

1. Mr. is 25 years old patient diagnosed with type I DM. He miss to take insulin and he enter to DKA? Which acid-base imbalance this patient develops?
- A. Respiratory alkalosis
 - B. Respiratory acidosis
 - C. Metabolic acidosis
 - D. Metabolic alkalosis

ANSWER: C

2. Fluid volume shift occur in all of the following, **except**?

- A. Extensive burns
- B. Lymphatic obstruction
- C. Hypoalbuminemia
- D. Hyperalbuminemia

ANSWER: D

3. For a male client in the oliguric phase of acute renal failure (ARF), which nursing intervention is most important?

- A. Encouraging coughing and deep breathing
- B. Promoting carbohydrate intake
- C. Limiting fluid intake
- D. Providing pain-relief measures

ANSWER: C

4. Nurse Fekadu do ECG and appreciate prominent U wave, which electrolyte imbalance basically indicates?

- A. Hyperkalemia
- B. Hypocalcemia
- C. Hypercalcemia
- D. Hypokalemia

ANSWER: D

5. A 29 years old female patient presents with rapid heartbeat, weight loss, heat intolerance, increased sweating and elevated thyroid hormone, what is the diagnosis of this patient?

- A. Hyperthyroidism
- B. Hypothyroidism
- C. Hyperparathyroidism
- D. Hypoparathyroidism

ANSWER: A

6. Which one of the following is **wrong** about acute renal failure?

- A. Rapid decline in GFR
- B. Asymptomatic
- C. Irreversible
- D. urine output < 400 ml/d is frequent

ANSWER: C

7. A client with chronic heart failure takes a daily thiazide diuretic. The client asks the nurse why it is important to consume potassium-rich foods on a daily basis. What is the best response?

- A. Consuming potassium-rich foods replaces what is lost in your urine.
- B. Potassium promotes the excretion of excess fluid by your kidneys.
- C. Potassium improves the therapeutic action of your diuretic.
- D. Your thiazide diuretic may cause significant hypocalcemia

ANSWER:A

8. A type of anemia happens due to loss of blood volume, which results in fewer blood cells?

- A. Iron deficiency anemia
- B. Hemolytic anemias
- C. Hypovolemic anemia
- D. Pernicious anemia

ANSWER: C

9. A 30 years old male patients presents to you with a complaint of fever, GI discomfort and vague Para umbilical pain that progress to right lower quadrant. On physical examination, palpating the left lower quadrants, causes pain in right lower quadrant and rebound tenderness. Laboratory finding reveals raised WBC? What is the Medical diagnosis of this patient?

- A. Peritonitis
- B. Small bowel obstruction
- C. Cholecystitis
- D. Appendicitis

ANSWER: D

10. A 29 years old female patient presents with hypothermia, hypotension, and hypoventilation. What is the diagnosis of this patient?

- A. Hyperthyroidism
- B. Hypothyroidism
- C. Hyperparathyroidism
- D. Hyperparathyroidism

ANSWER: B

In a 29-year-old female client who is being successfully treated for Cushing's syndrome, the following finding is seen now, which clinical feature is **not** improved?

- A. Decline in blood glucose
- B. Decline in serum sodium
- C. Increased oiliness of the skin
- D. Increased potassium
- E. Answer- C

11. Pick **wrong** finding regarding laboratory finding for respiratory acidosis?

- A. $\text{pH} < 7.35$
- B. $\text{PaCO}_2 > 42\text{mmHg}$
- C. HCO_3^- decreased (compensatory)
- D. HCO_3^- increased (compensator)

ANSWER:C

12. Which of the following electrolyte imbalance leads to cellular dehydration?

- A. Hyponatremia
- B. Hypomagnesemia**
- C. Hypocalcemia
- D. None

ANSWER:A

13. Which one of the following is **nothall mark features** related to nephritic syndrome?

- A. Hypoalbuminemia
- B. periorbital edema
- C. Hyperalbuminemia
- D. Hyperlipidemia

ANSWER:C

14. Mr. John, 28 year's old male patient presents with a complaint of flank pain, dysuria, and high grade fever 38.50°C that lasts for 1 week duration. He has CVA tenderness and his laboratory result reveal raised bacteria in the urine. What was his possible diagnosis?

- A. Acute pyelonephritis
- B. Nephrotic syndrome

- C. Chronic Pyelonephritis
- D. Acute glomerulo nephritis

ANSWER: A

15. A 25 years old male patient diagnosed with renal failure and laboratory finding reveals raised renal function test. Choose wrong finding regarding the case?

- A. Azotemia
- B. Hyperkalemia
- C. Hypokalemia
- D. Hypocalcemia

ANSWER: D

16. For a male client in the oliguric phase of acute renal failure (ARF), which nursing intervention is most important?

- A. Encouraging coughing and deep breathing
- B. Promoting carbohydrate intake
- C. Limiting fluid intake
- D. Providing pain-relief measures

ANSWER: C

17. Because of impaired urine elimination, the nurse is concerned about which potential skin problems that will require additional team planning for the client in renal failure?

- A. Extreme oiliness
- B. Loss of skin turgor
- C. Pronounced itching
- D. Reduced perspiration

ANSWER: C

18. Daniel 50 years old male patient was a known diabetic patient who have follow up at Assela Hospital diabetic clinic. He missed his appointment and left medication for two days and his RBS raised at which he enter to DKA. What clinical features we cannot appreciate mainly on this patient?

- A. Nausea, Vomiting
- B. Rapid / Acidotic breathing
- C. Sunken eye

D. Respiratory acidosis

ANSWER: D

19. Clients with aplastic anemia experience all the following features, **except**?

- A. Weakness and fatigue
- B. Unusual bleeding
- C. Ecchymosis
- D. Blood cell count shows sufficient number of leukocyte

Answer- D

20. The patient with gastroesophageal reflux disease should be taught:

- A. To avoid coffee, tea, or other caffeine-containing beverages.
- B. To take histamine 2 blockers, such as ranitidine, as directed.
- C. To avoid acidic foods such as citrus or tomato.
- D. All of the above

ANSWER: D

21. A 40 years old male patients presents to our hospital with clinical feature of edema on the leg, hepatomegaly on palpation and ascites. For this patient which type of heart failure typical manifest with this presentation?

- A. Left side heart failure
- B. Backward failure of the right heart
- C. Backward failure of the left heart
- D. Forward failure of left heart

ANSWER:B

22. Mr. Bedada enter to the clinic with a clinical feature of abdominal cramp and headache. The nurse takes vital sign and appreciate tenderness on palpation. What component of nursing process applied on this patient?

- A. Assessment
- B. Diagnosis
- C. Planning
- D. Implementation

ANSWER: A

23. Alison is being treated for hyperthyroidism. In reviewing her lab results, you would expect to see:

- A. Diminished thyroid hormone.
- B. Elevated thyroid hormone.
- C. Diminished parathyroid hormone.
- D. Elevated parathyroid hormone

ANSWER: A

Patients with nephrolithiasis or kidney stones need to increase fluid intake. This is to:

- A. Concentrate the urine.
- B. Help flush the stones through the urinary tract.
- C. Crystallize the struvite from the renal tubules.
- D. Break down the stones into smaller pieces that will more easily pass through the urinary tract.

ANSWER: B

24. Bernadette's morning lab results have just come in. Her serum potassium level is currently 7.4 mEq/L. You recognize this as:

- A. Hypokalemia.
- B. Hyperkalemia
- C. Hypocalcemia.
- D. Hypercalcemia

ANSWER: B

25. Mr. Daniel 56 years old man, suspected to have intestinal obstruction. X-ray of the abdomen was taken and it revealed that, there is volvulus which means...?

- A. strangulating the intestine
- B. telescoping the intestine
- C. necrosis of the intestine
- D. twisting of the intestine

ANSWER: D

26. A female client has a serum calcium level of 7.2 mg/dl. During the physical examination, nurse Noah expects to assess:

- A. Trousseau's sign.
- B. Homans' sign.
- C. Hegar's sign.
- D. Goodell's sign

ANSWER: A

27. During assessment of the patient with acromegaly, the nurse would expect the patient to report

- A. Infertility
- B. dry, irritated skin
- C. undesirable changes in appearance
- D. an increase in height of 2 to 3 inches per year

ANSWER: C

28. Daniel 40 year's old patient presents to you with a complaint of dysphagia, chest pain and heartburn. Barium swallow reveal dilation of the esophagus and endoscopy visualize tight cardiac and food remains in the esophagus. What is possible diagnosis of this patients?

- A. Diaphragmatic Hernia
- B. Esophageal Hernia

- C. Achalasia
 - D. Gastritis
- ANSWER:C

29. A patient is admitted to the hospital in thyrotoxic crisis. On physical assessment of the patient, the nurse would expect to find
- A. Hoarseness and laryngeal stridor
 - B. Bulging eyeballs and arrhythmias
 - C. Elevated temperature and signs of heart failure
 - D. Lethargy progressing suddenly to impairment of consciousness
- ANSWER: C

30. A patient with SIADH is treated with water restriction and administration of IV fluids. The nurses evaluates that treatment has been effective when the patient experiences
- A. increased urine output, decreased serum sodium, and increased urine specific gravity
 - B. increased urine output, increased serum sodium, and decreased urine specific gravity
 - C. decreased urine output, increased serum sodium, and decreased urine specific gravity
 - D. decreased urine output, decreased serum sodium, and increased urine specific gravity

ANSWER: B

31. A 10 years child is admitted to Adama Hospital with chief complaints of jerking movements around the face and arm. The nurses assess the child and he had repeated attacks of pharyngitis for which he was treated at the health center three weeks ago. A throat examination showed he had an increased Titer test and had inflammation in different parts of his joints. The possible diagnosis of this child will be:
- A. heart failure
 - B. rheumatoid heart disease
 - C. rheumatic fever
 - D. infective endocarditi
- ANSWER:C

32. A term newborn 2560 grams is admitted at 4 hours with a heart rate of 200/ minute, cold extremities, and capillary fill time of 4 seconds. The resident shifts the baby inside the neonatal intensive care unit and starts oxygen and other resuscitative measures. On asking about the history of the relatives, the resident realizes that the neonate was born by caesarian section done for placenta Previa and the mother had come with severe bleeding per vaginum to the hospital, and the baby was referred after birth to the NICU. Which is the most appropriate NEXT STEP for management?
- A. Give IV normal saline 10 mL/kg over 60 minutes
 - B. Give IV normal saline/ ringer lactate 10 mL/kg over 10 minutes
 - C. Arrange blood 10mL/kg and transfuse over 30 minutes
 - D. Immediately give 20 mL/kg O negative packed RBC over 30 minutes.
- ANSWER:C

33. During resuscitation of a newborn, you have established adequate ventilation with an endotracheal tube and your colleague has begun chest compressions for a heart rate under 60 bpm. Nevertheless, after 60 seconds, the heart rate has not increased. What is the most appropriate next step in management?*

- A. Administer epinephrine
- B. Discontinue chest compressions
- C. Start dopamine infusion.
- D. Stimulate the newborn

ANSWER:A

34. Failure of the lung to fully expand during inhalation due to lack of Surfactant deficiency?

- A. Meconium Aspiration Syndrome
- B. Hyaline Membrane Disease
- C. Transient hypoxia
- D. Chemical pneumonitis

ANSWER:B

35. Assume you are working at NICU ward and Neonate Chaltu was admitted 3 days back. Since, she is in distress and she was in Incubators, resuscitation equipment's was also used to save her life. After two days, she started showing the following symptoms Lethargic, inactive and unresponsive, Failure to suck, HSM, Cyanosis and body temperature as 38 degree Celsius. What is the possible diagnosis of the above neonate?

- A. Early onset of neonatal sepsis
- B. Late neonatal sepsis
- C. Severe pneumonia
- D. All are correct

ANSWER:B

36. The neonatologist nurse assesses the neonate after 3 days and finds the following manifestations, poor sucking stupor, hypotonia, seizures, obligatory tonic neck reflexes, and bilirubin levels exceed 30 mg/dL. What is the diagnosis of this neonate?

- A. Physiologic jaundice
- B. Kernicterus
- C. Breastfeeding jaundice
- D. pyloric stenosis-related jaundice

ANSWER:B

37. A term infant is placed under a radiant warmer, the skin is dried, and the oropharynx and nose are suctioned. After tactile stimulation, there is a minimal respiratory effect, dusky color, and a heart of 86 beats/minute. Bag/mask ventilation is performed for 30 seconds with 100% oxygen at a rate of 40-60 breath/minute. The heart rate increases to 100 beats/minute. One of the following is the best step for managing the neonate in the above case.

- A. perform endotracheal intubation
- B. continue bag/mask ventilation at a rate of 20 to 40 beats/minute
- C. continue ventilation and begin chest compressions

D. observe for spontaneous respiration and discontinue ventilation

ANSWER:D

38. Assume 2-year age-old child comes to you with cough, vomiting everything, lethargic and sudden onset of fever on your assessment. The RR for the child is 46 beat per minute.

What is the possible diagnosis and management for this child?

- A. Bronchial asthma and adrenaline
- B. Severe pneumonia and refer after pre-referral treatment
- C. Pneumonia and treat at health facility
- D. Acute nasopharyngitis and counsel the mother.

ANSWER:B

39. Tigist is the second child has diarrhea for 5 days. There is no blood in the stool. She is irritable and her eyes are sunken. The nurse offered fluids and the child drinks eagerly. The nurse measure Anthropometric and record Z-score of <-3 and classify Tigist's illness as she has severe malnutrition. The nurse of the ward focus on how to stop diarrhea prior to other care. What is the best to stop diarrhea of the following options?

- A. Provision of 75ml/kg of ORS
- B. Provision of RESOMAL solutions
- C. Initiate ringer lactate by 100ml/kg
- D. Zinc supplement with home remedy based care.

ANSWER:B

40. Which one of the following is NOT a reported benefit of breastfeeding?

- A. Promotes mother–infant bonding.
- B. Decreased intelligence quotient (IQ).
- C. Decreased risk of asthma and eczema in infants predisposed to these conditions.
- D. Reduced likelihood and severity of cows' milk protein allergy.

ANSWER:B

41. One of the following is not true about Good attachment means during breast-feeding.

- A. The baby's mouth should wide open
- B. Lower lips everted out
- C. More areola seen on upper part of breast
- D. mother should support the whole body of the baby

ANSWER:D

42. Baby Hana was born by use forceps. Accidentally pressure from forceps affected nerves around face and the affected side of the face droops and the infant is unable to close the eye tightly on that side. Assume you are the midwives assigned to assess the problem of this neonate. What is the possible diagnosis for the above case?

- A. Brachial plexus palsy
- B. Torticollis
- C. Subcutaneous fat necrosis
- D. facial palsy

ANSWER:D

43. A mother feeds her baby 4 times over 24 hours, and switch from one breast to other before completing from the first breast. This mother brought her 5 months with loss of weight and irritability. What is the possible problem of this infant?

- A. Feeding difficulty
- B. Feeding problem
- C. Good position
- D. Good attachment

ANSWER:B

44. Neonate of 10 days age from HIV positive mothers was brought to clinic. DNA test of neonate indicated the neonate is HIV positive. The classification for HIV is _____

- A. HIV unlikely
- B. Exposed HIV infection
- C. Confirmed HIV positive
- D. Symptomatic Confirmed HIV infection

ANSWER:C

45. Child developed primarily stiffed body, and after 15sec, abnormal body movement. What is possible dx of this child?

- A. Tonic-clonic
- B. Clonic only
- C. Subtile
- D. Absence seizure

ANSWER:A

46. Rhythmic jerky movements of extremities that can migrate to other body part is ____

- A. Clonic
- B. Tonic
- C. Myoclonic
- D. Subtle

ANSWER:A

47. A 10 years child came to OPD with chief complain of polyuria, weight loss and polyphagia. On assessment, Blood glucose level of child is 500mg/dl. If the weight of child is 30-35kg. The amount of fluid required to resuscitate is _____

- A. 300-700 ml 0.9% Ns
- B. 300-700 ml Ringer lactate/NS
- C. 300ml 0.9% NS and 5% dextrose
- D. 700ml of 0.9% Ns and ringer lactate

ANSWER:A

48. Assume target population (pt) for immunization in a *kebele* with a total population of 8,000. The percentages of surviving infants in the population is 3.4%. Using target coverage rate (ic) expressed as a decimal number = 90% and wastage factor (wf) = 1.33, calculate Annual pentavalent vaccine needs.

- A. 326
- B. 651
- C. 272
- D. 544

ANSWER:B

49. Assume the number of children's took BCG is 450, measles 150, DPT 1 350, and DPT 3 250. Calculate dropout rate of DPT?

- A. 44%
- B. 29%
- C. 80%
- D. 66

ANSWER:B

50. A 2 hours neonate admitted to hospital with **chonal atresia**. As nurse, what the first prioritized nursing care for this neonate.

- A. secure IV line and provide fluid as recommended
- B. insert NG tube and start formula feeding
- C. initiate breast feeding as much as possible
- D. advise mother how to feed breast

ANSWER:A

51. Assume a 3-year-old child comes to your health center with a chief complaint of difficulty of breathing. His parents told him that he had difficulty sleeping with a pillow and had repeated attacks of upper respiratory infections. ECO cardiographs show he has vegetation and is enlarged. The nurses assess and determine if he has heart failure. What is the name of this kind of heart failure?

- A. left side heart failure
- B. right side heart failure
- C. myocardial infarction
- D. Rheumatic heart disease.

ANSWER:A

52. A four-year-old child has repeated attacks of streptococcal pharyngitis for which he is not treated. After 3 weeks he developed purposeless jerky, repetitive movements particularly involving the hands and face. What is the possible diagnosis of this child?

- A. Poly arthritis
- B. carditis
- C. Sydenham's Chorea
- D. Subcutaneous Nodules

ANSWER:C

53. Eight years old child was assessed and proved as he has Rheumatic fever. The nurse planned to suppress the acute inflammatory process and administered **Aspirin**. As a

nurse, it is not allowed to leave this child alone. What is the main concern that makes not to leave the patient alone?

- A. because the disease is severe and emergency
- B. fear of developing bed sore
- C. because fever can kill the child
- D. fear of nasal bleeding

ANSWER:D

54. A 10 years child comes to Adama Hospital with clinical presentations of pronounced weight loss, stunting, hypertension, edema, and **GFR** below 60ml/min/1.73m²/ for more than 3 months. What is the possible diagnosis of this child?

- A. nephrotic syndrome
- B. post-streptococcal infections
- C. acute renal failure
- D. chronic renal failure

ANSWER:D

55. Nurse Mathewos prescribed amoxicillin 500mg PO TID for 7 days. From this prescription Amoxicillin and PO respectively shows.

- A. drug and route
- B. time and documentation
- C. dose and route
- D. documentation and time

ANSWER:A

56. Nurse administer ceftriaxone for patient for seven days. After 2 days patient developed rash, redness, swelling and itching over the skin. This scenario shows to nurse.

- A. drug side effect
- B. drug-drug interaction
- C. drug is working effectively
- D. desired effect of the drug

ANSWER:A

57. One of the following medication administration is dangerous.

- A. Parenteral
- B. oral
- C. rectal
- D. ophthalmic

ANSWER:A

58. Assume that nurse want to administer TAT 3000 IU IM stat after skin test. The nurse should wait for _____minute before administration of drug.

- A. 30 minute
- B. 20-25 minute
- C. 0-10 minute
- D. 10-15 minute

ANSWER:D

59. One Liter (1L) of ringer lactate was prescribed for patient over 8 hours. calculate drop rate for this case _____

- A. 42 drop/minute
- B. 20 drop/minute
- C. 125 drop/minute
- D. 4 drop/minute

ANSWER:A

60. The nurse prescribed 5mg/kg of gentamycin. (Weight of child 8kg and gentamycin preparation is 80mg/2cc). The amount to be given is _____

- A. 1cc
- B. 0.5cc
- C. 2cc
- D. 0.25cc

ANSWER:A

61. One of the following is wrong nurse action during IV fluid administration.

- A. irrigate in clot & hematoma
- B. Remove air from IV tube before connecting
- C. Balance input & out put
- D. Calculate amount to be administered.

ANSWER:A

62. Before the actual discharge occurs, what must the nurse ensure?

- A. The patient is well enough to go home.
- B. The patient has not been overly medicated.
- C. The patient understands the discharge instructions.
- D. The patient has adequate transportation.

ANSWER:C

63. Not nursing responsibility during patient admission?

- A. Check for orders that a patient need to be discharged

- B. Perform base line assessment
- C. Make introduction and orient the patient
- D. Check for orders of admission

ANSWER:A

64. Usually **unclaimed dead** bodies are kept in store for how many times?

- A. 48 hrs
- B. 24hrs
- C. 12hrs
- D. 72hrs

ANSWER:D

65. Recorded dead body data is required if the family of the patient have payment through Medicare or insurance agencies. This is called _____

- A. Data used as for research
- B. Data will be used for education
- C. Data will used for reimbursement
- D. Data used for legal issue

ANSWER:C

66. Which is the first step for nurse during postmortem care?

- A. Start tying
- B. Injection of formalin
- C. Tagging dead body
- D. Informing the families

ANSWER:D

67. Stiffening of the dead body accompanying the depletion of adenosine triphosphate in the muscle fibers is determined by?

- A. Rigor mortis
- B. Algor mortis
- C. Liver mortis
- D. Blue skin

ANSWER:A

68. The process by which a foreign substance in the body; is converted by enzymes into a less active and harmless agent that can be easily excreted is _____

- A. Excretion
- B. Transportation
- C. Biotransformation
- D. Route of absorption

ANSWER:C

69. Which of the following is **not the purpose/correct** of intradermal drug administration?

- A. diagnostic determinations
- B. desensitization
- C. Immunization.
- D. massage the site after injection

ANSWER:D

70. If repeated injections are given, the nurse should rotate the site of the injection so that each succeeding injection is about 5 cm away from the previous one. What is a common route?

- A. Sub-cutaneous
- B. Intravenous
- C. Intra-dermal
- D. PO

ANSWER:A

71. For which of the following patient back care is not recommended?

- A. heart disease
- B. Diabetic
- C. Hypertension
- D. Aneurysm

ANSWER:D

72. the pattern of crutches and weak leg for-ward together, then weight-bearing leg forward

- A. two-point gate
- B. three point gate
- C. four point gate
- D. swimming through

ANSWER:B

73. Proper nutrition in hospitalized clients is necessary for the following, except.

- A. wound healing
- B. reduction in morbidity
- C. causes protein energy problem
- D. decrease length of hospital stay

ANSWER:

74. Not advantages of suturing?

- A. To approximate wound edges until healing occurs
- B. To speed up the healing of the wound
- C. Increase the chance of infection
- D. For esthetic purposes

ANSWER:C

75. Not nursing responsibility for the patient presented with a wound.

- A. Clean all wound with hydrogen peroxide
- B. Dress wound accordingly
- C. Suture if indicated
- D. Provide antibiotics to prevent complications

ANSWER:A

76. prolonged inadequate quality and quantity of sleep, either of the REM or the NREM type

- A. sleep deprivation
- B. sleep apnea
- C. insomnia
- D. narcolepsy

ANSWER:C

77. Acute pain related to surgical procedure as evidenced by patient verbalization. Acute pain is _____

- A. Nursing plan
- B. Problem of patient
- C. Nursing assessment
- D. Etiology

ANSWER:B

78. Which age categories sleep less

- A. Elders
- B. Infants
- C. Children
- D. Neonate

ANSWER:A

79. Types of communication used to treat patient is _____

- A. Group
- B. Intra
- C. Therapeutic
- D. Inter

ANSWER:C

80. Babies born to mothers with some deficient conditions during pregnancy may have possibility of developing serious congenital malformations. Which of the following supplementary drug to mothers during pregnancy could help to protect babies from developing neural tube defect?

- A. Vitamin B6
- B. Iron
- C. Folic acid
- D. Zinc

ANSWER: C

81. A laboring mother is advised to take Anti D Ig within 72 hours following delivery to prevent the possibility of rhesus incompatibility in subsequent pregnancies. In which of the following conditions that Rh incompatibility is expected to occur?

- A. When Rh negative mother carries Rh positive fetus from Rh positive father
- B. When Rh positive mother carries Rh negative fetus from Rh negative father
- C. When Rh negative mother carries Rh positive fetus from Rh positive father
- D. When Rh positive mother carries Rh positive fetus from Rh negative father

ANSWER: C

82. Azeb is a 30 years old primigravida lady who came for the complain of a severe headache and epigastric pain for 3 days duration. GA is 36 weeks from LMP. While performing physical examination BP was 149 by 95mmhg. Otherwise, she never has history of raised blood pressure. The laboratory finding showed a positive 2 albumin in urine and organ function tests are in normal range and platelet counts is 95000. Which of the following is the right management for Azeb?

- A. Administer antihypertensive drug
- B. Administer steroid
- C. Administer magnesium sulfate
- D. Expectant management until term

ANSWER: C

83. A 32 weeks of gestation pregnant woman comes to the health center with dark clotty vaginal bleeding and abdominal pain following a falling accident. There is abdominal tenderness, and fetal heartbeats difficultly heard. What is the likely cause of the bleeding?

- A. Placenta abruption
- B. Placenta previa
- C. Vasa previa
- D. Abortion

ANSWER: A

84. A woman came with a post term pregnancy and unfavorable cervix. The duty team decided to terminate the pregnancy, which of the following alternative has to be used first?

- A. Misoprostol
- B. Mifepristone
- C. Membrane stripping
- D. Oxytocin

ANSWER: C

85. Assume Mr. Alemu attended delivery of neonate at Asella referral hospital. During neonatal assessment during first minute heart rate is less than 100, body pink and extremities blue, respiration is absent, limb and no reflex activity. What is the APGAR score of this neonate?

- A. 2
- B. 3
- C. 4
- D. 6

ANSWER: A

86. The following are the correct management of cardiac patient during intrapartum period except?

- A. Shorten the second stage by instrumental deliver
- B. Cesarean section is a preferred mode of delivery
- C. Conduct labor and delivery in lateral position
- D. Do not use Ergometrine in the third stage of labour

ANSWER: B

87. Which of the following maneuver is used to deliver the shoulders in assisted breech delivery?

- A. BurnMarshall maneuver
- B. Lovset maneuver
- C. Pinardmaneuver
- D. MauriceauSmellieVeit maneuver

ANSWER: B

88. Atype of lochia characterized by pink in colour and discharged during the next 5 to 9 days is ___?

- A.Lochia rubra
- B.Lochia serosa
- C.Lochia alba
- D.White lochia

ANSWER: B

88. Which of the following is a postpartum complication, whichis characterized by offensive vaginal discharge and fever?

- A. Postpartum psychosis
- B. Puerperal sepsis
- C. Postpartum blues
- D. Thrombophlebitis

ANSWER: B

89. Within how much days emergency contraceptive pills are most effective?

- A. Within 3 days

- B. Within 15days
- C. Within 7days
- D. Within10 days

ANSWER: A

89. Which of the following perineal tear involve rectal mucosa?

- A. First degree tear
- B. Second degree tear degree tear
- C. Third degree tear
- D. Fourth degree tear

ANSWER: D

90. Which of the following is true about option B Positive HIV care

- A. Put all pregnant women on ART for stage I and CD4 less than 500
- B. Put all lactating women on ART for stage I and CD4 less than 500
- C. Put all lactating women on ART based on WHO staging and CD4
- D. Put all pregnant women on ART irrespective of WHO staging and CD4

ANSWER: D

91. A 40 years old gravida III Para I with history of previous abortion came with a chief complain of abdominal pain and vaginal bleeding for a day duration. She claims to be amenorrheic for the past 5 months and she is a known hypertensive patient. On physical examination you found the size of the uterus to be at the level of the umbilicus and BP 110 by 75mmhg, PR 92,Temp ATT. While performing vaginal examination you found a closed cervix and a minimal bleeding on the examining fingers. Which of the following is the most likely diagnosis?

- A. Incomplete abortion
- B. Complete abortion
- C. Threatened abortion
- D. Inevitable abortion

ANSWER: C

92. Based on latest DHS, what is the current neonatal mortality rate in Ethiopia?

- A. 30 death per 1000 livebirth
- B. 29 death per 1000 livebirth
- C. 33 death per 1000 livebirth
- D. 41 death per 1000 livebirth

ANSWER: A

93. Which of the following is the commonest site for tubal pregnancy?

- A. Isthmus
- B. Ampulla
- C. Infundibulum
- D. Interstitial

ANSWER: B

94. A 19 years old sexually active lady came to your gynecology OPD complaining of menstrual pain that dull and localized to the lower quadrants of the abdomen, nausea, vomiting and diarrhea. She skipped school due to the pain whenever the period comes.

What could be the 1st line therapy for her?

- A. Ibuprofen
- B. OCP
- C. Naproxen
- D. Aspirin

ANSWER: B

95. Which of the following types of HPV are responsible for about 70% of cervical cancer?

- A. 16 and 18
- B. 45 and 31
- C. 16 and 35

D. 18 and 45

ANSWER: A

96. Which of the following is the best screening tool to screen cervical cancer in low resource countries like Ethiopia?

- A. Colposcopy
- B. Human Papillomavirus Testing
- C. Visual inspection with acetic acid and naked eye
- D. A traditional Pap smear

ANSWER: C

97. According to Ethiopian demographic and health survey report (EDHS 2016), the **MMR** (maternal death per 100,000 live births) is___?

- A. 673
- B. 676
- C. 412
- D. 422

ANSWER: C

98. Which one of the following is abnormal finding during neonatal examination?

- A. Arms and legs are flexed
- B. 40 breaths in 1 minute
- C. 120 beats in 1 minute
- D. The anterior fontanel swells when the baby sleeps

ANSWER: D

99. Which one of the following is not among the causes of first delay?

- A. Failure to recognize complication sign
- B. Poor skills of health providers
- C. Cost consideration
- D. Gender influence

ANSWER:

100. Which one of the following **is not** the cause for unsafe abortion?

- A. Self induced method.
- B. access to safe services
- C. Delay in seeking procedure
- D. Unskilled provider

ANSWER: B

101. Suppose a 28 years old women comes to Adama Medical College family planning unit to receive a family planning method which is inserted to upper arm and protects from pregnancy for a maximum of three years. Based on the information above which family planning method can be provided for this woman?

- A. Norplant
- B. Implant
- C. Implanon
- D. Jedelle

ANSWER: C

102. Mekdes is a 23 years old lady who wishes to use a calendar family planning method. Based on her 6 months menstrual cycle, her shortest and longest cycle is 28 and 32 days respectively, which of the following is her fertile period?

- A. From 10th upto 21th days of the cycle
- B. From 8th upto 17th days of the cycle
- C. from 7th upto 14th days of the cycle
- D. From 14th upto 18th days of the cycle

ANSWER: A

103. Which of the following condition influences the suitability of IUCD for a particular woman?

- A. Past history of abortion
- B. History of vaginal candidiasis
- C. History of pregnancy
- D. Current pelvic infection

ANSWER: D

104. A 37 year old man came to out patient family planning clinic after vasectomy was done for him. You counseled him to use condom or avoid intercourse for effectiveness of the method. For how long must he use condom or avoid sex?

- A. One month
- B. Two month
- C. Three month
- D. Four month

ANSWER: C

105. A pregnant woman took a second dose of tetanus diphtheria vaccine on today. When does next dose has to be provided?

- A. Immediately on today
- B. After four weeks
- C. After one year
- D. After six months

ANSWER: D

106. Which of the following complication of instrumental delivery is more common with ventous delivery?

- A. Cephalohematoma
- B. Brachial plexus injury
- C. Facial nerve injury
- D. Anterior eye injury

ANSWER: A

107. A woman whose GA of 39 week is admitted in hospital. Her physical examination shows strong uterine contraction, intra uterine fetal death, severs CPD, no previous cesarean scar, cervix is fully dilated and effacement is 100 percent. Which option is most appropriate for the management of this labour?

- A. Forceps

- B. Laparotomy
- C. Vacuum
- D. Caesarean section

ANSWER: D

108. A much-feared outcome of Alzheimer's disease is

- A. functional psychosis.
- B. paranoia.
- C. general paresis.
- D. senile dementia.

ANSWER: D

109. Multiple personality or dissociative identity disorder often begins

- A. as a result of combat exhaustion.
- B. in adulthood as a response to unremitting phobias.
- C. as a consequence of post-traumatic stress disorders.
- D. in childhood as a result of unbearable experiences.

ANSWER: D

110. A person who has an extreme lack of self-confidence and who allows others to run his or her life is said to have a(n) _____ personality.

- A. Dependent
- B. Narcissistic
- C. Paranoid
- D. Antisocial

ANSWER: A

111. Because there is considerable overlap among the types of schizophrenia, and because patterns of behavior shift over time, many patients are simply classified as suffering from _____ schizophrenia.

- A. Borderline
- B. Atypical
- C. Mixed
- D. Undifferentiated

ANSWER: D

112. Retreat from reality by hallucinations and delusions and by social withdrawal typically characterizes

- A. somatoform disorders.
- B. anxiety disorders.
- C. psychotic disorders.

D. personality disorders.

ANSWER: C

113. Persons who suffer from paraphilias are categorized as having

- A. somatoform disorders.
- B. generalized anxiety.
- C. sexual disorders.
- D. personality disorders.

ANSWER: C

114. Statistical approaches to abnormality define as "abnormal" those who

- A. show evidence of loss of contact with reality.
- B. are unhappy, withdrawn, and depressed.
- C. deviate from typical or average patterns of behavior.
- D. are disabled by anxiety.

ANSWER: C

115. Sensory experiences that occur in the absence of a stimulus are called

- A. illusions.
- B. hallucinations.
- C. delusions.
- D. affect episodes.

ANSWER: B

116. If a person suffering from schizophrenia has an identical twin, that twin

- A. will have a great percent chance of becoming schizophrenic.
- B. will be no more likely than anyone else to become schizophrenic.
- C. is almost sure to become schizophrenic.
- D. is also likely to have more than one personality.

ANSWER: A

117. The most severe psychological disorder is

- A. personality disorder.
- B. psychosomatic illness.
- C. anxiety disorder.
- D. psychosis.

ANSWER: D

118. Persons who are excessively narcissistic, dependent, or antisocial are characterized as having

- A. somatoform disorders.
- B. generalized anxiety.
- C. conversion reactions.
- D. personality disorders.

ANSWER: D

119. Mood disorders are those in which the person may

- A. experience severe depression and threaten suicide.
- B. exhibit symptoms suggesting physical disease or injury but for which there is no identifiable cause.
- C. exhibit behavior that is the result of an organic brain pathology.
- D. experience delusions and hallucinations.

ANSWER: A

120. A core feature of all abnormal behavior is that it is

- A. culturally absolute.
- B. learned.
- C. maladaptive.
- D. dependent on age.

ANSWER: C

121. DSM stands for

- A. diagnostic schedule of medicine.
- B. diagnostic and statistical manual.
- C. depressive scale modalities.
- D. doctor of surgical medicine.

ANSWER: B

122. Conversion disorder and hypochondriasis are classified as

- A. physio-mental disorders.
- B. somatoform disorders.
- C. psychosomatic disorders.
- D. somatization disorders.

ANSWER: C

123. An unusual state called "waxy flexibility" is sometimes observed in _____ schizophrenia.

- A. borderline
- B. disorganized
- C. catatonic
- D. paranoid

ANSWER: C

124. Irrational and very specific fears that persist even when there is no real danger to a person are called

- A. anxieties.

- B. phobias
- C. dissociation
- D. obsessions.

ANSWER: B

125. False beliefs that are held even when the facts contradict them are called

- A. fantasies.
- B. hallucinations.
- C. illusions.
- D. delusions.

ANSWER: D

126. The most widely used system of psychological classification today is

- A. the Freudian Psychoanalytic System (FPS).
- B. found in the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).
- C. the system designed by Emil Kraepelin and Eugen Bleuler.
- D. to be found in the Federal Uniform Code of Psychopathology (UCP).

ANSWER: B

127. Which of the following is classified as a mood disorder?

- A. bipolar disorder
- B. multiple personality disorder
- C. delusional disorder
- D. dissociative disorder

ANSWER: A

128. The distinction between obsessions and compulsions is the distinction between

- A. engaging in behaviors that are merely inconvenient and those that are severely disruptive.
- B. having positive and negative feelings toward an object or event.
- C. thoughts that are evidence of neurosis or those that are evidence of psychosis.
- D. having repetitious thoughts or engaging in repetitious actions.

ANSWER: D

129. Hearing voices that are not really there would be called

- A. hallucination.
- B. delusion.
- C. auditory regression.
- D. depressive psychosis.

ANSWER: A

130. A person who is preoccupied with fears of having a serious disease suffers from

- A. a conversion reaction.
- B. hypochondriasis.
- C. a traumatic disorder.
- D. an obsession.

ANSWER: B

131. One of the most important ways to prevent pressure ulcers is to:

- A. Do regular skin assessments
- B. Provide support surfaces for all
- C. Identify at-risk individuals
- D. Do regular repositioning

ANSWER: C

132. W/roKemila has fallen from stares . she has bruised brain with possible surface hemorrhages

This patient is unconscious for considerable period and with cool, pale skin. Most probably this patient has

- A. Concussion
- B. Contusion
- C. Confusion
- D. All

ANSWER: B

133 . Ato Kemal has come to Hospital with Bloody conjunctivae , CSF rhinorrhea, headache, vertigo and alter level of consciousness .

- A. Vitiligo
- B. Scalp injury
- C. Skull fracture & Brain injury
- D. Skull fracture

ANSWER: C

134. Minor head injury without significant sequelae with temporary loss of brain function with no apparent structural damage to the brain after head injury can be

- A .Paralysis
- B. Concussion
- C. Contusion
- D. Confusion

ANSWER: B

135. Which one is the first measure must be taken by NURSES for patient with head injury?

- A. Maintain an open airway
- B. Improve cognitive functions
- C. Maintain body temperature
- D. Promoting adequate nutrition

ANSWER: A

136. Is a break in the continuity of the bone which cover brain caused by forceful trauma with or without damage to the brain is called

- A. Cerebral laceration
- B. Skull fracture
- C. Scalp injury
- D. Head injury

ANSWER: B

137. To promote adequate nutrition professionals must NOT do

- A. Feed the patient via NG-tube
- B. If there is abdominal distention perform gastric aspiration
- C. Continue parenteral feeding until swallowing reflex returns
- D. Check dressing and cast

ANSWER: D

138. A victim has lost a lot of blood through a deep cut in his leg. He is breathing fast and seems pale and restless. He is probably....

- A. Having a stroke.
- B. Having a heart attack.
- C. In shock.
- D. Choking.

ANSWER: C

139. A victim has a large piece of glass sticking out of her leg. You should....

- A. Leave the glass in her leg and control the bleeding.
- B. Call your local emergency phone number.
- C. Remove the glass and then control the bleeding.
- D. Both a and b

ANSWER: D

140. You feel a sudden sharp pain in the bottom of your left foot. You look at the bottom of your left shoe and see a tack sticking in your shoe. What type of wound do you probably have?

- A. Avulsion
- B. Bruise
- C. Scrape

D. Puncture

ANSWER: D

140 Which should be part of your care for a severely bleeding open wound?

- A. Allow the wound to bleed in order to minimize infection.
- B. Apply direct pressure and elevate the injured area. (If no broken bones)
- C. Use a tourniquet to stop all blood flow.
- D. Both b and c

ANSWER: D

141. What should you do if you think a victim has serious internal bleeding?

- A. Apply heat to the injured area.
- B. Call your local emergency phone number for help.
- C. Place the victim in a sitting position.
- D. Give fluids to replace blood loss.

ANSWER: B

142. Which is the first step when caring for bleeding wounds.

- A. Apply direct pressure with a clean or sterile dressing.
- B. Apply pressure at the pressure point.
- C. Add bulky dressings to reinforce blood soaked bandages.
- D. Elevate the wound.

ANSWER: A

143. How can you reduce the risk of disease transmission when caring for open, bleeding wounds?

- A. Wash your hands immediately after giving care.
- B. Avoid direct contact with blood.
- C. Use protective barriers such as gloves or plastic wrap.
- D. All of the above.

ANSWER: D

144. You have tried to control a victim's bleeding with direct pressure and elevation, but the bleeding doesn't stop. Where would you apply pressure to slow the flow of blood to a wound on the forearm?

- A. Outside the arm midway between the shoulder and the elbow.
- B. On the inside of the elbow.
- C. Inside the arm between the shoulder and the elbow.
- D. Any of the above will slow the flow of blood.

ANSWER: C

145. When giving care for external bleeding, what should you do first?

- A. Elevate the injury.
- B. Apply direct pressure.
- C. Apply a loose dressing.
- D. Apply a tourniquet

ANSWER: B

146. When caring for a victim with a bloody nose you would not.....

- A. Apply an ice pack to the bridge of the nose.
- B. Apply pressure to upper lip just beneath nose.
- C. Have the victim sit with head tilted slightly backward while pinching the nostrils together.
- D. Have the victim sit with head tilted slightly forward while pinching the nostrils together.

ANSWER: D

147. Which symptoms would indicate internal bleeding?

- A. Rapid weak pulse, Excessive thirst.
- B. Skin that feels cool or moist, or looks pale or bluish.
- C. Tender, swollen, bruised, or hard areas of his body, such as the abdomen.

D. All of the above.

ANSWER: D

148. Blood ooze from wound which can be controlled by easily with direct pressure

- A. capillary bleeding
- B. Arterial bleeding
- C. Venous bleeding
- D. All

ANSWER: A

149. The two most common sources of internal bleeding may be

- A. Fractured extremities
- B. Injured or damaged internal organ
- C. Infection
- D. Shock
- E. A & B

ANSWER: E

150. In a cardiac centre of a hospital, more than 5000 children and other cardiac patients are on waiting list for cardiac surgery because of resource scarcity. Along with solving the budget constraint and training of health professionals, the hospital management team decided to give service in their order of visit. What is the most appropriate principle applied in this context?

- A. Autonomy
- B. Beneficence
- C. Non-maleficence
- D. Justice

ANSWER:D

151. A 25 years old female patient admitted to a medical ward with the diagnosis of hepatitis B. After nurse's counselling, the patient agreed to share her information to her family. Which ethical principle applied by the nurse?

- A. Advocacy

- B. Privacy
- C. Confidentiality
- D. Informed consent

ANSWER:C

152. A researcher obtained an informed consent from an HIV positive study participant for the purpose of estimating the prevalence of the diseases in a country. The researcher used the same data to identify list of HIV positive individuals in the community for another study without asking their consent gain. What ethical principle is breached?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-maleficence

ANSWER:B

153. A 35 year old woman was brought to a laboratory reception unit to give blood sample for pre-operative assessment. The woman was not fully conscious when she reached in the sample collection site. Then the laboratory technicians need permission or consent of the patient before drawing the blood sample. What type of consent he has to use to draw a blood sample based on this scenario?

- A. Patient autonomy
- B. Informed consent
- C. No need of consent
- D. Surrogate decision maker

ANSWER:D

154. A laboratory manager needs to distribute packs gloves for different units of laboratory. The glove was not enough to distribute according to their need. The manager studied

the consumption and future need and decided who deserves more resources to realize the best outcomes of the laboratory units. Which principle of ethics is the manager most concerned?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-malefeasance

ANSWER:A

155. A 25 year old pregnant woman visited an ANC clinic. The laboratory professional weighed the mental stability and consciousness of the woman to give blood sample and decided that she can give consent by herself regardless of any other representative. What type of consent is appropriate to take for the above case?

- A. Assent
- B. Verbal consent
- C. Written consent
- D. Surrogate decision making

ANSWER:B

156. A 35 years old male, athlete admits with pain in Right knee, investigations reveal that he is suffering from end stage osteosarcoma, and needs extensive chemotherapy, radiotherapy to treat the cancer. Athlete's father died with osteosarcoma in his fifties. His mother died due to breast cancer in her forties. He has a living will that if he ever diagnosed with cancer, not wanted to be aggressively treated by chemotherapy, radiotherapy and other therapies. Physician is consistent to treat him with all sorts of treatment possible to cure his cancer. Physician is exercising:

- A. Act of Paternalism
- B. Ideal beneficence

- C. Beneficence
- D. Surrogate decision maker

ANSWER:C

157. A 32 year old female patient came to hospital with sudden onset of sharp knife pain like chest pain accompanied by her family. On assessment, she pronounced dyspnea and central cyanosis. The physician and nursing team in charge performed emergency surgery. The patient and her family presented the case to court because she developed iatrogenic pleural effusion as complication. What nursing action most likely prevents accusation?

- A. Keep fidelity
- B. Take consent
- C. Keep veracity
- D. Assess patient capacity

ANSWER:B

158. A nursing instructor is teaching nursing students about principles of ethics in health care, and she tells them that the most important principle to observe while taking care of patient is doing no harm. The principle of Ethics she described here:

- A. Respect for autonomy
- B. Beneficence
- C. Justice
- D. Nonmaleficence

ANSWER:D

159. A nurse does not check patient's blood pressure admitted with severe hypertension and patient suffers Cerebral Vascular Accident (CVA) and irreversible hemiplegia due to raised blood pressure. This shows:

- A. Negligence
- B. Ethical dilemma
- C. Malpractice
- D. Beneficence

ANSWER:A

160. Discharging the patient without the permission of the physician/nurse in charge is called:

- A. Discharging with good Progress
- B. Discharging against medical
- C. Discharging by Referral
- D. Discharging by death

ANSWER:B

161. A hospital implemented the health intervention measure which is intended to reduce impairments, disability and suffering caused by health problems. The hospital also wants to patients' adjustment to not curable conditions. What level of disease prevention will be considered for this intervention?

- A. Primordial prevention
- B. Primary prevention
- C. Secondary prevention
- D. Tertiary prevention

ANSWER:D

162. The public health nurse who does blood pressure screening and related health education is conducting activities in the level of:

- A. Primary prevention

- B. Secondary prevention
- C. Tertiary prevention
- D. focused prevention

ANSWER:A

163. The health educator who teaches proper body mechanics for bending and lifting is conducting activities in the level of

- A. Primary prevention
- B. Secondary prevention
- C. Tertiary prevention
- D. Focused prevention

ANSWER:A

164. A 22 year old male client wants to start an exercise program, but he says he doesn't have the time or money to attend a fitness club. The nurse recognizes the behaviour-specific cognitions and affect variable of

- A. Perceived self-efficacy
- B. Perceived barriers to action
- C. Interpersonal influences
- D. Situational influences

ANSWER:B

165. The challenges faced, and the long term advantages of change, influence which part of the Health Belief model?

- A. Perceived benefits
- B. Perceived barriers
- C. Perceived efficacy
- D. Perceived severity

ANSWER:A

166. When the conditioned stimulus is no longer paired with the unconditioned stimulus, what occurs?

- A. Extinction
- B. Reversal
- C. Shaping
- D. Generalization

ANSWER:A

167. In operant conditioning, what is the tendency for a response to happen only when a particular stimulus is present?

- A. Stimulus generalization
- B. Shaping
- C. Response acquisition
- D. Stimulus discrimination

ANSWER:D

168. Classical conditioning usually works best if which of the following occurs?

- A. The conditioned stimulus is presented just after the unconditioned stimulus
- B. The conditioned stimulus is presented just before the unconditioned stimulus
- C. The conditioned stimulus is presented many hours after the unconditioned stimulus
- D. All of the above are equally effective

ANSWER:B

169. What is the tendency to respond to a new stimulus as if it's the original conditioned stimulus?

- A. Stimulus discrimination
- B. Stimulus generalization
- C. Higher-order conditioning
- D. Spontaneous recovery

ANSWER:B

