

1. Sara is a 3 months pregnant mother came to health center ANC for checkup. On history you found that she has 4 children, 2 previous abortion and 1 still birth.

This finding shows you that she is Gravida (G)_____and Para(P) _____.

A. G7,P4

C.G7,P5

B. G6,P4

D. G8,P5

Halima is a 24 year women is on her 5th pregnancy and has a history of abortion in the 4th pregnancy and the first pregnancy was a twin. She is considered to be

A. G4 P3

C. G5 P3

B. G5 P4

D. G4 P4

2. On abdominal palpation you found that the fundal height of the uterus is half way between the umbilicus and symphysis pubis. Her gestational age will be ____ weeks

A. 12

B. 16

C. 20

D. 24

3. What is the most common complication associated with too rapid delivery in precipitate labor?

- a. Pitting edema of the baby's scalp
- b. Dural or subdural tears in fetal brain tissue
- c. Premature separation of the placenta
- d. Prolonged retention of the placenta

- 4. The nurse has just completed emergency delivery of a term infant. What is the priority nursing concern at this time?
 - a. Controlling hemorrhage in the mother
 - b. Removing the afterbirth
 - c. Keeping the infant warm
 - d. Cutting the umbilical cord

5. A woman spontaneously delivers a baby girl who is immediately handed to the nurse. Which action is of highest priority for the nurse?

- a. Do an Apgar assessment.
- b. Check neonatal heart rate.
- c. Apply identification bracelets.
- d. Clear the nasopharynx.

- 6. In abdominal examination of a pregnant woman, the 3rd Leopold's Maneuver is
 - A. Deep pelvic palpation
 - B. Pawlick's Grip
 - C. Fundal height palpation
 - D. Lateral palpation

7. The long axis of the fetus in relation to the axis of mother uterus is known as: -

- A. Presentation
- B. Position
- C. Lie
- D. Attitude

- 8. Which of the following is the most important nursing assessment during the fourth stage of labor?
- a. Bonding behaviors
- b. Distention of the bladder
- c. Ability to relax
- d. Knowledge of newborn behavior

9. Kedija is a 20 years old primi-gravida mother comes to hospital with labor pain and progressive uterine contractions, on vaginal examination you find that the cervix is 4 fingers dilated.

Then the expected cervical dilation in centimeters is :-

- A. 2 cm
- B. 4 cm
- C. 6 cm
- D. 8 cm

10. Aster is a 28 years old gravida IV and para III mother who came to the health center with labor pain . While performing vaginal examination you felt the vagina is hot and dry. This finding may indicate that Aster may have:-

- A. Obstructed labor
- B. Pre-eclampsia
- C. Infection or sepsis
- D. Previous scar

11. Hana 25 years old female at term came to labor ward with active first stage of labor . On vaginal examination you find a spongy feeling between your fingers and the presenting part.

This may indicate Hana may have :-

- A. Obstructed labor
- B. Abruptio placentae
- C. Placenta previa
- D. Vasa previa

- 12. The nursing care plan for a woman who has placenta abruptio should include careful assessment for signs and symptoms of which of the following?
 - a. Jaundice
 - b. Hypovolemic shock
 - c. Impending convulsions
 - d. Hypertension

- 13. A woman is admitted with suspected placenta previa. What test does the nurse expect will be done to confirm the diagnosis?
- a. Vaginal exam (PV)
- b. Non-stress test
- c. Oxytocin challenge test
- d. Ultrasound

- 14. Magnesium sulfate is ordered for a client who is hospitalized for pregnancy-induced hypertension (PIH). What effects would the nurse expect to see as a result of this medication?
- a. CNS depression
- b. Decreased gastric acidity
- c. Onset of contractions
- d. Decrease in number of bowel movements

- 15. Fatuma is 24 years old woman came to your Labour and Delivery ward with complaining of bleeding during pregnancy. Then, which of the following signs and symptoms is associated with placenta previa?
- A. Painless bright red bleeding
- B. Hard, tender uterus
- C. Concealed bleeding
- D. Rigid abdomen

- 16. Bleeding from the genital tract in late pregnancy after 28th weeks of gestation and before the onset of labour is known as :-
 - A. Antepartum Hemorrhage
 - B. Post partum Hemorrhage
 - C. Abortion
 - D. None of the above

- 17. The nurse is caring for a laboring woman who has a history of rheumatic heart disease. How should the nurse position her during labor?
 - a. Supine
 - b. Semi-recumbent
 - c. Side-lying
 - d. Sitting

- Disseminated intravascular coagulation (DIC) can occur in _____. This happens because when the placenta becomes damaged and detaches from the uterine wall, large amounts of _____ are released into mom's circulation, leading to clot formation and then clotting factor depletion.
- A. Placenta previa, fibrinogen
- B . Abruptio placentae, fibrinogen
- C. Placenta previa, platelets
- D. Abruptio placentae, thromboplastin

- 19. A 25 year old patient at 32 weeks of amenorrhea was brought to maternity ward of Hiwot Fana Specialized university hospital . These are the clinical findings on the physical examination. She was Pale ,dehydrated, RP: 120/ min; BP 90/60 mmHg; delay in the capillary refilling time; . Abd: Fundal height 36 cm, tenderness, and uterus hard, no Fetal Heart heard. Vaginally: scanty blood coming through the canal, reddish area around the cervix was noticed. Which among the following is the most likely diagnosis?
- A. Placental abruption
- B. Placenta praevia type IV. .
- C. Cervical carcinoma
- D. Severe placental abruption with IUFD

20. Which area of health teaching will a new mother be most responsive to during the taking in phase of the postpartum period?

- a. Family planning
- b. Newborn care
- c. Community support groups
- d. Perineal care

- 21. A 20 years old woman came to Gyne OPD with complaints of fever and abnormally increased vaginal discharge. On physical examination she has lower abdominal and cervical motion tenderness. Your diagnosis will be
 - A. Uterine myoma
 - B. Pyelonephritis
 - C. Pelvic Inflammatory Disease
 - D. Placenta previa

- 22. A 30 years old G5P4 mother is admitted in labor room with complaining of 32 weeks gestation, mild vaginal bleeding and abdominal pain. On physical examination, her blood pressure is 140/100 mmHg, abdomen is tense, tender and hard. Fetal heart sounds are not audible. What is the most likely diagnosis?
 - A. Placenta praevia
 - B. Abruptio placenta
 - C. Preterm labour
 - D. Urinary tract infection

23. A condition by which the fetus which is died and is retained with its placenta in the uterus is known as :-

- A. Threatened abortion
- B. Missed abortion
- C. Inevitable abortion
- D. Incomplete abortion

24. One of the following anti-TB drugs can cause congenital deafness to the new born If given to the mother during pregnancy:

- A. Rifampicin
- B. Streptomycin
- C. Ethambutol
- D. Pyrazinamide

- 25. A 20 years old primigravida comes in emergency at 32 weeks of gestation. She is complaining of blurring of vision, gross edema. On examination her B.P is 160/115 mm Hg. What is the most likely diagnosis of this mother?
- A. Hypertension
- B. Preeclampsia
- C. Eclampsia
- D. Thyroid disease

- 26. A 17-year-old, pregnant woman was brought to maternity ward, because was found to have generalized convulsion at the central market. (positive finding) unconscious, pale, BP 156/100 mmHg, hyperreflexia, urine dipstick for protein +. V/E Cervix effaced, dilated 4cm, station -1. How do you manage this patient?
- A. General measure, prophylactic antibiotic and immediate C/section
- B. General measures, antihypertensive, MgSO₄, resuscitation of the patient, fetal assessment and emergency c/section.
- C. General measures, antihypertensive, anticonvulsant and augmentation.
- D. General measures, BP control, fit control, mother stabilization and conservative management.

- 27. A G2 P1 with previous cesarean section due to obstructed labour comes for first antenatal visit at 34 weeks of gestation. She is seeking advice for home delivery this time. What will be the most dangerous complication in her case if we allow her to deliver at home by untrained birth attendant?
 - A. Prolonged latent phase
 - B. Arrest in second stage of labour
 - C. Delayed progress in active phase of labour
 - D. Rupture uterus

- 28. A 28 years old woman come to health center with complaints of vulval pruritus, dyspareunia, and whitish curd like vaginal discharge. Your diagnosis will be:-
 - A. Candidiasis
 - B. PID
 - C. Urethritis
 - D. Folliculitis

- 29. Mrs Sofia is a 20 years old women with known history of schizophrenia and found to be 16 weeks pregnant. She is considered to physically and mentally unable to care for the would-be born child and therapeutic abortion is considered.
- Which method of therapeutic abortion will be used for Mrs Sofia ?
 - A. Dilatation and curettage (D&C)
 - B. Prostaglandin preparation
 - C. Manual vacuum aspiration (MVA)
 - D. Caesarian Section

30. Mary is a 16 years old full term gravida I para 0 teen ager presented with prolonged labour and she is diagnosed to have CPD and is in the second stage of labor with live fetus.

The management of Mary will be :-

- A. Spontaneous vaginal delivery
- B. Oxytocin augmentation
- C. Caesarian section
- D. Instrumental delivery