

Sleep and Rest

Fundamental of nursing-II

By Abbas .M (Bsc, Msc-In Nursing)

Sleep & Rest

- Rest refers to a state of relaxation and calmness, of both mental and physical.
- The nurse should try to understand what activities and ,
- environments the client defines as **restful** .

Ctd'...

- Sleep refers to a state of altered consciousness during
- In which an individual experiences minimal physical activity and a general slowing of the body's physiological processes.

Ctd'...

- **The sleeper is unresponsive to most stimuli.**
- **It differs from unconsciousness in that the individual can be aroused by, the introduction of direct stimuli, such as a sudden noise.**
- **During sleep the metabolic rate decreases**
- **and resulting in a reduction in all component of vital sign .**

Ctd'...

- Rest and sleep are considered as essential to conserve energy, prevent fatigue,
- promote healing and relieve tension.

Ctd'...

- Adequate rest and sleep are important
- in Promoting general health states ,
- Ensuring recovery from illness,Aid in healing process.
- Increase ability to learn and concentration and recalling knowledge.

STAGES OF SLEEP

- **Quiet sleep-** During the deep states of quiet sleep:-
- Blood supply to the muscles is increased Energy is restored
- Tissue growth and repair occur
- Important hormones are released for growth and development.

Ctd'...

- **Active sleep:-**
- During active sleep:
- our brains are active and dreaming occurs.
- Our bodies become immobile
- Breathing and heart rates are irregular.

ctd'....

- All individuals require certain periods of calm and lesser activity
- so that their bodies can regain energy and rebuild stamina.
- The need for rest and sleep varies with age, developmental level, health status, activity level, and cultural norms.

Normal sleep patterns and it's requirements

- Newborn from -16-18 hours / day
- Infants: from -12-14 hours/ day
- Toddlers from -10-12 hours/ day
- Preschool from -11-12 hours/ day
- School-Age from -8- 12 hours/ day
- Adolescents from -8 -10 hours/ day
- Adult from -6-8 hours/ day
- Elders from -6 hours/ day

Sleep and rest disturbance

- This will imply :-
- Irritable, anxiety and stress.
- Fatigue.
- Seek assistance from physician.
- Poor concentration.
- Difficulty in making decisions.

Factors Affecting Sleep

- Psychological stress
- Lifestyle and habits
- Physical activity and exercise
- Dietary habits
- Environmental factors
- Medications, Cultural Norms

Alteration in Sleep Patterns

- sleep pattern disturbance is defined as:
- The state in which an individual experiences ,
- or at risk of experiencing a change in the quantity or quality of his or her rest and sleep pattern,
- Which is related to the person's biological and emotional needs.

Ctd'...

- Sleep disturbances can take many forms and are quite **common**.
- Alterations in sleep patterns are generally viewed as either :-
 1. primary sleep disorders :-those in which the sleep alteration is the **fundamental problem**

Ctd'...

2.Secondary sleep disorders:-

- Those in which the alteration has a **medical or clinical cause,**
- That results in or contributes to the sleep alteration

Ctd;....

- The most common sleep alterations include,
 - a) insomnia,
 - b) hypersomnia or narcolepsy,
 - c) sleep apnea,
 - d) sleep deprivation and Parasomnias
 - e) and parasomnias.

Insomnia

- Refers to the chronic inability to sleep or inadequate quality of sleep due to sleep prematurely ended
- Or interrupted by periods of wakefulness.
- Insomnia is not a **disease**, but it may be a manifestation of many **illnesses**

Ctd'....

- Treatment for insomnia is best
- directed at **modifying** those **factors** and
- behaviors that are causing insomnia.
- It is impossible to force sleep.

Hypersomnia or Narcolepsy

- Hypersomnia is an alteration in sleep pattern characterized by **excessive sleep**,
- especially in the daytime.
- Persons suffering from hypersomnia often feel that they cannot get enough sleep at night,
- and therefore they sleep very late into the morning and nap several times throughout the day.

Ctd'...

- manifests as sudden uncontrollable urges to fall asleep during the daytime.
- Causes of hypersomnia can be **physical** or **psychological**;
- treatment depends on addressing the underlying cause.

Treatment

- avoiding substances or activities that cause sleepiness, taking short daytime naps,
- or taking prescribed stimulant medications.

Sleep Apnea

- Sleep apnea refers to periods of sleep during which airflow stops for 10 seconds or more.
- Sleep apnea gives rise to complications as a result of **oxygen desaturation**
- And retention of carbon dioxide .

Ctd'....

- Short-term consequences may include
- cognitive impairment (including memory changes),
- personality changes, and impotence

Ctd'...

- A major problem is **daytime sleepiness**,
- which may interfere with functional abilities such as **driving** and **working with machine**.
- If untreated, it can result in the following
 - Hypertension
 - Cardiac arrhythmias

Ctd'....

- Right-sided congestive heart failure
- Cerebral vascular accident (stroke)
- Cognitive dysfunction
- Death

Ctd'...

- The first line of defense against apnea is:
- treating its **cause which may be :-**
- emotional,
- cardiac, or
- respiratory alteration

Sleep Deprivation

- Sleep deprivation is a term used to describe prolonged, inadequate quality and quantity of sleep,
- Sleep deprivation can result from age, prolonged hospitalization, drug and substance use,
- illness, and frequent changes in lifestyle patterns

Ctd'...

- Sleep and dreaming have a restorative value,
- Necessary for mental and emotional recovery,
- And enhance the ability to cope with emotional problems.

Ctd'...

- Therefore, sleep deprivation can **cause symptoms**
- ranging from irritability, hypersensitivity, and confusion to
- Apathy, sleepiness, and diminished reflexes.
- Treating or minimizing the factors that cause the sleep deprivation is the most **effective solution.**

Parasomnias

- Parasomnias refer to sleep alterations resulting from
- “an activation of physiological systems
- at inappropriate times during the sleep-wake cycle”

Ctd'....

- Somnambulism (sleepwalking), sleep talking, bed wetting,
- and bruxism (teeth grinding) are the most common parasomnias.
- Treatment for parasomnias varies, and care should be focused on helping the client,
- And family understand the disorder and its potential safety risks.

Nsg Assessment

- When discussing a client's sleep and rest patterns,
- The nurse should try to understand
- what activities and environments does the client defines as restful.

Ctd'...

1. Subjective data

- **Sleep Habits:** How would you rate the quality of your sleep?
- **Special Problems:** Do you ever experience difficulty with falling asleep?
- **Remaining asleep?** Do you ever feel fatigue after a sleep period?

Ctd'...

- Sleep Aids:
- What helps you to fall asleep?
- Medications?
- Reading?
- Relaxation technique?
- Watching TV?
- Listening to music?

Ctd'...

2. Objective Data

1. Observation for behaviors of sleep deprivation

Observe behavior

- a. Yawning
- b. Dozing (sleep slightly) during day
- c. Irritability
- d. Short attention span

Ctd'...

- **Nursing Diagnosis**

- Sleep Pattern Disturbances which may be related:-

- to body temperature,

- Depression, drugs ,

- anxiety, medical illness ,etc

Ctd'

- Again these can also be Evidenced by
- Dissatisfaction with sleep.
- Awakening earlier or later than desired
- Less than age normal total sleep time
- Etc

Ctd'....

- **Planning**

- Goal will be made to promote normal,
- restful sleep for clients ,
- having sleep disturbances and/ or difficulties.

Ctd'...

Nursing Interventions:-

- to Promote Sleep Prepare a restful environment
- Promote bedtime spirituals (religious ceremony)
- Promote relaxation and comfort
- Use night light
- Provide privacy

Ctd'...

- Use medications to produce sleep(if..)
- Encourage patient to void before sleep
- Remove any irritants against patients skin such as,
 - moist or wrinkled sheets and etc....

Ctd'...

- Make an environment conducive to sleep and rest by:
- Ensuring privacy and dignity
- Controlling room temperature
- Ensuring good ventilation and etc ...

Ctd'...

- Speaking quietly and avoiding unnecessary conversation(discussion)
- in the vicinity(area) of resting clients
- Reducing environmental noise such as
- telephones, alarms, machinery, and movement of equipment

Ctd'...

- Prioritizing care needs
- Minimize stress and anxieties
- Administer pain relievers as prescribed,
- and evaluate effectiveness
- Making sure the bed/chair is clean, dry and comfortable

Strategies to promote relaxation and facilitate sleep and rest

- Encourage client to empty bladder before retiring(go to bed)
- Reduce fluid intake prior to retiring
- Encourage the client to avoid caffeine products,
- such as tea, coffee and chocolate.

Ctd'...

- Hot milky drinks can helps to induce sleep
- Encourage the client to avoid alcohol
- Discourage the client from eating immediately prior to retiring ,
- and eating heavy meals late at night

Ctd'...

- Discourage inappropriate daytime napping
- Encourage daily exercise in keeping with client's abilities
- Provide appropriate daytime mental and physical stimulation

Ctd'...

- Educate the client regarding the need for a balance between **activity** , **rest**, and **correct** any **misconceptions**
- Provide cutaneous **stimulation** such as offering a warm bath, cold compress or massage

Ctd'...

- Employ appropriate relaxation techniques as Complementary therapies should be used with caution
- Administer sedation as prescribed and monitor effect.

Ctd'...

- This should be an intervention of **last resort** as many sedatives,
- hypnotics and anxiolytics have undesirable side effects and can be addictive
- Monitor client's pattern, quantity and quality of sleep and rest

Promoting Rest and Sleep

- Rest period
- Nonessential tasks
- Night time/early ambulation bathing
- Lab. work
- Cluster activity
- Visitor control
- Interventions

Comfort Measures for Promoting Sleep

- ♥ Administer hygiene measures for clients on bed rest
- ♥ Loose fitting nightwear
- ♥ Remove or change any irritants against the client's skin (moist ,drainage tubes)
- ♥ Position and support dependent body parts,
- ♥ protect pressure points and aid muscle relaxation

Comfort Measures for Promoting Sleep....

- ♥ Provide caps and socks for older clients and those prone to cold
- ♥ Void before bedtime
- ♥ Analgesics or sedatives 30 min. H.S.
- ♥ Bedtime massage/backrub
- ♥ Comfortable mattress and a clean dry bed!

Noisy Nightshift

- Close doors to clients' rooms/work areas if possible
- Telephone/paging equipment
- Noisy footwear
- Equipment's
- Bedside monitors
- TV/radio
- Conversations

Client's Environment

- Chairs
 - Straight back post surgery
 - Lounge chair
- Lighting
 - Over bed
 - Night light
 - Call light
- Over bed table
- Bedside table

Reading assignment

Physiology and stage of sleep

THE END

THANK YOU FOR YOUR ATTENTION //



Thank You