

Abbreviation	Definition	Abbreviation	Definition
aa	Affected area	BW	Body weight
a.c.	Before meals	C	Centigrade
ABW	Actual body weight	c or $\bar{c}$	With
ad	Up to	cap	Capsule
a.d.*	Right ear	cc*	Cubic centimeter
a.m.	Morning	cr, crm	Cream
amp	Ampule	d	Day
APAP*	Acetaminophen	disc, D.C.*, d/c*	Discontinue
Aq	Water	disp	Dispense
a.s.*	Left ear	div	Divide
ASA	Aspirin	DOB	Date of birth
ATC	Around the clock	DS	Double strength
a.u.*	Each ear	d.t.d.	Give as such doses
b.i.d.	Twice a day	DW	Distilled water
b.i.w.	Twice a week	D5NS	Dextrose 5% in normal saline
BMI	Body mass index	D5½NS	Dextrose 5% in ½ normal saline (0.45% NaCl)
BSA	Body surface area	D5W	Dextrose 5% in water

EC	Enteric coated	OTC	Over the counter
elix.	Elixir	o.u.*	Each eye
e.m.p.	As directed	oz	Ounce
F	Fahrenheit	p or per	By
fl or fld	Fluid	p.c.	After meals
ft.	Make	PCN	Penicillin
g or Gm	Gram	p.m.	Afternoon or evening
gr	Grain	p.o.	By mouth
gtt, gtts	Drop, drops	post	After
h, hr, or °	Hour	PPM	Parts per million
HCTZ*	Hydrochlorothiazide	pr	Rectally
h.s.*	At bedtime	pre-op	Before surgery
IBW	Ideal body weight	p.r.n.	As needed
ID	Intradermal	pulv.	Powder
IM	Intramuscular	q	Every
inj.	Injection	q.d.*	Every day
IU*	International units	q.i.d.	Four times a day
IUD	Intrauterine device	q.o.d.*	Every other day
IV	Intravenous	q.s.	Sufficient quantity
IVP	Intravenous push	q.s. ad	A sufficient quantity to make
IVPB	Intravenous piggy back	s or s̄	Without
jt or j-tube	Jejunostomy tube	sc*, sq*, subq*, or subcut	Subcutaneous
KVO	Keep vein open	Sig.	Write on label
L	Liter	SL	Sublingual
LE	Lower extremities	sol.	Solution
LR	Lactated Ringer's injection	ss*	One half
M ² or m ²	Square meter	stat.	Immediately
mcg or µg*	Microgram	supp.	Suppository

mg	Milligram	tab	Tablet
min	Minute	tal. dos.	Such dose
ml or mL	Milliliter	tbsp.	Tablespoon
MOM	Milk of magnesia	t.i.d.	Three times a day
mOsm or mOsmol	Milliosmole	tinc	Tincture
MR	May repeat	t.i.w.*	Three times a week
MRX_	May repeat _ times	top	Topically
NG or NGT	Nasogastric or nasogastric tube	tsp.	Teaspoon
No. or no.	Number	U* or u*	Unit(s)
noct.	Night	u.d.* or utdict	As directed
non rep. or N.R.	Do not repeat or no refills	UE	Upper extremities
NPO	Nothing by mouth	ung.	Ointment
NS	Normal saline (0.9% NaCl)	vag.	Vaginally
½ NS	Half-strength normal saline (0.45% NaCl)	vol.	Volume
NTG	Nitroglycerin	w/	With
o.d.*	Right eye	w.a.	While awake
oint.	Ointment	w/o	Without
o.s.*	Left eye	x	Times