

Date: 08/24/2015

Alloted time: 70 Mins

Addis Ababa University College of Health Sciences

Public Health Nutrition FINAL examination for Medical students

Name:

Abrar Sherif

ID.No

MDR B695/02

Part-I

Choose the best answer and write the letter of your choice in the answer sheet provided in the last page

E

1. What should be the goal of intervening in maternal nutrition?

- A. To improve maternal micronutrient status
- B. To improve maternal height and weight
- C. To provide nutritional supplementation
- D. To Provide therapeutic feeding
- ☒ E. A&B
- F. None

C

2. Of the following one is the best approach to assess causes of primary under nutrition.



- Dietary survey
- Clinical method ✓
- Biochemical method ✓
- D. Anthropometric measurements
- E. A&D
- F. None

Dietary

C

3. All can be considered as underlying causes of under nutrition, except.

- A. Living in unsanitary environment.
- B. Inadequate access for health service.
- C. Lack of adequate access for food at a household level.
- ☒ D. Cultural factors that prohibit proper feeding.
- E. A&C
- F. None

E

4. Iron supplementation for all pre pubertal and adolescent women is recommended when the prevalence of iron deficiency anemia in the general population is more than:

- A. 50%
- ☒ B. 40%
- C. 45%
- D. 30%
- E. 10%

F. None

F 5. One of the following does not accompany nutrition transition.

- A. Increased prevalence of non-communicable diseases
- B. Sedentary life style
- C. Decreasing life expectancy
- D. Decreasing fertility rate
- E. Decreasing cost of food
- F. D&E

F 6. Which one of the following micro nutrients play a leading role in decreasing maternal morbidity and mortality.

- A. Iron
- B. Vitamin A
- C. Iodine
- D. Carbohydrates
- E. A&B
- F. All except D

B 7. From the following dietary reference intakes one indicates the median nutritional requirement of a group of people.

- A. Adequate intake
- B. Estimated average requirement
- C. Recommended nutrient intake
- D. Recommended dietary intake
- E. C&D
- F. A&B

EA

C 8. Deworming is one of the commonest interventions to avert:

- A. Iodine deficiency
- B. Vitamin A deficiency
- C. Iron deficiency
- D. Protein deficiency
- E. Zinc deficiency
- F. None

B 9. The range of intake that is considered sufficient to prevent nutritional deficiency while avoiding toxicity is between.

- A. EAR & RNI
- B. RNI & UL
- C. EAR & UL

AVD.

EAR - RNI

- D. A & B
- E. None

A 10. Of the following one is not true about carbohydrates.

- ☒ A. The human body uses trioses and hexoses for metabolism
- B. Glycogen is the storage form of glucose in animals
- C. Starch is a form of poly saccharide
- D. Dextrin is intermediate product of starch digestion
- E. Constitutes majority portion of diet of the poor
- F. None

hexose
(fructose)

A 11. Of the following measures of protein quality one estimates the amount of amino acids that is retained in the body.

- ☒ A. Net protein utilization
- B. Chemical score
- C. Daily requirement
- D. Estimated upper intake level
- E. A&B
- F. All of the above

K

A 12. A tool that contains a list of food items consumed in the area

- ☒ A. Food frequency questionnaire
- B. Dietary history
- C. Twenty four hour recall
- D. Estimated food record
- E. Weighed food record
- F. None

A 13. It is a measurement which indicates body composition

- ☒ A. Waist to hip ratio
- B. Height
- C. Weight
- D. Head circumference
- E. None

A 14. The waist to hip ratio cut off point which indicates an increased risk for non-communicable diseases in males is.

- A. >0.85
- B. >90
- C. >1.5
- ☒ D. >1
- E. >2

F 20. Of the following one is not an ideal site to take skinfold measurement

- A. Mid axillary
- B. Sub-scapular
- C. Over the biceps
- D. Over the triceps
- E. Suprailiac
- ☒ F. None

A 21. Which one of the following components of energy requirements demand the highest daily energy expenditure in humans.

- ☒ A. Basal metabolism
- B. Thermic effect of food
- C. Physical activity
- D. Thermoregulation
- E. All require equal energy expenditure

D 22. Of the following one is available carbohydrate.

- A. Starch
- B. Hemicellulose
- C. Fiber
- D. Sugars
- ☒ E. A & D
- F. B & C

A 23. The most deficient of all the essential amino acids in a protein content of a given food is termed as

- ☒ A. Limiting Amino acids
- B. Chemical score
- C. Net protein utilization
- D. In complete proteins
- E. Deficient proteins
- F. None

A 24. How many essential amino acids exist in nature

- ☒ A. 8
- B. 12
- C. 20
- D. 10

E. 15
F. None

B 26. Observation of mucus membranes may be an important indicator for vitamin A deficiency.
☒ True ☐ B. False

A 26. Protein deficiency may prevent the release of vitamin A from the liver resulting in a low serum level of vitamin A.
☒ A. True ☐ B. False

A 28. Vitamin A deficiency is the leading cause of blindness in people below the age of 21 years.
☒ A. True ☐ B. False

A 28. Egg and milk are among the best sources of organic (heme) iron.
☒ A. True ☐ B. False

B 29. Improving the vitamin A status of individuals could also help improving other nutrients.
☒ True ☐ B. False

B 30. The unit for measuring the adult body mass index (BMI) is kilogram divided by kilometer squared (kg/km^2).
☐ A. True ☒ B. False

A 31. Marasmic children usually have poor appetite.
☐ A. True ☒ B. False

B 32. Under nutrition does not affect cognitive development.
☐ A. True ☒ B. False

B 33. The most vulnerable children to develop marasmus are children above 3 years of age.
☐ A. True ☒ B. False

B 34. Stunting usually represents an acute state of malnutrition.
☐ A. True ☒ B. False

E 36. Which of the following statement is False?

- A. Women who were malnourished as infants are more likely to give birth to malnourished babies
- B. Low birth weight is a risk factor for underweight
- C. Low birth weight is a risk factor for stunting
- D. all
- E. None

D 33. Increases in weight can be achieved within a woman's reproductive life by _____?

- ☒ Increasing caloric intake
- ☐ Reducing energy expenditure
- ☐ Increasing birth intervals
- D. All
- E. A and b

D 34. Which of the following is optimal behaviour to Improve Women's Nutrition?

- A. Exclusive breast feeding for six months
- B. Reduce workload during pregnancy
- C. Reduce workload during lactation
- D. All
- E. B and c

D 35. Which intervention is effective in Improving Women's Micronutrient Status?

- A. Parasite control
- B. Fortification
- C. Supplementation
- ☒ All
- E. B and c

E 36. Which of the following is/are important cause/s of IUGR in developing countries?

- ☒ small maternal size at conception
- ☒ low gestational weight gain
- ☐ Cigarette smoking during pregnancy
- ☒ All
- E. A and b

B 37. Which one is not true about anthropometric nutritional assessment method?

- A) It is widely used in the assessment of protein-energy-malnutrition
- ☒ It can identify specific nutrient deficiencies
- C) It can be used to monitor periodic changes in growth and/or body composition
- D) The procedures can assist in the identification of mild to moderate malnutrition.
- E) None of the above

D 38. Which of the following is less likely addressed by cross sectional nutritional survey?

- A) Establishing baseline nutritional data,
- B) Ascertaining overall nutritional status of a population

- C) Describing extent of existing nutritional problems
D) Identifying cause of malnutrition
E) None of the above
- C 39. Continuous monitoring of nutritional status of selected population groups is
A) Nutritional survey B) Nutritional screening C) Nutritional surveillance D) None
- E 40. All are true about Anthropometric Indices except
A) Weight for height identify children suffering from current or acute under-nutrition
B) Weight for age identifies both chronic and/or acute under-nutrition
C) Height for age identifies chronic malnutrition
D) All of the above
E) None of the above
- B 41. Preferred method for assessing adult nutritional status is
A) Mid upper arm circumference
B) Body mass Index
C) Height-for-age
D) Weight-for-age
E) None of the above

WAST

H/S
WASTING

WAST

WAST