

Answers

1. A
2. E
3. E
4. D
5. B
6. D
7. B
8. A, correct answer:
III, IV, VI
9. F
10. E
11. C
12. D
13. D
14. E
15. B
16. C
17. 1.5 MJ/day (360
kcal/day)
18. B
19. B
20. D
21. D
22. D
23. C
24. E
25. D
26. A
27. E
28. E
29. B
30. D
31. B
32. D
33. E
34. D
35. D
36. A
37. -1.98, mild
38. C
39. B
40. C
41.

I. Empower women, men, families, and communities to recognize pregnancy-related risks, and to take responsibility for developing and implementing appropriate responses to Them

II. Ensure access to a core package of maternal and neonatal health services, especially in rural areas where health facilities are limited

III. Create an environment supportive to safe motherhood and newborn health

42. E
43. D
44. C
45. A
46. D
47. A
48. A
49. D
50. IEC, FP, perinatal care, Tx of STIs, Tx of infertility, Safe abortion, referral for FP and dx & Tx of further complications, discouragement of harmful traditional services
51. D
52. C
53. E
54. B

55. Parenteral antibiotics, parenteral oxytocic drugs, parenteral sedatives for eclampsia, manual removal of placenta, removal of retained products, assisted vaginal delivery

- 56.
- I. Decide number, spacing and timing of children
 - II. Attain highest standard of sexual and reproductive health
 - III. Decision concerning reproduction

57. B
58. E
59. GATHER
60. A
61. C
62. B
63. B
64. D
65. C
66. C
67. B
68. B
69. E
70. A
71. D
72. A
73. D
74. E
75. B
76. E
77. D
78. B
79. B

80. D

81. A

82. 87.59%, mild

83. E

84. A

85. D

86. Prematurity and
SGA, SGA