



Complete
Mark 1.00 out of 1.00
Flag question

Gastric glands are:

- ☐ a. Simple straight tubular
- ☐ b. Simple coiled tubular
- ☐ c. Compound tubular
- ☐ d. Simple acinar
- ☒ e. Simple branched tubular

Question 2

Complete
Mark 1.00 out of 1.00
Flag question

Which of the following mark(s) best ileum from the rest of small intestinal region?

- ☒ a. Peyer's patches
- ☐ b. Plicae semilunaris
- ☐ c. Absence of teniae coli
- ☐ d. Presence of long and broad leaf like villi
- ☐ e. Plicae circularis

Question 3

Complete
Mark 1.00 out of 1.00
Flag question

Ring like rhythmical contractions and relaxation of the intestine is _____.

- ☐ a. Mass movement
- ☐ b. Retro propulsion
- ☐ c. Migrating Motor Complex
- ☒ d. Segmentation contraction
- ☐ e. Peristalsis

Question 4

Complete
Mark 1.00 out of 1.00
Flag question

The type of epithelium lining the pharynx is:

- ☐ a. Transitional
- ☒ b. Stratified squamous non-keratinized
- ☐ c. Simple columnar
- ☐ d. Stratified squamous keratinized
- ☐ e. Simple cuboidal



- c. Gastrin
- d. Motilin
- e. Secretin

Question 51

Complete

Mark 1.00 out of 1.00

Flag question

A patient was diagnosed with pancreatitis due to a reflux of bile into the pancreatic duct caused by a gallstone. The stone is likely to be lodged at the:

- a. Cystic duct
- b. Common bile duct
- c. Right hepatic duct
- d. Common hepatic duct
- e. Hepatopancreatic ampulla

Question 52

Complete

Mark 1.00 out of 1.00

Flag question

Dietary fat, after being processed, is extruded from the mucosal cells of the gastrointestinal tract into the lymphatic ducts in the form of:

- a. Chylomicrons
- b. Triglycerides
- c. Free fatty acids
- d. Diglycerides
- e. Monoglycerides

Question 53

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following cell type is correctly matched with its secretion?

- a. Chief cells – HCL
- b. Parietal cells – Pepsinogen
- c. G-cells – secretin
- d. D-cells – Gastrin
- e. Enterochromafin like cells – Histamine

Question 54

Complete

Which of the following condition most likely stimulate the secretion of cholecystokinin from the duodenum?

Question 52

Complete

Mark 1.00 out of 1.00

Flag question

Dietary fat, after being processed, is extruded from the mucosal cells of the gastrointestinal tract into the lymphatic ducts in the form of:

- a. Chylomicrons
- b. Triglycerides
- c. Free fatty acids
- d. Diglycerides
- e. Monoglycerides

Question 53

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following cell type is correctly matched with its secretion?

- a. Chief cells – HCL
- b. Parietal cells – Pepsinogen
- c. G-cells – secretin
- d. D-cells – Gastrin
- e. Enterochromatin like cells – Histamine

Question 54

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following condition most likely stimulate the secretion of cholecystekinin from the duodenum?

- a. Elevated H⁺ the gut
- b. Elevated acid in the gut
- c. Elevated glucose in the gut
- d. Elevated pH in the gut
- e. Elevated fat in the gut

Question 55

Complete

Mark 1.00 out of 1.00

Flag question

Regarding vermiform appendix the following statements are correct except:

- a. It is attached to the cecum
- b. It has a blind free end and attached end.

Question 54

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following condition most likely stimulate the secretion of cholecystokinin from the duodenum?

- a. Elevated H^+ in the gut
- b. Elevated acid in the gut
- c. Elevated glucose in the gut
- d. Elevated pH in the gut
- e. Elevated fat in the gut

Question 55

Complete

Mark 1.00 out of 1.00

Flag question

Regarding vermiform appendix the following statements are correct except:

- a. It is attached to the cecum
- b. It has a blind free end and attached end.
- c. It is surgically removed by way of McBurney's incision
- d. Its common position is behind the cecum (retrocecal)
- e. Its arterial supply arises from right colic artery

Question 56

Complete

Mark 1.00 out of 1.00

Flag question

The anastomotic artery running along the border of the large intestine is called the:

- a. Ileocolic
- b. Marginal
- c. Vasa rectae
- d. Jejunal
- e. Arcade

Mark 1.00 out of 1.00

Flag question

- ... responsible for the migrating motor complex is
- a. Gastric inhibitory peptide
 - b. Secretin
 - c. Gastrin
 - d. Motilin
 - e. Cholecystokinin

Question 47

Complete

Mark 1.00 out of 1.00

Flag question

The effect of parasympathetic stimulation on the GIT include:

- a. Constriction of the pylorus
- b. Contraction of the internal anal sphincter
- c. Contraction of the gallbladder
- d. Contraction of the external anal sphincter
- e. Vasoconstriction on GI blood vessels

Question 48

Complete

Mark 1.00 out of 1.00

Remove flag

In Porto-systemic anastomosis, the systemic tributary around the umbilicus is the:

- a. Paraumbilical vein
- b. Azygos vein
- c. Lower intercostals vein
- d. Esophageal vein
- e. Inferior superficial epigastric vein

Question 49

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following is the transporter protein responsible for the entry of glucose into the intestinal enterocytes?

- a. Glut-2
- b. Glut-1
- c. Glut-5
- d. SGLT-2
- e. SGLT-1

- c. Fungiform
- d. Filiform
- e. Foliate

Question 36

Complete

Mark 1.00 out of 1.00

Flag question

True about pancreatic secretion,

- a. Enzyme secretion is enhanced by the hormone CCK
- b. It contains enzymes which produce large amounts of monosaccharides from disaccharides
- c. Bicarbonate secretion is enhanced by the hormone gastrin
- d. The digestive enzymes function optimally at a very low pH
- e. Enzyme-rich juice is secreted in response to a fall in duodenal pH

Question 37

Complete

Mark 1.00 out of 1.00

Flag question

Gastric emptying is primarily controlled by,

- a. The volume of chyme in the stomach
- b. The conditions and factors from large intestine
- c. Conditions and factors in the duodenum
- d. The type of food in the stomach
- e. The degree of fluidity of the chyme in the stomach

Question 38

Complete

Mark 0.00 out of 1.00

Remove flag

Which of the following statement is not true regarding ion absorption and secretion in the intestine?

- a. Calcium absorption depends on vitamin D
- b. Bicarbonate ion is absorbed from small intestine in the form of carbonic acid
- c. Aldosterone increases potassium secretion
- d. Aldosterone increases sodium reabsorption
- e. Na^+/K^+ pump contributes for sodium absorption

Question 39

Complete

Mark 1.00 out of

Following the return from the physiological umbilical herniation the cecum and appendix initially occupy:

- a. Midabdominal position behind the umbilicus

Question 39

Complete

Mark 1.00 out of 1.00

Remove flag

Following the return from the physiological umbilical herniation the cecum and appendix initially occupy:

- a. Midabdominal position behind the umbilicus
- b. Left upper quadrant of the abdominal cavity
- c. Subhepatic position in the upper right quadrant of the abdominal cavity
- d. Left lower quadrant of the abdominal cavity
- e. Right lower quadrant of the abdominal cavity

Question 40

Complete

Mark 0.00 out of 1.00

Flag question

You were asked to assist in a surgical operation on a young patient to treat an ulcer in the first part of the duodenum. You would expect that the surgeon will approach the ulcer by doing an anterior abdominal wall incision in the following region:

- a. Left inguinal
- b. Hypogastric
- c. Right hypochondriac
- d. Left hypochondriac
- e. Epigastric

Question 41

Complete

Mark 1.00 out of 1.00

Flag question

The reason why the main pancreatic duct and common bile duct drain together as the hepatopancreatic duct is:

- a. The development of the liver from the dorsal pancreatic bud.
- b. The two ducts fused because of the rotation of duodenum.
- c. The development of the ventral pancreatic bud from the hepatic bud.
- d. The two ducts fused because of the rotation of the pancreas.
- e. The development of the liver from the ventral pancreatic bud.

Question 42

Complete

Mark 0.00 out of 1.00

Flag question

The large intestine as opposed to the small intestine has:

- a. More number of villi
- b. More developed Plicae circularis
- c. More developed mucosa associated lymphoid tissue
- d. No serosa

Question 41

Complete

Mark 1.00 out of 1.00

Flag question

The reason why the main pancreatic duct and common bile duct drain together as the hepatopancreatic duct is:

- a. The development of the liver from the dorsal pancreatic bud.
- b. The two ducts fused because of the rotation of duodenum.
- c. The development of the ventral pancreatic bud from the hepatic bud.
- d. The two ducts fused because of the rotation of the pancreas.
- e. The development of the liver from the ventral pancreatic bud.

Question 42

Complete

Mark 0.00 out of 1.00

Flag question

The large intestine as opposed to the small intestine has:

- a. More number of villi
- b. More developed *Plicae circulares*
- c. More developed mucosa associated lymphoid tissue
- d. No serosa
- e. No microvilli

Question 43

Complete

Mark 1.00 out of 1.00

Flag question

The type of epithelium lining the surface of stomach where digestion is taking place in an acidic environment is:

- a. Stratified squamous keratinized
- b. Transitional
- c. Simple columnar
- d. Stratified squamous nonkeratinized
- e. Simple cuboidal

Question 44

Complete

Mark 1.00 out of 1.00

Flag question

A patient was admitted with symptoms of bowel obstruction. Further examination revealed that the obstruction was caused by the nutcracker-like compression of the bowel between the mesenteric artery and the abdominal aorta. The compressed bowel is most likely _____ part of the duodenum.

- a. 3rd
- b. 1st
- c. 4th
- d. 1st and 2nd
- e. 2nd

- d. Gastrin stimulates bicarbonate secretion from the pancreas
- d. Gastrin stimulates acid secretion
- e. Parasympathetic stimulation has inhibitory effect at the sphincters

Question 32

Complete

Mark 1.00 out of 1.00

Flag question

Stomach bed is formed by all, except:

- a. Left kidney
- b. Spleen
- c. Gall bladder
- d. Left suprarenal
- e. Pancreas

Question 33

Complete

Mark 1.00 out of 1.00

Flag question

The principal paracrine secretion involved in the inhibitory feedback regulation of gastric acid secretion is:

- a. Histamine
- b. Gastrin
- c. Somatostatin
- d. Secretin
- e. Acetylcholine

Question 34

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following is true regarding gastric secretion?

- a. Distension of the stomach reduces gastric secretion
- b. HCl secretion requires energy
- c. It is reduced by the hormone gastrin
- d. It starts only when food enters the stomach
- e. Acetylcholine decreases gastric secretion

Question 35

Complete

Mark 1.00 out of 1.00

Which papillae of the tongue are found in a single row in front of sulcus terminalis and have a large number of taste buds on their sides?

- a. Both Filiform and foliate
- b. Circumvallate

Question 27

Complete

Mark 1.00 out of 1.00

Flag question

A surgeon performing an appendectomy was unable to identify the base of the appendix due to massive adhesions in the peritoneal cavity. Eventually he identified the cecum and was able to localize the base of the appendix. What anatomical structure(s) on the cecum would he have used to find the base of the appendix?

- ☐ a. Semilunar folds
- ☐ b. Omental appendages
- ☐ c. Ileal orifice
- ☐ d. Haustra coli
- ☐ e. Teniae coli

Question 28

Complete

Mark 1.00 out of 1.00

Flag question

The parotid of adult humans is:

- ☐ a. Compound purely mucous
- ☐ b. Simple branched purely mucous
- ☐ c. Compound purely serous
- ☐ d. Simple mixed mucous and serous, but mainly serous
- ☐ e. Compound mixed mucous and serous, but mainly serous

Question 29

Complete

Mark 0.00 out of 1.00

Remove flag

The posterior boundary of the epiploic foramen (of Monro) is the:

- ☐ a. First part of the duodenum
- ☐ b. Common bile duct
- ☐ c. Hepatic portal vein
- ☐ d. Inferior vena cava
- ☐ e. Caudate lobe of the liver

Question 30

Complete

Mark 0.00 out of 1.00

Flag question

Which of the following substances must be further digested before it can be absorbed by specific carriers in intestinal cells?

- ☐ a. Tripeptides
- ☐ b. Amino acids
- ☐ c. Lactose

Complete
Mark 0.00 out of 1.00
Remove flag

The posterior boundary of the epiploic foramen (of Monro) is the:

- a. First part of the duodenum
- b. Common bile duct
- c. Hepatic portal vein
- d. Inferior vena cava
- e. Caudate lobe of the liver

Question 30
Complete
Mark 0.00 out of 1.00
Flag question

Which of the following substances must be further digested before it can be absorbed by specific carriers in intestinal cells?

- a. Tripeptides
- b. Amino acids
- c. Lactose
- d. Galactose
- e. Fructose

Question 31
Complete
Mark 1.00 out of 1.00
Flag question

Which of the following statement is not true regarding regulation of gastrointestinal activities:

- a. Sympathetic stimulation decreases GI motility
- b. Motilin secretion is reduced during fasting
- c. Secretin increases bicarbonate secretion from the pancreas
- d. Gastrin stimulates acid secretion
- e. Parasympathetic stimulation has inhibitory effect at the sphincters

Question 32
Complete
Mark 1.00 out of 1.00
Flag question

Stomach bed is formed by all, except:

- a. Left kidney
- b. Spleen
- c. Gall bladder
- d. Left suprarenal
- e. Pancreas

Question 18

Complete

Mark 1.00 out of 1.00

Flag question

Muscularis mucosae of the esophagus is:

- a. Found as outer longitudinal and inner circular layers
- b. Contains myenteric plexus
- c. Well-formed than the muscularis externa
- d. Made by both skeletal and smooth muscles
- e. Found as longitudinal layer with no circular layer

Question 19

Complete

Mark 1.00 out of 1.00

Flag question

A radiological examination of a patient revealed a large tumor in the quadrate lobe of the liver. During the surgical removal of the tumor, one of the vessels that needs to be clamped to effectively control bleeding is the:

- a. Left gastric artery
- b. Right gastric artery
- c. Cystic artery
- d. Left hepatic artery
- e. Right hepatic artery

Question 20

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following structures of the teeth and periodontium is easily brittle and could shatter?

- a. Cementum
- b. Enamel
- c. Periodontal ligaments
- d. Alveolar bone forming tooth socket
- e. Dentin

Question 21

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following is not related to the spleen?

- a. Pancreas
- b. Colon
- c. Stomach
- d. Duodenum

- c. Periodontal ligaments
- d. Alveolar bone forming tooth socket
- e. Dentin

Question 21

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following is not related to the spleen?

- a. Pancreas
- b. Colon
- c. Stomach
- d. Duodenum
- e. Diaphragm

Question 22

Complete

Mark 1.00 out of 1.00

Flag question

A surgeon needs to construct a bypass/shunt between the veins of the portal and caval systems to circumvent insufficient drainage through the natural porto-caval anastomoses. Which plan is likely to be successful?

- a. Superior mesenteric vein to splenic vein
- b. Left colic vein to middle colic vein
- c. Splenic vein to left renal vein
- d. Coronary vein to right gastroepiploic vein
- e. Inferior mesenteric vein to splenic vein

Question 23

Complete

Mark 1.00 out of 1.00

Flag question

Cells of gastric glands that produce gastric intrinsic factor which binds to vitamin B12 are:

- a. Paneth cells
- b. Zymogenic cells
- c. Oxyntic cells
- d. Basket cells
- e. Mucous neck cells

Question 24

Which of the following is true regarding slow wave electrical potentials in gastrointestinal smooth muscle?

Question 24

Complete

Mark 0.00 out of 1.00

Flag question

Which of the following is true regarding slow wave electrical potentials in gastrointestinal smooth muscle?

- a. They occur at the same frequency in all portions of the GIT
- b. Influx of sodium induces depolarization
- c. Influx of potassium results in repolarization
- d. They are true action potentials
- e. They can cause muscle contraction in all parts of GIT

Question 25

Complete

Mark 1.00 out of 1.00

Flag question

Regarding to the development of the hind gut, which of the following statements is wrong?

- a. Is supplied by the inferior mesenteric artery.
- b. Forms a loop with its convexity directed ventrally same as duodenum.
- c. Is not involved in the physiological umbilical herniation.
- d. Does not give rise to the distal part of anal canal.
- e. Has a dilated distal part known as cloaca which is divided later by urorectal septum into urogenital sinus ventrally and anorectal canal dorsally.

Question 26

Complete

Mark 1.00 out of 1.00

Flag question

Stellate or fat storing cells of Ito found in the liver are located:

- a. In the space of Disse
- b. In the space of Moll
- c. In the liver sinusoids
- d. At the portal canal
- e. Around the central vein

Question 27

Complete

Mark 1.00 out of 1.00

Flag question

A surgeon performing an appendectomy was unable to identify the base of the appendix due to massive adhesions in the peritoneal cavity. Eventually he identified the cecum and was able to localize the base of the appendix. What anatomical structure(s) on the cecum would he have used to find the base of the appendix?

- a. Semilunar folds
- b. Omental appendages
- c. Ileal orifice

Question 18

Complete

Mark 1.00 out of 1.00

Flag question

Muscularis mucosae of the esophagus is:

- a. Found as outer longitudinal and inner circular layers
- b. Contains myenteric plexus
- c. Well-formed than the muscularis externa
- d. Made by both skeletal and smooth muscles
- e. Found as longitudinal layer with no circular layer

Question 19

Complete

Mark 1.00 out of 1.00

Flag question

A radiological examination of a patient revealed a large tumor in the quadrate lobe of the liver. During the surgical removal of the tumor, one of the following arteries should be ligated to control bleeding is the:

- a. Left gastric artery
- b. Right gastric artery
- c. Cystic artery
- d. Left hepatic artery
- e. Right hepatic artery

Question 20

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following structures of the teeth and periodontium is easily brittle and could shatter?

- a. Cementum
- b. Enamel
- c. Periodontal ligaments
- d. Alveolar bone forming tooth socket
- e. Dentin

Question 21

Complete

Mark 1.00 out of 1.00

Flag question

Which of the following is not related to the spleen?

- a. Pancreas
- b. Colon
- c. Stomach
- d. Duodenum

Question 14

Complete

Mark 1.00 out of 1.00

Flag question

The stomach has the ability to absorb:

- a. Calcium ion
- b. Fat soluble vitamins
- c. Vitamin B12
- d. Alcohol
- e. Amino acids

Question 15

Complete

Mark 1.00 out of 1.00

Flag question

The following structures pass through porta-hepatic except:

- a. None of the above
- b. Right and left hepatic ducts
- c. Right and left hepatic veins
- d. Right and left portal veins
- e. Right and left hepatic arteries

Question 16

Complete

Mark 0.00 out of 1.00

Flag question

Which of the following is correct for distinguishing between parotid and exocrine portion of pancreas?

- a. Pancreas has centroacinar cells and many striated ducts, while parotid has none of these.
- b. Parotid has centroacinar cells and many striated ducts, while pancreas has none of these.
- c. Parotid has intercalated ducts, while pancreas none of these.
- d. Pancreas has centroacinar cells, while parotid has many striated ducts.
- e. Parotid has centroacinar cells, while pancreas has many striated ducts.

Question 17

Complete

Mark 1.00 out of 1.00

Remove flag

All are the mechanisms for absorption of sodium from the small intestine, except:

- a. Co-transport with galactose
- b. Co-transport with glucose
- c. Co-transport with potassium
- d. $\text{Na}^+ - \text{H}^+$ exchanger
- e. Co-transport with amino acids

- d. The stomach rotates, so that the distal end moves to the right and up
- e. The site of junction of foregut and midgut is the 2nd part of duodenum just proximal to the entrance of common bile duct

Question 13

Complete

Mark 1.00 out of 1.00

Flag question

Which one of the following is the major chemical digestive activity that takes place in the stomach?

- a. Digestion of starch
- b. Digestion of disaccharides
- c. Digestion of fats
- d. Digestion of cellulose
- e. Digestion of proteins

Question 14

Complete

Mark 1.00 out of 1.00

Flag question

The stomach has the ability to absorb:

- a. Calcium ion
- b. Fat soluble vitamins
- c. Vitamin B12
- d. Alcohol
- e. Amino acids

Question 15

Complete

Mark 1.00 out of 1.00

Flag question

The following structures pass through porta-hepatic except:

- a. None of the above
- b. Right and left hepatic ducts
- c. Right and left hepatic veins
- d. Right and left portal veins
- e. Right and left hepatic arteries

Question 16

Complete

Mark 0.00 out of 1.00

Flag question

Which of the following is correct for distinguishing between parotid and exocrine portion of pancreas?

- a. Pancreas has centroacinar cells and many striated ducts, while parotid has none of these.
- b. Parotid has centroacinar cells and many striated ducts, while pancreas has none of these.
- c. Parotid has intercalated ducts while pancreas none of these.



Complete
Mark 1.00 out of 1.00
Flag question

ring like rhythmical contractions and relaxation of the intestine is _____

- a. Mass movement
- b. Retro propulsion
- c. Migrating Motor Complex
- d. Segmentation contraction
- e. Peristalsis

Question 4
Complete
Mark 1.00 out of 1.00
Flag question

The type of epithelium lining the pharynx is:

- a. Transitional
- b. Stratified squamous non-keratinized
- c. Simple columnar
- d. Stratified squamous keratinized
- e. Simple cuboidal

Question 5
Complete
Mark 1.00 out of 1.00
Flag question

The origin of electrical slow wave activity in gastrointestinal tract smooth muscle is:

- a. The interstitial cells of Cajal
- b. The messiner's plexus
- c. Postganglionic parasympathetic fibers
- d. Preganglionic sympathetic fibers
- e. The myenteric plexus

You are logged in as [MINTESNOT WONDIMAGEGN DEMEKE ug](#) (Log out)
MTG



DELL

e. The myenteric plexus

Question 6

Complete

Mark 1.00 out of 1.00

Flag question

If a hernia enters into the scrotum, it is most likely:

- a. Direct inguinal hernia
- b. Hiatal hernia
- c. Femoral hernia
- d. Indirect inguinal hernia
- e. Obturator hernia

Question 7

Complete

Mark 1.00 out of 1.00

Flag question

The following statements are correct for duodenum except:

- a. Its length is 25 Cm (10 inches).
- b. Its 4th part is developed from the mid-gut.
- c. Its upper part contains plica-circulans
- d. Its 2nd part is related to the anterior surface of the Rt. Kidney.
- e. its 2nd part is supplied by both superior and inferior pancreaticoduodenal artery.

Question 8

Complete

Mark 0.00 out of 1.00

Flag question

All of the following substances use Na^+ dependant transport, except

- a. Fructose
- b. Glucose
- c. Galactose
- d. Tryptophan
- e. Proline

Question 9

Complete

Mark 1.00 out of 1.00

Flag question

A new mother calls the pediatrician because she is concerned that her infant defecates after every meal. Which of the following is the

- a. Peristaltic reflex
- b. The gastrocolic reflex
- c. The colonoileal reflex



DELL

Complete
Mark 1.00 out of 1.00
Flag question

- a. Peristaltic reflex
- b. The gastrocolic reflex
- c. The coloioilal reflex
- d. The enterogastric reflex
- e. The gastroileal reflex

Question 10
Complete
Mark 1.00 out of 1.00
Flag question

Which of the following is correct regarding saliva secretion?

- a. Parasympathetic stimulation has both inhibitory and stimulatory effect
- b. Primary saliva is hypotonic to plasma
- c. The acinar cells secrete isotonic saliva
- d. The ductal cells secrete hypertonic saliva
- e. Maximal salivation decreases the tonicity of saliva

Question 11
Complete
Mark 1.00 out of 1.00
Flag question

The brush border enzyme of the small intestine digest:

- a. Proteins
- b. Lactose
- c. Glucose
- d. Galactose
- e. Starch

Question 12
Complete
Mark 1.00 out of 1.00
Flag question

Regarding the development of gut:

- a. The epithelium of the GIT is entirely of endodermal origin.
- b. The midgut is temporarily connected to the yolk sac by the allantois
- c. The primitive gut tube is in open communication with the chorionic cavity during the 4th week
- d. The stomach rotates, so that the distal end moves to the right and up
- e. The site of junction of foregut and midgut is the 2nd part of duodenum just proximal to the entrance of common bile duct