

1 Which one of the following is not the principle of nutrient reference level?

- a. Refer to the average daily need over a reasonable period of time
- b. Refer to level of intake needed to maintain health in already healthy individuals
- c. It takes in to account possible interaction that involve nutrients and other dietary components
- d. Based on typical dietary pattern of a country
- e. None

2 Which one of the following statement is true?

- a. Basal metabolism rate represents 45-70% of daily total energy expenditure
- b. Basal metabolism is the most variable and account for the largest component of daily energy expenditure
- d. Dietary reference intakes provide qualitative advice to the public about diet and chronic disease prevention
- e. None

3 One of the following is a fundamental shift and a new approach in the reproductive health progress milestones.

- a. Adapting and expanding population policies
- b. Removing the barrier to family planning services to the level possible
- c. Considering the reproductive health care as a right of individuals or couples
- d. Control of female fertility
- E. All of the above

4 Which one of the following approach is not the limitation of risk/traditional approach of antenatal care service provision?

- a. It considers every pregnancy is at risk
- b. Majority who are considered as high risk may not develop complication at the end
- c. Risk factors identification is not a good indicator for complication
- d. Unnecessary utilization of resources for those who are considered as high risk

5 Respecting cultural diversity contributes less to adolescent and youth program success.

- a. True
- b. False

6 Which one of the following is not true about magnitude of maternal mortality?

- a. Maternal death is a global problem particularly in developing countries
- b. Sub-Saharan countries account for 99% of maternal mortality
- c. Maternal mortality is the leading cause of female adult death
- d. Average maternal mortality declined from 1990 to 2015 by 2.3% annually

7 Which one of the following measure is not included in the preventive measure for Iodine deficiency disorder?

- a. Universal salt iodization
- b. Water iodization
- c. Deworming
- d. Health education
- e. None

8 Which one of the following is not correct about WHO classification of public health problem of xerophthalmia?

- a. Bitot spot of $> 0.5\%$
- b. Corneal scar of $> 0.05\%$
- c. Corneal xerosis of $> 0.01\%$
- d. Night blindness of $> 5\%$
- e. None

9 In a nutritional survey, a female child of 90cm weighs 9kg. Note: The median weight for a female child of height 90cm is 13.7kg. Calculate percent of median?

- a. 65.6%, Moderate wasting
- b. 72.4%, Mild wasting
- c. 72.4%, Mild stunting
- d. 65.6% Sever wasting

10 Which of the following statements is not true?

- a. Good nutrition is not intuitive.
- b. People cannot tell when their children are becoming malnourished because it is not easy to detect for an untrained eye.
- c. Improving nutrition is pro-poor and increases the income-earning potential of the poor.
- d. All
- e. None

11 The waist to hip ratio cut off point which indicates an increased risk for non-communicable diseases in male is:

- a. > 0.85
- b. > 0.87
- c. > 1`
- d. > 2

12 Which of the following statement is correct?

- a. Zinc deficiency could be a cause of night blindness
- b. Zinc supplementation can reduce child morbidity and mortality
- c. A rate of stunting > 20% is indicative of zinc deficiency
- d. In areas where high Phytate diet is a staple zinc deficiency might be common
- e. All
- f. None

13 According to 2016 Ethiopian demographic and health survey (EDHS), the modern contraceptive prevalence rate is

- a. 42
- b. 29
- c. 36
- d. 27
- e. 35

14 Which of the following statement is not correct?

- a. Nutrition survey is important to establish base line data
- b. Nutrition survey is important to understand the extent of nutritional problem

- c. Nutrition survey is suitable to identify high risk group
- d. Nutrition survey is important to formulate prediction based on current trend
- e. All are correct

15 Choose the most common family planning delivery strategies in Ethiopia.

- a. Community based distribution
- b. Commercial retail sales
- c. Clinic based services
- d. Outreach based services
- e. All of the above strategies are equally applicable

16 Which of the following shows a recent acute malnutrition?

- a. Weight for age $< -2SD$
- b. Weight for height $< -2SD$
- c. Height for age $< -2SD$
- d. Weight for age $< -3SD$

17 Which of the following is not true about direct causes of maternal mortality?

- a. It results from obstetric complications
- b. It is related with incorrect treatment during pregnancy and delivery
- c. Accounts for 80% of maternal death
- d. It considers aggravated disease conditions by physiologic effects of pregnancy

18 Which one of maternal health services is most improved from 2000 to 2016 based on EDHS report?

- a. Birth attended by skilled birth attendant
- b. Postnatal care
- c. Antenatal care
- d. All of the above

19 Which one of the following causes of maternal mortality took short time for death of a mother?

- a. Haemorrhage
- b. Obstructed labour

- c.Abortion
- d.Hypertensive disorder of pregnancy

20 One of the following statements is not true?

- a.HAZ of $< -2SD$ indicates a chronic malnutrition
- b.waterloo classification helps to differentiate sever forms of malnutrition
- c.For intervention with immediate impact, we should use WHZ score
- d.Prevalence of stunting is usually higher than prevalence of Wasting
- e.All are correct

21 One of the following is not among the rationale for investing on adolescent and youth health?

- a.Available funding opportunities for adolescent and youth health and particularly that of reproductive health services
- b.Existence of real and perceived barriers that limit access and utilization of RH information and services
- c.The magnitude of adolescents and youths which can't be ignored
- D.Favourable time for transferring knowledge, information and skills about sexuality, relationships, marriage, and parenting

22 Which one measures the volume of program activity among the family planning indicators

- a.Contraceptive prevalence rate
- b.Ever use of contraceptive
- c.Total fertility rate
- d.Couple years of protection
- e.Method mixes

23 All are true about Anthropometric Indices except

- a.Weight for height identify children suffering from current or acute under-nutrition
- b.Weight for age identifies both chronic and/or acute under nutrition
- c.Height for age identifies chronic malnutrition
- d.All
- e.None of the above

24 One of the following intervention is not recommended to avoid third level delay for maternal health service utilization.

- a. **Availing necessary equipment , logistics and supplies in the health facility**
- b. **Improving quality of services**
- c. **Improve health provider mix for comprehensive service provision**
- d. **None of the above**

25 In a nutrition survey, a male child of 84cm weighs 9.3kg. Note: the reference median weight for boys of height 84cm is 11.7kg and the standard deviation is 0.908. What is the Z-score?

- A.- 2.6, Moderate wasting
- b.- 2.6, Moderate stunting
- c.-1.8, Mild wasting
- d.1.8, Normal

26 Which of the following is not a true statement about adolescence and adolescents?

- a.Physical maturation occurs earlier in girls than in boys
- b.Puberty is a marker of the period of adolescence
- c.psychological and emotional readiness occur earlier than physical maturation
- d.The duration and defining characteristics of adolescence may vary across time, culture, and socio-economic situations

27 Family planning is the same with birth control.

- a.False
- b.True

28 The prevalence of stunting at birth, in the first six month of life and in the age between 6-24 months respectively is

- a.10%, 10%, 50%
- b.50%, 20%, 20%
- C.20%, 20% 50%
- d.10%, 20%, 50%

29 A lactating woman should do the following except

- a.Breast feed at least 10 times day and night over 24 hr
- b.Exclusively breast feed for the first 24 month

- c. Exclusively breast feed for the first 6 month
- d. Should empty one breast before switched to the other breast
- e. All are correct

30 Which one of the following is correctly matched?

- a. Keratomalacia- ulceration and perforation of the cornea
- b. corneal ulceration- reversible structural change of the cornea
- c. corneal xerosis- corneal surface becomes dry
- d. Bitot's spot- the earliest structural change
- e. A and B
- f. A and C

31 Which one of the following is WRONG about the objective of reproductive health care?

- a. providing information about reproductive health services
- b. to enable and support decision making in family planning and childbearing
- c. to meet the reproductive health need of people based on the available resources
- d. none

32 One of the following is not true about marasmus

- A. Old man face
- B. No Fatty liver
- c. Weight for age between 60-80%
- d. Common among children less than 2 years of age

33

Which one of the following statement is true?

- a. A 20% of low hemoglobin for age is considered mild
- b. A higher PH level enhance iron absorption
- c. Heme Iron is found from plant source
- d. All
- e. None

34 Which of the following statement is False?

- a.Unrestricted exclusive breastfeeding result in ample production
- b.Vitamin B, C and A content of breastmilk is not sensitive to dietary intake
- c.Energy density of breast milk remain remarkably stable except in extreme deprivation
- d.All are correct

35 Which of the following indicator directly informs the impact of family planning service?

- a.Contraceptive prevalence
- b.Maternal mortality ratio
- c.Total fertility rate
- d.Perinatal Mortality Rate
- e.All of the above

36 One of the following is not true about Adolescent sexual reproductive health

- a.it refers to the ability of young people to remain free from unwanted pregnancy and unsafe abortion
- b.it refers to the ability of young people to remain free from STDs/HIV/AIDS
- c.It focuses more on adolescents and youths' sexual violence and coercion.
- d.It refers to the physical and emotional well-being of adolescents

37 Adolescent development underlies prevention of health problems.

- a.False
- b.True

38 Family planning development and progress in Ethiopia was smoother unlike other countries since it was adopted from and supported by international non-governmental organizational.

- a.False
- b.True

39 Early and unprotected sexual intercourse are the major causes of adolescent Sexual and Reproductive Health problems.

- a.True
- b.False

40 One of the following service is not expected to be provided for newborn with in the first 24 months of birth.

- a.Initiation of exclusive breast feeding
- b.Advice on family planning
- c.Cord care
- D.Immunization

41 Which of the following statement is true?

- a.Inadequate intake is immediate cause of iron deficiency anemia
- b.Intervention using medicinal tablet is recommended where anemia prevalence is > 40
- c.Primary school children are at high risk of iron deficiency anemia
- d.All
- e.None

42 Which one of the following models of delivery service approaches is not considered as successful?

- a.Attended by health professional at basic emergency obstetric care facilities
- b.Attended by non-health professional at home
- c.Attended by health professional at health facility
- D.Attended by health professional at home

43 One of the following is not a characteristic feature of adolescence.

- a.rapid physiological changes and vulnerability to physical, psychological and environmental influences
- b.progresses from the point of initial appearance of 20 sexual characteristics to sexual maturity
- c.of transition from the state of total socio – economic dependency to relative independence
- d.all of the above
- e.None of the above

44 Immediate cause of malnutrition includes

- a. preferential feeding
- b. Infections causing loss of appetite
- c. policies that govern the social sectors
- d. Immunization and early treatment

45 All of the following obstetric care services are provided at health center level for complications in labour and delivery except

- a. Neonatal resuscitation
- b. Perform manual removal of placenta
- c. Administer parenteral oxytocic drugs
- d. Caesarean sections

46 Which one of the following is the severest form of IDD?

- a. Reduced IQ
- b. Adynamia
- c. Goiter
- d. cretinism
- e. None

47 Which one of the following is delay mothers to seek skilled birth attendant during labour and delivery?

- a. Socioeconomic and cultural factors
- b. Perceived accessibility of the services
- c. Perceived Quality of Care
- d. All of the above

48 Which of the following does not show a correct nutrition & development cycle?

- a. Direct losses in productivity from poor physical status
- b. Indirect losses from poor cognitive development
- c. Losses due to increased health care costs
- d. None

49 One of the following statements is not true

- a. The income–malnutrition relationship is far above expectation.
- b. When gross national product (GNP) in developing countries doubles changes in underweight rates are modest.
- c. Under nutrition and micronutrient malnutrition are direct indicators of poverty.
- d. None
- e. All

50 One of the following focused antenatal care approach is not changed for the 2016 WHO model.

- a. Number of antenatal care visit/contact
- b. Respectful, individualized, person-centred care at every contact
- c. Category of interventions and recommended services
- d. Weeks of pregnancy for each visit

51 Which of the following statement is false about changes in maternal weight gain?

- a. Fat is stored most rapidly during mid-pregnancy while the fetus is small
- b. Approximately 5% of the total weight gain occurs in the first 10–13 weeks of pregnancy
- c. The amount of weight gain required during pregnancy to achieve a favorable outcome is also influenced by pre-pregnancy bodyweight
- d. Weight gain of 0.9 kg per week is the target during the second and the third trimester is the target for women with normal pre pregnancy BMI.
- e. None

52 Dietary reference level which is used when there is no enough data about Estimated Average Requirements and established based on animal experiments is called

- a. Recommended dietary allowance
- b. Upper tolerable intake
- c. Lower tolerable intake
- d. Adequate intake
- e. None

53 Which one of the following is incorrectly matched about indicators and their interpretation?

- a. WFA < 60% = Sever wasting
- b. BMI < 16 Kg/m² = Sever chronic energy deficiency
- c. MUAC < 12.5 = Wasting
- d. All
- e. None

54 Which one is the most effective method of controlling Iodine deficiency disorder

- a. Fortification
- B. Supplementation
- c. Iodized salt
- d. All

55 Which of the following statement is true about maternal vitamin A supplementation?

- a. Vitamin A supplementation is given as a routine ANC service
- b. In areas where the prevalence of night blindness is > 5% 25,000 IU of vitamin A daily is recommended
- c. Vitamin A supplementation for mothers within 45 days of delivery is recommended
- d. B and C
- e. None

56 Which one of the following is the global target of maternal mortality ratio reduction for 2030?

- a. Reduce it annually by 27%
- b. Reduce it 70/100,000 live birth
- c. 75% for each country
- d. Reduce it by 50%

57 Which one of the following statement is true?

- a.TGR of >10% indicates endemic area
- B.The major source of iodine is animal origin
- c.School children are at increased risk of Iodine deficiency
- d.Consumption of contaminated water could be a risk factor for IDD
- e.All
- f.None

58 Which of the following statement is not true?

- a.Nutrition sensitive interventions are those intervention that addresses immediate cause of malnutrition
- b.Building an enabling environment help address basic causes of malnutrition
- c.Feeding behavior and stimulation is categorized under nutrition specific intervention
- d.Water and sanitation is among nutrition sensitive intervention
- e.All are correct

59 At what time should breastfeeding be initiated?

- a.Immediately after the breast starts to express milk
- b.Immediately after the child wants to take breast milk
- c.Within one hour after delivery
- d.Within 4-12 hours after delivery
- e.A and B

60 Young people don't make economic and political demands.

- a.True
- b.False

61 Out of the three nutritional assessments, which one is less comprehensive?

- a.Nutrition screening
- b.Nutrition surveillance
- c.Nutrition survey

62 Adolescence is a time of opportunities and risks.

- a.True
- b.False

63 Which one is not true about anthropometric nutritional assessment method?

- a.It is widely used in the assessment of protein-energy malnutrition
- b.It can identify specific nutrient deficiency
- c.It can be used to monitor periodic changes in growth and/or body composition
- D.The procedures can assist in the identification of mild to moderate malnutrition
- e.None of the above

64 One correctly described the contraceptive prevalence trends in Ethiopia according to EDHS evidences (from 2000 to 2016).

- a.The contraceptive prevalence rate increased significantly from 2005 to 2011 than from 2000 to 2005
- b.The contraceptive prevalence rate showed significant increase from 2000 to 2005 compared with 2005 to 2011
- c.The contraceptive prevalence rate rose dramatically from 2011 to 2016 than another period.
- d.The contraceptive prevalence rate rose dramatically from 2000 to 2005 than another period.

65 Which of the following statements is not true?

- a.Undernourished children are likely to become short adults
- b.Child's weight for age is best predictor of human capital
- c.Stunting is linked to cognitive development
- d.Stunting in the first two years is related with reduced economic productivity

66 Which of the following statement is not true

- a. Preterm babies have higher risk of mortality compared to infants who are small for gestational age.
- b. Term infants with low birth weight are 10 times more likely to die compared to infants with birth weight between 3000-3499 gram.
- c. Preterm is the commonest cause of low birth weight among developing country
- d. Stress and anxiety is one causes of low birth weight among developed country
- e. All are correct

67 Which one of the following statement is not correct?

- a. Vitamin A deficiency has a direct relationship with severity and complication of child hood infections.
- b. Provitamins are found from animal source
- c. Vitamin A deficiency is associated with poor growth and poor school performance
- d. All
- e. None

68 Case A The following are the dietary reference values of Vitamin A for non-pregnant mothers in reproductive age

- A. Upper tolerable intake = 3,000 µg/day of preformed vitamin A
- B. Recommended dietary allowance = 700 µg RAE/day of vitamin A
- C. Estimated Average Requirements = 500 µg RAE/day of vitamin A

Based on the above information, the Vitamin A needs for 50% of non-pregnant women population is meet by?

- a. A
- b. B
- c. C

69 Case A The following are the dietary reference values of Vitamin A for non pregnant mothers in reproductive age

- A. Upper tolerable intake = 3,000 µg/day of preformed vitamin A
- B. Recommended dietary allowance = 700 µg RAE/day of vitamin A
- C. Estimated Average Requirements = 500 µg RAE/day of vitamin A

Which one meet only the need for 2- 3% of the non-pregnant population but exceeds the need of 97% of non-pregnant population

- a.A
- b.C
- c.B

70 Intake that poses no risk of adverse health effect

- a.Safe intake
- b.Lower reference intake
- c.Estimated average requirement
- d.Upper level of safe intake

71 One of the following is not an immediate cause to adolescent health problems

- a.Poor access to education and health services;
- b.Unsafe and unsupportive environment
- c.Unemployment and limited work opportunities
- d.Inadequate information and skills

72 Which one of the following define late maternal death?

- a.Death of a woman from obstetric causes more than 42 days but less than one year after the termination of pregnancy
- b.Death occurred during pregnancy or within 42 days after termination of pregnancy
- c.It includes deaths from causes and complications related with pregnancy only occurred in the uterus
- d.Maternal death that does not occur by incidental and accidental causes

73 Which of the following is not an underlying cause of malnutrition?

- a.insufficient health services
- b.inadequate access to food
- c.inadequate care for children and women
- d.inadequate dietary intake

74 Which one of the following is a risk factor for Iron deficiency anemia?

- a.Infection
- b.Low intake of phytates
- c.Deworming
- d.Increased intake of poly phenols
- e.A and D
- f.None

75 Which one of the following maternal mortality indicator calculated from female adult female deaths due to maternal causes?

- a.Life time risk of maternal death
- b.Proportion of maternal death
- c.Maternal mortality rate
- d.Maternal mortality ration

76 The pre-ICPD period was characterized by EXCEPT

- a.targeted the demography
- b.attained fertility regulation through holistic approach
- c.designed population policies
- d.reduced the population growth by family planning
- e.none

77 One of the following sources of data is more reliable and valid for measurement of maternal mortality?

- a.Health service data
- b.Vital registration system data
- c.Hospital based studies data
- d.None of the above

78 Which one of the following statement is true?

- a.Children tend to breastfeed less often when complementary foods are introduced
- b.Optimal complementary feeding depends only on what is fed
- c.The use of cups to feed liquids is most likely to result in transmission of infection that use of bottles with teats
- d.The actual amount (weight or volume) of food required depends on the energy density of the food
- e.A and D
- f.B and D

79 Case A The following are the dietary reference values of Vitamin A for non-pregnant mothers in reproductive age

- A. Upper tolerable intake = 3,000 µg/day of preformed vitamin A
- B. Recommended dietary allowance = 700 µg RAE/day of vitamin A
- C. Estimated Average Requirements = 500 µg RAE/day of vitamin A

Based on the above information, which one would you use for the non-pregnant women who came to your OPD for nutritional counselling?

- a.A
- b.B
- c.C

80 If you want to apply an anthropometric measurement for children with age greater than 24 months, which one is not valid measure of the nutritional status

- a.Head circumference forage Z score
- B.Height for age Z score
- c.Weight for height Z score
- d.Weight for age Z score