

1. A tumor infiltrating the foramen ovale might result in which of the following:
 - A. paralysis of tensor palatini muscle
 - B. sensory loss to the skin of the forehead
 - C. sensory loss to the skin of tip of the nose
 - D. paralysis of masseter muscle
 - E. sensory loss to the hard palate
2. In surgical removal of malignant parotid tumor, the facial nerve may be cut resulting in paralysis of one of the following muscles:
 - A. tensor palatine
 - B. masseter
 - C. stylopharyngeus
 - D. orbicularis oris
 - E. anterior belly of digastric
3. Clinically, four muscles are important in evaluating the integrity of the facial nerve. One of these muscles that **raises the eyebrows in surprise or horror** is:
 - A. Buccinator muscle ... (this helps to whistle or suck)
 - B. Orbicularis Oculi muscle... (this helps to close eyelids)
 - C. Orbicularis oris... (this helps to smile)
 - D. Frontalis muscle
 - E. Procerus muscle
4. A patient comes to the Out Patient Department (OPD) with a left sternocleidomastoid muscle paralysis. Which one of the following symptoms might she also have?
 - A. Weakness in turning her head to the left
 - B. Weakness in turning her head to the right
 - C. Inability to protrude her tongue
 - D. Inability to elevate the shoulder
 - E. Inability to close the eyelid
5. The pterion is an important landmark to indicate position of:
 - A. Anterior branch of superficial temporal artery.
 - B. Posterior branch of superficial temporal artery

C. Anterior branch of middle meningeal artery.

D. Posterior branch of middle meningeal artery

6. A thrombosis that occurs in the cavernous sinus could result in each of the following EXCEPT

A. inability to close the eyelids

B. lack of adduction of eyeball

C. lack of abduction of eyeball

D. inability to constrict pupil

E. loss of accommodation

7. An ER patient is unable to speak after being hit in the neck with a pool cue (club) in a bar fight. Examination reveals paralysis of most the laryngeal muscles that normally maintain speech. What nerve was damaged?

A. Glossopharyngeal nerve

B. Hypoglossal nerve

C. Superior laryngeal nerve internal branch

D. Recurrent laryngeal nerve

E. C2- ventral ramus

8. The most frequent clinical complaint in facial nerve peripheral palsy is an inability to close the eye on the affected side. However, difficulty with eating is also frequently reported. Which of the following statements explains best the difficulty with eating?

A. Innervation to the lateral pterygoid muscles is lost resulting in an inability to lower the mandible.

B. Innervation to the buccinator muscle which aids in chewing is lost.

C. Innervation of the masseter muscle, which is a powerful closer of the mandible, is lost resulting in a weak bite.

D. Innervation of the maxillary teeth is lost.

E. Innervation of the mandibular teeth is lost.

9. A tumor that compresses the neural contents within the right jugular foramen might result in:

A. loss of pain from anterior two-thirds of the tongue

B. inability to protrude tongue

C. sensory loss from skin overlying the zygomatic bone

D. inability to elevate the shoulder

E. inability to close the eye lid

10. At your first surgical experience of tonsillectomy, you encounter profuse bleeding from the tonsillar bed. Which of the following vessels may be responsible for post-tonsillectomy hemorrhage?

- A. Inferior thyroid artery
- B. Tonsilar branch of facial artery**
- C. Tonsilar branch of maxillary artery
- D. Lingual artery
- E. Tonsilar branch of superior thyroid artery

11. A lesion of the lingual nerve immediately after it receives the chorda tympani nerve could result in each of the following, EXCEPT

- A. loss of sublingual gland secretion
- B. sensory loss from lower teeth**
- C. loss of taste from anterior two-thirds of the tongue
- D. loss of submandibular gland secretion
- E. none of the above

12. Which one of the following is caused by **failure of the maxillary prominence** to merge with its **corresponding lateral prominence**?

- A. Isolateral cleft palate
- B. Bilateral cleft lip
- C. Oblique facial cleft**
- D. Cleft upper jaw
- E. None of the above

13. Innervation to the rotator cuff muscle that laterally rotates the arm is provided by which of the following?

- A. Suprascapular nerve**
- B. Median nerve
- C. Ulnar nerve
- D. Upper and Lower subscapular nerve
- E. Dorsal scapular nerve

14. Which nerve is probably damaged if a person CAN NOT abduct beyond 15 degrees?

- A. Axillary nerve**

- B. Radial nerve
- C. Musculocutaneous nerve
- D. Median nerve
- E. Ulnar nerve

15. Loss of supination of a flexed forearm suggests a lesion of the:

- A. Radial nerve
- B. Musculocutaneous nerve**
- C. Ulnar nerve
- D. Median nerve
- E. Axillary nerve

16. Injury to one of the following nerves causes paralysis of primary hip abductors.

- A. Sciatic
- B. Superior gluteal
- C. Inferior gluteal**
- D. Inferior cuneal

17. Fracture to the head of fibula results the following except

- A. Loss of cutaneous innervations between 1st and 2nd toes
- B. Loss of eversion of the foot
- C. Loss of cutaneous innervations on the plantar surfaces of foot**
- D. Foot drop

18. With regard to the type of collagens present in the matrix of the different types of cartilage:

- A. Fibrous cartilage has type I, hyaline cartilage has type II, and elastic cartilage has type III.
- B. Both fibrous and hyaline cartilages have type I, while elastic cartilage has type II collagen
- C. Fibrous cartilage has type I while both hyaline and elastic cartilages have type II collagen**
- D. Fibrous cartilage has type II, hyaline cartilage has type I and elastic cartilage has type III
- E. Both the fibrous and hyaline cartilages have type II, while elastic cartilage has type I collagen.

19. Endochondral bone ossification includes all of the following except:

- A. A cartilage model
- B. Proliferation and enlargement of cartilage cells

C. CHONDROCYTES CHANGING TO OSTEOCYTES

D. Calcification of cartilage matrix

E. Interruption of blood supply to the area of ossification

20. Which type of bone cells are involved in bone ossification?

A. OSTEOCLASTS

B. Osteoblasts

C. Osteocytes

D. a & b

e. All

21. A newly deposited uncalcified bone matrix is known as:

A. Osteon

B. Perosteoid

C. OSTEOID

D. Haversian system

E. Cancellous bone

22. Which one of the following is **not true** regarding the cardiac muscles?

A. Divides and anastomose but the cytoplasm remains separated.

B. Intercalated discs correspond to the zone of the Z lines

C. HAVE PERIPHERALLY LOCATED NUCLEI.

D. Sarcoplasmic reticulum makes diad with T-tubule but not triad

E. Cannot regenerate

23. Region of cross-linking between myosin molecules in a sarcomere is represented by:

A. A-BAND

B. I band

C. M line

D. H band

E. Z line

24. The body of **first lumbar vertebrae** is formed by:

A. The 12th thoracic sclerotome alone

B. The cranial half of the 1st thoracic sclerotome alone

C. FUSION OF CAUDAL HALF OF THE 12TH THORACIC WITH THE CRANIAL HALF OF THE 1ST LUMBAR SCLEROTOMES

D. Fusion of caudal half of the 1st lumbar with the caudal half of the 2nd lumbar sclerotomes

E. The 1st lumbar sclerotome alone

25. The ribs develop from:

A. somatic layer of lateral mesoderm

B. Visceral layer of lateral mesoderm

C. PARAXIAL MESODERM

D. Intermediate mesoderm

E. Neural crest

27. Which one of the following is true regarding **hypomeric** and **epimeric** divisions of the myotomes?

- A. Epimers are larger and located dorsally as compared to the hypomers
- B. EPIMERS ARE SMALLER AND LOCATED DORSALLY AS COMPARED TO THE HYPOMERS.**
- C. Epimers are innervated by the dorsal branches of the ventral primary rami of spinal nerves
- D. Hypomers give rise to the extensors of the back
- E. None of the above.

28. The **ventral mesodermal** condensations of the limb bud give rise to which type of muscles?

- A. Extensors and pronators
- B. Extensors and abductors
- C. Extensors and adductors.
- D. FLEXORS AND PRONATORS.**
- E. Flexors and abductors.

29. Keratinocytes show mitotic activity in which layer of epidermis?

- A. Stratum lucidum and stratum spinosum
- B. Stratum basale and stratum granulosum
- C. Stratum basale, stratum granulosum and stratum basale
- D. Stratum basale and stratum spinosum
- E. All layers, except stratum corneum**

30. An eosinophilic layer of dead cells that contain densely packed filaments embedded in an electron dense matrix and may be absent in thin skin is:

- A. Stratum lucidum**
- B. Stratum spinosum
- C. Stratum corneum
- D. Stratum basale
- E. Stratum granulosum

31. Which one of the following is not true about melanocytes?

- A. Are less in number in white skinned than dark skinned races.**
- B. Produce melanin pigments which are dark brown pigments
- C. Have capacity of mitotic activity

D. Appear as cells with elongated nuclei surrounded by a clear cytoplasm under light microscopy following H & E staining.

E. Have long irregular and branched processes.

32. The most inner layer of the inner epithelial root sheath of hair follicle that anchors the hair in its root is:

- A. Cortex B. Huxley's layer C. Cuticle D. Henle's layer E. Medulla

33. Outer coat of hair follicle is:

A. Made by connective tissue continuous with the dermis

B. Equivalent to stratum corneum

C. Equivalent to the most inner layer of epidermis

D. Absent from fine hairs

E. None of the above

34. Body of the nail or hair plate is equivalent to:

A. Epidermis, dermis, and subcutaneous tissue of the skin

B. Epidermis and dermis of the skin

C. All layers of epidermis with no dermis

D. Stratum corneum of epidermis

E. Stratum germinativum of epidermis

35. Which one of the following is derived from the second pharyngeal arch?

A. The sphenomandibular ligament

B. The stapes of auditory ossicles

C. The incus of auditory ossicles

D. The zygomatic bone

E. The malleus of auditory ossicles

36. Which one of the following is not derived from the mandibular pharyngeal arch?

A. Mylohyoid

B. Tensor palatine

C. muscles of mastication

D. Tensor tympani

E. None of the above

37. Which one of the following doesn't arise from pharyngeal pouches?

A. Stroma of palatine tonsil

B. External auditory canal

C. Middle ear cavity

D. Parathyroid gland and thymus

E. Eustachian tube

38. Which prominences form the cheek and lateral portions of upper limb

A. Maxillary

B. Mandibular

C. Lateral nasal

D. Median nose

E. All of the above

39. Which prominence forms the bridge of the nose?

A. Lateral nasal

B. Medial nasal

C. Frontonasal

D. Maxillary

E. Mandibular

40. Structure that is not developed in the third brachial arch?

A. Lesser cornu of hyoid bone

B. Glossopharyngeal nerve

C. Common carotid artery

D. Stylopharyngeal artery

E. None of the above

41. Structures derived from endoderm of the 4th pharyngeal pouch include:

A. Thymus gland

B. Palatine tonsil

C. Thyroid gland

D. Para follicular cells of thyroids

E. None of the above