

Name _____

ID No _____

1. Regarding chest Inspection, One of the following is not included
- a. Symmetry of the chest
 - b. Chest movement
 - ☒ c. Diaphragmatic Excursion
 - d. Retractions

2. Auscultation of friction rub over an enlarged spleen suggests
- a. Splenic abscess
 - b. Splenic tumor
 - ☒ c. Splenic infarction
 - d. Splenic granuloma

3. Elicitation of a fluid thrill in a 50 year old female patient with grossly distended abdomen without flank fullness and shifting dullness suggests
- a. Ascites
 - b. Haemo-peritoneum
 - ☒ c. Ovarian cyst
 - d. Pregnancy

4. 1990 (EC). Menilik II Hospital, Addis Ababa. Bleeding duodenal ulcer. Discharged symptom free after transfusion of 2 units of blood. The above description is an example of
- a. Chief Complaint
 - b. Past Medical History
 - ☒ c. Previous Admission
 - d. None of the above

5. Causes of pleuritic chest pain include
- a. Ischemic heart disease
 - ☒ b. TB pleurisy of tuberculosis
 - ☒ c. Pneumothrax
 - d. All

6. Which of the following is least commonly used technique of examination in cardiovascular system?
- ☒ a. Percussion
 - b. Inspection
 - c. Palpation
 - d. Auscultation
 - e. None

Name _____

ID No _____

7. False negative findings in breast palpation may be due to
- a. Small mass in a large breast
 - b. Inherent properties of breast tissue
 - c. Poor technique
 - d. Failure to listen to the patient
 - e. All of the above
8. In the evaluation of breasts which of the following is the appropriate first stage
- a. Inspection of the breasts
 - b. Palpation of the breasts
 - c. Exposure of the breasts
 - d. Positioning of the patient
 - e. Talking to the patient
9. Which of the following is most informative in abdominal examination?
- a. Inspection
 - b. Auscultation
 - c. Palpation
 - d. Percussion
10. Tenderness upon superficial palpation of the abdomen is most likely due to:
- a. Inflammatory conditions of the peritoneum
 - b. Intra abdominal malignancy
 - c. Peritoneal carcinomatosis
 - d. Ascitic collection in the peritoneal cavity
11. True about jaundice includes:
- a. Sclera has high affinity to bilirubin because of its location and appearance.
 - b. Presence of bilirubin in urine is a sign of direct hyperbilirubinemia
 - c. Gilbert's syndrome is a congenital defect of enzyme leading to conjugated hyperbilirubinemia
 - d. Most common cause of icteric sclera is Carotenoderma
12. Select the inappropriate interviewing technique
- a. Empathy
 - b. Sympathy
 - c. Confrontation
 - d. Free speech
 - e. Summarization

13. Loss of what percentage of weight is considered significant

a. 3%

b. 5%

☒ c. 10% *in 3 months (our setup)*

d. 15%

14. What are the properties that we should consider when analyzing pain

a. Main site

b. Radiation

c. Quality

d. Severity

☒ e. All of the above

15. All of the following are accessible lymphnodes except

a. Axillary lymph node

b. Cervical lymphnode

☒ c. Hilar lymph node

d. Epitrochlear lymph node

e. Femoral lymph node

16. Which one of the following statements is true about Fever

a. Fever is a manifestation of infections only ☒

☒ b. Fever is defined as an elevated oral temperature $\geq 37.8^{\circ}\text{C}$ ☒

c. Fever is the same as hyperthermia ☒

d. Axillary temperature is the best estimate of core body temperature ☒

17. Which of the following should be included in the HPI of a patient with fever

a. Travel to malaria endemic area ☒

b. Chest pain ☒

c. Headache ☒

☒ d. All of the above


18. Concerning facial nerve examination

☒ a. Nuclear or infranuclear lesions are of the same degree in upper and lower parts of the face ☒

b. Differential facial weakness is suggestive of infranuclear weakness ☒

☒ c. Lower face weakness preserving the upper face is typical of facial nerve lesion at the parotid gland ☒

d. If a crossed lesion i.e. left facial palsy and right hemiparesis in a patient are found, the localization of the lesion is at the cortex of the brain ☒



1. Supranuclear ☒ Upper face spared
2. Infranuclear (nucleus) ☒ The whole face affected

19. The following are meningeal signs except
- a. Nuchal rigidity
 - b. Kerning's sign
 - ☒ c. Babinski's sign
 - d. Brudzinski's sign
20. The classic pupillary abnormality of neurosyphilis is
- a. Miotic pupil
 - ☒ b. Argyll Robertson pupil
 - c. Adie's tonic pupil
 - d. Unilateral dilated pupil
21. Which of the following indicates a renal parenchymal cause for HEMATURIA?
- a. Presence of clots in the urine
 - ☒ b. Flank pain accompanying hematuria
 - ☒ c. Continuous hematuria
 - d. Dysuria
22. Which one of the following clinical symptoms/signs may be an early indicator of the presence of Chronic Kidney Disease (CKD)?
- ☒ a. Oliguria
 - b. Nocturia
 - c. Anemia
 - d. Bone pain
23. Which one of the following physical signs indicates that heart failure is the most likely cause for the generalized edema a patient has presented with?
- a. Huge ascites
 - ☒ b. +4 pretibial edema
 - c. Periorbital puffiness
 - d. Hepatomegaly
24. The elevation of the plasma levels one of the following hormones serves as an indicator of the severity of congestive heart failure.
- a. Angiotensin II
 - ☒ b. Arginine vasopressin (AVP)
 - c. Endothelin
 - ☒ d. Brain natriuretic peptide (BNP)