

Anatomy – upper limb mcq

Regarding pronator teres which of the following statements is correct?

- a) It forms the lateral border of the cubital fossa
- b) It arises from the coronoid process and lateral epicondyle.
- c) The Median nerve passes deep to both heads
- d) It's medial border forms the medial boundary of the cubital fossa
- e) It is the most lateral of the superficial flexors of the forearm

Regarding the radial nerve which statement is incorrect?

- a) It passes anterior to the lateral epicondyle of the humerus
- b) Injury to the radial nerve from fracture of the shaft of the humerus will result in wrist drop
- c) Injury to the deep radial nerve in the mid forearm will prevent extension only at the MCPJs
- d) Sensory loss from injury to the superficial radial nerve will usually result in loss of sensation over the entire thumb
- e) It is the larger terminal branch of the posterior cord of the brachial plexus

Regarding the blood supply of the forearm

- a) The radial artery is the larger of the terminal branches of the brachial artery
- b) The radial artery runs under brachialis as it leaves the cubital fossa
- c) The radial artery has just one named branch proximal to the carpal braches
- d) Ligation of the radial artery at its origin will significantly reduce blood flow through the posterior interosseous artery.
- e) The radial artery has no involvement in the elbow anastomotic network

The median nerve

- a) Supplies flexor carpi ulnaris and half of flexor digitorum profundis
- b) Gives rise to most of its braches in the upper arm
- c) Gives rise to the common interosseous nerve which divides anterior to the radial head
- d) May be compressed between the two heads of pronator teres
- e) Enters cubital fossa lateral to the brachial artery

- 1) E
- 2) D
- 3) C
- 4) D