

Question 1

Which one of the following is not concerning the magnitude of maternal mortality?

- a. It is dropping from the previous estimate to now
- b. Needs large sample size to estimate its burden
- c. Maternal mortality is a worldwide
- d. Its burden is the same for all African regions

Question 2

In a nutrition survey, a male child of 84cm weighs 9.3kg. Note: the reference median weight for boys of height 84cm is 11.7kg and the standard deviation is 0.908. What is the Z-score?

- a. 2.6
- b. 1.8
- c. - 2.6
- d. -1.8

Question 3

Which of the following statement is true about maternal vitamin A supplementation?

- a. Vitamin A supplementation is given as a routine ANC service
- b. In areas where the prevalence of night blindness is < 5% 25,000 IU of vitamin A daily is recommended
- c. Vitamin A supplementation for mothers within 45 days of delivery is recommended
- d. B & C
- e. None

Question 4

One of the following is not true about MDSR (Maternal Death Surveillance and Response)

- a. Maternal deaths in communities notified to the appropriate authorities within 48 hours
- b. Triangulating data sources is important to avoid duplication in reporting
- c. It involves a continuous surveillance and action cycle
- d. Notification of maternal death does not include zero death

Question 5

The waist to hip ratio cut off point which indicates an increased risk for non-communicable diseases in male is:

- a. > 0.85
- b. > 0.87
- c. > 2
- d. > 1

Question 6

The risk of unwanted pregnancy and unplanned parenthood in adolescents is high.

- a. True
- b. False

Question 7

Which one of the following is not correct about WHO classification of public health problem of xerophthalmia?

- a. Bitot's spot of > 0.5%
- b. Corneal scar of > 0.05%
- c. Corneal xerosis of > 0.01%
- d. Night blindness of > 5%
- e. None

Question 8

Family planning is the same with family planning.

- a. False
- b. True

Question 9

All of the following obstetric cares are given only for complications in labor and delivery except

- a. Obstetric First Aid
- b. Essential Obstetric Care
- c. Emergency Obstetric Care
- d. None of the above

Question 10

Which of the following indicator directly informs the impact of family planning service?

- a. Contraceptive prevalence
- b. Maternal mortality ratio
- c. Total fertility rate
- d. Perinatal Mortality Rate
- e. All of the above

Question 11

Which of the following statements is not true?

- a. Stunting is linked to cognitive development
- b. Child's weight for age is best predictor of human capital
- c. Undernourished children are likely to become short adults
- d. Stunting in the first two years is related with reduced economic productivity

Question 12

Which one of the following is the major cause of maternal mortality at global and national level?

- a. Sepsis and
- b. Hypertensive disorder of pregnancy and obstructed labor
- c. Abortion
- d. Hemorrhage

Question 13

A 19-year-old pregnant girl is considered as an adolescent pregnancy.

- a. True
- b. False

Question 14

The pre-ICPD period was characterized by EXCEPT

- a. targeted the demography
- b. attained fertility regulation through holistic approach
- c. designed population policies
- d. reduced the population growth by family planning
- e. none

Question 15

Which one of the following is not the principle of nutrient reference level?

- a. Refer to the average daily need over a reasonable period of time
- b. Refer to level of intake needed to maintain health in already healthy individuals
- c. It takes in to account possible interaction that involve nutrients and other dietary components
- d. Based on typical dietary pattern of a country
- e. None

Question 16

In Ethiopia, the age of first sexual intercourse for girls and boys is nearly equal.

- a. False
- b. True

Question 17

Which of the following statement is true?

- a. Inadequate intake is immediate cause of iron deficiency anemia
- b. Intervention using medicinal tablet is recommended where anemia prevalence is ≥ 40
- c. Primary school children are at high risk of iron deficiency anemia
- d. All
- e. None

Question 18

Which of the following does not show a correct nutrition & development cycle?

- a. Direct losses in productivity from poor physical status
- b. Indirect losses from poor cognitive development
- c. Losses due to increased health care costs
- d. None

Question 19

Which of the following causes of maternal mortality takes short time for death of a mother?

- a. Sepsis
- b. Preeclampsia
- c. Hemorrhage
- d. Obstructed labor

Question 20

Short answer

Write down short answers for the following questions. (6 Points)

1. Indicate the age limit in years for the following
 - a. Child: 0 to 18 years
 - b. Early Adolescence: 10 to 14 years
 - c. Late Adolescence: 15 to 19 years
 - d. Adolescent: 10 to 19 years
 - e. Youth: 15 to 24 years
 - f. Young people: 10 to 24 years

Question 21

In a nutritional survey, a female child of 90cm weighs 8kg. Note: The median weight for a female child of height 90cm is 13.7kg. Calculate percent of median?

- a. 67%
- b. 58.4%
- c. 80%
- d. 52%

Question 22

One of the following statements is not true

- a. The income–malnutrition relationship is far above expectation.

- b. When gross national product (GNP) in developing countries doubles changes in underweight rates are modest.
- c. Under nutrition and micronutrient malnutrition are direct indicators of poverty.
- d. None
- e. All

Question 23

Which one measures the volume of program activity among the family planning indicators

- a. Method mixes
- b. Contraceptive prevalence rate
- c. Couple years of protection
- d. Total fertility rate
- e. Ever use of contraceptive

Question 24

One of the following is not true about marasmus

- a. No Fatty liver
- b. Common among children less than 2 years of age
- c. Old man face
- d. Weight for age between 60-80%

Question 25

In Ethiopia, based on recent DHS studies, nearly half of adult women has initiated sexual intercourse during their early adolescence age.

- a. False
- b. True

Question 26

One of the following interventions is recommended for all causes of maternal health complications.

- a. Treatment of anemia
- b. Having skilled birth attendant
- c. Replacement of fluids
- d. Urgent referral

Question 27

Which one of the following is correctly matched?

- a. Keratomalacia- ulceration and perforation of the cornea
- b. corneal ulceration- reversible structural change of the cornea
- c. corneal xerosis- corneal surface becomes dry
- d. Bitot's spot- the earliest structural change
- e. A& B

f. A& C

Question 28

Which of the following statements is not true?



Good nutrition is not intuitive.

- b. People cannot tell when their children are becoming malnourished because it is not easy to detect for an untrained eye.
- c. Improving nutrition is pro-poor and increases the income-earning potential of the poor.
- d. All
- e. None

Question 29

Which one is not true about anthropometric nutritional assessment method?

- a. It is widely used in the assessment of protein-energy malnutrition
- b. It can identify specific nutrient deficiency
- c. It can be used to monitor periodic changes in growth and/or body composition
- d. The procedures can assist in the identification of mild to moderate malnutrition
- e. None of the above

Question 30

Which one of the following is not included in the definition of maternal mortality?

- a. Cause of maternal death related to or aggravated by pregnancy or its management
- b. Death occurred during pregnancy or within 42 days after termination of pregnancy
- c. Maternal death occurred by incidental and accidental causes
- d. All sites and duration of pregnancy considered equally

Question 31

Which of the following statement is correct?

- a. Zinc deficiency could be a cause of night blindness
- b. Zinc supplementation can reduce child morbidity and mortality
- c. A rate of stunting $\geq 20\%$ is indicative of zinc deficiency
- d. In areas where high Phytate diet is a staple zinc deficiency might be common
- e. All
- f. None

Question 32

Which one of the following is the severest form of IDD?

- a. Reduced IQ
- b. Adynamia
- c. Goiter
- d. cretinism

- e. None

Question 33

One correctly described the contraceptive prevalence trends in Ethiopia according to EDHS evidences (from 2000 to 2016).

- a. The contraceptive prevalence rate showed significant increase from 2000 to 2005 compared with 2005 to 2011
- b. The contraceptive prevalence rate rose dramatically from 2000 to 2005 than another period.
- c. The contraceptive prevalence rate increased significantly from 2005 to 2011 than from 2000 to 2005
- d. The contraceptive prevalence rate rose dramatically from 2011 to 2016 than another period.

Question 34

Which of the following is not an underlying cause of malnutrition?

- a. inadequate access to food
- b. insufficient health services
- c. inadequate dietary intake
- d. inadequate care for children and women

Question 35

All are true about Anthropometric Indices except

- a. Weight for height identify children suffering from current or acute under-nutrition
- b. Weight for age identifies both chronic and/or acute under nutrition
- c. Height for age identifies chronic malnutrition
- d. All
- e. None of the above

Question 41

Which one of the following antenatal care services expected to be provided for the pregnant women at third visit?

- a. Calculate LNMP and EDD
- b. Advice on family planning and breastfeeding
- c. Perform physical examination
- d. Assess fetal well being

Question 42

Which of the following shows a recent acute malnutrition?

- a. Height for age < -2SD
- b. Weight for height < -2SD
- c. Weight for age < -2SD
- d. Weight for age < -3SD

Question 43

Which one of the following measures is not included in the preventive measure for Iodine deficiency disorder?

- a. Universal salt iodization
- b. Water iodization
- c. Deworming
- d. Health education
- e. None

Question 44

In Ethiopia, based on recent DHS studies, nearly half of adult men has initiated sexual intercourse during their late adolescence age.

- a. False
- b. True

Question 45

Which one of the following is a risk factor for Iron deficiency anemia?

- a. Infection
- b. Low intake of phytates
- c. Deworming
- d. Increased intake of poly phenols
- e. A & D
- f. None

Question 46

Dietary reference level which is used when there is no enough data about Estimated Average Requirements and established based on animal experiments is called

- a. Recommended dietary allowance
- b. Upper tolerable intake
- c. Lower tolerable intake
- d. Adequate intake
- e. None

Question 47

Growth of external genitalia is considered as sign of secondary sexual characteristics in boys, but not in girls.

- a. False
- b. True

Question 48

One of the following is a fundamental shift and a new approach in the reproductive health progress milestones.

- a. Adapting and expanding population policies
- b. Removing the barrier to family planning services to the level possible
- c. Considering the reproductive health care as a right of individuals or couples
- d. Control of female fertility
- e. All of the above

Question 49

Immediate cause of malnutrition includes

- a. Immunization and early treatment
- b. policies that govern the social sectors
- c. Infections causing loss of appetite
- d. preferential feeding

Question 50

Which one of the following is WRONG about the objective of reproductive health care?

- a. providing information about reproductive health services
- b. to enable and support decision making in family planning and childbearing
- c. to meet the reproductive health need of people based on the available resources
- d. none

Question 51

What was the maternal mortality ratio of Ethiopia according to EDHS 2016?

- a. 676 maternal deaths/100,000LBs
- b. 871 maternal deaths/100,000LBs
- c. 412 maternal deaths/ 100,000LBs
- d. 573 maternal deaths/100,000LBs
- e. None of the above

Question 52

Family planning development and progress in Ethiopia was smoother unlike other countries since it was adopted from and supported by international non-governmental organizational.

- a. False
- b. True

Question 53

Which one of the following statements is true?

- a. Basal metabolism rate represents 45-70% of daily total energy expenditure

- b. Basal metabolism is the most variable and account for the largest component of daily energy expenditure
- c. Dietary reference intakes provide qualitative advice to the public about diet and chronic disease prevention
- d. None

Question 54

According to 2016 Ethiopian demographic and health survey (EDHS), the modern contraceptive prevalence rate is

- a. 29
- b. 42
- c. 36
- d. 27
- e. 35

Question 55

Which one of the following statements is true?

- a. TGR of > 10% indicates endemic area
- b. The major source of iodine is animal origin
- c. School children are at increased risk of Iodine deficiency
- d. Consumption of contaminated water could be a risk factor for IDD
- e. All
- f. None

Question 56

Which one of the following is component of comprehensive emergency obstetric care services?

- a. Administer parenteral oxytocic drugs
- b. Perform Caesarean sections
- c. Perform manual removal of placenta
- d. Perform assisted vaginal delivery

Question 57

One of the following maternal health service utilization is very low in Ethiopia

- a. Delivery conducted by skilled birth attendant
- b. Postnatal care
- c. Family planning
- d. Antenatal care

Question 58

Which one of the following is not the strategy of Safe Motherhood Initiative?



Use of family planning to avoid unwanted pregnancy

- b. Considering every pregnancy is at risk
- c. Promote use of traditional birth attendants for high-risk pregnant women
- d. Recommend risk assessment during pregnancy by trained non-physicians

Question 59

Which of the following statement is false about changes in maternal weight gain?

- a. Fat is stored most rapidly during mid-pregnancy while the fetus is small
- b. Approximately 5% of the total weight gain occurs in the first 10–13 weeks of pregnancy
- c. The amount of weight gain required during pregnancy to achieve a favorable outcome is also influenced by pre-pregnancy bodyweight
- d. Weight gain of 0.5 kg per week is the target during the second and the third trimester is the target for women with normal pre pregnancy BMI.
- e. None

Question 60

Choose the most common family planning delivery strategies in Ethiopia.

- a. Community based distribution
- b. Commercial retail sale
- c. Clinic based services
- d. Outreach based services
- e. All of the above strategies are equally applicable

Question 61

Which one of the following statements is not correct?



- a. Vitamin A deficiency has a direct relationship with severity and complication of child hood infections.
- b. Provitamins are found from animal source
- c. Vitamin A deficiency is associated with poor growth and poor school performance
- d. All
- e. None

Question 62

Intake that poses no risk of adverse health effect

- a. Lower reference intake
- b. Upper level of safe intake
- c. Safe intake
- d. Estimated average requirement

Question 63

The prevalence of stunting at birth, in the first six month of life and in the age between 6-24 months respectively is

- a. 10%, 10%, 50%
- b. 20%, 20%, 50%
- c. 10%, 20%, 50%
- d. 50%, 20%, 20%

Question 64

One of the following is not the reason why safe motherhood issue to be concerned

- a. Majority of causes of maternal death are not prevented
- b. Benefit of maternal survival for child health
- c. Social and economic contribution of women in addition to reproduction
- d. Magnitude of the problem

Question 65

One of the following is delaying the woman to decide to seek care for maternal health services?

- a. Perceived severity
- b. Staff attitudes
- c. Unavailability of drugs
- d. All of the above