

Health promotion and Disease Prevention

Question 1

Answer saved

Marked out of 1.0

A woman who uses infant formula milk occasionally for her baby can use lactational amenorrhea as contraceptive method.

Select one:

- ☐ a. FALSE
- ☐ b. TRUE

Question 2

Answer saved

Marked out of 1.0

Which one of the following is incorrectly matched about indicators and their interpretation?

Select one:

- ☐ a. WFA < 60% = Sever malnutrition
- ☐ b. BMI < 16 Kg/m² = Sever chronic energy deficiency
- ☐ c. MUAC < 12.5 = Wasting
- ☐ d. All
- ☐ e. None

Question 3

Answer saved

Marked out of 1.0

Maternal death, as considered in maternal mortality ratio, don't count/include such deaths:

Select one:

- ☐ a. A high school teenager girl died of abortion complication,
- ☐ b. A pregnant woman who died of eclampsia,
- ☐ c. A pregnant woman who died of severe malaria and severe anemia,
- ☐ d. A postpartum woman who died of sepsis after 1 month of delivery
- ☐ e. None of the above

Question 4

Answer saved

Marked out of
1.0

One of the following was not the characteristic of the pre Cairo international conferences of population and development:

Select one:

- ☐ a. Setting demographic targets,
- ☐ b. Designing population policies,
- ☐ c. Applying FP to restrain population growth,
- ☐ d. Emphasizing the rights & health of women,
- ☐ e. None of the above

Question 5

Answer saved

Marked out of
1.0

Sexual health and reproductive health have same meanings.

Select one:

- ☐ a. TRUE
- ☐ b. FALSE

Question 6Not yet
answeredMarked out of
1.0

The waist to hip ratio cut off point which indicates an increased risk for non-communicable diseases in male is:

Select one:

- ☐ a. > 0.85
- ☐ b. > 0.87
- ☐ c. > 2
- ☐ d. > 1

Question 7

Answer saved

Marked out of
1.0

If you want to apply an anthropometric measurement for children with age greater than 24 months, which one is not valid measure of the nutritional status

Select one:

- ☐ a. Weight for age Z score
- ☐ b. Head circumference for age Z score
- ☐ c. Weight for height Z score
- ☐ d. Height for age Z score

Question 8Not yet
answeredMarked out of
1.0

Below are list of disease conditions and complications causing neonatal and child deaths

- I. Pneumonia
- II. Diarrhea
- III. Sepsis
- IV. Birth asphyxia
- V. Congenital anomalies
- VI. Prematurity
- VII. Injury

Which of the above three conditions are the leading cause of neonatal deaths?

Select one:

- ☐ a. I, III, & VI
- ☐ b. I, IV, & VI
- ☐ c. II, III. & IV
- ☐ d. III, IV, V
- ☐ e. III, V, VI

Question 9

Answer saved

Marked out of
1.0

Which one of the following is correctly matched?

Select one:

- ☐ a. Keratomalacia- ulceration and perforation of the cornea
- ☐ b. corneal ulceration- reversible structural change of the cornea
- ☐ c. corneal xerosis- corneal surface becomes dry
- ☐ d. Bitot's spot- the earliest structural change
- ☐ e. A and B
- ☐ f. A and C

Question 10

Answer saved

Marked out of
1.0

According to the recent EDHS survey, in Ethiopia _____ & _____ are the two commonly used modern contraceptive methods:

Select one:

- ☐ a. Oral pills & Injectables
- ☐ b. Injectables & IUDs
- ☐ c. Oral pills & condoms
- ☐ d. Estrogen & Progestrone based oral pills
- ☐ e. Injectables & Implants

Return to: 0 

Health promotion and Disease Prevention

Question 11

Answer saved

Marked out of 1.0

Which one of the following measure is not included in the preventive measure for IDD?

Select one:

- ☐ a. Universal salt iodization
- ☐ b. Water iodization
- ☐ c. Deworming
- ☐ d. Health education
- ☐ e. None

Question 12

Answer saved

Marked out of 1.0

The gold standard source of data for estimating MM worldwide is

Select one:

- ☐ a. Census
- ☐ b. VA
- ☐ c. Survey
- ☐ d. VR

Question 13

Answer saved

Marked out of 1.0

The ultimate aim of skilled delivery care is to

Select one:

- ☐ a. Provide Clean care
- ☐ b. Provide Safe care
- ☐ c. Detect and treat common delivery related complications
- ☐ d. All

Question 14

Answer saved

Marked out of
1.0

Among 10-14 years old adolescent females which of the following is the leading cause of death?

Select one:

- ☐ a. Pregnancy related complications
- ☐ b. Diarrheal disease
- ☐ c. Road traffic injury
- ☐ d. Drowning
- ☐ e. Lower respiratory infection

Question 15

Answer saved

Marked out of
1.0

One of the following factors has contributed to the launch of the Safe Motherhood Initiative (SMI) in 1987?

Select one:

- ☐ a. MM was equally the problem of public health importance for all nations
- ☐ b. MM was unacceptably massive in 1980s
- ☐ c. MM was used to be hard to measure
- ☐ d. MM was the most talking points in pre-1980s

Question 16

Answer saved

Marked out of
1.0

The Cairo ICPD recommendations were formulated for countries for ----- years.

Select one:

- ☐ a. Ten
- ☐ b. Fifteen
- ☐ c. Twenty
- ☐ d. Twenty five
- ☐ e. None

Question 17

Answer saved

Marked out of
5.0

Many women in low-income countries do not seek antenatal care until the second trimester, in this situation what is recommendation for these women in order to reach the total energy cost of pregnancy? **(5points)**

Question 18Not yet
answeredMarked out of
1.0

The following issues are collectively termed as process indicators:

Select one:

- ☐ a. LFR, SBA, PNC and PM
- ☐ b. ANC, SBA, and PNC
- ☐ c. ANC, PNC and MMRatio
- ☐ d. MMRate, LFR and PM

Question 19Not yet
answeredMarked out of
1.0

Which one of the following statement is not correct?

Select one:

- ☐ a. Vitamin A deficiency has a direct relationship with severity and complication of child hood infections.
- ☐ b. Provitamins are found from animal source
- ☐ c. Vitamin A deficiency is associated with poor growth and poor school performance
- ☐ d. All
- ☐ e. None

Question 20

Not yet
answered

Marked out of
1.0

The following reason does not cause us to draw huge attention to women, which one?

Select one:

- ☐ a. Their huge number
- ☐ b. Being susceptibility to wide range of problems
- ☐ c. They being voiceless
- ☐ d. None

Return to: 0 

Health promotion and Disease Prevention

Question 21

Not yet
answered

Marked out of
1.0

Which of the following statement is true?

Select one:

- ☐ a. Inadequate intake is immediate cause of iron deficiency anemia
- ☐ b. Intervention using medicinal tablet is recommended where anemia prevalence is > 40
- ☐ c. Primary school children are at high risk of iron deficiency anemia
- ☐ d. All
- ☐ e. None

Question 22

Not yet
answered

Marked out of
1.0

Which one of the following is the severest form of IDD?

Select one:

- ☐ a. Reduced IQ
- ☐ b. Adynamia
- ☐ c. Goiter
- ☐ d. cretinism
- ☐ e. None

Question 23

Not yet
answered

Marked out of
1.0

According to the recent national survey (EDHS), one of the following is correct about Ethiopia's maternal mortality:

Select one:

- ☐ a. Four maternal deaths occur for every thousand live births
- ☐ b. Four maternal deaths occur for every ten thousand live births
- ☐ c. Four hundred maternal deaths occur for every hundred thousand live births
- ☐ d. A and C
- ☐ e. None of the above

Question 24

Not yet
answered

Marked out of
1.0

Which of the following statements is not true?

Select one:

- ☐ a. Good nutrition is not intuitive.
- ☐ b. People cannot tell when their children are becoming malnourished because it is not easy to detect for an untrained eye.
- ☐ c. Improving nutrition is pro-poor and increases the income-earning potential of the poor.
- ☐ d. All
- ☐ e. None

Question 25

Not yet
answered

Marked out of
1.0

Which of the following statement is not correct?

Select one:

- ☐ a. Nutrition survey is important to establish base line data
- ☐ b. Nutrition survey is important to understand the extent of nutritional problem
- ☐ c. Nutrition survey is suitable to identify high risk group
- ☐ d. Nutrition survey is important to formulate prediction based on current trend
- ☐ e. All are correct

Question 26

Not yet
answered

Marked out of
1.0

Compared to the earlier international conferences, the Cairo ICPD brought a significant paradigm shift in the issues of health, population and development.

Select one:

- ☐ a. TRUE
- ☐ b. FALSE

Question 27

Not yet
answered

Marked out of
1.0

Which of the following statement is correct?

Select one:

- ☐ a. Zinc deficiency could be a cause of night blindness
- ☐ b. Zinc supplementation can reduce child morbidity and mortality
- ☐ c. A rate of stunting >20% is indicative of zinc deficiency
- ☐ d. In areas where high Phytate diet is a staple zinc deficiency might be common
- ☐ e. All
- ☐ f. None

Question 28

Not yet
answered

Marked out of
1.0

One of the following is not important to fulfil the need and right of adolescents

Select one:

- ☐ a. Family
- ☐ b. Teachers
- ☐ c. Religious leaders
- ☐ d. All
- ☐ e. None

Question 29

Not yet
answered

Marked out of
1.0

Which of the following intervention prevents both pneumonia and diarrhea?

Select one:

- ☐ a. Hib vaccination
- ☐ b. Measles vaccination
- ☐ c. Rota virus vaccination
- ☐ d. PCV vaccination

Question 30

Not yet
answered

Marked out of
1.0

One of the following is included under sexual and reproductive health problem of adolescents

Select one:

- ☐ a. Unwanted pregnancy
- ☐ b. Unsafe abortion
- ☐ c. HIV
- ☐ d. All
- ☐ e. None

Return to: 0 

Health promotion and Disease Prevention

Question 31

Not yet answered

Marked out of 1.0

Which one of the following is not classified under the category of obstetric death?

Select one:

- ☐ a. Abortion caused death
- ☐ b. Deaths of women by profuse bleeding after fall down accident
- ☐ c. Deaths to a woman due to hypertension that for the first time have occurred during gestation
- ☐ d. Mortality of a woman by infection introduced during the process of delivery
- ☐ e. None

Question 32

Not yet answered

Marked out of 1.0

One of the following is the reason for focusing on adolescent

Select one:

- ☐ a. They are high in number
- ☐ b. High unmet need of adolescents services
- ☐ c. Less decision making power of adolescents
- ☐ d. All
- ☐ e. None

Question 33

Not yet answered

Marked out of 1.0

All are true about Anthropometric Indices except

Select one:

- ☐ a. Weight for height identify children suffering from current or acute under-nutrition
- ☐ b. Weight for age identifies both chronic and/or acute under nutrition
- ☐ c. Height for age identifies chronic malnutrition
- ☐ d. All
- ☐ e. None of the above

Question 34

Not yet
answered

Marked out of
1.0

Can be considered as the cause of adolescent SRH problem

Select one:

- ☐ a. Early marriage
- ☐ b. Peer pressure
- ☐ c. Lack of knowledge on contraceptive use
- ☐ d. All
- ☐ e. None

Question 35

Not yet
answered

Marked out of
1.0

Below are list of disease conditions and complications causing neonatal and child deaths

I. Pneumonia

II. Diarrhea

III. Sepsis

IV. Birth asphyxia

V. Congenital anomalies

VI. Prematurity

VII. Injury

Which of the above three conditions are the leading cause of deaths during post-neonatal period and before 5 years of age?

Select one:

- ☐ a. III, V, VI
- ☐ b. II, IV, & VII
- ☐ c. II, III. & IV
- ☐ d. I, II, & III
- ☐ e. III, II, & VII

Question 36

Not yet
answered

Marked out of
1.0

One of the following statements is not true

Select one:

- ☐ a. The income–malnutrition relationship is far above expectation.
- ☐ b. When gross national product (GNP) in developing countries doubles changes in underweight rates are modest.
- ☐ c. Under nutrition and micronutrient malnutrition are direct indicators of poverty.
- ☐ d. None
- ☐ e. All

Question 37

Not yet
answered

Marked out of
5.0

In a nutrition survey, a male child of 84cm weighs 9.9kg. **(5 points)**

- A. Calculate a z-score for this child.
- B. Comment on his result?

The reference median weight for boys of height 84cm is 11.7kg and the standard deviation is 0.908.

Question 38

Not yet
answered

Marked out of
1.0

Which of the following falls outside the definitions children's health?

Select one:

- ☐ a. Enabling children to realize their potential
- ☐ b. Enabling children to satisfy their needs
- ☐ c. Empowering children to nurture their special talents
- ☐ d. Capacitating children to effectively interact with their environment

Question 39

Not yet
answered

Marked out of
1.0

The health benefit of family planning is only for females.

Select one:

- ☐ a. TRUE
- ☐ b. FALSE

Question 40

Not yet
answered

Marked out of
1.0

In an area where there is no Vital Registration (VR) system and Verbal Autopsy (VA), what we basically estimate is

Select one:

- ☐ a. MM (maternal mortality)
- ☐ b. Late MM
- ☐ c. Pregnancy related MM
- ☐ d. All can be estimated easily

Return to: 0 

Health promotion and Disease Prevention

Question 41

Not yet
answered

Marked out of
4.0

Discuss the three interrelated set of strategies required to have improved maternal health (4 points)

Question 42

Not yet
answered

Marked out of
1.0

Which one of the following statement is true?

Select one:

- ☐ a. TGR of > 10% indicates endemic area
- ☐ b. The major source of iodine is animal origin
- ☐ c. School children are at increased risk of Iodine deficiency
- ☐ d. Consumption of contaminated water could be a risk factor for IDD
- ☐ e. All
- ☐ f. None

Question 43

Not yet
answered

Marked out of
1.0

On average an Ethiopian woman have 4 to 5 children during her life span. Which indicator is needed to verify this statement:

Select one:

- ☐ a. Crude birth rate (CBR)
- ☐ b. Sex specific birth rate
- ☐ c. Gravidity & parity
- ☐ d. Total fertility rate (TFR)
- ☐ e. None of the above

Question 44

Not yet
answered

Marked out of
1.0

Which of the following is not an underlying cause of malnutrition?

Select one:

- ☐ a. inadequate access to food
- ☐ b. inadequate care for children and women
- ☐ c. inadequate dietary intake
- ☐ d. insufficient health services

Question 45

Not yet
answered

Marked out of
1.0

If the current trend of neonatal mortality reduction continues how long willit take sub-Saharan African countries to reach same level of neonatal mortality rate developed regions had in 2015 despite new interventions?

Select one:

- ☐ a. 110 years
- ☐ b. 55 years
- ☐ c. 210 years
- ☐ d. 220 years

Question 46

Not yet
answered

Marked out of
1.0

Maternal Death Surveillance and Response (MDSR) aims to

Select one:

- ☐ a. Cut down the preventable portion of MM
- ☐ b. Improve maternal health
- ☐ c. Fight morbidities among mothers
- ☐ d. All

Question 47

Not yet
answered

Marked out of
1.0

Which of the following shows a recent acute malnutrition?

Select one:

- ☐ a. Weight for height < -2SD
- ☐ b. Weight for age < -2SD
- ☐ c. Weight for age < -3SD
- ☐ d. Height for age < -2SD

Question 48

Not yet
answered

Marked out of
1.0

Immediate cause of malnutrition includes

Select one:

- ☐ a. Infections causing loss of appetite
- ☐ b. policies that govern the social sectors
- ☐ c. preferential feeding
- ☐ d. Immunization and early treatment

Question 49

Not yet
answered

Marked out of
1.0

Provision of legal identify to all children including registration at birth is one of the targets of Sustainable Development Agenda. Which SDG agenda does this target fall under?

Select one:

- ☐ a. Survive
- ☐ b. Develop
- ☐ c. Thrive
- ☐ d. Transform

Question 50

Not yet
answered

Marked out of
5.0

List five components of reproductive health care service. **(5 points)**

Health promotion and Disease Prevention

Question 51

Not yet
answered

Marked out of
1.0

Which of the following does not show a correct nutrition & development cycle?

Select one:

- ☐ a. Direct losses in productivity from poor physical status
- ☐ b. Indirect losses from poor cognitive development
- ☐ c. Losses due to increased health care costs
- ☐ d. None

Question 52

Not yet
answered

Marked out of
1.0

One of the following is not true about marasmus

Select one:

- ☐ a. Old man face
- ☐ b. No Fatty liver
- ☐ c. Weight for age between 60-80%
- ☐ d. Common among children less than 2 years of age

Question 53

Not yet
answered

Marked out of
1.0

Which of the following is not a rationale for focusing on child health?

Select one:

- ☐ a. High mortality and morbidity burden
- ☐ b. Early child development is foundational
- ☐ c. The child population is too big
- ☐ d. Health is basic human right
- ☐ e. Causes of death are hardly preventable

Question 54

Not yet
answered

Marked out of
1.0

One of the following statements is not true?

Select one:

- ☐ a. HAZ of $< -2SD$ indicates a chronic malnutrition
- ☐ b. waterloo classification helps to differentiate sever forms of malnutrition
- ☐ c. For intervention with immediate impact, we should use WHZ score
- ☐ d. Prevalence of stunting is usually higher than prevalence of Wasting
- ☐ e. All are correct

Question 55

Not yet
answered

Marked out of
4.0

List down the 7 basis essential obstetric care **(4 points)**

Question 56

Not yet
answered

Marked out of
5.0

Describe the reproductive rights. And list three human rights that can be affected by reproductive rights. **(5 points)**

Question 57

Not yet
answered

Marked out of
1.0

In addition to effective contraception use, hormonal contraception methods are effective in protecting women from STI/STDs.

Select one:

- ☐ a. TRUE
- ☐ b. FALSE

Question 58

Not yet
answered

Marked out of
1.0

Which one of the following is a risk factor for IDA?

Select one:

- ☐ a. Infection
- ☐ b. Low intake of phytates
- ☐ c. Deworming
- ☐ d. Increased intake of poly phenols
- ☐ e. A and D
- ☐ f. None

Question 59

Not yet
answered

Marked out of
5.0

Describe the "**GATHER**" counselling approach in FP. **(5 points)**

|

|

|

|

Question 60

Not yet
answered

Marked out of
1.0

What is in the heart of today's ANC approach?

Select one:

- ☐ a. Birth Preparedness and Complication Readiness
- ☐ b. History taking and thorough physical examination
- ☐ c. Number of visits
- ☐ d. All

Return to: 0 

Health promotion and Disease Prevention

Question 61

Not yet answered

Marked out of 1.0

The proportion of births attended by the skilled birth attendants according to the 2016 EDHS report is

Select one:

- ☐ a. 26%
- ☐ b. 64%
- ☐ c. 28%
- ☐ d. 34%

Question 62

Not yet answered

Marked out of 1.0

One of the following statements is not true?

Select one:

- ☐ a. HAZ of $< -2SD$ indicates a chronic malnutrition
- ☐ b. waterloo classification helps to differentiate sever forms of malnutrition
- ☐ c. For intervention with immediate impact, we should use WHZ score
- ☐ d. Prevalence of stunting is usually higher than prevalence of Wasting
- ☐ e. All are correct

Question 63

Not yet answered

Marked out of 1.0

According to the recent national survey (EDHS), one of the following is correct about Ethiopia:

Select one:

- ☐ a. Nearly a quarter of married women use modern contraceptive,
- ☐ b. Nearly a third of married women use modern contraceptive,
- ☐ c. Nearly half of married women use modern contraceptive,
- ☐ d. Nearly two third of married women use modern contraceptive,
- ☐ e. None of the above

Question 64

Not yet
answered

Marked out of
1.0

Dietary reference level which is used when there is no enough data about Estimated Average Requirements and established based on animal experiments is called

Select one:

- ☐ a. RDA
- ☐ b. Upper tolerable intake
- ☐ c. Lower tolerable intake
- ☐ d. Adequate intake
- ☐ e. None

Question 65

Not yet
answered

Marked out of
1.0

Changing social relationships, expectations, roles and responsibilities is majorly seen in

Select one:

- ☐ a. Childhood
- ☐ b. Adulthood
- ☐ c. Adolescents
- ☐ d. All

Question 66

Not yet
answered

Marked out of
1.0

Which of the following statement is true about maternal vitamin A supplementation?

Select one:

- ☐ a. Vitamin A supplementation is given as a routine ANC service
- ☐ b. In areas where the prevalence of night blindness is $>5\%$ 25,000 IU of vitamin A daily is recommended
- ☐ c. Vitamin A supplementation for mothers within 45 days of delivery is recommended
- ☐ d. B and C
- ☐ e. None

Question 67

Not yet
answered

Marked out of
1.0

Early pregnancy and abortion related deaths are two of the major conditions that lead to under estimation of MM

Select one:

- ☐ a. False
- ☐ b. True

Question 68

Not yet
answered

Marked out of
1.0

The claim that all pregnant women must be attended by Skilled Birth Attendant is built up on the fact that

Select one:

- ☐ a. All women have the right to get the highest possible quality of care
- ☐ b. Major obstetric complications are usually unpredictable
- ☐ c. Skilled birth attendants are easily accessible
- ☐ d. Skilled birth attendants such as physicians are highly valued professionals
- ☐ e. All

Question 69

Not yet
answered

Marked out of
1.0

The Cairo ICPD recommendation identified the following as favorable conditions for fostering RH:

Select one:

- ☐ a. Women empowerment
- ☐ b. Gender equity
- ☐ c. Male participation
- ☐ d. A and B only
- ☐ e. all of the above

Question 70

Not yet
answered

Marked out of
1.0

One of the following is false about adolescents

Select one:

- ☐ a. All adolescents have the same needs
- ☐ b. Adolescents have changing needs
- ☐ c. A boy adolescent is physically and psychologically different from a girl adolescent
- ☐ d. None

Return to: 0 

Health promotion and Disease Prevention

Question 71

Not yet
answered

Marked out of
1.0

Which one is not true about anthropometric nutritional assessment method?

Select one:

- ☐ a. It is widely used in the assessment of protein-energy malnutrition
- ☐ b. It can be used to monitor periodic changes in growth and/or body composition
- ☐ c. The procedures can assist in the identification of mild to moderate malnutrition
- ☐ d. It can identify specific nutrient deficiency
- ☐ e. None of the above

Question 72

Not yet
answered

Marked out of
1.0

Intake that poses no risk of adverse health effect

Select one:

- ☐ a. Upper level of safe intake
- ☐ b. Safe intake
- ☐ c. LRI
- ☐ d. EAR

Question 73

Not yet
answered

Marked out of
1.0

Which one of the following is not correct about WHO classification of public health problem of xerophthalmia?

Select one:

- ☐ a. Bitot spot of $> 0.5\%$
- ☐ b. Corneal scar of $> 0.05\%$
- ☐ c. Corneal xerosis of $> 0.01\%$
- ☐ d. Night blindness of $> 5\%$
- ☐ e. None

Question 74

Not yet
answered

Marked out of
1.0

Adolescents friendly service can provide all except

Select one:

- ☐ a. Safe abortion services
- ☐ b. Nutrition service
- ☐ c. All
- ☐ d. Contraception
- ☐ e. None

Question 75

Not yet
answered

Marked out of
1.0

Which of the following statements is not true?

Select one:

- ☐ a. Stunting is linked to cognitive development
- ☐ b. Child's weight for age is best predictor of human capital
- ☐ c. Undernourished children are likely to become short adults
- ☐ d. Stunting in the first two years is related with reduced economic productivity

Question 76

Not yet
answered

Marked out of
1.0

Which one of the following statement is true?

Select one:

- ☐ a. A 20% of low hemoglobin for age is considered mild
- ☐ b. A higher PH level enhance iron absorption
- ☐ c. Heme Iron is found from plant source
- ☐ d. All
- ☐ e. None

Question 77

Not yet
answered

Marked out of
1.0

Which of the following is true about child mortality burden and trend?

Select one:

- ☐ a. With current rate of mortality reduction, the world will meet SDG targets
- ☐ b. Reduction in neonatal and post-neonatal U5 mortality is proportional
- ☐ c. There is no disparity in neonatal mortality burden across the different regions of the world
- ☐ d. 75 – 80% stillbirths occur in Southern Asia and Sub-Saharan Africa
- ☐ e. The rate of mortality reduction among under-5 and 5-9 is the same

Question 78

Not yet
answered

Marked out of
1.0

Lactational amenorrhea contraceptive method is effective only for the first six months after delivery.

Select one:

- ☐ a. FALSE
- ☐ b. TRUE

Question 79

Not yet
answered

Marked out of
1.0

Which of the following mother has low likelihood of having stillbirth?

Select one:

- ☐ a. Mother from impoverished neighborhood
- ☐ b. Mother who use modern contraceptive
- ☐ c. Older mother
- ☐ d. Mother living in rural area

Question 80

Not yet
answered

Marked out of
1.0

Criteria for Adolescents friendly service include

Select one:

- ☐ a. Welcoming
- ☐ b. Good listener
- ☐ c. Non judgmental
- ☐ d. All
- ☐ e. None

Health promotion and Disease Prevention

Question 81

Not yet
answered

Marked out of
1.0

Adolescents are within the age range of

Select one:

- ☐ a. 10-19
- ☐ b. 15-24
- ☐ c. 10-24
- ☐ d. All

Question 82

Not yet
answered

Marked out of
5.0

The median weight for a female child of height 90cm is 13.7kg.

In a nutritional survey, a female child of 90cm weighs 12kg. **(5 points)**

A. Calculate percent of median?

B. Comment on his result.

Question 83

Not yet
answered

Marked out of
1.0

One of the following is not considered as natural contraception method:

Select one:

- ☐ a. Abstinence
- ☐ b. Standard calendar method
- ☐ c. Rhythmic calendar method
- ☐ d. Coitus interruptus
- ☐ e. None of the above

Question 84

Not yet
answered

Marked out of
1.0

Which one of the following statement is true?

Select one:

- ☐ a. Basal metabolism rate represents 45-70% of daily total energy expenditure
- ☐ b. Basal metabolism is the most variable and account for the largest component of daily energy expenditure
- ☐ c. Dietary reference intakes provide qualitative advice to the public about diet and chronic disease prevention
- ☐ d. None

Question 85

Not yet
answered

Marked out of
1.0

One of the following is important for adolescents to grow and develop in good health

Select one:

- ☐ a. Information
- ☐ b. Safe and supportive environment
- ☐ c. Counselling
- ☐ d. All
- ☐ e. None

Question 86

Not yet
answered

Marked out of
5.0

What are the two cause of LBW and which one is the
commonest in developing country? **(5Points)**

|

|

|

|

Return to: 0 