

MCQs
In
Upper limb

By

DR.KHALID MILAD

Select the most appropriate answer:

1. one of the following is true for radial nerve :

- a-Branch of the medial cord
- b-Anterior interosseous nerve its branch in forearm
- c-Passing in spiral groove with profunda brachii artery
- d-Supply the flexor carpiradialis

C

2-Regarding the cubital fossa, all of the following are true except :

- a-Contains the biceptal tendon
- b-Contain the medial nerve medially
- c-The floor is formed by the brachioradialis and supinator muscles .
- d-The apex is formed by meeting the pronator teres and brachioradialis .

C

3-The following are contents of deep group of anterior compartment of forearm except :

- a-Flexor pollicis longus
- b-Pronator teres
- c-Pronator quadrates
- d-Supinator muscles

B

4-Regarding the radial artery all of the following are true except :

- a-Lies deep to brachioradialis muscle in upper part of forearm .
- b-Gives the common interosseous artery.
- c-Lies lateral to the tendon of flexor digitorum Superficialis .
- d-Can be felt in lower part of forearm .

B

5-Pronator quadrates muscle is supply by ;

- a-Ulnar-nerve , b-Median nerve c-Radial nerve d-Anterior interosseous nerve

D

6-Regarding flexor Carpi radialis muscle all of the following ;

- a-Most Lateral muscle in the superficial layer of flexor compartment of forearm .
- b-Take origin from the common flexor origin .
- c-Flexor and abductor of the wrist
- d-Innervated by the median nerve

ALL true

7-The structure which pass behind the medial epicondyle is :

- ☒ a-Ulnar nerve b-Ulnar artery c-Radial artery d-Radial nerve

8-The Lateral half of the flexor digitorum profundus is supply by:

- a-Ulnar nerve only ☒ b-Median nerve only
c-Both ulnar and median nerves d-Posterior interosseous nerve

9-All the following are true for median nerve except :

- a-Has two root from Lateral and medial cords.
☒ b-Posterior interosseous nerve it is branch in forearm .
c-Passing between two heads of pronator teres .
d-Enters the palm by passing behind the flexor retinaculum .

10-Regarding the cubital fossa ,All the following are true except :

- a-Contains the brachial artery
b-Contains the radial nerve
☒ c-The floor is formed mainly by the brachialis muscle
d-The apex is formed by meeting the pronator teres and brachioradialis

11-All the following are contents of anterior compartment of forearm except:

- a-Flexor digitorum superficialis .
b-Pronator teres
c-Pronator quadratus
☒ d-Brachioradialis

12-Regarding the radial artery all the following are true except :

- a-Lies deep to brachioradialis muscle in upper part of forearm
☒ b-Gives the superficial palmar arch
c-Lies medial to the tendon of flexor carpi radialis in the distal forearm
d-Can be felt in snuff box

13-Regarding flexor carpi ulnaris muscle all the following are true except :

- ☒ a-Most medial and in the deep layer of flexor

b-Take origin from the olecranon and posterior border of the ulna

c-Flexor and adductor of the wrist

d-Innervated by the ulnar nerve

14-The structure which pass behind the medial epicondyle is :

☒ a-Ulnar nerve 2-Ulnar artery 3-Radial artery d-Radial nerve

15-Regarding the forearm ,all the following are true except :

a-The radius is lateral

b-The interosseous membrane is a thin membrane

☒ c-Flexor muscles arranged in two groups

d-All the extensor superficial muscles group are take origin from lateral epicondyle

16-The deltoid muscle can perform all the following action around the shoulder ,except :

a-Flexion

b-Medial rotation

c-Lateral rotation

☒ d-Initiate Abduction

17-Only one structure from the following is lateral to the third part of axillary artery :

a-Ulnar nerve

b-Axillary vein

c-Medial cutaneous nerve of arm

☒ d-Musculocutaneous nerve

18-The shoulder joint can be adducted by all the following Except:

☒ a-Supraspinatus

b-Latissimus dorsi

c-Teres major

d-Teres minor

19-The following are anterior to the shoulder joint ,except:

a-subscapularis muscle

b-Axillary muscle

c-Coracoacromial ligament

☒ d-Brachial plexus

20-Pain may be referred to the shoulder region in the gallbladder disease ,by the following nerve :

☒ a-Supraclavicular nerve

b-Axillary nerve

c-Subscapular nerve

d-Nerve to subclavius

21-All the following branches of brachial artery anastomoses around elbow joint ,except :

- a-Profunda artery
- b-Superior ulnar collateral artery
- ☒ c-Nutrient artery
- d-Inferior Ulnar collateral artery

22-Regarding the arm region all the following statements are true except :

- a-Ulnar nerve has no branches in the anterior compartment of the arm
- b-Radial nerve in the spiral groove ,accompanied by the profunda vessels
- ☒ c-Median nerve crosses the brachial artery from medial to lateral ,
- d-tendon of biceps is lateral to brachial artery in the lower part

23-The content of the cubital fossa ,arranged from lateral to medial as the following :

- ☒ a-Median nerve ,Bifurcation of brachial artery ,Tendon of biceps, Radial nerve ,
- b-Bifurcation of brachial artery ,tendon of biceps ,Axillary vessels ,radial nerve
- c-Radial nerve ,tendon of biceps ,Bifurcation of brachial artery ,median nerve.
- d-Radial nerve, tendon of biceps ,median nerve ,ulnar nerve.

24-The structure in the roof of cubital fossa ,protects brachial artery during vein puncture is :

- a-Superficial nerves
- ☒ b-Bicipital aponeurosis
- c-Radius bone
- d-Skin

25-Distal end of radius bone is all the following ,Except :

- a-Styloid process projects distally from its Lateral margin
- b-Ulnar notch medially ,articulates with round head of ulna
- c-Articulates with the Scaphoid and lunate bones at wrist joint
- ☒ d-Posteriorly grooved on medial side by tendon of extensor carpi radialis

26-Regarding the elbow joint ,only one statement is true:

- a-Triceps and anconeus muscles are the elbow flexors
- b-Distal radioulnar joint communicates with the elbow joint
- c-Carrying angle disappears when elbow joint is fully extended

☒ d-Strength of proximal radioulnar joint depends on annular ligaments

27-Stability of elbow joint depends mainly on all the following ,except :

a-Wrench –shaped articular surface of olecranon

b-Muscular support

☒ c-Pulley-shaped trochlea of humerus

d-Medial and lateral ligaments

28-The direct articulation parts in wrist joint are all the following except:

a-Radius

b-Articular disc

c-Scaphoid

☒ d-Ulna

29-Flexor and extensor carpi ulnaris can do another movement around the wrist joint:

a-Abduction

☒ b>Adduction

c-Rotation

d-All of the above

30-Regarding the thumb ,all are true ,except ;

☒ a-Flex the thumb occur in anteroposterior plane

b-Metacarpal bone of the thumb functions independently

c-Trapezium articulates with the metacarpal bone of the thumb

d-Rotation is possible movement in the saddle joint of the thumb

31-Compartment of the forearm is separated by all the following except:

a-Lateral intermuscular septum

b-Interosseous membrane

☒ c-Medial intermuscular septum

d-Attachment of deep fascia and ulna

32-One from the following sensory nerves is radial nerve branch :

^{lower}
☒ a-Lateral cutaneous nerve of forearm

b-Medial and ulnar nerves

c-Ulnar nerve

d-Median and radial nerves

33-About the Palmar Aponeurosis one statement is true:

a-Thickening of the hypothenar part of deep fascia of the palm

b-Attached to the distal border of the extensor retinaculum

☒ c-Receives the insertion of the palmaris longus tendon

d-Deep to the long flexor tendons

- a-Opponens b-Flexion c-Extension d-Adduction

37-The unique muscles connecting flexor tendons with extensor tendons are :

- 38-Injury to median nerve at elbow cause all the following except :**

- 39-If the small muscle of hand are lost except thenar muscles and first 2 Lumbricals.the nerve injury in:**

- 40-The following clinical signs are related to the accompanied nerve ,Except :**

- 41-Allen's test (hand clinical examination)used to test the :**

- a-Hand arterial supply
- b-Hand motor innervations
- c-Hand sensory innervations
- d-Thumb function



- 7

c-Anterior branch of inferior ulnar collateral A

☒ d-Posterior branch of inferior ulnar collateral A

43-Injury of the long thoracic nerve is causing :

a-Drop the wrist b-Claw hand finger ☒ c-Wedge of scapula d-Extension

44-Abduction of the arm done by the following except :

a-Start the action of the supra spinatus muscle up to 15 degree

b-Deltoid muscle up to 90 degree

☒ c-Above the 90 degree by rotation of scapula backward

d-Is Opposite by adduction

45-Basilic vein is terminated to :

☒ a-Axillary vein

b-Subclavian vein

c-Cephalic vein

d-Brachial vein

46-Anterior axillary fold is formed by :

a-Latissimus dorsi

☒ b-Pectoralis major

c-Pectoralis minor

d-Serratus anterior

47-The breast is lies on the following muscles except :

a-Pectoralis major
abdominis

b-serratus anterior

☒ c-Latissimus dorsi

d-External Oblique

48-Delto-pectoral groove has the following structure except:

a-Branch of the thoracoacromial artery

b-Cephalic vein

☒ c-Basilic vein

d-Lymph node

49-Extensor expansion of the finger formed by following except :

☒ a-Extensor carpi ulnaris

b-Palmar interosscous

c-extensor digitorum

d-Lumbricalis

50-All the following arteries are sharing in the anastomosis around the scapula except :

- ☒ a-Posterior circumflex humeral artery
- b-Suprascapular artery
- c-Circumflex subscapula artery
- d-Subscapular artery

51.All of the following are lymphatic drainage of the breast except :

- a-Anterior group of axillary lymph nodes
- b- **Apical** group of axillary lymph nodes
- ☒ c-Lateral group of axillary lymph nodes
- d-Central group of axillary lymph node

52-One of the following is branch of the third part of axillary artery :

- a-Internal thoracic artery
- b-Lateral thoracic artery
- c-Thoraco-acromial artery
- ☒ d-Subscapular artery

53-Biceps brachialis has the following action except:

- a-flexion of the shoulder joint
- b-Flexion of the elbow joint
- ☒ c-pronation of the forearm
- d-Supination of forearm

54-Clavi pectoral fascia is piercing by the following structure except:

- a-Acromial branch of the thoracoacromial artery
- b-Cephalic vein
- ☒ c-Basilic vein
- d-Lateral pectoral nerve

55-All the following are true for deltoid muscle except :

- a-Has origin in the clavical ,acromial and spine process of scapula
- b-Innervated by axillary nerve

c-Powerful adductor muscle

d-Multipennate fibers

