

EPIDEMIOLOGY

- ___ 1. In a recent survey, investigators found that the prevalence can be represented Disease A was higher than the prevalence of Disease B. The incidence and seasonal pattern of both diseases are similar. Explanations consistent with this observation include:
- A. patients recover more quickly from Disease A than from Disease B
 - B. patients recover more quickly from Disease B than from Disease A
 - C. patients die more quickly from Disease A than from Disease B
 - D. patients die more quickly from Disease B than from Disease A
 - E. A and C are correct
 - F. B and D are correct
- _?_ 2. If 10 cases of Disease A occur during a 2-year period in a stable population of 50,000 people, then the person-time rate of Disease A in that population is approximately:
- A. 10 cases/ 5,000 person-years
 - B. 10 cases/ 25,000 person-years
 - C. 10 cases/ 49,990 person-years
 - D. 10 cases/ 50,000 person-years
 - E. 10 cases/ 100,000 person-years
 - F. none of the above
- ___ 3. To investigate the association between Kawasaki syndrome (KS) and carpet shampoo, investigators conducted a case-control study with 100 cases and 200 controls. Among the children with KS, 50 gave a history of recent exposure to carpet shampoo. Among those without KS, 50 gave a history of recent exposure to carpet shampoo. For this study, the odds ratio is:
- A. 1.0
 - B. 1.5
 - C. 2.0
 - D. 3.0
 - E. can not be calculated from the information given
- d ___ 4. Bar charts may be distinguished from histograms at a glance because:
- A. bar charts are not used for time series data
 - B. histograms are used to display discrete data
 - C. bar charts are based on area under the curve
 - D. histograms do not have spaces between consecutive columns
 - E. none of the above
- ___ b _ 5. The primary difference between a surveillance system and a survey is:
- A. a surveillance system is population-based
 - B. a surveillance system is ongoing
 - C. a surveillance system cannot assure confidentiality
 - D. a survey is generally cheaper
 - E. none of the above
- ___ 6. The ability of a surveillance system to detect the cases it is intended to detect is referred to as:
- A. predictive value positive
 - B. representativeness

ifcity

C. sensitivity

D. spec

E. none of the above

___ 7. Which one of the following is **NOT** a disease having animal reservoir?

A. Rabies

B. Typhoid fever

C. Brucellosis

D. Anthrax

E. Non of the above

___ a_ 8. Which of the following is the objectives of PRIMARY PREVENTION ?

A. to prevent exposure, and prevent disease.

B. to stop or slow the progression of disease

C. to limit the impact of damage

D. for early detection and treatment of disease.

E. none of the above

___ 9. The time course of a disease outbreak is usually displayed as a/ an:

A. secular trend

B. seasonal trend

C. epidemic curve

D. endemic curve

E. none of the above

___ b_ 10. The primary difference between an experimental and observational study is:

A. the investigator is 'blinded' in an experimental study but not in an observational study

B. the investigator controls the subject's exposure in an experimental study but not in an observational study

C. the investigator controls the subject's outcome in an experimental study but not in an observational study

D. experimental studies are conducted with animals; observational studies are conducted with humans

E. none of the above