

The largest lymphatic duct (thoracic duct) begins

☐

a.

Infront of L4 vertabrae

☐

b.

Infront of T12 vertabrae

☒

c.

At left venous angle

☐

d.

Infront of L2 vertabrae

☐

e.

Infront of T10 vertabrae

All of the following features can be seen on inspection of abdominal wall except

☒

a.

Umbilical folds

☐

b.

Caput medusae

☐

c.

Striae gravidarum:

☐

d.

Linea nigra

☐

e.

None

Abeba, a 25-year-old 4th year medical student was studying in the library until lunch hour. As she was going home for lunch, she felt a smell of delicious 'doro wot' from the nearby restaurant. She then entered into the Hotel and ate the 'doro wot' following which her gastric juice secretion increased tremendously. This phase of gastric juice secretion is most likely:

☐

- a.
Gastric phase of salivary juice secretion
☐
- b.
Intestinal phase of gastric juice secretion
☒
- c.
Gastric phase of gastric juice secretion
☐
- d.
Cephalic phase of pancreatic juice secretion
☐
- e.
Cephalic phase of gastric juice secretion

Correct about salivary secretion

- ☐
- a.
At highest flow rate it is most dissimilar with plasma
☒
- b.
Halitosis in sjogren syndrome
☐
- c.
The secretion is increased profusely by sympathetic action
☐
- d.
Saliva is hypertonic than plasma
☐
- e.
Contains mucin for amylase action

The nutrient type absorbed into the lymphatic system is

- ☐
- a.
Vitamin A
☐
- b.
Short chain fatty acid

☐

c.

Triglyceride

☐

d.

Amino acid

☒

e.

A and C are correct

A persistence of the vitelline duct may result in all except:

☐

a.

umbilical fistula

☐

b.

Meckel's diverticulum

☒

c.

urachus (allantoic duct)

☐

d.

ileal diverticulum

☐

e.

vitelline cyst

The cloacal membrane is derived from:

☐

a.

ectoderm

☐

b.

mesoderm

☐

c.

endoderm

☒

d.

both endoderm and ectoderm

☐

e.
yolk sac membrane

Select the false statement about CHPS?

☒

a.
Caused by closure of gastric inlet by cardiac sphincters muscle

☐

b.
More common in males than females

☐

c.
Presents with projectile vomiting

☐

d.
Visible abdominal peristalsis

Patients with portal hypertension can present with all of the following signs and symptoms except

☐

a.
Esophageal varices

☐

b.
Caput medusa

☐

c.
Hemorrhoid

☒

d.
None

Which of the following cells of the gastric glands are large and acidophilic?

☐

a.
Basket cells.

☐

b.
Zymogenic cells.

☐

c.

Paneth cells.

☐

d.

Mucous neck cells.

e. oxyntic cell

☒

In infants, defecation often follows a meal. The cause of colonic contraction in this situations is

a.

The gastroileal reflex

b.

The enterogastric reflex

c.

The gastrocolic reflex

d.

Increased circulating levels of CCK

e.

Increased circulating levels of somatostatin

Which component of GIT is least adapted for protection of GIT mucosa from assault?

a.

Somatostatinergic secretion

b.

Histaminergic secretion

c.

Serous layer of the GIT

d.
Secretion of Brunner's gland

e.
Electrolyte secretion

The caudal limb of the primary intestinal loop forms many intestine structures except:

a.
proximal 2/3 rd of transverse colon

b.
cecum

c.
distal ileum

d.
ascending colon

e.
vitelline duct

Brunner's glands are distinguishing glands of which region of the GIT?

a.
Jejunum

b.
Appendix

c.
Ileum

d.
Large intestine

e.
Duodenum

Which of the following is NOT true about defecation reflex

- a.
Reflex centers are located in the sacral spinal cord
- b.
Initiated by accumulation of feces in the rectum by mass movement
- c.
Involves relaxation of distal colon
- d.
The external sphincter relaxes
- e.
ENS has minimal role as compared to ANS

If the IVC is compressed by the large fetus or by any other pathology at L4vertebrae, by which of the following vein can the blood reach in to the heart?

- a.
Paraumbilical veins
- b.
Azygo vein
- c.
Inferior epigastric veins
- d.
All

Which region of stomach has relatively short gastric glands, but long and branching pits?

- a.
Pylorus
- b.
Fundus
- c.
Cardia
- d.
Body and fundus
- e.
Cardiac and fundus

A patient deficient in trypsin inhibitor has a primary abnormality that involves the:

- a.
Small intestine epithelium
 - b.
Salivary glands
 - c.
Pancreas
 - d.
Liver
 - e.
Oxyntic cells of the stomach
- The bare areas of the liver is found on

- a.
Anterior surface

- b.
Posterior surface
- c.
Superior surface
- d.
Inferior surface

Which of the following statements is not correct regarding the histology of the oral cavity?

- a.
Enamel has no collagen fibers.
- b.
The collagen bundles present in the periodontal ligaments of the teeth have high turnover.
- c.
The extrinsic muscles of the tongue are skeletal, while the intrinsic muscles are smooth in type.
- d.
The transitional border of the lip has no gland, unlike the outer or inner surfaces.
- e.
Circumvallate papillae of the tongue are located in front of the sulcus terminalis in a single row.

The superior mesenteric and splenic veins unite to form the portal vein posterior to which of the following structure?

- a.
right kidney
- b.
neck of the pancreas

- c.
pylorus of stomach
- d.
Spleen
- e.
2nd portion of the duodenum

Muscularis mucosae of the esophagus is:

- a.
Contains myenteric plexus
- b.
Well-formed than the muscularis externa
- c.
Found as outer longitudinal and inner circular layers
- d.
Found as longitudinal layer with no circular layer
- e.
Made by both skeletal and smooth muscles

Select the true statement about stomach

- a.
Lesser curvature gives attachment to gastrophrenic ligaments
- b.
Anteriorly it relates to the diaphragm
- c.
Lesser curvature is shorter and forms left border of stomach

d.
None

Long-chain fatty acids absorbed by the intestinal mucosa cells reach the blood in the form of:

- a.
micelles
- b.
free fatty acids
- c.
monoglycerides
- d.
chylomicrons
- e.
diglycerides

Which of the following regions of the digestive tract does not contain submucosal glands?

- a.
Stomach
- b.
Duodenum
- c.
Anal canal
- d.
Oral cavity

e.
Esophagus

Identify the correctly matched gastric cell with its secretions

a.
Enterochromaffin cells..... Histamine

b.
Chief cells..... HCl

c.
D cells Intrinsic factor

d.
G-cells Secretin

e.
Parietal cells Pepsinogen

Glands in the large intestine are:

a.
Simple straight tubular

b.
Compound tubular

c.
Simple coiled tubular

d.
Simple branched tubular

e.

Simple acinar

During development of the gut which one is wrong?

- a.
the urorectal septum may fail to divide the cloaca, causing a rectovaginal fistula in the female
- b.
failure of the intestinal loops to return into the abdominal cavity leads to omphalocele
- c.
the stomach rotates, causing the left vagal trunk to innervate its anterior wall
- d.
gut rotation and fusion of mesentery to the dorsal body wall makes the spleen retroperitoneal
- e.
umbilical fistula derived from retention of the urachus

The transport of glucose through basolateral membrane of intestine into blood capillary is by:

- a.
GLUT2 transporter
- b.
Primary active transport
- c.
H⁺ - glucose symport
- d.
GLUT5 transporter

e.

Na⁺ - glucose symport

Which of the following is correct for distinguishing between parotid and exocrine portion of pancreas?

a.

Pancreas has a few mucous producing glandular cells, while parotid is purely serous.

b.

Parotid has centroacinar cells, while pancreas has many striated ducts.

c.

Pancreas has centroacinar cells and many striated ducts, while parotid has none of these.

d.

Parotid has centroacinar cells and many striated ducts, while pancreas has none of these.

e.

Pancreas has centroacinar cells, while parotid has many striated ducts.

The portal lobule of the liver is demarcated by:

a.

Portal triad at the center and central veins of 6 classic liver lobules at the corners.

b.

Sides of two adjacent classic lobules at the center, and alternate corners by central veins and portal canals of two adjacent classic lobules.

c.

Central vein at the center and portal triads of 3 classic liver lobules at the corners.

d.
Central vein at the center and portal triads of 6 classic liver lobules at the corners.

e.
Portal triad at the center and central veins of 3 classic liver lobules at the corners.

One of the following is true about ascending colon.

a.
Narrower than caecum

b.
Crosses right lateral cutaneous nerve of thigh

c.
It gets blood supply from right colic artery

d.
All

One of the following is incorrectly matched item

a.
Short gastric artery – celiac trunk

b.
Left gastroepiploic artery –splenic artery

c.
Right gastroepiploic artery –gastroduodenal artery

d.
Right gastric artery—common hepatic artery

Bile canaliculi are lined by:

- a.
Simple columnar epithelium
- b.
Hepatocytes
- c.
Cholangiocytes
- d.
Simple squamous epithelium
- e.
Partly by hepatocytes and partly by cholangiocytes

If large portion of small intestine is removed due to medical reason, a person will develop a short bowel syndrome. As a result he may suffer from all of the following EXCEPT;

- a.
Muscle wasting
- b.
Constipation
- c.
Steatorrhea
- d.
Megaloblastic anemia
- e.
Osteomalacia

Identify the INCORRECT statement about large intestine

- a.
Provides enzymatic digestion of organic nutrients
- b.
Site of diverticulosis and diverticulitis
- c.
Provides microbial digestion and vitamin synthesis
- d.
Site for haustral contractions that decrease transit time of ingesta
- e.
Provides absorption of water and electrolytes

The type of epithelium lining the surface of stomach where digestion is taking place in an acidic environment is:

- a.
Simple columnar
- b.
Stratified squamous keratinized
- c.
Stratified squamous nonkeratinized
- d.
Simple cuboidal
- e.
Transitional

One of the following is false about liver H

- a.
Posterior of right limb is formed by fissure of IVC

- b.
Anterior part of right limb is formed by fossa of gallbladder
- c.
Posterior part of left limb is formed by fissure of falciform ligament
- d.
Anterior part of left limb is formed by fissure of ligamentum teres

The major lipolytic enzyme in the man

- a.
Are mainly found in gastric juice
- b.
are mainly the Enteric lipases
- c.
Are potentiated by bile salt
- d.
Are activated by enterokinase
- e.
Require a PH greater than 10 for optimal activity

After secretion into the duodenum, the enzyme trypsinogen is converted into its active form, trypsin, by:

- a.
Procarboxypeptidase
- b.
Enterokinase
- c.
Co-lipase

d.
Trypsin inhibitor

e.
HCL and trypsin already found in the duodenum

One of the following is the 2nd most common presentation of appendix.

a.
Pre-ileal

b.
Retrocaecal

c.
Subcaecal

d.
Pelvic