

1. What is the cut-off rate for fast breathing in a child who is 7 weeks old?
 - a) 60 breaths per minute or more
 - b) 40 breaths per minute or more
 - c) 50 breaths per minute or more
 - d) 30 breaths per minute or more
 - e) 70 breaths per minute or more
2. In a sick child, which of the following signs are not considered as a general danger sign?
 - a) unable to drink or breast-feed
 - b) vomiting everything
 - c) severe chest indrawing
 - d) lethargy or unconsciousness
 - e) all are general danger signs
3. What are the four main symptoms for which every sick child should be asked and assessed?
 - a) malnutrition, cough, vitamin A status, ear problem
 - b) cough, diarrhea, malnutrition, ear problem
 - c) anemia, fever, diarrhea, ear problem
 - d) cough, diarrhea, fever, ear problem
 - e) None of the above
4. What is a child's classification or diagnosis if he is 10 months old, has had a cough that lasted two days, has a breathing rate of 46 breaths per minute and chest indrawing?
 - a) no pneumonia: cough or cold
 - b) pneumonia
 - c) severe pneumonia or very severe disease
 - d) very severe febrile disease
 - e) None of the above
5. A 3-year-old child has fever and diarrhea. He has no danger signs and no cough. The child has had diarrhea for 2 weeks. The child is restless and irritable, but is not drinking eagerly. His eyes are not sunken. A skin pinch goes back slowly. He does not have a cough or difficult breathing. He does not have an ear problem. The child should be classified as:
 - a) Severe dehydration
 - b) Some dehydration and Severe persistent diarrhea
 - c) Persistent diarrhea
 - d) Severe dehydration and Persistent diarrhoea
 - e) Dysentery

6. A child with fever plus any general danger sign should be classified as:
- a) Malaria
 - b) Very severe febrile disease
 - c) Severe pneumonia
 - d) Mastoiditis
 - e) Severe and complicated measles
7. If a child has had ear pain and pus draining from the ear for 10 days, you will classify this child as having:
- a) acute ear infection
 - b) Mastoiditis
 - c) chronic ear infection
 - d) not enough signs to classify this child
 - e) None
8. Which sick children should be checked for malnutrition and anemia?
- a) only children with feeding problems
 - b) only children who are not breastfed
 - c) only children who are younger than 12 months old
 - d) only children with diarrhea
 - e) all children brought to the health facility
9. Feeding should be assessed in a child who:
- a) need urgent referral
 - b) is classified as having anemia or Moderate acute malnutrition
 - c) is less than 3 years old and does not need urgent referral
 - d) is classified as having persistent diarrhean
 - e) None
10. A 12 month old boy with persistent diarrhoea, has MUAC of 11.4 cm and no edema; which of the following are correct action for this child?
- a) Child should be considered to have moderate acute malnutrition
 - b) Child should be admitted to OTP program in the nearby HP or HC
 - c) Health worker should do appetite test with RUTF
 - d) Child should be admitted to inpatient service
 - e) None of the above