

Module 15 test one and two

1. Characteristics of stable malaria vs unseasonal malaria, stable malarias are:
 - A. Makes the Area prone to malaria epidemics
 - B. Population low level of immunity
 - C. Affects mostly children
 - D. Its epidemiology fluctuates from week-to-week month to month
2. The usual infective stage of malaria to man is the Select one:
 - A. gametocytes
 - B. sporozoites
 - C. schizonts
 - D. merozoites
 - E. C and D
3. Infective form of plasmodium for human form of mosquito is
 - A. Merozoite
 - B. Schizont
 - C. Sporozoite
 - D. sporocyst
 - E. gametocyte
4. Trichinosis which one is incorrect
 - A. encysted larva ingestion causes dz
 - B. are larviparous
 - C. Same host acts as definitive and intermediate host
 - D. males die after fertilization
5. Found on veins of bladder
 - A. S. H
 - B. S. M
 - C. S. I
 - D. S. J
 - E. S. masoni
6. What is the cause of febrile episode of malaria
 - A. rupture of hepatocyte
 - B. rupture of RBC
 - C. infection of hepatocyte by sporozoite
7. HbG S is protective against which specie of plamodoium
 - A. P.o
 - B. P.v
 - C. P.f
 - D. P.m
 - E. P.k
8. The unique pathological condition of P. falciparum?
 - A. Vascular obstruction
 - B. Hepatomegaly
 - C. Leukopenia

- D. Anemia
E. Fever
9. RBC cytoadherence in the microvasculature such as in cerebral brain is characteristic of which malaria species
- A. P.o
B. P.f
C. P.m
D. P.v
10. Babesia its definitive host is
- A. dog
B. pig
C. human
D. tick
E. anopheles mosquito
F. none
11. Hydatid treatment and Laboratory diagnosis correct
- A. ■ Histological examination to find larvae
B. ■ X – ray examination to find adult
C. ■ Examination of cystic fluid for brood capsules and eggs protoscolices
12. Terminal spine in the egg of which species defines is in which species
- A. haematobium
B. S. buski
C. S. mekongi
D. S. intercalatum
13. Ingestion egg infected water plant intestinal fluke
- A. F buski
B. F hepatica
C. F gigantica
D. paragonimus westermani
14. Infective stage for the bladder cancer is the of S. haematobium which stage of its life cycle
- A. adult stage
B. ova
C. sporocyst
D. egg inflammatory reaction
15. Relapse of P. ovale and vivax is due to
- A. sporozoite
B. schizont
C. merozoite
D. oocyte
16. Podoconiosis clinical manifestation all are correct except
- A. mostly limited below the knee ankle and leg
B. Caused by immune response to certain minerals
C. Non parasitic elephantiasis
D. is except has no hydrocele is due to immune reaction its one of clinical manifestation

17. The most pathogenic stage of drancunculus medinesnsisi is
 - A. Microfilarie
 - B. Adult form
 - C. Ova
 - D. Sporocyst
18. Most common manifestatuion of elepahntiasusiks
 - A. Hydrocele ,
 - B. Elephantiasis
 - C. Lymphedema
 - D. chyluria
19. Ovulvusi manifestation laeast likely to occure in Eth
 - A. oocorercoma
 - B. oncho nodule
 - C. blidness
 - D. nodule
 - E. dermatitis s
20. Lymphatic filarais which is false
 - A. W. banchrofin is the sole cause
 - B. Are larviparoios
21. Not caused by microfilaria in ovulvulus in onchocerciasis
 - A. nodule
 - B. Onchocercoma
 - C. Blindness
 - D. Dermatitis
 - E. Lymphheadenitis
22. Tx and of loalosais includes all except
 - A. ahigh protein and rest regimen(o.volvulus)
 - B. give antihistamine (corticosteroids)
 - C. pruritic of wrist and panful aedema of hand
23. Not calaber swelling
 - A. non pitting swollen areas of wrists & ankles skin
 - B. painful/may be painless
 - C. Itchy and pruritic of wrist of hand
 - D. sub conjunctival swelling
24. Chikungunya correct statement
 - A. 50 become asysptomatic
 - B. Symptoms usually disappear between the first- and second-week post-infection.
 - C. 1/4th of the cases evolve to a chronic phase, in which debilitating arthralgia persists for months or even years.
25. Which one of the following does not correctly describe pathophysiology of Actinomycosis?
 - A. Anaerobes help to reduce oxygen tension
 - B. Polymicrobial infection enhance invasiveness of Actinomyces species
 - C. Destruction of host tissues provides nutrients

- D. Actinomycosis infections are polymicrobial
 - E. Presence of foreign bodies prevents the spread of infection
26. Actinomyces with polymicrobial infection ,
- A. anaerobes cause low oxygen tension
 - B. the microbes decrease the toxin production
 - C. the microbes do not help to form capsule
 - D. actinomyces breaks the mucosal barrier
27. Nocardia infections are
- A. endogenous infection
 - B. modified acid fast stained
 - C. cause cutaneous and pneumonia (inhalational) infections
28. Brucella are
- A. G -ve non-motile coccobacillus bacteria
 - B. G -ve non-motile coccobacillus fungi
29. Which one of the following is incorrect about Anthrax
- A. Cutaneous anthrax is the less severe form of anthrax
 - B. The bacteria that causes anthrax is aerobic, spore forming gram positive bacteria
 - C. Anthrax is a contagious disease
 - D. Capsule and exotoxins are the two most important virulence factors of B anthracis
 - E. The causative agent, B. anthracis forms non hemolytic colonies on blood agar
30. Wool sorter's infection is synonym for
- A. cutaneous anthrax
 - B. pulmonary anthrax
 - C. GIT anthrax
31. Intermittent increase and decrease of body temperature in brucella
- A. undulant brucellosis
 - B. undulant fever
32. Which one is incorrect Compared with
- A. an individual who is not infected with HIV, a person infected with HIV has a 10 times increased risk of developing TB
 - B. with DM increase 10% each year
33. TB Risk factors and relative probability of being infected Which is
34. correct about tuberculin skin test,
- A. means active virus found on the patient
 - B. means cannot get infected
 - C. read after 12 hours (48-72)
 - D. redness/ erythema is also referred to report additional to size
35. TB incorrect mycobacterium tuberculosis
- A. can't grow without cell
 - B. CO₂ tension enhances growth
 - C. are obligatory aerobic
 - D. has high lipid content cell wall
36. Yellow fever are correct except

- A. hepatocyte infection lead to shock
 - B. immunoglobulin vaccine give life long immunity Neutralizing antibodies endure for life and provide complete protection from disease
 - C. Mortality from severe stage (hepatorenal failure)
 - D. is There is no antiviral drug therapy
 - E. threat of a yellow fever outbreak exists wherever A. aegypti is present.
37. Dengue virus correct
- A. one antigenic type
 - B. pathogenesis is due to extracellular permeability
38. Which one of the following is not correct about Tularemia?
- A. Rabbit is a reservoir for the bacteria
 - B. Causative agent has low infectious dose
 - C. Causative agent may be exploited for bioterrorism
 - D. Causative agent can be killed by macrophages
 - E. Causative agent is highly virulent
39. Tularemia are correct except
- A. killed by macrophages
 - B. causative are bioterrorist
 - C. have slow infection dose
 - D. high virulence
40. Rabies virus bot correct
- A. NTV is Safe , (cause post-vaccine encephalitis)
 - B. moves by retrograde to CNS
 - C. after CNS saliva infection
 - D. after manifestation invariably death follows
 - E. death one of its salient clinical feature is hydrophobia
41. Saprophytic form of coccidioidomycosis
- A. arthroconidia
 - B. spherule
 - C. tuberculate chlamydospore
42. Which of the following is the saprophytic form of coccidioidomycosis?
- A. Arthroconidia
 - B. Coccidioidin
 - C. Spherule
 - D. Macroconidia
 - E. Hyphae
43. Blastomyces dermatitidis is characterized by
- A. Buds attached to the parent cell by narrow base
 - B. Forms hyphae and conidia at 37°C
 - C. Yeast cells attached to the parent cell by a broad base
 - D. has multiply budding yeast cells
 - E. Forms spherules in tissue or at 37°C
44. Blastomyces dermatitidis
- A. are broad based budding

- B. multiple based budding
45. Normal flora fungi causes disseminated diseases in immunocompromised
- A. Aspergillosis
 - B. cryptococcosis
 - C. candida
 - D. mucormycosis
46. Does not contribute to emerging and re-emerging diseases
- A. mostly are DNA viruses
 - B. bats /animals are most
 - C. are deadly mostly affect immunocompromised
 - D. no herd immunity
47. Which one of the following is not correct about emerging diseases?
- A. Majority are due to DNA viruses
 - B. Genomic mutation of the etiology contributes for emergence
 - C. Immunocompromised host contributes for emergence
 - D. Tending to be more deadly to humans
 - E. Majority cross from animals to humans
48. All are true about SARS-CoV-2 except
- A. feco-oral transmission are established
 - B. asymptomatic are infectious
 - C. contact of mucous by hand is infection
 - D. no aerosol transmission documented in community
 - E. droplet infection is the major MOT
49. Pick the incorrect statement concerning SARS-CoV-2's mode of transmission?
- A. Hand- to-mucous contact transmission is well established
 - B. Droplet transmission is the most common way
 - C. Asymptomatic cases can transmit the virus
 - D. Fecal-oral transmission is established
 - E. Airborne transmission is not well established in the community
50. Meds treatment is
- A. worm removal
51. Used during WW1 as biological warfare bioterrorism
- A. Burkholderia mallei
 - B. Francisella tularensis
52. Which one of the following organism was used as a biological weapon during World War I?
- A. Lassa virus
 - B. Hantavirus
 - C. Burkholderia mallei
 - D. Francisella tularensis
 - E. Pasteurella canis