

30. A nurse working in intensive care unit inserted nasogastric tube for parental feedings for 16 years old male patient. The nurse started the procedure by collecting necessary equipment.

What is the most likely final action of this procedure?

(A) Confirm tube size

(B) Insure tube patency

(C) Checking bowel sound

(D) Secure the tube in place

31. A 50-year-old male patient was admitted to surgical unit after surgical management of large bowel obstruction with colostomy. The patient was physiologically stable during admission.

What is the priority nursing intervention that should be done immediately in the above scenario?

(A) Check the surgical dressing

(B) Monitor vital signs

(C) Monitor input and output

(D) Start colostomy irrigation

32. A 53-year-old female patient was admitted to the medical ward with medical diagnosis of chronic liver disease. A nurse took admission assessment and identified 'Fatigue related to decreased energy and the disease process'.

What is the most likely outcome criteria the nurse should set to this patient?

(A) Decide her nutritional requirements

(B) Report strategies to preserve power

(C) Perform activities of daily living

(D) Report increased level of comfort

33. A 37-year-old female patient was admitted to orthopedic unit due to hip fracture. The nurse in charge identified 'Impaired physical mobility related to fractured hip' a nurse place pillow between legs while doing patient positioning.

What is the most likely expected outcome after pillow placement?

(A) Actively participates in exercise regimen

(B) Uses ambulatory aids correctly and safely

(C) Participates in progressive ambulation program

(D) Cooperate in position changes and increased independence

34. A 30-year-old male patient presented to emergency orthopedic unit with complaint of severe pain, tenderness, edema & abnormal joint movement.

What is the most likely diagnosis for this patient?

(A) Fourth degree sprain

(B) Third degree sprain

(C) Second degree sprain

(D) First degree sprain

35. A 35-year-old male client was admitted to orthopedics emergency unit with soft tissue injury at right upper extremity.

What is the initial nursing intervention immediately after admission?

(A) Promote rest

(B) Apply cold packs

(C) Encourage elevation

(D) Apply elastic bandage

71. A nurse working at post anesthesia care unit recognized tachycardia immediately after surgery. The nurse recognized it as the earliest indicator of hemorrhage and shock.

What is the primary responsibility of the nurse?

- (A) Administer anti pain and continue as usual
- (B) Re-assess the patient and identify the cause
- (C) Check vital sign and report to the physician
- (D) Document findings and wait for 30 minutes

72. A nurse working at central sterilization department received surgical equipments immediately after surgery. She was requested to prepare the equipment for the next case.

What is the initial activity to be done by the nurse?

- (A) Decontamination
- (B) Cleaning
- (C) High level disinfection
- (D) Sterilization

73. An emergency patient admitted with chief complaint of sharp chest pain and Thoracentesis is performed to obtain a specimen of pleural fluid for diagnosis of TB. The lab result shows drainage purulent, Glucose equal to serum levels, increased white blood cells and increased RBC.

Which lab result best predict presence of induced pleural effusion of the above case?

- (A) Drainage appearance
- (B) Glucose level
- (C) RBC level
- (D) WBC level

74. Rescue personnel brought a 19-year-old female patient to the emergency department after a motorcycle crash. She has facial contusions and lacerations, a fractured sternum, three fractured ribs, a hemothorax, a dislocated hip, a fractured pelvis, and multiple minor lacerations.

What additional data you need as priority assessment criteria?

- (A) Airway and breathing
- (B) Consciousness level
- (C) Body fluid status
- (D) Vital sign

75. A 39-year-old male patient was admitted due to motorcycle crash. He has facial contusions and lacerations, a fractured sternum, three fractured ribs, and hemothorax. Initial findings include sharp pain over chest, an unobstructed airway with absent breath sounds in the right basilar lung field, tachypnea with 32 shallow respirations per minute, BP is 94/70 mmHg; skin is cool, bloody and clammy; heart rate is 110 bpm and peripheral pulses are intact.

What appropriate nursing intervention should be done first?

- (A) Stop hemorrhage
- (B) Ventilation support
- (C) Initiate IV resuscitation
- (D) Provide morphine sulphate

71. A nurse working at post anesthesia care unit recognized tachycardia immediately after surgery. The nurse recognized it as the earliest indicator of hemorrhage and shock.

What is the primary responsibility of the nurse?

- (A) Administer anti pain and continue as usual
- (B) Re-assess the patient and identify the cause
- (C) Check vital sign and report to the physician
- (D) Document findings and wait for 30 minutes

72. A nurse working at central sterilization department received surgical equipments immediately after surgery. She was requested to prepare the equipment for the next case.

What is the initial activity to be done by the nurse?

- (A) Decontamination
- (B) Cleaning
- (C) High level disinfection
- (D) Sterilization

73. An emergency patient admitted with chief complaint of sharp chest pain and Thoracentesis is performed to obtain a specimen of pleural fluid for diagnosis of TB. The lab result shows drainage purulent, Glucose equal to serum levels, increased white blood cells and increased RBC.

Which lab result best predict presence of induced pleural effusion of the above case?

- (A) Drainage appearance
- (B) Glucose level
- (C) RBC level
- (D) WBC level

74. Rescue personnel brought a 19-year-old female patient to the emergency department after a motorcycle crash. She has facial contusions and lacerations, a fractured sternum, three fractured ribs, a hemothorax, a dislocated hip, a fractured pelvis, and multiple minor lacerations.

What additional data you need as priority assessment criteria?

- (A) Airway and breathing
- (B) Consciousness level
- (C) Body fluid status
- (D) Vital sign

75. A 39-year-old male patient was admitted due to motorcycle crash. He has facial contusions and lacerations, a fractured sternum, three fractured ribs, and hemothorax. Initial findings include sharp pain over chest, an unobstructed airway with absent breath sounds in the right basilar lung field, tachypnea with 32 shallow respirations per minute, BP is 94/70 mmHg; skin is cool, bloody and clammy; heart rate is 110 bpm and peripheral pulses are intact.

What appropriate nursing intervention should be done first?

- (A) Stop hemorrhage
- (B) Ventilation support
- (C) Initiate IV resuscitation
- (D) Provide morphine sulphate

66. Newly diagnosed adult type II Diabetes mellitus patient was admitted to a medical ward. The nurse formulated a nursing diagnosis 'knowledge deficit related to medication administration' and the nurse's goal was 'patient will safely administer all ordered medications before discharge'.

Which outcome criteria need improvement from the nurse's plan?

- (A) Verbalize understanding of desired and adverse effects of medications.
- (B) Be able to monitor blood glucose levels to determine medication.
- (C) State signs, symptoms, and treatment of hypoglycemia.
- (D) Request insulin 30 minutes after taking meal on daily basis.

67. You are attending examination of a patient with peritonitis using drainage tube which is done by physician. The drainage is characterized by pale yellow, watery, and fluid from a blister.

Which wound drainage is described in the above case?

- (A) Serous
- (B) Purulent
- (C) Sanguinous
- (D) Serosanguineous

68. After road traffic accident a patient has above knee cast. During assessment of distal extremity for neuro-vascular status, the result shows minimal swelling, discomfort and color change, prompt capillary refill response and decreased sensations.

Which sign best indicates poor prognostic factor?

- (A) Minimal Swelling
- (B) Minimal discomfort
- (C) Minimal sensations
- (D) Minimal color change

69. A 29-year-old male patient was scheduled for emergency surgery for management of acute abdomen secondary to appendicitis. A pre operative nurse needs to prepare the patient for surgery.

What psychological component should be assessed first?

- (A) Level of understanding for pre & postoperative instructions
- (B) Coping strategies for common health problems
- (C) Emotional state related to current diagnosis
- (D) Roles and responsibilities at home

70. A 70-year-old male patient was admitted to surgical unit with diagnosis of benign prostatic hyperplasia. He was scheduled for surgery. The patient reported to the nurse that he didn't clearly understand the overall care outcome to sign informed consent.

Who is responsible to solve this problem at preoperative period?

- (A) The nurse
- (B) The surgeon
- (C) The assistant surgeon
- (D) The anesthetist

56. A 30-years-old electrician with major electrical burn injury was admitted to a hospital. The burn injury has affected the anterior trunk. The nurse in charge devoted to keep safety of patient and promote healing of injury in short time.

What action is less important in maximizing safety and healing?

- (A) Apply baby powder between lower limb
- (B) Apply sand bag between lower limb
- (C) Apply bed cradle to protect wound
- (D) Apply cotton ring over occipute

57. A 22-year-old male trauma patient was admitted to the emergency department. His vital sign is Temperature cold to touch; Respiratory rate 12/min, Pulse rate 130/min; Blood pressure 70/50 mmHg. On physical examination, an open wound with active bleeding and lethargic in appearance was observed.

What reason best describes cold body of this patient?

- (A) Decreased peripheral blood circulation
- (B) Activation of anterior hypothalamus

- (C) Increased Peripheral vasodilatation
- (D) Exposing patient unnecessarily

58. A menopause woman with medical diagnosis of hyperthyroidism was admitted to medical ward. She is agitated and diaphoretic with some neural symptoms. Vital sign: Temperature = 38°C , Respiratory rate = 19/min, Pulse rate = 105/min and Blood pressure = 110/70 mmHg.

What do you interpret from this particular client?

- (A) Euthermia; tachycardia; Tachypnea; hypotension
- (B) Hypothermia; tachycardia; eupnea; hypertension
- (C) Hyperthermia; tachycardia; eupnea; normo-tension
- (D) Hyperthermia; tachycardia; tachypnea; normo-tension

Eupnea

59. A Patient with medical diagnosis of Polycythemia in which hematocrit is increased was admitted to medical ward. Allopurinol is given to prevent gouty attacks. Vital sign: Temperature = 37.2°C , Pulse rate bounding = 76/min, Respiratory rate = 19/min and Blood pressure 130/100 mmHg.

Which factor most likely influences value of this blood pressure measurement?

- (A) Medication
- (B) Blood viscosity
- (C) Peripheral resistance
- (D) Circulating blood volume

60. Upon assessing patient's IV site before administering medication, the nurse noted that IV site has been swollen, cold and patient complaining of pain.

Which one is the priority nursing measure?

- (A) Irrigate the line with distilled water
- (B) Increase the height of IV stand
- (C) Maintain IV on extremity
- (D) Stop IV fluid

61. A nurse working in surgical ward collecting equipment for cleaning septic wound. The nurse needs to use first surgical forceps for wound dressing.

What is the most likely purpose of the forceps for the above scenario?

- (A) To remove inner cover of wound
- (B) To clean the wound before dressing
- (C) To dry cleaned wound before dressing
- (D) To apply new dressing for final dressing

62. On home visiting service, you identified 75-year-old urine incontinent, dehydrated and neglected woman with bed sore over buttock. The wound is characterized by serous drainage which extends to subcutaneous tissue with no sign of infection over it.

What is the immediate risk factor for pressure ulcer of the above woman?

- (A) Age
- (B) Dehydration
- (C) Urine Incontinence
- (D) Serous drainage from wound

63. You are providing enema pre-operatively for client before bowel surgery. Client vital sign: Temperature 37.2°C , Pulse rate bounding 79/min, Respiratory rate 18/min, Blood pressure 140/89mmHg. You selected warm tap water for your procedure.

What is the mechanism of action of this enema fluid?

- (A) Isotonic with simple expansion of colon
- (B) Irritation to the colon to cause sense of defecation
- (C) Hypotonic with simple expansion of colon
- (D) Hypertonic with absorption of fluid from colon

64. In routine morning care for client with colostomy, you observed that content coming from colostomy opening was fluid. The patient reported that he always has similar appearance of discharge.

What is the most likely type of colostomy in this patient?

- (A) Sigmoid colostomy
- (B) Ascending colostomy
- (C) Transverse colostomy
- (D) Descending colostomy

65. A 75-year-old male patient with type II diabetes mellitus was counseled for home management. He reports that, he is physically active and engages in routine physical exercise. The nurse identified that client has decreased vision, decreased peripheral sensation and dribbling of urine.

What is the most priority nursing diagnosis of the patient?

- (A) Risk for pressure ulcer related to decreased peripheral sensation
- (B) Risk for constipation related to active physical exercise
- (C) Risk for infection related to dribbling of urine
- (D) Risk for injury related to decreased vision

95. A 32-year-old rabies infected woman was admitted to intensive care unit with loss of consciousness, generalized spasm, wound and convulsions. Vital signs finding shows that:

- Blood pressure 90/50 mmHg
- Pulse rate 45/min
- Respiratory rate 10/min
- Temperature: 35.7°C

What is the first line management of this patient?

- (A) Giving Human rabies immune globulin
- (B) Giving wound care
- (C) Sedating the patient with barbiturates
- (D) Supporting the airway

96. A researcher wants to study the prevalence of stunting and associated factors among school children's in a certain district.

What is the most appropriate anthropometric measurement to be used by the researcher?

- (A) Weight for age
- (B) Height for age
- (C) BMI with age
- (D) Weight for height

97. A mother brought a child to pediatric OPD with complaint of cough for one day duration. On examination, the child can walk with hands held, understand simple commands like wave bye-bye, cruise round furniture, casting of objects and have fluctuating moods.

What is the actual age of this child?

- (A) 9 month
- (B) 12 month
- (C) 15 month
- (D) 18 month

98. A two year-old child was brought to under-five OPD with complaint of no day time sleepiness for five days. The mother reported that the child was agitated if awakened, displaced from bed and has sleep walking.

What is the best management for this child?

- (A) Administer PO anxiety medications
- (B) Parental education and reassurance
- (C) Provide sleep medication at night time
- (D) Investigate further for anxiety disorders

99. A three year-old child was brought to the health center with a complaint of oedema of both feet. On physical examination, the nurse noticed that the child had grade-II oedema of both feet. On appetite test, the baby passed the test.

What is the most likely diagnosis of this infant?

- (A) Moderate acute malnutrition
- (B) Complicated severe acute malnutrition
- (C) Uncomplicated severe acute malnutrition
- (D) Complicated moderate acute malnutrition

90. A 26-year-old man with multiple sexual partners visited the health center by complaining burning sensation on urination and acute onset of profused and purulent urethral discharge. Testicles are tender and swollen.

What is the possible diagnosis of this patient?

- (A) Candidiasis
- (B) Gonorrhea
- (C) Herpes genitalis
- (D) Trichomoniasis

91. A woman came to the hospital with white, thick and curd like discharge from vagina. She has also burning sensation during urination.

What is the most likely diagnosis of this patient?

- (A) Candidiasis
- (B) Gonorrhea
- (C) Trichomoniasis
- (D) Urethritis

92. A 32-year-old man came to clinic with complaint of urethral discharge and pain during urination. The nurse confirmed presence of purulent discharge by milking the urethra.

What is the priority nursing activity for the above case?

- (A) Testing him for HIV
- (B) Educate him to use condom
- (C) Call, notify and manage his partner
- (D) Tell him to abstain from sex until cured

93. In one governmental farm, anthrax killed more than three cows. The local and national public health authorities planned to prevent the spread of the disease.

Which strategy is vital to prevent the transmission?

- (A) Mass vaccination of animals in the area
- (B) Bury animals believed to have died of anthrax
- (C) Use occupational health attire before any contact
- (D) Isolate and control exposed zone and individuals

94. A woman was bitten by rabid bid Fox while gathering sticks and woods in the jungle. After three weeks, she started to develop headache, fever, nausea and a tingling sensation at the site of the bite. Based on history and manifestations, a nurse suspected rabies.

Which clinical phase of rabies is most likely seen on this patient?

- (A) Excitatory phase
- (B) Neurologic phase
- (C) Paralytic phase
- (D) Prodromal phase

66. Newly diagnosed adult type II Diabetes mellitus patient was admitted to a medical ward. The nurse formulated a nursing diagnosis 'knowledge deficit related to medication administration' and the nurse's goal was 'patient will safely administer all ordered medications before discharge'.

Which outcome criteria need improvement from the nurse's plan?

- (A) Verbalize understanding of desired and adverse effects of medications.
- (B) Be able to monitor blood glucose levels to determine medication.
- (C) State signs, symptoms, and treatment of hypoglycemia.
- (D) Request insulin 30 minutes after taking meal on daily basis.

67. You are attending examination of a patient with peritonitis using drainage tube which is done by physician. The drainage is characterized by pale yellow, watery, and fluid from a blister.

Which wound drainage is described in the above case?

- (A) Serous
- (B) Purulent
- (C) Sanguinous
- (D) Serosanguineous

68. After road traffic accident a patient has above knee cast. During assessment of distal extremity for neuro-vascular status, the result shows minimal swelling, discomfort and color change, prompt capillary refill response and decreased sensations.

Which sign best indicates poor prognostic factor?

- (A) Minimal Swelling
- (B) Minimal discomfort
- (C) Minimal sensations
- (D) Minimal color change

69. A 29-year-old male patient was scheduled for emergency surgery for management of acute abdomen secondary to appendicitis. A pre operative nurse needs to prepare the patient for surgery.

What psychological component should be assessed first?

- (A) Level of understanding for pre & postoperative instructions
- (B) Coping strategies for common health problems
- (C) Emotional state related to current diagnosis
- (D) Roles and responsibilities at home

70. A 70-year-old male patient was admitted to surgical unit with diagnosis of benign prostatic hyperplasia. He was scheduled for surgery. The patient reported to the nurse that he didn't clearly understand the overall care outcome to sign informed consent.

Who is responsible to solve this problem at preoperative period?

- (A) The nurse
- (B) The surgeon
- (C) The assistant surgeon
- (D) The anesthetist

85. A 34-year-old man was diagnosed with paucibacillary leprosy. He did not have any information about the disease. He requested the nurse to tell him about the manifestations of the disease.

What is the best explanation to be given by the nurse to this patient?

- (A) One to five skin lesions and one nerve trunk enlarged
- (B) Decreased skin sensation and 6-12 Painless lumps
- (C) More than 10 skin lesions and two nerves affected
- (D) Six or more skin lesions and >1 nerve affected

86. A 42-year-old immune suppressed patient with uncontrolled diabetes mellitus is admitted to a medical ward with aseptic meningitis.

What are the first line combined drugs should be prescribed by the physician to this patient?

- (A) Benzyl penicillin, Chloramphenicol, Ampicillin, Meropenem
- (B) Ceftriaxone, Vancomycin, Ampicillin, Gentamycin
- (C) Ceftazidime, Vancomycin, Dexamethasone, Ampicillin
- (D) Ceftazidime, Acyclovir, Ampicillin, Vancomycin

87. A 34-year-old man with fever, marked weight loss and splenomegally was admitted to medical ward. One month ago, he had travel history to the warm part of the country where sand flies are prevalent.

What is the most likely diagnosis of this patient?

- (A) Babesiosis
- (B) Leishmaniasis

- (C) Onchocerciasis
- (D) Trypanosomiasis

88. A 32-year-old patient came to OPD with painless ulcer on the tips of the nose and submandibular lymph node enlargement. The lesion persisted for more than six months. Lab finding showed Leshmania Aetiopica.

What is the first line drug treatment for this patient?

- (A) Miltefosine
- (B) Paromomycin

- (C) Meglumin Antimoniate
- (D) Sodium Stibogluconate

89. The nurse plans to work on prevention of relapsing fever and typhus in the nearby prison.

Which preventive action is the priority?

- (A) Changing clothes and bedding frequently
- (B) Drying clothes and bedding in sunlight
- (C) Treating louse-infested clothes with chemicals
- (D) Washing body and clothes regularly

51. A 50-year-old diabetic patient with uncontrolled glycemia was brought to the Emergency Department with sudden loss of vision after 12 years of disease onset. The nurse in charge suspected a complication.

What is the most likely complication expected for the above scenario?

- (A) Diabetic ketoacidosis
- (B) Diabetic nephropathy
- (C) Diabetic neuropathy
- (D) Diabetic retinopathy

52. A client who spent one week in the ward refused to stay and wanted to be discharge by his own decision. The nurse in charge tried to explain about the process of care to keep patient on treatment. Unfortunately, the patient escaped without fulfilling formal procedure from the ward.

What nursing action takes priority in the above scenario?

- (A) Record the fact on the form and have another health professional witness
- (B) Search for the clients care giver around hospital to sign the form
- (C) Notify the physician, nurse in charge, and agency administration
- (D) Call the police officer of the hospital and city administration

53. After 10-days of inpatient care, a client is going to be discharged today. The nurse obtained authorization for discharge from the physician in charge.

What should be the nurse's next plan of action for this case?

- (A) Discharge client
- (B) Write discharge summary
- (C) Notify business personnel
- (D) Provide discharge instruction

54. A 65-year-old client with laboratory result of low white blood count was admitted to medical ward. The nurse in charge worn mask, washed hand on entry to room and kept client in private closed room with specially monitored room pressure. The nurse advised the client to wear mask during transportation from one room to another.

What principle of standard precaution method is applied in the above scenario?

- (A) Air born precautions
- (B) Contact precautions
- (C) Droplet precautions
- (D) Standard precautions

55. A nurse was performing wound dressing of infected wound in emergency ward and needs to decontaminate all the medical instruments used.

What is the duration of decontamination the nurse should wait?

- (A) 5 minutes
- (B) 8 minutes
- (C) 10 minutes
- (D) 12 minutes

80. A nurse was accused of medication error committed during the care of an admitted patient. The nurse was punished for giving inappropriate drug to the patient without following medication administration guidelines.

What is the most appropriate initial action the nurse should take to prevent this error?

- (A) Checking right medication label
- (B) Checking the right client
- (C) Checking the right dose
- (D) Checking right route

81. A male prisoner patient came to a clinic with the complaint of abrupt onset of fever, malaise, anorexia, and abdominal discomfort. He has visible jaundice. He strongly criticizes the prison site for the poor hygiene of the drinking water. The lab finding showed that his IgM has abnormally increased.

What is the possible diagnosis for this client?

- (A) Hepatitis A
- (B) Hepatitis B
- (C) Hepatitis C
- (D) Hepatitis D

82. A man is washing clothes of cholera patient at stream water. He is washing without using any protective equipment. A nurse planned to train on the measures of controlling the transmission of the disease.

What is the priority control measure the nurse should take?

- (A) Disinfect the contaminated clothes
- (B) Avoid contamination of drinking water
- (C) Boil drinking water from the stream source
- (D) Isolate the victim and wash hands frequently

83. A woman brought her four-year-old boy to the clinic because of long round worms which measure 19 cm long coming out of the child's anus with stool.

What is the best treatment option for this boy?

- (A) Albendazole 400 mg PO stat
- (B) Albendazole 800 mg PO stat
- (C) Mebendazole oral suspension 500 mg/5ml
- (D) Mebendazole oral suspension 250 mg/2.5 ml

84. A 36-year-old HIV positive woman is living with her husband who was diagnosed with smear positive pulmonary TB one week ago. She is asymptomatic for TB. The nurse at the health center traced her as household contact.

What should be the first action of the nurse to prevent TB transmission?

- (A) Give her isoniazid preventive therapy
- (B) Conduct appropriate clinical evaluation
- (C) Start first line anti-TB treatment regimen
- (D) Go home and screen other family members