

1. A patient has been receiving high-dose corticosteroids and broad-spectrum antibiotics for treatment of traumatic injury and secondary infection. The nurse plans care knowing that the patient is most susceptible to

- a. Aspergillosis
- b. Histoplasmosis
- C. Coccidioidomycosis
- d. Candidiasis

2. Which of the following measures is used to assess the strength of association between exposure and disease in a case-control study?

- a. Attributable risk
- b. Odds ratio
- c. Relative risk
- D. Hazard ratio

3. During assessment of a patient the nurse notes an area of red, sharply defined plaques covered with silvery scales that are mildly itchy on the patient's knees and elbows. The nurse recognizes this finding as

- a. Actinic keratosis
- b. Psoriasis
- C Vertigo
- d. Seborrheic keratosis

4. A 35 year old mother brought a 4 month baby to under 5 OPD where you are working. When you evaluate the baby the baby is not well attached to breast & not suckling effectively, has more than 18 breastfeeds in 24 hours and didn't start other foods or drinks. Which of the following classification best describe this condition?

- a. Sever developmental delay
- b. No feeding problem and not underweight
- C No developmental delay
- d. Feeding problem or underweight

5. When planning care for the infant diagnosed with cleft lip and palate, which action would the nurse take in relation to the priority nursing diagnosis for this child?

- a. Temporarily refrain from having the baby breastfeed
 - b. Encourage mother to use false palate covering when feeding baby
 - c. Prevent the baby from vigorously crying
 - d. Burp the baby well throughout feedings
6. Which of the following statements about metabolic side effects of ART is false?
- a. Changes in body shape and size are often difficult for patients to accept
 - b. These are an annoying set of symptoms that are ultimately harmless
 - c. Lipid abnormalities include increases of triglycerides and decreases in high-density cholesterol
 - d. Insulin resistance and hyperlipidemia can be treated with drugs to control glucose and cholesterol
7. Which of the following terms best describes the synthesis of major features of decreasing risk of transmission of blood borne pathogens and body substance isolation?
- a. Universal Precautions
 - b. Protective Isolation
 - c. Medical Asepsis
 - d. Standard Precautions
8. Which one of the following is priority nursing intervention for HIV exposed neonate?
- a. Assess feeding and counsel
 - b. Link only neonate to ART clinic and start ART
 - c. Give Co-trimoxazole from 5 weeks
 - d. Give Co-trimoxazole from 4 weeks
9. Which of the following is a type of study design that compares the occurrence of a disease or condition in two or more groups?
- a. Case series
 - b. Cohort study
 - c. Cross-sectional study
 - d. Case-control study

10. One of your clients with mania begins dancing in the ward. When she twisted her skirt in front of the male clients, it was obvious she had no underwear on. You as a nurse distract her and take her to her room to put on underwear. Which of the following best describe your action?

- a. Avoid embarrassing the male clients who are watching.
- b. Minimize the client's embarrassment about her present behavior.
- c. Teach her about proper attire and hygiene.
- d. Keep her from dancing with other client

11. The nurse in X hospital is caring for a client who has been taking fluphenazine for 3 days. The client suddenly cries out, his neck twists to one side, and his eyes appear to roll back in the sockets. The nurse finds the following PRN medications ordered for the client. Which one should the nurse administer?

- a. Fluphenazine 2 mg PO tid PRN
- b. Haloperidol 5 mg IM PRN
- c. Benztropine 2 mg PO bid PRN
- d. Diphenhydramine, 25 mg IM, PRN

12. How can nurses incorporate creativity in promoting breastfeeding success?

- a. Providing formula supplementation without exploring other options
- b. Discouraging mothers from seeking lactation support or consulting professionals
- c. Restricting breastfeeding practices to a strict schedule and regimen
- d. Encouraging the use of different breastfeeding positions and techniques

13. A nurse is admitting a patient with congestive heart failure. Which of the following describe appropriate aspects of assessment?

- a. Apply oxygen therapy via nasal cannula.
- b. Appraise current therapeutic regimen at home.
- c. Define patient goals for increasing ability for self-care.
- d. Identify activities that exacerbate symptoms

14. The nurse demonstrates to a patient scheduled for a total hip replacement that it is important after surgery to avoid:

- a. Sleeping on the abdomen

b. Bearing weight on the affected leg after 4 to 6 weeks

C Sitting with the legs crossed

d. Abduction exercises of the affected leg

15. Which term would the nurse use to document wound drainage that is thick, odorous, and green?

a. Serosanguineous

b. Serous

c. Sanguineous

d. Purulent

16. A five year female child brought to Health center from low malaria risk area with fever, convulsion and lethargic.

Which of the following is the best category of this child?

a. severe complicated malaria

b. Fever: no malaria

C very severe febrile disease

d. Malaria

17. Which one of the following is the most important for the nurse to assess the client's with anxiety beginning treatment with lorazepam?

a. Use of coping mechanisms.

b. Motivation for treatment.

C. Family and social support.

d. Use of alcohol.

18. Which shistosoma species known to cause Urinary Tract Infection that leads to blood in the urine?

a. Shistosoma Mansoni

b. Shistosoma Intercalatum

c. Shistosoma Japonicum

d. Shistosoma Heamatobium

19. Which one of the following best defined as an exercise in which the students act the parts of the different people and so begin to experience some of the feeling of these people?

a. Demonstration

b. Role play

c Case study

d. Group Discussion

20. A 25 year old male patient with Obsessive Compulsive Disorder is presented with ritualistic hand washing that last several hours each day. The skin on his hands is red and cracked, has minor bleeding. Which one of the following is the nursing goal for this client?

a. Eliminating the hand washing rituals

b. Providing good skin care

C. Decreasing the time spent washing hands

d. Providing milder soap for hand washing

21. Nurse Gemechu is working at under 5 OPD in which he faced new born infant with the following condition Skin and eyes yellow and baby is < 24 hrs old. Which of the following is the best term to describe her symptoms?

a. Severe Jaundice

b. Jaundice

C Mild jaundice

d. No jaundice

22. When providing care for a patient with thrombocytopenia, you should instructs the patient to:

a. Be careful when shaving with a safety razor

b. Dab his or her nose instead of blowing

C Continue with physical activities to stimulate thrombopoiesis

d. Avoid aspirin because it may mask the fever that occurs with thrombocytopenia

23. After surgery, Ms. Hanna is having difficulty voiding. Which nursing action would most likely lead to an increased difficulty with voiding?

a. Pouring warm water over Ms. Hanna's fingers

- b. Having Ms. Hanna ignore the urge to void until her bladder is full
- c. Using a warm bedpan when Ms. Hanna feels the urge to void
- d. Stroking Ms. Hanna's leg or thigh

24. Which of the following describes an odds ratio of 1?

- a. The exposure is not associated with the outcome
- b. The exposure is associated with an increased risk of the outcome
- c. The exposure is associated with a decreased risk of the outcome
- d. There is no association between the exposure and the outcome

25. Captopril may be administered to a client with HF because it acts as a:

- a. Vasopressor
- c. Potassium-sparing diuretics
- b. Vasodilator
- d. Volume expander

26. A 28 years mother has a 4 weeks baby on exclusive breast feeding. The mother is reactive to HIV antibody test. The baby also tested for HIV status and the result is Positive antibody test, but DNA PCR is negative. Which of the following is the best classification of this neonate?

- a. HIV infected
- b. HIV infection unlikely
- c. HIV exposed
- d. HIV status unknown

26. Mr. Tadese is a head nurse at a health center. His major focus as head is building and maintaining relationships among staffs who are working under his unit. Which one of the following best described the role of Mr. Tadese?

- a. individual roles
- b. Social-emotional roles
- c. Task roles
- d. Procedural roles

27. In planning nursing interventions to increase bladder control in the patient with urinary incontinence, the nurse includes:

- a. Clamping and releasing a catheter to increase bladder tone
- b. Teaching the patient biofeedback mechanisms to suppress the urge to void
- C. Teaching the patient to use Kegel exercises
- d. Counseling the patient concerning choice of incontinence containment device

28. When explaining the action of a hypertonic solution enema, the nurse incorporates which of the following as the basis for action?

- a. Bowel mucosa irritation
- b. Osmosis of water into colon
- C. Softening of fecal contents
- d. Diffusion of water out of colon

29. Mr. Abera is receiving continuous tube feedings through a small-bore nasogastric tube. Which method is the most accurate in verifying correct tube placement?

- a. Measurement of the gastric aspirate PH
- b. Measurement of the amount of residual
- C. Radiographic examination
- d. Auscultatory method

30. A two days male newborn comes to X hospital by chief complain of cough. He has excessive secretions coming out of nose and constant drooling of foamy saliva. After evaluation he diagnosed by Oesophageal Atresia. Which of the following is the first priority nursing diagnosis of above newborn case scenario?

- a. Risk for deficient fluid volume.
- b. Risk for Impaired nutrition.
- C. Risk for impaired breathing.
- d. Risk for altered comfort related to disease process.

31. 72-year-old woman is taking several medications on a regular basis. Which of the following categories of drugs would be most likely to increase her surgical risk?

- a. Sedatives
- b. Antacids
- C. Anticoagulants
- d. Laxatives

32. Which one of the following is a teaching method in which 2-4 people discuss a question/problem for a short period of time followed by plenary feedback?

- a. Case study
- b. Demonstration
- C. Buzz groups
- d. Group Discussion

33. Mrs.X is a 28-year-old woman who wants a contraceptive method that provides protection against sexually transmitted infections (STIs in addition to preventing pregnancy. Which of the following methods offers dual protection?

- a. Condom
- b. Intrauterine device (IUD)
- C. Diaphragm
- d. Emergency contraception (morning-after pill)

34. The nurse is teaching an adult patient how to administer antibiotic eardrops. Instructions should include which of the following?

- a. Cool the drops so that they decrease swelling in the canal
- b. Placement of a cotton wick to assist in administering the drops is not recommended
- c. Be careful to avoid touching the tip of the dropper bottle to the ear
- d. Keep the head tilted for 5 to 7 minutes after administering the drops to prevent them from running out of the ear canal

35. Which of the following actions demonstrates the nurse's commitment to providing compassionate care during the postpartum period?

- a. Focusing solely on physical health assessments and ignoring emotional needs
- b. Dismissing the mother's emotional distress as common postpartum blues

C. Showing genuine interest and actively listening to the concerns of the new mother

d. Rushing through postpartum education and discharge planning

36. nurse is assessing respiratory function on a patient with a pneumothorax by asking him to say "99" while palpating the chest. This is a technique used to determine which of the following?

a. Excursion

b. Fremitus

C Friction rub

d. Egophony

37. Which of the following best illustrate the change theory that viewed change as sequential (occurring in a step-by-step fashion ?

a. Havelock Change Model

b. Nonlinear change theories

C Linear change theories

d. Lewin's Force-Field Model of Change

38. patient is hypotensive related to extensive bleeding in surgery. Which type of fluid would be most appropriate to administer to this patient?

a. Hypertonic 3% saline (NaCl

b. Hypotonic D5W + 1½ NaCl

C Colloid total parenteral solution

d. Isotonic lactated Ringer's

39. The nurse is teaching an adult patient how to administer antibiotic eardrops. Instructions should include which of the following?

a. Cool the drops so that they decrease swelling in the canal

b. Placement of a cotton wick to assist in administering the drops is not recommended

C Be careful to avoid touching the tip of the dropper bottle to the ear

d. Keep the head tilted for 5 to 7 minutes after administering the drops to prevent them from running out of the ear canal

40. A patient has recently been diagnosed with early stages of breast cancer. Which of the following is most appropriate intervention for the nurse to focus on?

a. Discussing the patient's past experiences with her grandmother's cancer

b. Maintaining patient's hope

C Preparing a will and advance directives

d. Discussing replacement child care for patient's children

41. patient with a stable, closed traumatic fracture of the humerus has a temporary splint with bulky padding applied with an elastic bandage. The nurse suspects compartment syndrome when the patient experiences:

a. Pain at the fracture site

b. Muscle spasms of the lower arm

C. Pain when the nurse passively extends the fingers

d. Increasing edema of the limb

42. A patient who has fallen and injured his wrist carefully cradles it with the other hand. The patient is demonstrating which of the following responses to pain?

a. Behavioral

b. Affective

C. Physiologic

d. Involuntary

43. Which of the following is an example of a strategic leadership role for a nurse leader?

a. Providing direct patient care.

b. Conducting research studies

C Developing and implementing organizational goals

d. Training new nursing staff

44. A 28-year-old woman who is 6 months pregnant presents to the emergency department with a temperature of 38.2° C (100.8° F) and complains of shaking chills and pain on her right side, which she locates by pointing to the area above her right iliac crest. During the examination she is tender to percussion at the junction of the lower ribs and the thoracic vertebrae. Urinalysis reveals WBC casts. Which of the following is the most likely diagnosis?

- a. Appendicitis
- b. Ectopic pregnancy
- c. Localized cystitis
- d. Pyelonephritis

45. Which one of the following is the diagnostic method of endemic typhus?

- a. Weil Felix agglutination test (serology)
- b. Widal test
- c. sputum smear
- d. blood film

46. A nurse is identifying nursing diagnoses that would be appropriate for an elderly patient with failure to thrive after the loss of a spouse. Which of the following would be most appropriate?

- a. Loneliness
- b. Impaired Sleep Pattern
- c. Impaired Self-Perception
- d. Ineffective Coping

47. Which action demonstrates proper technique for conducting a newborn assessment immediately after delivery?

- a. Performing the assessment while the baby is still wet and unclothed
- b. Delaying the assessment until the baby is clothed and in a warmer
- c. Using a non-sterile stethoscope to auscultate the baby's heart sounds
- d. Focusing solely on physical assessment and omitting the neurologic examination

48. During the postoperative period, the patient with an above-the-knee amputation should be instructed that the residual limb should not be routinely elevated because (2 extra

- a. Unnecessary movement of the extremity can cause wound dehiscence
- b. The flexed position can promote hip flexion contracture
- c. This position reduces the development of phantom limb sensation
- d. This position promotes clot formation at the incision site and thigh

49. An indication of a neurovascular problem noted during assessment of the patient with a fracture is:

- a. Exaggeration of extremity movement
- b. Petechiae on the head and upper thorax
- c. Decreased sensation distal to the fracture site
- d. Purulent drainage at the site of an open fracture

50. A 10-year-old male was recently admitted to hospital with signs of increased intracranial pressure (ICP). He presented with severe headaches, vomiting, and altered levels of consciousness. A CT scan was sent and it revealed a space-occupying lesion. Finally, brain tumor was diagnosed. What is the most appropriate nursing interventions for this patient?

- a. Elevating the head of the bed at 30 degree
- b. Administering antibiotics
- c. Administering antipyretics
- d. Encouraging deep breathing exercises

51. The nurse would increase the comfort of the patient with appendicitis by:

- a. Sitting the patient upright in a chair
- b. Having the patient lie prone
- c. Flexing the patient's right knee
- d. Turning the patient onto his or her left side

52. Which of the following symptoms is most commonly associated with left-sided heart failure?

- a. Hypotension
- b. Hepatic engorgement
- c. Arrhythmias
- d. Crackles

53. Which of the following is an example of a nurse leader demonstrating ethical behavior?

- a. Ignoring ethical guidelines and standards
- b. Advocating for patient rights and autonomy
- c. Engaging in dishonest practices for personal gain

d. Prioritizing their own interests over the well-being of patients

54. The physician has ordered an indwelling catheter inserted in a hospitalized male patient. The nurse is aware of which of the following considerations?

a. The catheter is inserted 2 to 3 cm. into the meatus

b. The male urethra is more vulnerable to injury during insertion

c. Since it uses a closed system, the risk for urinary tract infection is absent

d. In the hospital, a clean technique is used for catheter insertion

55. Nursing care for a patient with an indwelling catheter includes which of the following?

a. Irrigation of the catheter with 30 ml of normal saline solution every 4 hours

b. Disconnecting and reconnecting the drainage system quickly to obtain a urine specimen

c. Encouraging a generous fluid intake if not contraindicated by the patient's condition.

d. Telling the patient that burning and irritation are normal, subsiding within a few days

56. Which of the following diagnostic procedure is most commonly applied to determine the location of myocardial damage?

a. Cardiac enzymes

b. Electrocardiogram

c. Echocardiogram

d. Cardiac catheterization

57. Which of the following is defined as death of a woman while pregnant, or within 42 days of termination of pregnancy, from any cause related to, or aggravated by, the pregnancy or its management, but not from accidental or incidental death?

a. Indirect maternal death

b. Maternal death

c. Direct maternal death

d. Maternal mortality rate

58. Which of the following action demonstrates the nurse's creative problem-solving skills in a postpartum complication scenario?

- a. Seeking input from the healthcare team to explore different solutions
- b. Maintaining a rigid treatment plan without adjustments
- C. Dismissing alternative treatment options without consideration
- d. Ignoring the signs and symptoms of complications

59. patient is on a continuous epoprostenol infusion pump. The alarm goes off indicating an obstruction in the intravenous line downstream. The nurse should:

- a. Monitor for flushing and hypotension due to rebound from no medication and the short half-life of the drug
- b. Auscultate the lungs for pulmonary congestion
- C. Assess the central line immediately for any obstruction or accidental clamping of tubing
- d. Check vital signs and oxygen saturation

60. A researcher who is providing therapy for a person should not develop a close friendship with the client because such dual relationships can compromise the success of the therapy. This problem relates to which General Ethical Principle?

- a. Fidelity and responsibility
- b. Justice
- c. Beneficence and nonmaleficence
- d. Respect for people's rights and dignity

61 In research, what does the acronym "IRB" stand for?

- a. Institutional Review Board
- b. International Review Board
- C. Institute of Research and Biomedicine
- d. Investigative Research Bureau

62. Bladder training interventions in a male patient who has urinary incontinence after a stroke

- a. Keeping a urinal in place at all times
- b. Assisting the patient to stand to void
- C. Limiting fluid intake

d. Catheterizing the patient every 4

63. Mr. Dagnachew presented with ladder fashion fever, rose spot found on chest and abdomen. On physical examination he has small pallor, slightly raised blanching, and bradycardia. CBC finding showed leucopenia with relative lymphocytosis (WBC<4000/mm³. Which of the following is the best term differentiate his sign/symptoms?

a. Typhoid fever

b. Giardiasis

C. Dysentery

d. Typhus

64. A 24-year-old man who is in the hospital for treatment of a severe gram-negative infection subsequently becomes oliguric. He is also having difficulty hearing the hospital staff. Laboratory studies show elevated BUN and creatinine levels. Which of the following is the most likely cause of this patient's symptoms?

a. Chloramphenicol

b. Erythromycin

C. Gentamicin

d. Doxycycline

65. When a nurse assesses a patient for pain and offers a plan to manage the pain, which principle is used to encourage the nurse to monitor the patient's response to the pain?

a. Non-maleficence

b. Fidelity

C. Beneficence

d. Respect for autonomy

66. Mr. Molla has a fecal impaction. The nurse correctly administers an oil-retention enema by doing which of the following?

a. Mixing milk and molasses in equal parts for an enema

b. Administering a large volume of solution (500-1,000 mL)

C. Administering the enema while the patient is sitting on the toilet

d. Instructing the patient to retain the enema for at least 30 minutes

67. A 23-year-old woman who is 36 weeks pregnant presents to delivery ward. She complains of feeling generally unwell. Clinical examination reveals a 28-week size fetus. Her blood pressure was noted to be 120/90mmHg and on urine analysis proteinuria was present. During the clinical examination, she has a seizure. Which of the following is this history most suggestive of?

- a. Epileptic fit
- b. Pre-eclampsia

C. Eclampsia

- d. Hypoglycemia

68. The factor related to cerebral blood flow that most often determines the extent of cerebral damage from a the:

- a. Degree of collateral circulation
- b. Oxygen content of the blood
- c. Amount of cardiac output
- d. Level of carbon dioxide in the blood

69. Mr. Tolosa is the Metron at X hospital in one of the region where conflicts had affected the health of the community at large. Majority of the individuals experience or live with the consequences of the conflict and its resolution. Which one of the following best describe this history?

- a. perceived and felt conflict
- b. conflict aftermath
- C. Latent conflict
- d. Manifest Behaviour

70. Which of the following study designs is most suitable for investigating rare diseases?

- a. Prospective study
- b. Retrospective study
- c. Case-control study**
- d. Cross-sectional study

71. Administering or providing opioids to a dying patient who is experiencing moderate to severe pain:

- a. Will likely hasten the person's death.**

- b. May cause addiction.
- c. Will probably be ineffective.
- d. Is an appropriate nursing action

72. A patient with a comminuted fracture of the femur is to have an open reduction with internal fixation (ORIF) of the fracture. The nurse explains that ORIF is indicated when:

- a. Adequate alignment cannot be obtained by other nonsurgical methods.
- b. Cast would be too large to provide normal mobility.
- c. The patient is able to tolerate long-term immobilization.
- d. The patient cannot tolerate the discomfort of a closed reduction.

73. Which of the following actions demonstrates the nurse's creative approach to providing comfort during labor?

- a. Dismissing the use of alternative therapies for pain relief
- b. Restricting pain relief options based on traditional practices
- c. Limiting the use of position changes and movement during labor
- d. Offering a variety of non-pharmacological pain management techniques

75. A 10-year-old male child falls headfirst from a tree and lands on the ground. He experienced immediate loss of consciousness, vomiting, and have unequal pupil size upon regaining consciousness. What is the most appropriate nursing intervention for this patient?

- a. Keep the head elevated at a 30-degree angle
- b. Performing neurologic assessments.
- c. Stabilize the child's neck and spine.
- d. Administer antiemetic medications

76. Which of the following best describes your attitude towards patients who are non-compliant with their treatment plan?

- a. I try to understand the reasons behind the non-compliance and find alternative solutions.
- b. I become annoyed and judgmental towards these patients.
- c. I feel frustrated and give up on these patients, believing they will never change.

d. I dismiss these patients as uncooperative and give up on trying helping them.

78. Which one of the following is a vaccine preventable infection characterized by bluish-white or grey adherent membrane in the throat and tonsils which cannot wipe off with swab and bull neck?

a. Diphtheria

b. Measles

c. Pertusis

d. Shigellosis

79. Performing the assessment while the baby is still wet and unclothed When providing postpartum care, the nurse should instruct the patient on proper breastfeeding technique, which of the following best describe this technique?

a. Applying heat to the breasts before feeding to promote milk production

b. Taking the baby off the breast abruptly to avoid overfeeding

c. Offering the baby only one breast during each feeding session

d. Ensuring a deep latch with the baby's mouth covering the nipple and areola

80. To control the side effects of corticosteroid therapy, the nurse teaches the patient who is taking corticosteroids to:

a. Increase calcium intake to 1500 mg/day

b. Avoid abrupt position changes because of orthostatic hypotension

c. Obtain immunizations due to high risk of infections

d. Perform glucose monitoring for hypoglycemia

81. The most common response of the stroke patient to the change in body image is:

a. Intellectualization

b. Disassociation

c. Denial

d. Depression

82. A patient is admitted to the hospital in hypertensive crisis. The nurse recognizes that hypertensive urgency differs from hypertensive emergency in that:

- a. Hypertensive emergencies always require intra-arterial BP measurement.
- b. Hypertensive urgency is treated with rest and tranquilizers to lower the BP.

C. Hypertensive emergencies are associated with evidence of target organ damage.

- d. The BP is always higher in a hypertensive emergency.

83. A nurse is explaining the transmission of hepatitis "A" to new nursing students. Which of the following explanations of the chain of infection would be most correct?

- a. Infected human stool vehicle transmission gastrointestinal tract
- b. Contact transmission stool gastrointestinal tractinfected human**
- c. Infected human gastrointestinal tract contact transmission stool
- d. Infected human vector bome transmission blood nonintact skin

83. Which of the following best describes your attitude towards patients who are non-compliant with their treatment plan?

- a. I feel frustrated and give up on these patients, believing they will never change.
- b. I become annoyed and judgmental towards these patients.
- cl try to understand the reasons behind the non-compliance and find alternative solutions.
- d. I dismiss these patients as uncooperative and give up on trying helping them.

84. A one-day newborn referred to X hospital from Y health center due to defect of the abdominal wall that allows the abdominal contents to protrude outside the body and no covering over the bowel or contents. What is diagnosis of this newborn?

- a. Gastroschisis
- b. Omphalocele
- c. Hypertrophic Pylonc Stenosis
- d. Oesophagal Atresia

85.