


MCQs in




ORTHODONTICS



K Vijayalakshmi

Foreword

N Sendamaraai



JAYPEE

Published by

Jitendar P Vij

Jaypee Brothers Medical Publishers (P) Ltd

EMCA House, 23/23B Ansari Road, Daryaganj

New Delhi 110 002, India

Phones: +91-11-23272143, +91-11-23272703, +91-11-23282021, +91-11-23245672

Fax: +91-11-23276490, +91-11-23245683 e-mail: jaypee@jaypeebrothers.com

Visit our website: www.jaypeebrothers.com

Branches

- 202 Batavia Chambers, 8 Kumara Krupa Road, Kumara Park East
Bangalore 560 001, Phones: +91-80-22285971, +91-80-22382956, +91-80-30614073
Tele Fax: +91-80-22281761 e-mail: jaypeebc@bgl.vsnl.net.in
- 282 IIIrd Floor, Khaleel Shirazi Estate, Fountain Plaza
Pantheon Road, **Chennai** 600 008, Phones: +91-44-28262665, +91-44-28269897
Fax: +91-44-28262331 e-mail: jpmedpub@md3.vsnl.net.in
- 4-2-1067/1-3, Ist Floor, Balaji Building, Ramkote
Cross Road, **Hyderabad** 500 095, Phones: +91-40-55610020, +91-40-24758498
Fax: +91-40-24758499 e-mail: jpmedpub@rediffmail.com
- 1A Indian Mirror Street, Wellington Square
Kolkata 700 013, Phone: +91-33-22451926 Fax: +91-33-22456075
e-mail: jpbcal@cal.vsnl.net.in
- 106 Amit Industrial Estate, 61 Dr SS Rao Road, Near MGM Hospital
Parel, **Mumbai** 400 012, Phones: +91-22-24124863, +91-22-24104532, +91-22-30926896
Fax: +91-22-24160828 e-mail: jpmedpub@bom7.vsnl.net.in

MCQs in Orthodontics

© 2006, K Vijayalakshmi

All rights reserved. No part of this publication should be reproduced, stored in a retrieval system, or transmitted in any form or by any means: electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the author and the publisher.

This book has been published in good faith and belief that the material provided by author is original. Every effort is made to ensure accuracy of material, but the publisher, printer and author will not be held responsible for any inadvertent error(s). In case of any dispute, all legal matters to be settled under Delhi jurisdiction only.

First Edition : 2006

ISBN 81-8061-582-0

Typeset at JPBMP typesetting unit

Printed at Sanat Printers

Contents

1. Growth and Development	1
2. Development of Dentition	20
3. Dental Occlusion and Stomatognathic System	26
4. <u>Diagnosis, Classification of Malocclusion, Model and Cephalometric Analysis</u>	<u>29</u>
5. <u>Etiology of Malocclusion</u>	<u>68</u>
6. Biomechanics of Tooth Movement and Anchorage .	81
7. Appliances	
a. Removable Appliance	91
b. Functional Appliance	101
c. <u>Fixed Appliance</u>	<u>105</u>
d. Orthopedic Appliance	114
8. <u>Preventive Orthodontics</u>	<u>117</u>
9. Interceptive Orthodontics	122
10. Surgical Orthodontics	124
11. Retention and Relapse	127
12. <u>Orthodontic Materials</u>	<u>130</u>
13. Treatment Plan, Extraction and Superimposition ..	136
<u>Bibliography</u>	<u>139</u>

5. In mixed dentition, which of the following may be considered as a self-correcting problem with age?
A. Unilateral loss of primary canine
B. Lack of interdental spaces
C. A distal step
D. Open bite
6. Order of eruption of permanent teeth in a maxillary arch is:
A. 6123457 B. 6124357
C. 6124537 D. 43
7. The change observed in the arch length of a child between 3 to 18 years is:
A. 4 mm B. 11 mm
C. 17 mm D. 34 mm
8. Occlusal adjustment is affected by all *except*:
A. Compensating curve
B. Cusp height
C. Incisal guidance
D. Postural position
9. After eruption of teeth, root is completed within:
A. 1½ to 2½ years
B. 2½ to 3½ years
C. 5 to 6 years
D. 5 to 6 months
10. Intercanine width is increased with:
A. Eruption of lateral incisors
B. Eruption of permanent canines
C. Eruption of premolars
D. Eruption of peg laterals
11. The first tooth bud develops when embryo length is:
A. 6 mm B. 10 mm
C. 12 mm D. 20 mm

5 C 6 C 7 A 8 D 9 B 10 B 11 A

22 MCQs in Orthodontics

12. In an ideal occlusion, which of the following is undesirable?
A. Crown angulation
B. Crown inclination
C. Curves of Spee
D. Spacing
13. Dental age is most closely related to:
A. Dental maturation
B. Resorption of roots
C. Diffuse calcification of pulp
D. Chronological age
14. Which of the following is a self-correcting anomaly?
A. Ugly duckling stage B. Deep bite
C. Retrusion D. Protrusion
15. Primate spaces in the primary dentition are seen between:
A. Maxillary central and lateral incisor and mandibular central incisor and canine
B. Maxillary primary first molar and second molar and mandibular canine and first primary molar
C. Maxillary primary lateral incisor and primary canine and mandibular primary canine and first primary molar
D. None of the above
16. Primate spaces are a characteristic of:
A. Most mammalian species
B. Human beings
C. Frogs
D. Cows and men
17. The leeway space is:
A. More in maxilla than in mandible
B. More in mandible than in maxilla
C. Equal in maxilla and mandible
D. None of the above

12 D 13 D 14 A 15 C 16 A 17 B

18. A distal step in the deciduous dentition leads to:
A. A normal relationship of permanent teeth
B. A class II malocclusion in permanent teeth
C. A class III malocclusion in permanent teeth
D. End to end relationship in permanent teeth
19. At what age is a child expected to have 12 erupted primary teeth and 12 erupted permanent teeth?
A. 4.5 years
B. 6.5 years
C. 8.5 years
D. 11.5 years
20. First permanent molars begin to calcify at:
A. 6 months of intrauterine life
B. At birth
C. Before deciduous incisors
D. About 1 year after birth
21. Before eruption, the position of permanent mandibular incisor buds relative to primary incisors is:
A. Superior and facial
B. Superior and lingual
C. Inferior and facial
D. Inferior and lingual
22. Tooth buds generally initiated after birth are:
A. The entire permanent dentition only
B. All permanent teeth and some primary teeth
C. First and second premolars and second and third molars only
D. All of the above
23. First evidence of calcification of 2nd premolar on radiographs is:
A. At birth
B. 3 to 4 months
C. 4 to 5 months
D. 1.5 to 2.5 years
-

18 B 19 C 20 B 21 D 22 C 23 D

24 MCQs in Orthodontics

24. **Apexogenesis is:**
A. Formation of the apical one-third of the root
B. Formation of the apical two-thirds of the root
C. Closure of the apical foramen in a developing tooth
D. Deposition of cellular cementum at the apex of the root
25. **During the transition from deciduous to permanent dentition, the overbite tends to decrease. This is mainly attributed to:**
A. Change in axial inclination of permanent incisors
B. Alveolar bone growth
C. Physiologic bite opening
D. All of the above
26. **Late mesial shift occurs due to:**
A. Presence of Leeway space
B. Primate spaces
C. Presence of unrestored caries on distal surface of teeth
D. In adults due to attrition of proximal surfaces of teeth
27. **Late mesial shift of mandibular permanent first molar is compensated by:**
A. Primate spaces
B. Median diastema
C. Leeway space
D. Wearing of proximal contacts of premolar teeth
28. **The calcification of the primary teeth begins about:**
A. 11th week of IUL
B. 12th week of IUL
C. 14th week of IUL
D. 16th week of IUL
29. **The time of initiation of first permanent molar formation (in utero):**
A. 2nd month
B. 4th month
C. 6th month
D. 7th month

24 C 25 C 26 A 27 C 28 A 29 B

Dental Occlusion and Stomatognathic System

1. The teeth which are most often referred to as the key to normal occlusion are:
 - A. First permanent molars
 - B. First permanent molars and second primary molars
 - C. Maxillary and mandibular second primary molars
 - D. Second permanent molars
2. The six keys to normal occlusion was introduced by:
 - A. Dewey
 - B. Andrews
 - C. Tweed
 - D. Angle
3. The pharyngeal muscle which forms a part of buccinator mechanism is:
 - A. Inferior constrictor
 - B. Middle constrictor
 - C. Superior constrictor
 - D. Palatopharyngeal
4. Leeway space helps in the development of normal occlusion by:
 - A. Allowing mesial drift of molars
 - B. Providing space for eruption of premolars
 - C. Providing space for eruption of permanent second molars
 - D. All of the above

28 MCQs in Orthodontics

- 11. What is Bennett movement?**
A. In the lateral shift of the mandible the articular disk moves toward the side of the working bite
B. A muscle develops its maximum force when all of its fibers are repeatedly stimulated to fire
C. The intensity of the contraction of any fiber is independent of the strength of the exciting stimulus, provided that the stimulus is adequate
D. None of the above
- 12. Who divided the deglutitional cycle into 4 phases?**
A. Fletcher
B. Meyer
C. Benninghoff
D. Enlow
- 13. The instrument most often used for evaluating the activity of orofacial muscles:**
A. Electromyograph
B. Holograph
C. Bronchoscope
D. Tonometer
- 14. Kinesiology is the study of movement resulting from action of:**
A. Muscles
B. Nerve
C. Blood vessel
D. Cells
- 15. Which period is considered as early mixed dentition period:**
A. Between 6 to 9 years
B. Between 7 to 12 years
C. Between 9 to 12 years
D. Between 8 to 12 years

11 A 12 A 13 A 14 A 15 A

Diagnosis, Classification of Malocclusion, Model and Cephalometric Analysis

DIAGNOSIS

1. **Brachycephalic individuals usually have:**
 - A. Narrow dental arches
 - B. Broad dental arches
 - C. Normal dental arches
 - D. B or C
2. **Carpel index denotes:**
 - A. Number of carpal bones
 - B. Skeletal maturity
 - C. Treatment prognosis
 - D. Incidence of malocclusion
3. **The best time to correct anteroposterior discrepancy in skeletal base is:**
 - A. Primary dentition period
 - B. Mixed dentition period
 - C. Permanent dentition period
 - D. Post-permanent dentition period
4. **Skeletal indicators of maturity are determined by:**

A. Chest X-ray	B. Periapical X-ray
C. Hand wrist X-ray	D. Occlusal X-ray
E. Combination of B and C	

1 B 2 B 3 B 4 E

11. The most difficult malocclusion to be corrected by orthodontic treatment is:
A. Class II division 2
B. Class III with horizontal growth pattern
C. Class III with vertical growth pattern
D. Bimaxillary protrusion
12. The ideal proportion of upper facial height to lower facial height is:
A. 45% : 55% B. 55% : 45%
C. 40% : 60% D. 50% : 50%
13. The anterior lower facial height is decreased in:
A. Skeletal open bite
B. Skeletal deep bite
C. Long face syndrome
D. Backward rotating mandible
14. Dental arch form is ultimately determined by the:
A. Facial type
B. Angle classification
C. Facial growth pattern
D. Balance between facial and intraoral musculature
E. Interaction of environmental influence on the genetic pattern
15. Most class III malocclusions are associated with:
A. Skeletal class III pattern
B. Deep bites
C. Open bite
D. Competent lips
16. Overclosure of the mandible occurs when there is extensive loss of posterior teeth. This can lead to:
A. Clicking of temporomandibular joint
B. Destruction of the articular disk
C. Crepitus
D. None of above

33. If there is no physiological spacing in between deciduous incisors, what is your investigation?
A. Intra-oral X-ray periapical view
B. Study model
C. Cephalogram
D. All of the above
34. What is the importance of VTO (Visualized treatment objective) in orthodontics?
A. As a clinical diagnostic aid to select a case for functional appliance
B. As a superimposition technique
C. As a soft tissue analysis
D. None of the above
35. Which VTO is required for functional appliance?
A. +ve VTO
B. -ve VTO
C. Difficult to elicit
D. None of the above
36. Which type of malocclusion is most often associated with fractured maxillary anterior tooth?
A. Class I
B. Class II div 1
C. Class II div 2
D. Class III
37. Apertognathia is:
A. Anterior cross bite
B. Open bite
C. Posterior cross bite
D. Prognathism
38. Difference between dental open bite and skeletal open bite is:
A. In dental, open bite present from canine to canine whereas in skeletal, open bite presents from 1st premolar to 1st premolar
B. Dental open bite can be corrected orthodontically whereas skeletal open bite by functional appliance in young age or by surgical orthodontics
C. Both are same (A and B)
D. None of A and B

36 MCQs in Orthodontics

39. Which is the supplemental diagnostic aid?
A. Study cast
B. Facial photographs
C. Cephalogram
D. Clinical examination
40. Appearance of sesamoid bone denotes:
A. Retardation of pubertal development
B. Onset of puberty
C. Peak growth
D. No significance
41. Initial ossification of the sesamoid and advanced ossification of the hook of the hamate denotes:
A. Peak growth
B. Onset of puberty
C. Decline of growth
D. No significance
42. When hand-wrist radiograph is not available what could be the other alternative to assess the pubertal growth spurt in your daily practice?
A. Periapical intra-oral mandibular radiograph
B. Orthopantomogram
C. Cephalogram
D. Bite wing films
43. Root completion of lower left canine denotes:
A. Decline of pubertal growth
B. Onset of puberty
C. Peak growth
D. Cessation of pubertal growth
44. Which is the indicator of pubertal growth spurt?
A. Sesamoid bone and hook of Hamate
B. Cervical vertebra
C. Left lower canine
D. All of the above

39 C 40 B 41 A 42 A 43 D 44 D

57. **Space closure is rapid following loss of which primary molars:**
A. Maxillary molars; closing within three months
B. Mandibular molars; closing within six months
C. Maxillary molars; closing within six months
D. Mandibular molars; closing within three months
58. **Deep skeletal bite will show which of the following features?**
A. High mandibular plane
B. Low mandibular plane
C. Deep palate
D. All of the above
59. **The two finger method of finding skeletal pattern is proposed by:**
A. Mills
B. Korkaus
C. Jacobson
D. Houston
60. **Mirror technique is advocated to check:**
A. Mouth breathing
B. Nasal airway technique
C. None of the above
D. Combination of 1 and 2
61. **Which type of mirror is used in mirror technique to elicit mouthbreather?**
A. Two surfaced steel mirror
B. One surfaced mirror
C. Concave mirror
D. None of the above
62. **Who described the patients into Ectomorphic, endomorphic and mesomorphic?**
A. Sheldon
B. Graber
C. Mills
D. Profitt

57 D 58 B 59 A 60 D 61 A 62 A

46 MCQs in Orthodontics

98. Angle's classification of malocclusion is based on which of the following principles?
- A. The most indicative irregularity of the teeth is in the anteroposterior direction
 - B. The maxillary first permanent molar is the key to occlusion
 - C. The curvature and size of the "line of occlusion" (greatest normal occlusal contact) is unique to each individual
 - D. None of the above
 - E. A, B and C are all correct
99. The terms "neutroclusion" for Angle's class I "distocclusion" for class II and "mesiocclusion" for class III were suggested by:
- A. Dewey Anderson
 - B. Angle himself later suggested these terms to simplify his classification
 - C. Lischer
 - D. Simon
100. The term "incompetent lips" refers to:
- A. Lips which do not seal when relaxed with the mandible in its rest position
 - B. Lips against which tongue thrusts during swallowing
 - C. Lips that contract during swallowing
 - D. Lips which have some dryness in the divisions
101. The divisions in Angle's class I were proposed by:
- A. Paul Simon
 - B. Martin Dewey
 - C. Edward H. Angle
 - D. Profitt
102. The skeletal classification of malocclusion was given by:
- | | |
|----------------|-----------------|
| A. Calvin Case | B. Martin Dewey |
| C. Paul Simon | D. EH Angle |

98 B 99 C 100 A 101 B 102 C

214. A boy has ANB angle of -4° facial angle of 98° , he is a case of:
A. True Cl-III
B. Pseudo Cl-III
C. Skeletal class II
D. Dental class II
215. Which of the following tells about maxillomandibular relationship?
A. SNA
B. SNB
C. IMPA
D. ANB
216. The anatomic porion is:
A. The upper edge of the shadow of the auditory canal seen on cephalometric films
B. The location of radiopaque marker placed on the rod that extends into the external auditory meatus as part of the cephalometric head positioning device.
C. Difficult to locate
D. Reliable cephalometric landmark
217. The machine porion is:
A. The upper surface of the ear rod of the cephalometric headholder
B. The upper surface of the shadow of the auditory canal on the cephalogram
C. Not reliable cephalometric landmark
D. Difficult to locate
218. Which is preferred in modern cephalometrics to establish true horizontal plane?
A. Natural head position-NHP
B. Anatomic head positioning
C. Frankfort plane
D. SN plane

214 B 215 D 216 A 217 A 218 A

219. For diagnostic purpose, careful evaluation of facial proportion both in front and profile view is also known as:
- A. Poor man's cephalometric analysis
 - B. Moyer's analysis
 - C. Bolton's analysis
 - D. Down's analysis
220. Which view of cephalogram is useful to diagnose facial asymmetry?
- A. Lateral cephalogram
 - B. AP view
 - C. PA view
 - D. Lateral cephalogram in mouth wide open
 - E. None of the above
221. Gonial angle with age tends to:
- A. Increase in size
 - B. Decrease in size
 - C. Increases upto 9 years and then decreases
 - D. Decreases upto 9 years and then increases
222. In a skeletal class III, the value of ANB will be:
- A. 10
 - B. +2
 - C. +4
 - D. +6
223. Anteroposterior relation of jaws is measured by:
- A. ANB angle
 - B. Angle between SN and FH plane
 - C. Angle between SN and mandibular plane
 - D. Facial angle
224. Sella turcica lies above:
- A. Pons
 - B. Frontal sinus
 - C. Foramen ovale
 - D. Sphenoidal sinus

219 A 220 B 221 B 222 A 223 A 224 D

Etiology of Malocclusion

1. **Abnormal thick Labial frenum results in:**
 - A. Midline diastema
 - B. Imbrication
 - C. Labial inclination of incisors
 - D. Cross bite
2. **Commonest cause of midline diastema is:**
 - A. Rotated canine
 - B. Mesiodens
 - C. Cleft palate
 - D. Traumatic Occlusion
3. **If the maxillary deciduous central incisors are over retained, the permanent central incisors erupt:**
 - A. Palatally
 - B. Buccally
 - C. Get transposed
 - D. Get impacted
4. **The most common local cause of malocclusion is:**
 - A. Hypothyroidism
 - B. Presence of supernumerary teeth
 - C. Early loss of primary teeth
 - D. Late eruption of permanent teeth
5. **Prolonged thumb sucking directly results in all of the following *except*:**
 - A. Deep over bite
 - B. Proclination of upper central incisors
 - C. Rotation of upper laterals
 - D. Constriction of maxilla

1 A 2 B 3 A 4 C 5 A

18. The main cause of bruxism is:
A. Physiological problems
B. Periodontal problems
C. Traumatic occlusion
D. Psychological factors
19. Multiple supernumerary and unerupted teeth are the characteristic features of:
A. Treacher Collins syndrome
B. Ectodermal dysplasia
C. Cleidocranial dysplasia
D. Crouzon's syndrome
20. The most common aetiology of non-skeletal anterior cross bite is:
A. Delayed eruption of permanent teeth
B. Over retention of primary incisors
C. Lack of space for the permanent teeth
D. None of the above
21. 'Beta hypothesis' deals with:
A. Long-face syndrome and its allied problem
B. It is a terminology coined by Dunlap to break deleterious oral habits, in children who are amenable to reason
C. Deals with pressure resorption and deposition
D. None of the above
22. Which of the following statements is correct regarding the "submerged" primary molar?
A. Tooth is non-vital
B. The tooth is ankylosed
C. There is no permanent successor
D. Resorption of roots is not evident
23. Bottle feeding may be stopped in a child by the age of:
A. 6 to 12 months
B. 12 to 18 months
C. 24 to 30 months
D. 30 to 36 months

18 D 19 C 20 B 21 B 22 B 23 B

72 MCQs in Orthodontics

24. As compared to other anterior teeth, a maxillary canine is often found impacted because:
- A. It erupts at the corner position where the arch turns abruptly
 - B. It is the last anterior tooth to erupt, leaving little space for it
 - C. Shape and length of its root prevent its proper alignment
 - D. Already erupted premolars are mesially inclined
25. Nursing bottle caries is least likely to involve:
- A. Maxillary incisors
 - B. Maxillary molars
 - C. Mandibular incisors
 - D. Mandibular molars
26. Cleft lip occurs due to disturbances in:
- A. 4 to 6 weeks of intrauterine life
 - B. 6 to 8 weeks of intrauterine life
 - C. 8 to 10 weeks of intrauterine life
 - D. 10 to 12 weeks of intrauterine life
27. In thumb-sucking posterior cross-bite occurs due to:
- A. Loss of the normal outward thrust of the tongue
 - B. Negative pressure within the oral cavity which causes
 - C. A and B
 - D. Loss of oral seal
28. If the maxillary central incisor is not erupted we must consider:
- A. Congenital absence
 - B. Presence of a supernumerary tooth
 - C. Eagle's syndrome
 - D. None of the above
29. Which test is suitable to elicit mouth breathers?
- A. Butterfly technique
 - B. On inspection
 - C. On intra oral clinical examination
 - D. None of the above

24 B 25 C 26 B 27 C 28 B 29 A

74 MCQs in Orthodontics

36. **Tongue thrusting habit is often exaggerated by:**
A. Enlarged tonsils
B. Tongue-tie
C. Cross bite
D. Prognathic mandible
37. **Cleft lip and palate is:**
A. Congenital
B. Heredity
C. Twin study
D. Chromosomal disorder
38. **Supernumerary tooth is:**
A. Congenital
B. Heredity
C. Dental anomaly
D. Combination of A and B
39. **Midline diastema is due to:**
A. Heredity
B. Peg shaped lateral
C. Tongue thrusting habit
D. Supernumery teeth
E. All of the above
40. **Which theory tells abnormal pressure habit?**
A. Equilibrium theory
B. Blood flow theory
C. Learning theory
D. Psychoanalytic theory
E. Combination of C and D
41. **What is "Trident of habit factors"?**
A. Duration, frequency and intensity
B. Heavy, low and light pressure
C. Intermittent, continuous and interrupted
D. None of the above

36 A 37 A 38 C 39 E 40 E 41 A

86 MCQs in Orthodontics

28. The first tissue to react when orthodontic forces are applied is:
A. PDL
B. Cementum
C. Dentine
D. Alveolar bone
29. During orthodontic tooth movement, periodontal ligament provides:
A. Osteoblasts
B. Osteoclasts
C. Both A and B
D. None of the above
30. The centre of rotation in bodily tooth movement is located:
A. Within the apical one-third
B. At the junction of apical, and middle one-third
C. At infinity
D. Within the middle one-third
31. In orthodontic treatment, rapid tooth movement will cause all *except*:
A. Devitalisation of teeth
B. Resorption of roots
C. Diffuse calcification of pulp
D. Ankylosis
32. In primary tooth intrusion is common because:
A. Alveolar bone is pliable
B. Vertical overlap of teeth
C. Lip protection to maxillary anterior
D. All of the above
33. An orthodontic tooth movement that takes place along the vertical axis is:
A. Tipping
B. Translation
C. Intrusion
D. Rotation

28 A 29 C 30 C 31 D 32 A 33 C

88 MCQs in Orthodontics

40. **Implants are used as anchorage control in orthodontics:**
A. False
B. True
C. Partly correct
D. Under research
41. **Which is the example for reciprocal anchorage?**
A. Midline diastema correction
B. Through the bite elastic
C. Using Cl-II elastics
D. Combination of A and B
42. **Which Law is applicable in anchorage?**
A. Sir Issac Newton Law
B. Canine Law
C. Wolf's Law
D. Coulomb's Law
43. **Which of the following is not a source of orthodontic anchorage?**
A. Bone
B. Root
C. Dentine
D. Skull
44. **Baker anchorage utilises:**
A. Intermaxillary anchorage with screws
B. Intramaxillary anchorage with screws
C. Intermaxillary anchorage with elastics
D. Intramaxillary anchorage with elastics
45. **Diastema closure involves:**
A. Stationary anchorage
B. Simple anchorage
C. Reciprocal anchorage
D. All of the above
46. **Occipital anchorage is related to:**
A. Couple force
B. Intraoral force
C. Intramaxillary force
D. Extraoral force
E. Tipping force

40 B 41 D 42 A 43 C 44 C 45 D 46 D

81. Which screw is used for Activator-III appliance?
A. Norwegian screw
B. Fan type screw
C. Universal standard screw
D. Hyrax screw
82. Which appliance is not suitable for skeletal CI-III malocclusion at the age of 8 years?
A. Frankel III
B. Activator III
C. Chin cap
D. Reverse pull headgear
E. Maxillary splint appliance
83. A disadvantage of the activator is that it cannot:
A. Help in correction of thumb sucking
B. Be used in cases with low facial height
C. Be used in skeletal maxillary protrusion
D. Utilize development during growth period

FIXED APPLIANCE

84. The first appliance system designed by Angle was:
A. Ribbon arch
B. E-Arch
C. Pin and tube
D. Edgewise appliance
85. The straight wire appliance was developed from which of the following techniques?
A. Begg
B. Edgewise
C. Ribbon arch
D. Tip edge
86. The force needed to move teeth in a fixed appliance is derived from:
A. Bands
B. Brackets
C. Archwires
D. Eyelets and cleats

81 A 82 E 83 A 84 B 85 B 86 C

87. Who is the father of straight wire appliance?
A. Andrews
B. Bennett
C. Wick Alexander
D. Begg
88. The edgewise appliance was introduced by:
A. PR Begg 1965
B. LF Andrews 1972
C. Edward Angle 1929
D. Johnson 1938
89. The apparatus used to measure the amount of pressure being exerted by a spring or elastic is known as:
A. Wanson Caliper
B. Vernier Caliper
C. Boley Gauge
D. Dontrix
90. The edgewise arch mechanism was introduced by Edward Angle in:
A. 1929
B. 1928
C. 1931
D. 1932
91. The following is not a component of fixed appliances:
A. Bands
B. Brackets
C. Open hellicie
D. Adam's clasp
92. Molar band material is normally:
A. 0.180×0.005 inch
B. 0.125×0.003 inch
C. 0.150×0.004 inch
D. 0.180×0.003 inch

- 116. If water or saliva contaminates with etched surface, what is the reaction on etched enamel?**
A. Remineralisation takes place
B. Fluoride ions get absorbed
C. Less etching and poor bond strength
D. All of the above
- 117. Which is the esthetic bracket?**
A. Ceramic brackets
B. Plastic brackets
C. Both A and B
D. None of the above
- 118. What is the choice of appliance suitable for skeletal Cl-II div.1 at the age of 14 years (Early permanent dentition)?**
A. Herbst appliance
B. Twin block appliance
C. Either A or B
D. Fixed appliance
- 119. What is TPA?**
A. Transpalatal arch
B. Treatment planning approach
C. Transverse plane articulation
D. None of the above
E. All of the above
- 120. Who demonstrated acid pretreatment in orthodontic practice?**
A. Buonocore
B. Miura
C. Retief
D. Smith
- 121. How much percentage of phosphoric acid used by Buonocore in 1955?**
A. 85% phosphoric acid
B. 65% phosphoric acid
C. 30% phosphoric acid
D. 36% phosphoric acid

116 D 117 C 118 C 119 A 120 A 121 A

- 128. Tooth positioner is a flexible rubber appliance used for:**
A. Splinting of mobile teeth
B. Expansion of the arches
C. Correction of minor rotation/crowding by applying a passive force
D. Altering the frenum attachment passively to achieve a favourable tooth position
- 129. Correction of a posterior cross bite by using elastics is an example of:**
A. Intermaxillary reciprocal simple single anchorage
B. Intermaxillary reciprocal simple compound anchorage
C. Intramaxillary reciprocal simple single anchorage
D. Intramaxillary reciprocal simple compound anchorage
- 130. Which of the following procedures is an example of corrective orthodontics?**
A. Early elimination of oral habits
B. Fixed appliances
C. Mandibuloplasty
D. Planned serial extractions
- 131. The bond formed between a bracket and the enamel is:**
A. Chemical
B. Mechanical
C. Ionic
D. Covalent
- 132. Orthodontic bands must:**
A. Allow a thick layer of cement to be placed
B. Overextend on to gingival for esthetics
C. Have an inherent springiness to adapt well
D. Always be available in preformed shapes and sizes
- 133. A Siamese bracket is used in:**
A. Begg's technique
B. Edgewise technique
C. Twin wire technique
D. All of the above

128 C 129 A 130 B 131 B 132 C 133 B

Preventive Orthodontics

1. When lower 2nd deciduous molar is prematurely extracted before the eruption of permanent 1st molar what could be the choice of space maintainer?
 - A. Distal shoe appliance
 - B. Bonded space maintainer
 - C. Band and loop space maintainer
 - D. Functional space maintainer
2. The following conditions should be met by space maintainers:
 - A. Maintain the desired mesiodistal space
 - B. Should not interfere with the eruption of the permanent tooth
 - C. Should not interfere with the vertical eruption of the adjacent teeth
 - D. All of the above
3. The band and loop space maintainer is most frequently used:
 - A. To maintain in the space of a primary first molar before eruption of permanent first molar
 - B. To maintain the space of permanent first molar
 - C. For maintaining bilateral posterior space when more than one tooth is lost
 - D. Both A and B

4. If the mandibular second premolar is missing, then the primary second molar should:
- A. Be extracted and space be maintained till the eruption of all permanent teeth
 - B. Be retained if patient has an acceptable occlusion with slight mesiodistal grinding and later on replaced by fixed bridge
 - C. Be extracted at an early age and be replaced by a fixed bridge
 - D. None of the above
5. In which of the following situations is space most difficult to manage?
- A. A 9.5-year-old patient with loss of maxillary first molar
 - B. A 5-year-old patient with loss of a primary maxillary central incisor
 - C. A 6-year-old with loss of primary first molar
 - D. A 5-year-old with loss of a primary mandibular second molar
6. When multiple maxillary deciduous teeth are extracted bilaterally before shedding time, what could be your space maintainer?
- A. Nance holding arch
 - B. Lingual arch
 - C. TPA
 - D. Lip pumber
7. When multiple mandibular deciduous teeth are extracted bilaterally before shedding time, what could be your space maintainer?
- A. Nance holding arch
 - B. Lingual arch
 - C. TPA
 - D. Lip Bumber

- 12. Which is active retainer?**
- A. After orthodontic treatment any minor correction can be carried during retention period
 - B. It prevents the relapse
 - C. It holds the corrected teeth in position
 - D. None of the above
- 13. The retention period for which the appliance is worn after completion of maxillary expansion is:**
- A. 1 month
 - B. 3 months
 - C. 5 months
 - D. 9 months

Orthodontic Materials

1. **Passivating effect is due to formation of:**
 - A. Iron oxide on austenitic steel
 - B. Chromium oxide on stainless steel
 - C. Bulky layer on polished acrylic
 - D. Nickel oxide on martensitic steel
2. **Orthodontic wires and bands are made from:**
 - A. Martensitic S.S.
 - B. Austenitic S.S.
 - C. Ferritic S.S.
 - D. Carbon steel
3. **Which of the following orthodontic wire has shape memory phenomena?**
 - A. Stainless steel
 - B. Elgiloy
 - C. Nitinol
 - D. Composite plastic wires
4. **Which wire is ideal for coffin spring?**
 - A. 19 gauge wire
 - B. 22 gauge wire
 - C. 21 gauge wire
 - D. 30 gauge wire
5. **Which wire is used for unravelling the crowding?**
 - A. Co-axial wire
 - B. Niti wire
 - C. Copper Niti wire
 - D. All of the above

1 B 2 B 3 C 4 A 5 D

17. **Bending of loop is almost impossible with one of the following wires:**
A. Nitinol wire
B. TMA wire
C. Stainless steel wire
D. Elgiloy wire
18. **Which of the following wires is more useful in space closure and finishing?**
A. Stainless steel wire
B. Nitinol wire
C. β -TMA wire
D. Elgiloy wire
19. **The cement generally used for cementing orthodontic bands:**
A. Zinc oxide cement
B. Glass ionomer cement
C. Zinc phosphate cement
D. Zinc silicophosphate cement
20. **Elgiloy is:**
A. Chrome cobalt alloy
B. Chrome gold alloy
C. Cobalt palladium alloy
D. Chrome palladium alloy
21. **Thickness of the band material used in molars is:**
A. 0.0003 inches
B. 0.03 inches
C. 0.005 inches
D. 0.05 inches
22. **Thickness and width of bands used in the molar area is:**
A. 0.005 x 0.180 inches
B. 0.005 x 0.150 inches
C. 0.005 x 0.125 inches
D. 0.05 x 0.150 inches

17 A 18 C 19 C 20 A 21 C 22 A

4. If the incisor overjet increases during treatment of a class-II div.1 case, you may be seeing:
 - A. Loss of anchorage
 - B. Loss of retention
 - C. Collapse of the lower arch
 - D. Unfavourable growth
5. The mother of a 10-year-old girl is concerned about a 1.5 mm diastema between the child's maxillary central incisors. A very fibrous maxillary frenum is present and inserts on the palatine papilla. When tension is applied to the frenum. The papilla blanches. The treatment of choice is to:
 - A. Perform a maxillary frenectomy to allow the teeth to close
 - B. Close the space orthodontically expecting the frenum to atrophy due to pressure
 - C. Postpone treatment until complete eruption of maxillary canines
 - D. Close the space with a composite/veneer.
6. Slow progress in molar uprighting in an adult patient is due to:
 - A. Overextended bands
 - B. Lack of anchorage control
 - C. The occlusion not being relieved
 - D. The stabilizing wire not being passive
7. Routine radiographic examination of a 6-year-old patient discloses a supernumerary tooth between maxillary central incisors. The dentist should:
 - A. Delay removal of the supernumerary tooth until its complete eruption
 - B. Wait until the child is 12 years of age to remove the supernumerary tooth
 - C. Remove the supernumerary tooth only if it develops acyst
 - D. Remove the supernumerary tooth as soon as possible without injury to the central incisors.

4 A 5 C 6 C 7 D

Bibliography

1. Clinical Sciences – Dentogist MCQs in dentistry by Sanjeev Raj Bhatia, 2nd and 3rd editions.
2. Dental Examination Review by Atul Soin.
3. MCQs in Dental Sciences by Dhirendra Srivastava.
4. MCQs in Dentistry (Clinical Sciences) by Dr. Vijay Pratap Singh.

Also available...


Textbook of Oral and Maxillofacial Surgery (Full Colour)

Neelima Anil Malik

732 Pages, Reprint 2005

ISBN 81-8061-012-8

Price: 825.00




Textbook of Orthodontics

Gurkeerat Singh

664 Pages, 2004

ISBN 81-8061-359-3

Price: 750.00




Textbook of Prosthodontics

Deepak Nallaswamy

864 Pages, 2003

ISBN 81-8061-199-X


Price: 795.00



MCQs in ORTHODONTICS

K Vijayalakshmi studied at Government Dental College, Chennai both BDS in 1977 and MDS in 1981 from Madras University. She was working at Rajah Muthiah Dental College & Hospital, Annamalai University for 20 years from 1981 to 2001. During her service at RMDC&H, she presented 15 papers both Scientific and Clinical papers at National and International Conferences. She published 12 papers in National Journals. She delivered Guest Lecture in Pedodontic Conference in 1994, another in PG convention 1999 at Davangere and many places at Madurai, Coimbatore, etc. She conducted several courses namely Tip edge, Surgical Orthodontics, Two Twin Block Courses one at RMDC&H and another one is Pedodontic Conference 1997 at Chennai. She is currently working as Principal at JKK Nataraja Dental College & Hospital, Komarapalayam from January 2003 to till date. She is a recipient of Capt Dr JS Gangadharan Memorial National Award and Second Best Scientific Paper Award given at IOS Conference held at Ooty in 1982.

JPB Rs. 80.00



81-8061-582-0

JAYPEE BROTHERS
MEDICAL PUBLISHERS (P) LTD
EMCA House, 23/23B Ansari Road, Daryaganj
New Delhi 110 002, India