

Medical Surgical Model Exam January, 2026

MCQ questions with answers related to MSN I

1. A patient with chronic gastritis is found to have a low hemoglobin level and elevated mean corpuscular volume (MCV). What is the most likely underlying cause?
A. Iron deficiency anemia
B. **Megaloblastic anemia**
C. Hemolytic anemia
D. Anemia of chronic disease
2. The nurse is advising a client who has recently undergone a hemorrhoidectomy, instructs to not use a sitz bath until at least 12 hours after the operation. This precaution is primarily to prevent which potential postoperative complication?
A. Constipation
B. **Hemorrhage**
C. Rectal spasm, D. Urinary retention.
3. The hospitalized client with GERD is complaining of chest discomfort that feels like heartburn following a meal. After administering an ordered antacid, the nurse encourages the client to lie in which of the following positions?
A. Supine with the head of the bed flat.
B. On the stomach with the head flat.
C. **On the left side, with the head of the bed elevated 30 degrees.**
D. On the right side, with the head of the bed elevated 30 degrees
4. A nurse is reviewing the results of a client's diagnostic tests to confirm the presence of ascites. Which of the following laboratory findings is commonly associated with ascites?
A. Elevated blood glucose levels, C. Increased white blood cell count
B. **Low albumin levels**
D. Elevated serum level protein
5. The nurse is providing care to a client with myxedema coma. Priority nursing care would include which of the following?
A. Measures to decrease heart rate
B. **Measures to decrease body temperature**
C. Measure to decrease blood pressure
D. Measures to maintain the airway.
6. For the first 72 hours after thyroidectomy surgery, the nurse would assess the female client for Chvostek's sign and Trousseau's sign because they indicate which of the following?
A. **Hypocalcemia**
B. Hypercalcemia
C. Hypokalemia
D. Hyperkalemia
7. A client is returning from a subtotal thyroidectomy for the treatment of hyperthyroidism. The immediate priority in assessing this client would include which of the following?
A. Assess for pain
B. Assess neurological status
C. Assess fluid volume status.
D. **Assess for respiratory distress.**
8. A male client has just been diagnosed with type 1 diabetes mellitus. When teaching the client and family how diet and exercise affect insulin requirements, the nurse should include which guideline?

- A. "You'll need more insulin when you exercise."
- B. "You'll need less insulin when you exercise."
- C. "You'll need less insulin when you increase your food intake."
- D. "Exercise have no role in insulin adjustment."
9. A 50-year-old man with Type 1 diabetes is learning about self-monitoring of blood glucose (SMBG). Which site and technique should he use for optimal blood glucose testing to minimize discomfort and improve accuracy?
- A. The side of the fingertip is rotated among different fingers
- B. The tip of the fingertip by using a deep puncture for more blood
- C. The side of the fingertip by avoiding rotating among different fingers
- D. Tip of the fingertip by applying pressure to ensure ample blood flow
10. A 50-year-old woman with diabetes is at a family gathering when she suddenly feels weak, confused, and faint. Her friends notice she is also sweating heavily. In her bag she has insulin and sugar, but not a glucometer. What should be the first action taken by her friends?
- A. Give her water
- B. Give her insulin
- C. Give her sugar
- D. Have her lie down and rest.
11. A 45-year-old pt with a history of T1D presents to the emergency department with signs of DKA. His serum potassium level is measured at 2.8 mEq/L. What is the most appropriate action regarding insulin therapy for this patient?
- A. Start insulin therapy alone immediately.
- B. Administer a bolus of insulin and monitor potassium levels.
- C. Administer 20-30 mEq of potassium IV and start insulin therapy
- D. Administer 20-30 mEq of potassium IV and hold insulin therapy.
12. A patient with a brain tumor is experiencing increased intracranial pressure. Which of the following nursing interventions is most appropriate?
- A. Administer oxygen via nasal cannula.
- B. Position the patient in a flat position
- C. Keep the patient's head elevated at 30 degrees
- D. Encourage the patient to perform deep breathing exercises
13. A patient with a recent spinal cord injury presents with flaccid paralysis and no sensation below the level of the injury. Which nursing diagnosis reflects the patient's condition regarding skin care?
- A. Impaired Skin Integrity
- B. Risk for Impaired Skin Integrity
- C. Altered Sensory Perception
- D. Acute Pain
14. A nurse is conducting an assessment on a patient who has sustained a traumatic brain injury. Which of the following assessment findings should the nurse prioritize?

- A. Patient's blood pressure
B. Patient's temperature
C. **Glasgow Coma Scale score**
D. Urine output
15. A 35-year-old patient presents with hoarseness and a sore throat. The nurse assesses the patient's vital signs and notes a low-grade fever. Which nursing intervention is most appropriate for this patient?
A. **Encourage the patient to engage in vocal rest.** C. Administer antibiotics as prescribed.
B. Suggest the use of throat irritants D. Immediate hospitalization for further evaluation.
16. A 72-year-old woman who recently underwent hip replacement surgery is hospitalized and develops a fever and cough on the fifth day of her hospital stay. A chest X-ray shows new infiltrates. What classification of pneumonia does she likely have?
A. Community-acquired pneumonia C. **Hospital-acquired pneumonia**
B. Aspiration pneumonia D. Chlamydia pneumonia
17. A patient diagnosed with chronic bronchitis is being evaluated for future management and treatment options. Which of the following strategies is most important in managing this condition?
A. Initiating long-term corticosteroid therapy C. **Smoking cessation**
B. Starting antibiotic prophylaxis D. Routine use of cough suppressants
18. A 68-year-old male patient is admitted with sharp chest pain that increases with inspiration and coughing. Upon assessment, you note diminished breath sounds on the affected side and a pleural rub upon auscultation. Which nursing diagnosis is most appropriate for this patient?
A. Impaired Gas Exchange C. **Acute Pain**
B. Ineffective Breathing Pattern D. Risk for Infection
19. A patient received burns to his entire back & left arm. Using the Rule of Nines, the nurse can calculate that he has sustained burns on what percentage of his body?
A. 9 % B. 18% C. **27%** D. 36%
20. A patient is admitted to burn intensive care unit with extensive full-thickness burns. The nurse is most concerned about the patient's:
A. Fluid & electrolyte status B. **Risk of infection** C. Body image D. Level of pain
21. A 60 k.gs patient came to the emergency department with a thermal burn accident in which 35% of the TBSA was affected. By taking 2 as a multiplying factor in the consensus formula for calculating the amount of fluid to be administered. How much fluid should be administered for this patient within the first 24 hours?
A. **8400 ml** B. 4200 ml C. 2100 ml D. 7000 ml
22. The nurse is caring for a patient with Glomerulonephritis. Which of the following interventions would the nurse recommend be included in the patient's plan of care?
A. Increase fluid intake C. **Decrease sodium intake.**
B. Increase potassium intake D. Decrease calcium intake.

23. A 22-year-old male patient came to the hospital with hematuria, edema, and excess protein in the urine. The doctor was admitted to the medical ward with a diagnosis of acute glomerulonephritis. After seven days of admission, he completed his drug regimen and planned to discharge the patient.

What is the most appropriate nursing discharge education?

- A. Emphasis on emotional support
- B. Educate to drink excess liquids
- B. Educate about restrictions of fluid and diet
- D. Inform about skin care

24. A patient with pyelonephritis once it detects it need protect from repeated attacks of urinary tract infection, encouraging of fluid intake and taking of proper medication, if it fails, the patient develops a complication. Which is the nurse most likely accept?

- A. Acute glomerulonephritis
- C. Nephritic syndrome
- B. Prostatitis
- D. Chronic renal failure

25. A 28-year-old man with a fractured humerus calls the nurse into his room. Upon assessment, the nurse finds the patient to be tachycardic, pale, and confused. The nurse suspects the patient may be experiencing which of the following complications?

- A. Disseminated intravascular coagulopathy
- C. Compartment syndrome
- B. Fat emboli
- D. Deep vein thrombosis

26. Case Scenario: A 68-year-old male patient, Mr. Demeke, is admitted to the JMC medical ward with shortness of breath, bilateral leg swelling, and fatigue. He has a history of hypertension and was recently non-compliant with his medication. On assessment, you find elevated blood pressure (160/100 mmHg), bilateral pitting edema, crackles on auscultation, and reduced urine output. Which of the following is the most appropriate objective data the nurse should collect during the assessment of Mr. Demeke?

- A. Patient states he feels tired all the time
- B. Patient expresses fear of being hospitalized
- C. Patient has bilateral pitting edema on the lower limbs
- D. Patient believes his condition is due to stress

27. A nurse is assessing a patient with a history of heart failure. Which of the following data is **objective**?

- A. Patient reports feeling dizzy
- C. Patient complains of shortness of breath
- B. Nurse notes crackles on lung auscultation
- D. Patient states he is tired

28. During a physical exam, the nurse observes a reddened area on the patient's sacrum. This is:

- A. Objective data
- B. Inference
- C. Secondary data
- D. Subjective data

29. Ms. Hana, a 50-year-old female, is 24 hours post-abdominal surgery. She reports pain rated 8/10, has a shallow respiratory pattern, and avoids moving due to discomfort.

Which nursing diagnosis best fits this situation?

- A. Acute pain related to the surgical incision
- B. Risk for constipation related to inactivity

- C. Impaired gas exchange related to the surgical site
- D. Risk for infection related to the hospital environment

30. Patient Abebe, 26 years old, newly diagnosed with HIV/AIDS, Nurse Aster wants to initiate antiretroviral therapy. What is a typical first-line Anti-retroviral therapy regimen for Abebe in Ethiopia?

- A. AZT/3TC/EFV
- B. TDF/3TC/NVP/EFV
- C. ABC/3TC/LPV/r
- D. d4T/3TC/NVP

31. Patient Elemo, 34 years, has a cough and night sweating. Nurse Dereje uses STAT-PAK, positive. ABON is positive, and SD BioLine, is negative for Elemo. How Nurse Dereje's next action, following the results, and interpret the results of Elemo:

- A. Report HIV positive
- B. Perform ABON rapid HIV test
- C. Report HIV Inconclusive and appoint after 14 days
- D. Perform Stat Pak, and ABON parallel rapid HIV test