

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

MIDWIFERY



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH-ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority and is essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and to the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPL

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, leadership and management, ethics, and Health Promotion as outlined in the Information Booklet 2024.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey—and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Midwifery

1. A 34-year-old primigravida mother has come to a hospital for initiation of antenatal care. A midwife on duty has taken her biographical data, history of her current pregnancy, past gynaecologic history, and family history.
What is the most important additional history needed to complete the history taking?
A. General medical history
B. Developmental history
C. Past obstetric history
D. Psycho-social history
ANSWER: A

2. A 34-year-old pregnant woman who wanted to know her gestational age came to a maternity clinic on 10/07/2016 E.C. Her last normal menstrual period had started on 05/05/2016 E.C. and completed on 09/05/2016 E.C.
What is the most accurate gestational age of the mother?
A. 8 weeks + 1 day
B. 8 weeks + 5 days
C. 9 weeks + 3 days
D. 9 weeks + 6 days
ANSWER: B

3. A pregnant woman has come to a health center for her first ANC visit. A midwife has provided her HIV counseling and testing services after history taking and physical examination.
What is the most appropriate pre-test counseling to be made in this case?
A. Reasons for recommending tests for pregnant women
B. Denial of the test may affect her access to all other services
C. Test results must be shared with her partner and children
D. Repeat test after three months is not needed if tested negative
ANSWER: A

4. A pregnant mother at the 37th week of gestational age came for a routine ANC visit. She had complained of severe chest discomfort for one month. On physical examination, her fundus was occupied by a ballotable, round, hard mass. The fetal heart sound was audible around the umbilicus.
What is the most likely diagnosis for this case?
A. Breech presentation
B. Cephalic presentation
C. Shoulder presentation
D. Compound presentation
ANSWER: A

5. A 22-year-old primigravida pregnant woman came to a health center for her first ANC contact at the 24th week of gestation and complained of feeling a tense and tingling sensation on her breast. She also had irregular brownish discolorations on her forehead, nose, and cheeks.
What is the most likely explanation in this case?
A. Pathological change of breast and skin
B. Physiological change in pregnancy
C. Minor disorder of pregnancy
D. Major disorder of pregnancy
ANSWER: B

6. A 38-year-old pregnant mother came to a hospital with complaints of abdominal pain and vaginal bleeding of five hours duration at the 38th week of pregnancy. She had a history of pregnancy-induced hypertension. Physical examination showed a blood pressure of 70/50 mmHg, a pulse rate of 75/min, and a respiration rate of 16/min. On abdominal examination, tenderness on palpation and a fetal heartbeat of 75/min were felt. On presentation, there was active, copious menstruation like bleeding in the vulva with an estimated amount of 500 ml blood loss.

What is the most likely late pregnancy bleeding in this case?

- A. Mild abruptio placentae
- B. Severe abruptio placentae
- C. Concealed abruptio placentae
- D. Moderate abruptio placentae

ANSWER: D

7. A pregnant mother has come to a clinic for a follow-up. A duty midwife noticed that her fundal height and gestational age calculated by a reliable last normal menstrual period were not coherent. On ultrasound investigation, the fetal ponderal index was found to be in the 8th percentile, and the single deepest vertical pocket of amniotic fluid was found to be less than one centimeter.

What is the most likely primary diagnosis of this case?

- A. Intrauterine growth restriction
- B. Intrauterine fetal death
- C. Oligohydramnios
- D. Wrong date

ANSWER: A

8. A woman came to a health center at the 24th week of her pregnancy with a complaint of inability to pass stool. She was diagnosed with constipation. On history, she did not have regular bowel habits and had minimal physical activity. Other potential medical and surgical causes were excluded.

What is the most appropriate initial step intervention in this case?

- A. Advise to take more refined foods
- B. Prescribe anti-hemorrhoid cream
- C. Advice to take high-fiber diets
- D. Prescribe castor oil

ANSWER: C

9. A pregnant woman has come to a hospital with complaints of shortness of breath, palpitations, and swelling of the legs. She has been diagnosed with polyhydramnios. On abdominal examination, the abdomen looks globular and full at the flank, and fundal height is more than the period of amenorrhea. On ultrasound, the deepest vertical pocket (DVP) is 10 cm.

What is the most effective prioritized management in this case?

- A. Find the underlying causes
- B. Advice prolonged bed rest
- C. Perform amniotic reduction
- D. Administer diuretics

ANSWER: B

10. A pregnant woman came to a hospital for routine follow-up at the 34th week of gestation from reliable LNMP and was diagnosed with IUGR. On physical examination, a fundal height of 30-week-size uterus was recorded. Her mid-upper arm circumference was found to be 20 cm, and her weight gain during pregnancy was insignificant during the whole pregnancy. On ultrasound scanning, there was no evidence of oligohydramnios and the ponderal index is around the 7th percentile.

What is the most effective management plan in this case?

- A. Administer low-dose aspirin
- B. Provide ready-to-use foods
- C. Terminate the pregnancy
- D. Reduce physical activity

ANSWER: B

11. A 32-year-old pregnant woman presented to a hospital for ANC contact at the 32nd week of gestational age and complained fatigue recently. Her vital signs showed a blood pressure of 90/60 mmHg, a pulse rate of 90/min, and a respiratory rate of 22/min. Ultrasound scanning revealed a dichorionic twin pregnancy with a reassuring biophysical profile.

What is the most appropriate management plan in this case?

- A. Schedule for frequent fetal surveillance
- B. Consider pregnancy termination
- C. Admit to hospital for follow-up
- D. Follow as a standard ANC client

ANSWER: A

12. A pregnant woman has come to a health centre for a second antenatal care visit at the 20th week of gestation following her first contact, which was one month before. On review of her previous record, the midwife has found negative results for HBV, HIV, and syphilis and a hemoglobin level of 9 gm/dl. The midwife has also confirmed that the woman had been adherent to iron-folic supplementation for the last month. The midwife is planning to further evaluate the condition of the woman.

What is the most appropriate evaluation to be undertaken in this case?

- A. Hemoglobin test
- B. HIV test
- C. HBV test
- D. Syphilis test

ANSWER: A

13. A 22-year-old G-III, Para-II mother has come to a hospital for routine ANC follow-up at the 34th week of gestational age. On assessment, the ultrasound record has revealed a twin pregnancy with monochorionic placentation. Otherwise, no other abnormal findings are detected.

When will the next most appropriate appointment be in this case?

- A. 35 weeks
- B. 36 weeks
- C. 37 weeks
- D. 38 weeks

ANSWER: A

14. A 24-year-old primigravida woman has come to a health center at the 26th week of gestation for the first ANC contact. A midwife has assessed her conditions and confirmed, based on laboratory results, that the woman is Rh-negative.

What is the most likely next step intervention in this case?

- A. Anti-D after 28 weeks
- B. Re-testing at 28 weeks
- C. Plan counseling at 28 weeks
- D. Partner testing before 28 weeks

ANSWER: D

15. A pregnant mother on ART came to a hospital with a complaint of pushing-down pain of six hours duration at her 35th week of gestation. On assessment, vital signs showed a blood pressure of 110/70 mmHg, a pulse rate of 82/min, a respiratory rate of 20/min, a temperature of 37.5 degrees Celsius. The membrane was intact, cervix was 6 cm dilated, and there were 3 contractions in 10 minute each lasting 40 seconds.

What is the most likely strong evidence of a risk factor for MTCT in this case?

- A. Poor immune status
- B. Maternal anemia
- C. Chorioamnionitis
- D. Prematurity

ANSWER: D

16. A pregnant woman came to a hospital at the 38th week of gestation with pushing-down pain of three hours duration. A midwife assessed her antenatal care record and found normal findings for all routine lab investigations that had been done at her 34th week of gestation.

What is the most likely laboratory test to be repeated in this case?

- A. HBsAg
- B. VDRL
- C. HIV
- D. Hgb

ANSWER: D

17. A pregnant mother at her 37th week of gestation has come to a hospital with a referral slip, as she was diagnosed as having "protracted active first stage of labor plus early term pregnancy." On assessment, cephalic fetal presentation, membrane rupture, and the fetal shoulder and the rest of the body are delivered at arrival to the hospital.

What is the most likely cardinal movement assessed at arrival in this case?

- A. Flexion
- B. Expulsion
- C. Extension
- D. Engagement

ANSWER: B

18. A pregnant woman came to a health center for her first ANC contact at the 22nd week of gestation. Following history taking and physical examination, a midwife conducted an HIV test for the woman and reported a positive on the first test (stat pack). The second (Abon) and the third (SD-bioline) tests were positive and negative, respectively.

What is the most appropriate interpretation of this finding?

- A. Invalid
- B. Reactive
- C. Inconclusive
- D. Non-reactive

ANSWER: C

19. A 32-year-old pregnant woman came to a hospital at her 37th week of gestation with a complaint of pushing-down pain of four hours duration. Upon examination, three contractions were observed in 10 minutes, each lasting 40 seconds; a fetal heartbeat of 144/min recorded; the membrane was intact; and the cervix was dilated 6 cm.

What is the most likely stage of labor in this case?

- A. First stage
- B. Second stage
- C. Third stage
- D. Fourth stage

ANSWER: A

20. A 24-year-old Gravida-II, Para-I woman presented to a health center with a complaint of a pushing-down pain of 12 hours duration. Physical examination showed two contractions in 10 minutes, each lasting 15-19 seconds, an intact membrane, and a cervix that admits the tip of a finger. After four hours, the re-examination result showed one uterine contraction in 10 minutes, each lasting 15 seconds, and no change in cervical dilation.

What is the most likely classification of labor in this case?

- A. False labor
- B. Prolonged labor
- C. Precipitated labor
- D. Latent first stage labor

ANSWER: A

21. A laboring mother, who is in the latent phase of the first stage labor, is tested for HIV and has become positive. Previously, she didn't know her HIV status and had never been tested. After brief counselling, the mother agreed to take ART.

What is the best appropriate time to start ART for this case?

- A. At the end of post-partum period
- B. As soon as possible during labor
- C. When cervix is fully dilated
- D. Immediately after delivery

ANSWER: B

22. A 25-year-old woman has come to a health centre with a complaint of pushing down pain of three hours duration. She had no antenatal care contact. On physical examination, feto-maternal conditions are stable with satisfactory progress of labor. Her HIV test result is positive.

What is the most appropriate safe practice for this case?

- A. Refer the mother to hospital for caesarean delivery
- B. Use double glove for any physical contact
- C. Advise the mother for mixed feeding
- D. Avoid invasive vaginal procedures

ANSWER: D

23. A 25-year-old laboring woman has been admitted and followed for the last 10 hours at a hospital. The partograph passes to the action line, the cervix is eight centimetres dilated, membrane is intact, fetal heart beat is 170 beats/minute, the contraction is two in 10 minutes each lasting 15-25 seconds, and the fetal head descent is 3/5.

What is the most appropriate next step in the management of this case?

- A. Caesarean delivery
- B. Augmentation of labor
- C. Feto-maternal follow-up
- D. Artificial rupture of membrane

ANSWER: A

24. A primigravida woman has come to a health centre with a complaint of pushing down pain of five hours duration at her 41st week of gestation. She had no history of antenatal care contact. She has leakage of watery fluid from her vagina of two hours duration.

What is the most likely next step to be performed in this case?

- A. Providing TT vaccination
- B. Perform physical examination
- C. Doing HIV testing and counselling
- D. Sending for laboratory investigations

ANSWER: B

25. A woman has been admitted to a health centre in the second stage of labor. Physical examination shows that the fetal head engagement of 2/5, fetal heart beat of 145 beats/min and uterine contraction of 4 in 10 minutes.

What is the appropriate management plan for this case?

- A. Refer the woman to a hospital
- B. Advice the mother to hold pushing
- C. Monitor fetal heart beat every 15 minutes
- D. Conduct the delivery using obstetric forceps

ANSWER: C

26. A midwife has followed a 28-year-old primigravida woman in the first stage of labor for four hours. On examination, three uterine contractions are recorded, each lasting 50 seconds and fetal heart rate of 140 beats/min. The cervicograph line crosses the alert line.

What is the most appropriate next evaluation to be done for this case?

- A. Maternal blood pressure
- B. Colour of meconium
- C. Head engagement
- D. Assess hydration

ANSWER: C

27. A multigravida woman was admitted to a hospital with a fully dilated cervix and vertex presentation. After two hours, the midwife reassessed the woman and found that station +1, blood pressure of 90/60 mmHg, and fetal heart rate of 156 beats/min. The midwife has planned to apply vacuum extractor to deliver the fetus.

What is the most likely indication of vacuum delivery in this case?

- A. Unstable maternal condition
- B. Non-reassuring fetal heart beat
- C. Prolonged second stage of labor
- D. Deep engagement of the fetal head

ANSWER: C

28. A 34-year-old woman presented to a hospital at her 39th week of pregnancy with pushing down pain of six hours duration. Physical examination result showed cervical dilatation of 6cm, vertex presentation, -2 station, and 3 contractions in 10 minutes each lasting 45 seconds. Ultrasound assessment revealed amniotic fluid index of 6 and estimated fetal weight of 4.6 kg. Emergency caesarean section was decided to be done after evaluation.

What is the most likely indication for caesarean section in this case?

- A. Macrocosmic fetus
- B. Oligohydraminous
- C. Polyhydraminous
- D. Mal presentation

ANSWER: A

29. A 26-year-old primigravida woman came to a health centre at her 34th week of gestation with a complaint of pushing down pain for six hours duration. She had ANC follow up and her record disclosed that she was non-reactive for HIV, syphilis, and normal urine analysis, and haemoglobin is 10gm/dl.

What is the most likely risk factor for preterm labor in this case?

- A. Anaemia
- B. Maternal age
- C. Urinary tract infection
- D. Socio economic status

ANSWER: A

30. A 34-year-old multigravida woman presented to a health centre at her 32nd week of gestation with complaints of exaggerated fetal movements and excessive maternal weight gain. The midwife wanted to diagnose the condition at the health center before referral to a hospital.

What is the most important assessment component to diagnose this case?

- A. Palpation of multiple fetal poles manually
- B. Determination of gestational age by LNMP
- C. Measurement of fundal height by tap meter
- D. Confirmation of family history of pregnancy

ANSWER: A

31. A 34-year-old pregnant woman came to a hospital with vaginal bleeding of two hours duration at her 34th week of gestation. On rapid initial assessment, she was restless and anxious. The midwife diagnosed her as shock and administered IV fluid.

What is the most likely sign/symptom to confirm the diagnosis in this case?

- A. Tachycardia
- B. Warm skin
- C. Bradypnea
- D. Polyuria

ANSWER: A

32. A 38-week pregnant woman came to a hospital with labor pain of 10 hours duration. Upon assessment, cervix is dilated 8 cm, membrane is ruptured, station is +1 and occiput is felt directly towards the sacrum.

What is the most appropriate position in this case?

- A. Left occiputo posterior
- B. Right occiputo posterior
- C. Direct occiputo posterior
- D. Direct Occiputo transverse

ANSWER: C

33. A 32-year-old multigravida woman presented to a health facility at 2:00 AM with pushing down pain of 10 hours duration. On admission, cervix was fully dilated and station was +1, a uterine contraction was 3 in 10 minutes each lasting 40 seconds. On re-assessment, station became +2 and adequate contraction but failed to give birth until 5:00 AM.

What is the mostly likely diagnosis of this case?

- A. Arrest of descent
- B. Failure of descent
- C. Protraction disorders
- D. Prolonged second stage

ANSWER: D

34. A 38-year-old Gravida III para II woman presented to a hospital at her 38th week of gestation with pushing down pain of 14 hours duration. On physical examination; a clear fluid leaked from vagina during coughing, blood pressure was 70/40mmHg, Pulse rate was 110 beats/min, respiratory rate was 24 breathes/min, and temperature was 39 degrees centigrade. The midwife assessed her conditions and administered IV fluid but hypotension failed to be corrected.

What is the most likely type of shock in this case?

- A. Hypo-volemic shock
- B. Cardiogenic shock
- C. Neurogenic shock
- D. Endotoxic shock

ANSWER: D

35. A 40-year-old Gravida-II, Para-I woman presented at her 36th week of gestation to a hospital for ANC follow up. Trans-abdominal ultrasound examination showed a double fold of amnion and chorion creates a broad, opaque white ring and ridge on the fetal surface.

What is the most likely placental abnormality in this case?

- A. Circummarginate placenta
- B. Circumvallate placenta
- C. Succenturiate placenta
- D. Multilobate placenta

ANSWER: B

36. A 32-year-old multi-Gravida mother presented to a hospital with pushing down pain of 12 hours duration. On physical examination, uterine size measured 38 weeks, cervix was fully dilated, and four uterine contractions were recorded in 10 minutes each lasting 45 seconds. Assessment result after two hours indicated that there was a recoil of the head back against the perineum and inadequate restitution following delivery of the head.

What is the most likely diagnosis in this case?

- A. Shoulder dystocia
- B. Brow presentation
- C. Compound presentation
- D. Persistent occiput transverse

ANSWER: A

37. A 34-year-old multi-Gravida woman has come to a health centre at her 32nd week of pregnancy from reliable LNMP with a complaint of excessive weight gain. The woman also concerned about exaggerated fetal movement during the current pregnancy. Abdominal examination revealed palpation of multiple fetal poles and uterus was found to be 36 weeks sized.

What is the most likely diagnosis in this case?

- A. Multiple pregnancy
- B. Macrocosmic fetus
- C. Polyhydraminous
- D. Uterine tumour

ANSWER: A

38. A 30-year-old Gravida-III, Para-II woman came to a hospital at 38 weeks of gestation with a complaint of pushing down pain of 12 hours duration. On assessment; cervix was fully dilated, vertex presentation, station was +2 and membrane is ruptured. Vacuum was applied to shorten the second stage of labor. The woman failed to deliver after three effective pulls following the application of the centre of the cup on the flexion point.

What is the most likely diagnosis in this case?

- A. Poor vacuum traction
- B. Failed vacuum application
- C. Ineffective vacuum pressure
- D. Inappropriate vacuum position

ANSWER: B

39. A midwife has decided to manage second stage of labor with vacuum extraction. The physical examination findings are vertex presentation with occipito-anterior position, 2/5 fetal head descent, and 148 fetal heart beats/min. The Midwife has assembled all necessary materials.

What is the most appropriate next step in the management of this case?

- A. Assess for descent
- B. Identify flexion point
- C. Apply the largest cup
- D. Check for contraction

ANSWER: B

40. A primigravida laboring woman is admitted to a hospital with obstructed labor. The cervix is fully dilated, fetal heart beat is 170 beats/min, and fetal head descent is 4/5.

What is the most appropriate management for this case?

- A. Destructive delivery
- B. Caesarean delivery
- C. Vacuum delivery
- D. Forceps delivery

ANSWER: B

41. A 28-year-old multigravida woman is admitted to a hospital. On physical examination, cervix is 8 cm dilated, frank breech presentation, fetal heart beat is 148 beats/min, uterine contraction is 4 times in 10 minutes each lasts 55 seconds, and membrane is intact.

What is the most appropriate next management for this case?

- A. Send for Emergency caesarean delivery
- B. Perform artificial rapture of membrane
- C. Follow the labor with partograph
- D. Provide prophylactic antibiotics

ANSWER: C

42. A 28-year-old gravid II, para I mother is admitted to a hospital in the first stage of labor. She has one lower uterine segment transverse caesarean scar performed for the indication of non-reassuring fetal heart rate. Cervical dilation and descent is progressing well, intact fetal membranes, the fetal heart beat is 130 beats/min, vertex presentation, right occipito-anterior position, and adequate uterine contraction. Ultrasound showed that the pregnancy is singleton with the estimated fetal weight of 3100 grams. No other pertinent finding is obtained.

What is the most appropriate next plan in the management of this case?

- A. Monitor feto-maternal condition
- B. Emergency caesarean section
- C. Perform amniotomy
- D. Augment the labor

ANSWER: A

43. A pregnant mother came to first ANC visit after five-months of amnhorea. On assessment, she did not have significant weight gain despite the pregnancy was confirmed to be five-month. She looked very thin and a midwife wanted to assess the nutritional status of the mother.

What is the most appropriate assessment in this case?

- A. Weight gain
- B. Body mass index
- C. Daily calorie intake
- D. Mid-Upper arm circumference

ANSWER: D

44. A 30-year-old pregnant woman came to the ANC clinic of a hospital for the first contact. On history, she reported as there was a family member confirmed to have liver disease. Now, she wants to understand the available test to confirm liver disease.
What is the most appropriate initial test in this case?
- A. Hepatitis B virus viral load
 - B. Hepatitis B e antigen (HBeAg)
 - C. Hepatitis B surface antigen (HBsAg)
 - D. Hepatitis B surface antibody (anti-HBs)
- ANSWER: C
45. A pregnant mother came to a hospital with a complaint of lesion on genital area. She reported that the lesions begin very small but disseminated up to her anus from time to time. On history, she reported as she had multiple sexual partners and poor in exercising safe sexual practice. On inspection, the lesions were papillary, cauliflower appearance, and clustered on the verge of labia and vagina.
What is the most appropriate assessment in this case?
- A. Genital Warts
 - B. Herpes Genitalis
 - C. Granuloma Inguinale
 - D. Molluscum contagiosum
- ANSWER: A
46. A 20-week pregnant mother came to a health centre for routine ANC visit. On history, she did not have significant fatigue or weakness. However, conjunctivas were slightly pallor with normal vital signs. On investigation haemoglobin level was 10g/dl and RBC was 3.2 million per cubic millimetre and peripheral smear shows normal morphology.
What is more likely type of anemia present in this case?
- A. Physiological anaemia
 - B. Iron deficiency anaemia
 - C. Folic acid deficiency anaemia
 - D. Vitamin B12 deficiency anaemia
- ANSWER: A
47. A pregnant mother came to the ANC clinic of a hospital visit at her 16th week of pregnancy with a complaint of dizziness and weakness of one week duration. All routine laboratory investigations were done and found to be normal; except her haemoglobin level was found to be 11g/dl.
What is the most appropriate management for this case?
- A. Iron sulphate in supplementary dose
 - B. Iron sulphate in therapeutic dose
 - C. Folic acid in a therapeutic dose
 - D. Prophylactic Mebendazole dose
- ANSWER: A
48. A 10-week Gravida-II Para-I woman has come to a hospital for her first visit. She is concerned about the health of her current pregnancy since her previous birth was complicated by neural tube defect. She has been screened for all routine laboratory investigations and found to be normal.
What is the most appropriate management in this case?
- A. Parenteral iron therapy
 - B. Folic acid in therapeutic dose
 - C. Prophylactic anthelmintic drug
 - D. Iron sulphate in therapeutic dose
- ANSWER: B

49. A midwife was counselling a 39-week pregnant woman birth preparedness plan. On her previous birth, she had experienced still birth while giving birth with the help of a traditional birth attendant. A midwife intended to provide client specific birth preparedness and complication readiness plan. What is the most appropriate plan to be given in focus in this case?
- A. Importance of skilled attendance at birth
 - B. How to get transport to health facility
 - C. Types of health facility to seek care
 - D. Aware about possible danger signs
- ANSWER: A
50. A midwife was counselling antenatal mother on issues related to nutrition. On assessment of daily nutritional intake, the woman reported as she consumes raw fruits, vegetables and meat. She also likes poultry and fish products in addition to taking vitamin B complex products. As she reported, she wants to have raw meat in her daily dish. What is the most appropriate point to be raised in counseling this case?
- A. Avoid raw meat
 - B. Avoid poultry and fish
 - C. Avoid raw vegetables and fruits
 - D. Avoid micronutrient supplements
- ANSWER: A
51. A multi gravid mother came to a health facility for ANC visit and received counselling about family planning options after delivery. Her history shows that she was not a fully breast fed due to her job and had pregnancy after 6 weeks of birth in her previous births. Currently, she wants to space giving birth and wants appropriate recommendation from a health professional. What is the most likely necessary counseling in this case?
- A. Lactation amenorrhea method
 - B. Post-partum family planning methods
 - C. Abstinence form sex for the first six months
 - D. Utilization of barrier family planning methods
- ANSWER: B
52. A pregnant mother came to ANC clinic for her third contact at her 32nd week of pregnancy. After history taking and necessary investigation, TT₂ vaccination was given to the woman and scheduled for the next TT vaccination. When is the most appropriate next schedule in this case?
- A. After 1 year
 - B. After due date
 - C. After 4 weeks
 - D. After 6 months
- ANSWER: D

53. A 25-year-old primigravida woman came to a hospital for the first ANC follow up at 16-week of gestation. On assessment, the woman was found to be positive for HIV test with no other pertinent findings. Following counselling, the midwife has decided to put the woman on TDT/3TC/DTG regimen at the first contact and appoints her for viral load assessment.

What is the most appropriate time to evaluate viral load for this case?

- A. One month
- B. Two months
- C. Three months
- D. Four months

ANSWER: C

54. A 25-year-old pregnant mother has come to a hospital OPD with a complaint of lower abdominal pain and mild vaginal bleeding of two days duration at her 8 weeks of gestational age. On physical examination, she has abdominal tenderness and rebound tenderness.

What is the most appropriate history to be asked for this woman?

- A. Frequent urination
- B. Shoulder tip pain
- C. Chest pain
- D. Headache

ANSWER: B

55. A 35-year-old woman at her 10th week of gestational age came to a health facility with a complaint of slight vaginal bleeding and unilateral lower abdominal pain of four-hour duration. She also reported that she felt shoulder tip pain and pain on defecation. Her history indicated that she had pelvic inflammatory diseases before a year.

What is the most likely diagnosis in this case?

- A. Abortion
- B. Ovarian torsion
- C. Ectopic pregnancy
- D. Hydatidiform Mole

ANSWER: C

56. A pregnant woman has come to a hospital with a complaint of lower abdominal pain at a six-week of gestation. On assessment, she is hemodynamically stable, the site of implantation is in fallopian tube, her serum β -hCG level is 4000 mIU/ml, the size of ectopic pregnancy mass is 3 cm, fetal cardiac activity is absent and she is happy to comply the post treatment follow-up.

What is the most appropriate initial management for this woman?

- A. Surgical management
- B. Treat her with Methotrexate
- C. Schedule for a follow-up visit
- D. Secure IV line with large bore cannula

ANSWER: B

57. A 14-year-old girl has come to a clinic with a complaint of pain during menstruation. The pain has been two years since menarche. Her menstrual cycle is monthly regular and moderate in amount. She has no history of sexual contact before.
What is the most appropriate history to be asked for this case?
- A. The onset of pain
 - B. History of oral pills
 - C. The duration of bleeding
 - D. abnormal vaginal discharge
- ANSWER: A
58. A 15-year-old adolescent girl came to a youth clinic with a complaint of delayed menstrual cycle. She had never seen her menstruation like her friends in the same age. On physical examination, she was healthy and well-nourished with well-developed secondary sexual characteristics. Trans-vaginal ultrasound revealed the absence of uterus.
What is the most likely diagnosis?
- A. Amenorrhea
 - B. Hypomenorrhea
 - C. Polymenorrhea
 - D. Oligomenorrhea
- ANSWER: A
59. A 12-year-old girl came to a hospital with a complaint of painful menstruation. She reported that midline cramping pain was started with menses. It became worse between 24 to 36 hours and resolved on the second day of menses. Her menarche had occurred before four months. Pelvic examination showed normal findings.
What is the first line therapy for the girl in this case?
- A. NSAID
 - B. Oral contraceptive pills
 - C. Depot medroxyprogesterone
 - D. Levonorgestrel-releasing IUD
- ANSWER: A
60. A 48-year-old woman has come to the gynaecology OPD of a hospital with a complaint of heavy vaginal bleeding of five months duration. She is feeling dragging pain or pressure in her pelvis.
What is the most likely additional history to help the diagnosis of this case?
- A. Menstrual history
 - B. Circumcision history
 - C. History of last sexual act
 - D. History of age at menarche
- ANSWER: A
61. A 33-year-old woman came to a hospital with a complaint of postcoital bleeding for the last three months. She stated that she had seen slight vaginal bleeding between menstrual cycles and yellowish mucous secretion. On physical examination, a soft, red pedunculated protrusion from the cervical canal at the external os was found. Otherwise, medical and gynaecological findings were normal.
What is the most likely diagnosis in this case?
- A. Cervicitis
 - B. Cervical polyp
 - C. Cervical cancer
 - D. Pedunculated myoma
- ANSWER: B

62. A midwife is monitoring a mother's condition who underwent gynaecologic surgery. The mother had spent 48 postoperative hours in bed, and she stated that it was difficult to walk with urinary catheter. Vital signs revealed blood pressure of 90/60 mmHg, pulse rate of 80 beat/min, respiratory rate of 18 breath/min and temperature of 37.8 degrees Celsius. Physical examination showed calf muscle tenderness, edema, and warmth of lower extremities.
What is the most likely result of the assessment based on this evaluation?
- A. Recovery
 - B. Hypovolemic shock
 - C. Pulmonary embolism
 - D. Deep vein thrombosis
- ANSWER: C
63. An 8-year-old girl was brought to a health facility by her mother with a complaint of early abnormal breast enlargement starting from her seven years.
What is the most appropriate assessment to be considered for this case?
- A. Assessment of breast trauma
 - B. Evaluation of FSH and LH level
 - C. Calculating her body mass index
 - D. Asking family history breast engorgement
- ANSWER: B
64. A 15-year-old adolescent has come to a health center with a complaint of missed menstrual cycle. She has never seen her menses yet. She states that her breast development started at her age four.
What is the most likely diagnosis in this case?
- A. Normal puberty
 - B. Delayed puberty
 - C. Premature puberty
 - D. Precocious puberty
- ANSWER: B
65. A 16-year-old adolescent presented to a hospital with a complaint of cyclic pelvic pain, and she had never seen menses. She reported that she had back pain, difficulty with defecation and urination. On physical examination, inspection of the vulva revealed a purplish-red membrane bulging outward.
What is the most likely definitive management in this case?
- A. Surgical
 - B. NSAIDs
 - C. Antibiotics
 - D. Cryotherapy
- ANSWER: A
66. A 28-year-old woman came to a hospital with a complaint of foul-smelling vaginal discharge of seven days duration. She was treated for pelvic inflammatory disease two weeks back.
What is the most appropriate examination to be performed for this case?
- A. Identify the colour of the discharge
 - B. Check for previous abdominal scar
 - C. Assess for swelling of the vulva
 - D. Determine hormonal level
- ANSWER: A

67. A 25-year-old woman presented to a health center with complaints of lower abdominal pain and foul-smelling vaginal discharge. She also stated that she had backache, nausea and vomiting. On assessment, she was febrile to touch; she had abdominal rebound tenderness; cervical motion tenderness; and palpable adnexal mass.
What is the most likely diagnosis?
A. Cervicitis
B. Peritonitis
C. Appendicitis
D. Pelvic inflammatory disease
ANSWER: D
68. A midwife on duty has been giving a care for a woman with first degree uterine prolapse. The midwife has planned to limit uterine prolapse progression and alleviate symptoms without obstructing normal discharge.
What is the most appropriate non-surgical management in this case?
A. Sit's bath
B. Pelvic rest
C. Kegel exercise
D. Vaginal pessary
ANSWER: C
69. A 20-year-old woman has been admitted to a clinic for post abortion care service. The Midwife has started to collect a history.
What is the most appropriate history to be collected in this case?
A. Last menstrual history
B. Past abortion history
C. Past Medical history
D. Past Surgical history
ANSWER: A
70. A 28-year-old woman has come to a clinic with a complaint of heavy clotted vaginal bleeding of two days duration at her 16th week of gestation.
What is the most appropriate assessment to be performed for this case?
A. Epigastric tenderness
B. Cervical dilatation
C. Vulval swelling
D. Perineal scar
ANSWER: B
71. A 19-year-old adolescent presented to a health center with complaints of lower abdominal cramping and slight vaginal bleeding of two hours duration. She was amenorrhic for the last four months. On physical examination, there was partial expulsion of products of conception through the cervix and the uterus size was smaller than expected date.
What is the most likely diagnosis?
A. Abortion
B. Molar pregnancy
C. Ruptured ectopic pregnancy
D. Unruptured ectopic pregnancy
ANSWER: A

72. A pregnant woman presented to a hospital with complaints of light vaginal bleeding and abdominal cramping at her 15th week of gestation. Physical examination revealed that the uterus corresponds to dates with closed cervix.

What is the most likely type of abortion in this case?

- A. Threatened
- B. Incomplete
- C. Inevitable
- D. Complete

ANSWER: A

73. A midwife on duty was following a 19-year-old married woman on her first day of post abortion. She wanted to become pregnant soon after having complete abortion. Her medical history indicated that she had taken TT2 vaccine before a month. She was screened for cervical lesion nine-months ago.

What is the most appropriate counselling to be considered for this case?

- A. Family planning
- B. Treatment for STD
- C. Tetanus prophylaxis
- D. Cervical cancer screening

ANSWER: A

74. A midwife was caring for a 19-year-old married adolescent on her 7th day of post abortion. She was using intrauterine contraceptive device. Her medical history indicated that she had taken TT1 vaccine a month before, and she was screened for cervical lesion 9 months before. Otherwise, medical investigation showed normal findings.

What is the most important service to be given in this case?

- A. Tetanus prophylaxis
- B. Cervical cancer screening
- C. HIV testing and counselling
- D. Treatment for sexually transmitted diseases

ANSWER: A

75. A woman has been admitted to a hospital with a complaint of continuous urine leakage from the vagina after two weeks of caesarean delivery.

What is the most appropriate assessment to be performed for this case?

- A. Anterior vaginal wall
- B. Caesarean wound site
- C. Abdominal distension
- D. Per-rectal examination

ANSWER: A

76. A 16-year-old nulliparous woman had obstructed labor and gave birth to a 3.2 kg alive neonate seven days ago. On postpartum follow up, she complained urinary incontinence but no urge to urinate for the last 36 hours. The midwife decided to do three swab tests since it was difficult to outline by the naked eye. Assessment finding showed that upper and lower swabs remained dry but the middle stained with dye.

What is the most likely diagnosis in this scenario?

- A. Urethrovaginal fistula
- B. Ureterovaginal fistula
- C. Vesicovaginal fistula
- D. Rectovaginal fistula

ANSWER: C

77. A woman with a surgical treatment for uterine mass has been admitted to gynaecology ward. On the second day after surgery, it is planned to provide wound care.
What is the most appropriate technique to be considered for this case?
- A. Aseptic precautions
 - B. Applying pressure
 - C. Immobilization
 - D. Positioning
- ANSWER: A
78. A midwife on duty was providing preoperative care for a woman with obstructed labour. A midwife planned to insert a bladder catheter with a better view of the urinary meatus and reducing the risk of catheter contamination.
What is the most appropriate position to insert the catheter in this case?
- A. Supine
 - B. Knee chest
 - C. Dorsal recumbent
 - D. Lateral with the right leg flexed
- ANSWER: C
79. A midwife on duty planned to insert a nasogastric tube for an infant with a difficulty of swallowing. A midwife wanted to determine the length of the plastic tube before insertion.
What is the most important land mark that determines how far to insert NG tube for this case?
- A. From the tip of nose to the tip of the ear lobe and then to the tip of the sternum
 - B. From the nose to the tip of the ear lobe and then to the xiphoid process
 - C. From the tip of upper lip to the tip of the ear lobe and then to the umbilicus
 - D. From the nose to the tip of the ear lobe and then to the left nipple
- ANSWER: B
80. A 35-year-old patient presented to a health center with a complaint of bleeding from wound site previously dressed at nearby health post. A midwife on duty assessed the wound and diagnosed as septic wound.
For what purpose is the wound care most likely important in this case?
- A. To apply pressure
 - B. To keep wound clean
 - C. Aids to prevent haematoma
 - D. Prevent pain, swelling and injury
- ANSWER: D
81. A pregnant woman has been brought by ambulance to a hospital. On arrival to the labor ward, she starts to convulse. The midwife prevents her from falling and performs rapid assessment. The woman is unconscious and no respiration effort identified.
What is the most appropriate initial management for this woman?
- A. Initiate fluid resuscitation
 - B. Bag and mask ventilation
 - C. Administer 4-6L nasal oxygen
 - D. Provide Loading dose of MgSo₄
- ANSWER: B

82. A midwife on duty accidentally grabs the radiant warmer with hand. The midwife calls for colleagues to help out with the burn injury.

What is the most appropriate initial management for the burn accident?

- A. Apply ice pack
- B. Apply ointments
- C. Flush with cool water
- D. Apply antiseptic solution

ANSWER: C

83. A 28-year-old newly married woman came to a hospital for preconception care. Her history revealed that she had repeated episodes of malarial attacks, felt fatigue, and had loss of appetite. She had pallor conjunctiva. Laboratory result included haemoglobin of 9 gm/dl, red blood cells count of 2million/mm³, Packed cell volume (PCV) 30 percent (normal value 39-42 percent), MCHC of 30 percent (normal value 32-36 percent), MCV of 75cubic micrometre (normal value 75-100 cubic micrometre), and MCH of 23pg (normal value 27-32 pg).

What is the most likely assessment of this patient?

- A. Iron deficiency
- B. Protein deficiency
- C. Folic acid deficiency
- D. Vitamin B12 deficiency

ANSWER: A

84. A 23-year-old woman came to a gynaecologic OPD with a complaint of forced sexual intercourse of two days duration. She feels depressed, has lack of appetite and disturbance of sleep pattern. The midwife requested for collecting semen for forensic evidence.

What is the most likely history to be identified for the laboratory investigation?

- A. The victim has a history of STIs
- B. The victim had sexual partners
- C. The victim is not sexually active
- D. The victim had used contraceptives

ANSWER: C

85. A 23-year-old woman came to a gynaecologic outpatient department claiming being raped two days back. She reported that she has been slapped, bitten and punched in the face. The midwife continued with the physical examination.

What is the most pertinent finding to be examined for this case?

- A. Abscess formation
- B. Healing recent scar
- C. Signs of pregnancy
- D. Marks of restraints

ANSWER: D

86. A 22-year-old para II mother came to a clinic for the preconception care. She had a previous history of IUGR, intrauterine death and premature birth. On physical examination, her height for age ratio was low, otherwise she had normal examination.

What is the most appropriate interpretation of her nutritional result?

- A. Chronically malnourished
- B. Acutely malnourished
- C. Adequately nourished
- D. Over nourished

ANSWER: A

87. A midwife was interviewing an elderly woman during community service. The woman emphasized that girls' virginity should be ensured and the future of a woman has to be secure, with a lot of children.

What is the most likely harmful traditional practice the woman was advocating?

- A. Early marriage
- B. Trafficking in women
- C. Marriage by abduction
- D. Female genital mutilation

ANSWER: A

88. A 28-year-old woman taking antihypertensive drug came to a preconception care unit. Her history revealed that she had been on antihypertensive medication for two years and it was poorly controlled. The midwife reviewed her medical record to develop care plan.

What is the most appropriate care plan for this woman?

- A. Counselling on exercise
- B. Optimize cardiac function
- C. Switch antihypertensive drug
- D. Refer for further monitoring

ANSWER: D

89. A 22-year-old woman came to STI clinic with a chief complaint of vaginal discharge of seven days duration. On history, she expressed the discharge as thin, homogenous whitish color with fishy odor. Her physical examination reveals dry congestion of the vulva with discharge. On speculum examination redness, contact bleeding from the cervix and endocervical discharge.

What is the most appropriate management for this woman?

- A. Ceftriaxone, Spectinomycin, Metronidazole and Clotrimazole
- B. Ceftriaxone, Azithromycin, Metronidazole and Clotrimazole
- C. Spectinomycin, Azithromycin, Doxycycline, and Metronidazole
- D. Spectinomycin, Azithromycin, Doxycycline, and Metronidazole

ANSWER: B

90. A 25-year-old para II woman has come to a health centre for preconception counselling. Her history reveals that she has been treated for TB for the past one month, and she was also treated for TB five years ago. On physical examination, she has no abnormal findings and is clinically well.

What is the most likely appropriate time for this woman to get pregnant?

- A. Right away
- B. After one month
- C. After three months
- D. After five months

ANSWER: D

91. A 22-year-old house wife came to a health center for family planning service. During exploration she revealed that she has planned to delay her next pregnancy for one year, and wanted to use combined oral contraceptives.

What is the most likely history that makes her eligible to use combined oral contraceptives?

- A. Stroke
- B. Tuberculosis
- C. Breast cancer
- D. Deep vein thrombosis

ANSWER: C

92. A midwife is counselling a 35-year-old mother on her six-week of postpartum period for family planning. Her history reveals that she is breastfeeding and wants to delay a pregnancy for at least five years. She is taking phenytoin for convulsion. The midwife has helped her to choose IUCD. What is the most important history for the recommendation of IUCD?

- A. Age
- B. Breastfeeding
- C. Medical history
- D. Postpartum period

ANSWER: C

93. A 25-year-old para-I mother on her six-day postpartum has come to a health center to use combined oral contraception (COC). Her history reveals that she is exclusively breastfeeding, is on antihypertensive medication for the last three years, and her mother was diagnosed for breast cancer. On physical examination, her blood pressure is 100/70 mmHg, and BMI is 22 kg/m². Based on this assessment, the midwife has decided that the woman is not eligible for COC. What is the most likely history that makes this mother ineligible for COC?

- A. Breastfeeding practice
- B. Reproductive history
- C. Medical history
- D. Family history

ANSWER: A

94. A 20-year-old primiparous woman on her 8th day of postpartum came to a family planning unit for counselling. During counselling, the woman expressed her need of non-hormonal temporary contraception method. The midwife recommended her to use lactational amenorrhea method. What is the most important history to recommend this method?

- A. Treatment of sexually transmitted infections
- B. Regularity of menstrual cycle
- C. Breastfeeding Practice
- D. Fertility awareness

ANSWER: C

95. A newly married couple came to a health centre for family planning service. During exploration, the couple revealed that they wanted a temporary backup method which would not be removed immediately after ejaculation.

What is the most likely barrier method recommended for this couple?

- A. Diaphragm
- B. Cervical cup
- C. Male condom
- D. Female condom

ANSWER: D

96. A 23-year-old primiparous mother on her three-week of postpartum has come to a health center for family planning service. Her history reveals that she had a stillbirth in the last pregnancy and wants to wait for one year till next pregnancy. She has epilepsy and has been on medication for the last two years. On physical examination the BMI is 20kg/m^2 .

What is the most likely contraceptive method this woman is eligible for?

- A. Progestogen-only pills
- B. Fertility awareness methods
- C. Combined oral contraception
- D. Intrauterine contraceptive device

ANSWER: C

97. A 21-year-old newly married woman came to a health centre for family planning service. During counselling, the woman expressed her need for standard day method. The midwife has continued the counselling focusing on the method.

What is the most appropriate counselling component to be included for the method?

- A. Following cervical section
- B. Menstrual cycle regularity
- C. Recording body temperature
- D. Discussing the consequences of each method

ANSWER: B

98. A 38-year-old woman has come to a family planning unit wanting to space her pregnancy for three years. Her history reveals that she smokes more than 15 cigarettes a day and wants to use hormonal contraceptives.

What is the most appropriate option method for this mother?

- A. Combined oral contraceptive
- B. Progestin only pills
- C. Copper based IUD
- D. Combined Patch

ANSWER: B

99. A 38-year-old woman and her husband came to a family planning unit for counselling. They worried about long-acting contraception might be hormonal, might need frequent visit, too expensive and might have long-term side effects. After exploration, the couple decided to use implanon.

What is the most important counselling component addressed in this case?

- A. The efficacy of the method
- B. The provider of the service
- C. The availability of the method
- D. The side effects of the method

ANSWER: D

100. A 47-year-old Gravida-V Para-III woman at her 40th week of gestation has come to a hospital to be admitted for elective caesarean delivery. During counselling, the woman had revealed that she does not want to have any more children and wants to switch from injectable contraception. The woman had previous caesarean delivery and a history of ectopic pregnancy. The midwife has counselled her for family planning methods.

What is the most appropriate contraceptive method for this woman?

- A. Implant
- B. Post placental IUD
- C. Combined injectable
- D. Bilateral tubal ligation

ANSWER: D

101. A 22-year-old woman came to a health centre for family planning service. On exploration, she expressed that she had never used contraception, and has heard about combined contraceptives pills side effects.

What is the most likely first step in the counseling of this woman?

- A. Possibility of switching to another method
- B. Address concerns through counselling
- C. The medical eligibility criteria
- D. The clinical management

ANSWER: B

102. A 22-year-old man came to family planning unit for condom refill. The midwife evaluated the proper utilization of male condom before refilling.

What is the most appropriate component in the utilization of condom?

- A. Reuse of condom
- B. Use in every sexual act
- C. Use of oil base lubrication
- D. Using with female condom

ANSWER: B

103. A 32-year-old para-III mother on her six-week of postpartum came to health centre for her third routine postnatal follow up. After reviewing her record the midwife took history regarding how she feels, her breastfeeding practice, vaginal bleeding and discharge, and fever.

What is the most likely prior history the midwife has missed?

- A. Danger sign
- B. Menstrual flow
- C. Family planning choice
- D. Complications during delivery

ANSWER: C

104. A 26-year-old primiparous mother on her third-day of postpartum came to a health center for follow up. She had no complaint except she has mood swings. She was using four sanitary pads per day, had breast pain which relieved with feeding. On physical examination; vital signs were stable, and had lower extremities oedema. After evaluation, the midwife has planned to admit the woman.

What is the most likely danger sign the midwife identified in this case?

- A. Breast pain
- B. Mood swing
- C. Leg swelling
- D. Vaginal bleeding

ANSWER: C

105. A 34-year-old HIV positive mother came to a health centre for follow up of her nine-month-old infant. On assessment, it was found that the infant had stopped breastfeeding after six months. The DNA/PCR test result came non-reactive.
What is the most likely HIV status of this infant?
A. HIV positive
B. HIV negative
C. HIV exposed
D. HIV unknown status
ANSWER: B
106. A 25-year-old woman on her third day of postpartum has been admitted to a postnatal unit after caesarean delivery. On planning for her discharge, the midwife has reviewed her record and found that she lived in malaria endemic area and had one Antenatal care visit. Laboratory finding indicates haematocrit of 33 percent.
What is the most appropriate preventive and promoting care for this woman?
A. Provide Tetanus toxoid vaccine
B. Advice to sleep under a bed net
C. Order Iron folate 2 tablet per day
D. Treat with broad-spectrum anti-helminthic
ANSWER: B
107. A 30-year-old para III mother on her third day of postoperative after caesarean delivery is planned to be discharged. Her record shows, she lives in highland area and she was treated with MgSo₄. Currently, she has stable vital signs, and BMI of 30 kg/m².
What is the most important counseling component that has to be included for this woman?
A. Use of bed net
B. Physical exercise
C. High calcium diet
D. Restricted salt intake
ANSWER: B
108. A 32-year-old para II mother on her seven-day postpartum came to a hospital for follow up. Upon evaluation, the midwife identified, firm uterus palpated midway between umbilicus and symphysis pubis.
What is the most possible explanation for the evaluation of this woman?
A. Uterine infection
B. Uterine Involution
C. Uterine decompression
D. Uterine Sub-involution
ANSWER: B
109. A 30-year-old para II mother has come to a hospital for follow up after six months of taking ART. Her history shows that she was diagnosed with HIV and started ART at labor. During evaluation, she is on WHO stage 3, has pulmonary TB, CD4 levels of 200 cells/mm³ and Plasma viral load of 4000 copies/ml.
What is the possible explanation for the treatment progress of this woman?
A. Viral failure
B. Clinical failure
C. Adherence failure
D. Immunological failure
ANSWER: B

110. A 25-year-old primiparous mother on her third-day of postpartum has come to a hospital with a complaint of excessive vaginal bleeding of one day duration. Physical examination result shows blood pressure of 90/50mmHg, pulse rate of 120 beat/min, profuse vaginal bleeding, sub-involute tender uterus, and no perineal tear observed.

What is the most appropriate next step in the assessment of this woman?

- A. Perform Ultrasound scan
- B. Perform bed side clotting test
- C. Perform speculum examination
- D. Perform manual exploration of uterus

ANSWER: A

111. A 35-year-old para III woman has come to a hospital with a complaint of fever of one day duration. On physical examination, blood pressure of 80/40mmHg, pulse rate of 125 beat/min, temperature of 38.5 degree Celsius are recorded. Laboratory investigation shows WBC level of 12,000 (normal level- 4,000-11,000), Haemoglobin level of 11.5gm/dl and urine ketone-negative, fibrinogen level of 500gm/dl (normal value 444-670gm/dl). The midwife has diagnosed the woman with septic shock.

What is the most important assessment that led the midwife to diagnose septic shock?

- A. Urine ketone
- B. WBC count
- C. Fibrinogen level
- D. Haemoglobin level

ANSWER: B

112. A 32-year-old primiparous mother has been admitted to postnatal unit with a diagnosis of shock. Her history reveals that she gave birth before two weeks at home. Physical examination results show blood pressure of 80/40mmHg, pulse rate of 125 beats/min, temperature of 38.5-degrees Celsius, tender uterus, minimal vaginal bleeding, and granulated perineal tear.

What is the most likely cause of shock in this case?

- A. Trauma
- B. Infection
- C. Haemorrhage
- D. Anaphylactic

ANSWER: B

113. A 24-year-old primiparous mother on her third-day of postpartum came to a hospital with a complaint of breast pain of one day duration. She has started breastfeeding. Physical examination results show body temperature of 38.1 degrees Celsius, pulse rate of 120 beats/min, and bilateral swollen breasts, but no palpable mass noticed.

What is the most likely diagnosis of this woman?

- A. Breast engorgement
- B. Breast abscess
- C. Breast cancer
- D. Mastitis

ANSWER: A

114. A 34-year-old para III mother on her seven-day of postpartum has come to a health center with a complaint of severe headache. She has no history of chronic disease. Physical examination reveals blood pressure of 150/90mmHg, and Pulse rate of 100 beat/min. Laboratory investigation shows +2 urine protein.

What is the most likely diagnosis for this woman?

- A. Mild Pre-eclampsia
- B. Postpartum pre-eclampsia
- C. Pre-eclampsia with severity feature
- D. Superimposed pre-eclampsia without severe features

ANSWER: C

115. A 23-year-old primiparous mother on her seven-day postpartum came to a health centre with a complaint of fever. Her history revealed that she was on antimicrobial therapy for the last four days for fever and had delivered with caesarean section. She also claimed to have chill and no other symptoms. Her temperature was 38.4 degrees Celsius.

What is the most likely diagnosis of this woman?

- A. Surgical site infection
- B. Parametrial phlegmon
- C. Persistent endometritis
- D. Septic thrombophlebitis

ANSWER: D

116. A 20-year-old primiparous mother has been transferred to a postnatal ward after caesarean delivery. Up on evaluation, a midwife has identified a blood pressure of 80/40 mmHg, pulse rate of 110 beat/min, boggy uterus and cold extremities.

What is the most likely diagnosis of this woman?

- A. Septic shock
- B. Cardiogenic shock
- C. Anaphylactic shock
- D. Hypovolemic shock

ANSWER: D

117. A team of midwives were managing a primiparous woman with vaginal bleeding secondary to uterine atony. They provided uterotonic agents, massage the uterus, applied bimanual uterine compression and aortic compression. Despite all these management, the bleeding was not controlled.

What is the most appropriate next management for this woman?

- A. Hysterectomy
- B. Uterine artery ligation
- C. Uterine balloon tamponade
- D. Uterine compression sutures

ANSWER: C

118. A multiparous mother has come to a hospital for postnatal follow up on her seven-day postpartum. Physical examination reveals a blood pressure of 170/110mmHg, pulse rate of 100 beat/minute and well contracted uterus. Urine analysis result indicates + 2 proteinuria.

What is the most appropriate management for this woman?

- A. Schedule for weekly follow up
- B. Admit and provide anticonvulsant drug
- C. Send home with PO antihypertensive drug
- D. Repeat BP measurement and urine analysis after 4 hours

ANSWER: B

119. A 35-year-old para III mother has been brought to a hospital with ambulance after 30 minutes of home delivery. Physical examination reveals blood pressure of 80/40mmHg, pulse rate of 120 beat/minute, cold clammy skin, boggy uterus and excessive vaginal bleeding. The midwife has mobilized the team and put her on shock position.

What is the most appropriate next step in the management of this woman?

- A. Start an IV infusion
- B. Catheterize the bladder
- C. Identify possible cause
- D. Send laboratory investigation

ANSWER: A

120. A 28-year-old woman is on her third-day of admission to a clinic with a diagnosis of endometritis. She is on IV Clindamycin phosphate and Gentamycin regimen. Her record indicates she was fever free for the last 48 hours. Current evaluation shows blood pressure of 110/90mmHg, and temperature of 36.6 degree Celsius.

What is the most appropriate next step in the management of this woman?

- A. Shift to oral Ampicillin and Metronidazole
- B. Discontinue IV Clindamycin phosphate and Gentamycin
- C. Shift to Ampicillin, Metronidazole, and Gentamycin regimen
- D. Do abdomino-pelvic ultrasound and appoint for weekly follow up

ANSWER: B

121. A 28-year-old woman gave birth at 37 weeks of gestation by spontaneous vaginal delivery in a health institution. During delivery, the neonate cried immediately. The neonate's examination revealed a heart rate of 90 beats/minute, flexion of an extremities, grimace for a reflex, and bluish colour of the whole body.

What is the Apgar score of the neonate?

- A. 3
- B. 4
- C. 5
- D. 6

ANSWER: C

122. A neonate was delivered at 36th week of gestation by caesarean section for an indication of Oligohydramnios. The assessment reveals; irregular respiration, pulse rate of 85 beats/minute, flaccid posture, grimace reflex, pink trunk and blue extremities.

What is the most appropriate Apgar score interpretation for the neonate?

- A. Low
- B. Severe
- C. Normal
- D. Moderate

ANSWER: D

123. A 29-year-old primiparous woman gave birth of a female neonate at 38-week of gestation by vaginal delivery two hours ago. The midwife counselled the mother about the importance of immunization.

What is the most appropriate vaccination to be given for the neonate?

- A. BCG
- B. DPT
- C. OPV-1
- D. Rota virus-1

ANSWER: A

124. A neonate was born at term, weighed 2.7 kg and had Apgar score of 8 at 1 minute of delivery. The neonate fed exclusive breast milk for 1 month. The weight for the current visit showed that the baby weighed 3.5 kg.

What is the most appropriate evaluation of the feeding status of the neonate?

- A. Difficult to decide
- B. Inappropriate feeding
- C. Adequate breast feeding
- D. Inadequate breast feeding

ANSWER: C

125. A 28-year-old woman gave birth of an alive neonate at 37-week of gestation at a health centre. The baby was brought back by the age of six-month, the physician performs physical examination to assess the neonatal reflex.

What is the most appropriate reflex assessment in this case?

- A. Moro reflex
- B. Rooting reflex
- C. Babinski reflex
- D. Grasping reflex

ANSWER: C

126. A 23-year-old woman gave birth to an alive neonate at 35-week of gestational age in a health center. Assessment results showed no respiration, pulse rate of 57 beat/minute, cyanotic, and flaccid posture. Resuscitation was initiated but the chest didn't rise with mask ventilation.

What is the most appropriate next assessment for this patient?

- A. Check for displaced tube
- B. Check patency of the air way
- C. Make sure the head in hyperextended position.
- D. Make sure tightness of the mask over the eye, nose and mouth

ANSWER: B

127. A 44-year-old woman gave birth to 1.8 kg alive neonate at 36-week of gestational age in a health centre. On assessment, the neonate had an upward slant of eyes, epicanthic folds, low set simple ears, flat occiput, short neck, single palmar crease and wide gap between first and second toes.

What is the most likely diagnosis of this case?

- A. Patau syndrome
- B. Down syndrome
- C. Noonan syndrome
- D. Edwards syndrome

ANSWER: B

128. A 28-year-old woman gave birth at 32-week of gestational age at a hospital. Assessment finding showed protrusion of abdominal contents to the right of umbilicus which are not covered with sac. What is the most likely diagnosis of this neonate?
- A. Omphalocele
 - B. Gastroschisis
 - C. Duodenal atresia
 - D. Hirschsprung disease
- ANSWER: B
129. A 28-year-old woman came to a hospital with a complaint of pus discharge from her neonate's eyes. On assessment, she had multiple sexual partners, no regular ANC follow up and gave birth to a neonate at home five days ago. What is the most appropriate management for the neonate?
- A. High dose acyclovir
 - B. Ceftriaxone 50mg/IM stat
 - C. Erythromycin PO for 14 days
 - D. One percent tetracycline ointment QID daily
- ANSWER: B
130. A 28-year-old woman brought her infant to a general hospital with a complaint of coughing and night sweating that started 15 days ago. On assessment, the infant had contact with known TB patient. What is the most appropriate diagnosis for the infant based on IMNCI?
- A. TB disease
 - B. TB infection
 - C. No TB infection
 - D. Difficult to rule out
- ANSWER: A
131. A woman who lived at high malaria risk area has brought her six-month-old infant to a health centre with a complaint of fever. Neonatal assessment shows axillary body temperature of 38-degree Celsius, poor sucking reflex, and stiff neck. What is the most appropriate management before referring the infant?
- A. Quinine
 - B. Paracetamol
 - C. Chloroquine
 - D. Oral coarthem
- ANSWER: A
132. A 24-year-old woman has brought her 10-week infant to a health centre for immunization. The infant has taken all vaccines expected to be taken for his/her age. What is the most appropriate vaccine for the infant at this age?
- A. BCG
 - B. OPV-2
 - C. Measles
 - D. Hepatitis B1
- ANSWER: B

133. A midwife working in a tertiary hospital admitted a labouring mother with breech presentation. He assisted the patient during spontaneous vaginal delivery with good outcome by using moral judgement.
What is the most likely professional ethics practiced in this case?
A. Teleology
B. Deontology
C. Intuitionism
D. Ethic of caring
ANSWER: A
134. A 22-year-old pregnant woman at her 6-week gestational age presented to a health centre with excessive vomiting of 3 days duration. Even though the midwife on duty tried to administer medication to stabilize her condition, the woman refused the treatment.
What is the most likely ethical principle employed by the midwife?
A. Justice
B. Autonomy
C. Beneficence
D. Non-maleficence
ANSWER: C
135. A labouring woman of 41-week gestational age presented to a general hospital with a complaint of pushing down pain. A midwife on duty diagnosed non-reassuring foetal heart rate and absent amniotic fluid which needs emergency caesarean section to save the life of the foetus.
What is the most likely factor that could influences woman's decision making in this case?
A. Urgency
B. Anxiety
C. Family
D. Age
ANSWER: A
136. A 22-year-old Primigravida woman came to a health centre for first antenatal care visit. Her name was similar with another patient appointed for family planning. Consequently, she was injected with Depo-Provera.
What is the most likely medico-legal issue occurred in this case?
A. Medical Competence
B. Medical Malpractice
C. Medico-Negligence
D. Medication error
ANSWER: D
137. A two-year-old female child was brought to health centre by her mother with excessive vomiting and bloody diarrhoea of six hours duration. A midwife on duty was watching football match in the duty room heard child crying. Then, the midwife discontinued watching the football match and curiously treated the child compassionately.
What is the most likely quality of compassionate care implemented in this case?
A. Motivation
B. Sympathy
C. Empathy
D. Judge
ANSWER: C

138. A 34-year-old multigravida labouring woman of 40-week gestational age presents to a health centre with complaints of pushing down pain and ruptured membrane of six-hour duration. A junior male midwife on duty admitted the woman to the facility. The woman informed the ward head that the midwife didn't demonstrate professional behaviour like other midwives. She, nonetheless, witnessed that he was punctual, hardworking and motivated during her stay. He was arriving late, looking depressed, tired and irritable.

What is the most likely action to overcome the threat in this case?

- A. Exercising in the club
- B. Taking to disciplinary
- C. Training of compassion
- D. Changing his profession

ANSWER: C

139. A 22-year-old labouring woman was brought to a nearby hospital by her mother-in law with complaints of pushing down pain and slight vaginal bleeding for the past 14 hours. The midwife on duty informed the woman to wait for another night duty midwife since the car was waiting to take her home.

What is the most likely disrespectful character practiced by the midwife in this case?

- A. Non-consented care
- B. Detention in facilities
- C. Abandonment of care
- D. Discrimination in facilities

ANSWER: C

140. A medical director of a rural health centre has assigned a nurse and a midwife during the night shift program, and a public health officer for a day time due to shortage of budget. A public health officer assigned for day time refused to work in maternal services as it was not her competency.

What is the most likely conflict of interest reflected in this case?

- A. Salary scale
- B. Shortage of budget
- C. Facility management
- D. Professional practice

ANSWER: D

141. A 22-year-old pregnant woman of 12-week gestational age has come to a health centre with excessive nausea and vomiting of 24 hours duration. Since the pregnancy is wanted and planned, she needs coping mechanism with the symptoms to have a healthy baby with her husband.

What is the most likely key step in establishing trust with this patient?

- A. Entertaining
- B. Information
- C. Appointment
- D. Communication

ANSWER: D

142. A 30-year-old woman presented to a general hospital from a rural health centre with a referral paper diagnosis of breech presentation. The midwife on duty found that the foetal heart rate was absent and there was excessive vaginal bleeding. The midwife asked for obstetric ultrasound and confirmed the foetus was not alive. The midwife revealed this finding to the woman. As a result, the woman cried and became not responsive.

What is the most appropriate issue to be considered when breaking bad news in this case?

- A. Age
- B. Setup
- C. Emotion
- D. Perception

ANSWER: C

143. A 26-day-old male neonate who was diagnosed for late neonatal sepsis was brought by his mother to a district hospital. A midwife on duty admitted the neonate and communicated with his colleagues regarding the treatment guideline and protocol in the facility. One of the colleagues informed him that the guideline recommends referral to a neonatologist, and suggested a tertiary hospital in their catchment area.

Why the communication within and across health care provider is needed in this scenario?

- A. To get acceptance
- B. To fulfil settled criteria
- C. For the better outcome
- D. For the need of the patient

ANSWER: C

144. A 20-year-old woman presented to a nearby health centre with a complaint of pain during sexual intercourse after her marriage of three months. A male midwife on duty started to ask medical information but the woman refused to communicate with him.

What is the most likely communication barrier faced in this case?

- A. Language
- B. Husband
- C. Age
- D. Sex

ANSWER: D

145. A researcher planned to explore the impact of religious beliefs on female genital mutilation. He collected data for his research through observation from the practices in the community. After data analysis, the researcher declared that community religious belief is associated with female genital mutilation significantly.

What type of research is most appropriate in the above scenario?

- A. Action research
- B. Applied research
- C. Qualitative research
- D. Quantitative research

ANSWER: C

146. A researcher wanted to conduct a study on the relationship between per capital salt consumption and magnitude of hypertension. Hence, he collected data on annual salt consumption from the entire population of two different districts and compared the proportion of hypertension.

What is the most appropriate study design to be used by the researcher?

- A. Ecological
- B. Case series
- C. Case report
- D. Cross-sectional

ANSWER: A

147. A midwife planned to conduct a study on the effect of health education during pregnancy to prevent low birth weight. She intended to conduct the research on 440 women who are in their antenatal care follow-up out of which 220 got health education and the rest did not receive.

What is the most appropriate study design for the above scenario?

- A. Cohort
- B. Case-control
- C. Interventional
- D. Cross-sectional

ANSWER: C

148. A gynaecologist planned to conduct a study on the problems experienced by pregnant women who had antenatal care follow-up at a hospital. The gynaecologist invited six pregnant women from both outpatient and inpatient departments to discuss and reach agreement on the problems they faced.

What is the most appropriate research type to realize this objective?

- A. Action
- B. Applied
- C. Qualitative
- D. Quantitative

ANSWER: C

149. A student planned to conduct a study for the partial fulfilment of the requirements of his degree. Hence, the student reviewed related literatures continuously and found additional burden and the severity problem of the research that was used for the student to rewrite the proposal.

What is the importance of rewriting the proposal in this case?

- A. To prevent duplication of the study
- B. To refine the statement of the problem
- C. To be familiar with different research method
- D. To get confidence because the research conducted

ANSWER: B

150. A midwife conducted a study to assess the prevalence of cervical cancer and its associated factors among reproductive age group. The midwife determined a sample size of 380 and drew 190 samples from rural and the remaining 190 samples from urban residents.

Which sampling method was used in the above scenario?

- A. Multistage
- B. Simple random
- C. Systematic random
- D. Stratified random

ANSWER: D

151. Midwifery students were evaluating the method of teaching of a lecturer for the 2016 E.C academic year. The students were requested to provide their level of satisfaction on the teaching method of the lecturer simultaneously. Then after, the students filled out and returned the documents to the one who requested them.

Which type of data collection tool was used in this case?

- A. Questionnaire
- B. Interview Guide
- C. Observation check-list
- D. Focus Group Discussion Guide

ANSWER: A

152. A researcher conducted a study on the factors associated with anaemia among school children. He collected data, analysed, and declared the results; but later another researcher conducted the study and found more valid results using larger sample size.

What is the most likely error committed by the first researcher in this case?

- A. Bias
- B. Chance
- C. Confound
- D. Negligence

ANSWER: B

153. A patient who was admitted had no progress in the morning round and a gynaecologist on duty ordered a new work-up but did not get any relevant finding. Finally, the gynaecologist decided to understand the patient's interest on the treatment to contrast with what he had in mind about the patient for further intervention.

Which component of evidence-based practice was applied in this case?

- A. Best and latest evidence
- B. Patient preference and value
- C. Clinical expertise and assessment
- D. Knowledge of psychosocial issues

ANSWER: B

154. A researcher understood why all pregnant women did not avail themselves for antenatal care follow-up at a nearby healthcare facility after gathering information from invited guests. During the data collection, the researcher assigned women representative to facilitate, and youth representative to record their discussion.

Which data collection method was used in this case?

- A. Observation
- B. Face-to-face interview
- C. Focus group discussions
- D. Using available information

ANSWER: C

155. A researcher planned to conduct a study on the timely breastfeeding initiation and associated factors among mothers who gave birth in a district. The midwife on duty checked to have consistent understanding between her and the mothers on the content of the questionnaire before the actual data collection.

Which data quality assurance technique was applied in this case?

- A. Providing training
- B. Conducting pre-test
- C. Performing supervision
- D. Checking data completeness

ANSWER: B

156. An Epidemiologist would like to assess the prevalence and associated factors of urinary tract infection among reproductive age group women. The Epidemiologist determined the presence or absence of the disease after urine sample was drawn via catheterization for laboratory examination, but five women experienced urinary tract injury caused by the sample collectors.

Which research ethical principle was violated in this case?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-Maleficence

ANSWER: D

157. A researcher is interested to assess maternal health need via face-to-face interview among key informants in a district. Then the researcher identified the problems and made conclusion for the whole communities after processing and analysing the data.

Which research ethical principle was violated in this case?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-Maleficence

ANSWER: A

158. A midwife who works in a hospital recorded a weight of 10 pregnant women who visited the hospital for ANC follow-up. The weights of the women were 56, 60, 49, 70, 64, 78, 50, 80, 66, and 74 in kilogram.

Calculate the median weight of the pregnant women who visited the hospital?

- A. 65
- B. 50
- C. 60
- D. 70

ANSWER: D

159. A graduate student conducted a study to know the knowledge, attitude, and practices of food choices among pregnant women who were attending ANC follow up in a hospital. The age of the study participants was 18, 59, 24, 42, 21, 23, 24, and 32 years. Calculate the interquartile range (IQR) age of the women?

A. 39.5
B. 24.0
C. 21.5
D. 18.0

ANSWER: D

160. A researcher conducted a study to know the prevalence of under nutrition among pregnant women attending ANC clinic in a health center. The investigator used mid upper arm circumference with the corresponding records of 21.2, 21.4, 21.6, 21.8, and 21.5.

What is the most appropriate measure of central tendency in this scenario?

A. Mean
B. Mode
C. Median
D. Geometric mean

ANSWER: A

161. A researcher conducted a study on college students to know the amount of time they spend for physical exercises. The researcher considered the data on time (in hours) that 100 college students devoted to leisure activities during a typical school week.

What is the most appropriate method of data presentation in this scenario?

A. Bar graph
B. Line graph
C. Histogram
D. Frequency polygon

ANSWER: C

162. A district health office conducted a survey to measure the death rate occurred in population specific to age groups of zero to four years, where the total number of populations of interest was 84,000 and the number of deaths specific to this age group was 1,000.

Calculate age-specific-death rate in this case?

A. 9.0
B. 16.0
C. 10.0
D. 11.9

ANSWER: D

163. A comprehensive specialized hospital conducted annual review meeting of the 2020 fiscal year specific to maternal and child health activities. The chief of the hospital opened the meeting recognizing that the number of mothers who gave live birth was 81453 which is the largest number ever. The coordinator of the MCH unit presented reports on each activity. From the report, the number of fetal deaths was 124.
Calculate fetal death rate in this scenario?
A. Two child deaths per 1000 live births
B. Three child deaths per 1000 live births
C. Four child deaths per 1000 live births
D. Five child deaths per 1000 live births
ANSWER: A
164. A nurse wants to determine the prevalence and associated factors of gestational diabetes mellitus among women admitted to the obstetrics wards of a hospital from September to November, 2024. In the hospital; 10, 12, and 15 cases were treated in September, October, and November respectively among the total of 200 women admitted.
Calculate the prevalence of gestational diabetes mellitus in September in this case?
A. 5 percent
B. 12 percent
C. 10 percent
D. 15 percent
ANSWER: A
165. An epidemiologist wants to conduct a study on short term clinical and pharmaceutical outcomes of pharmaceutical care services and associated factors among diabetes patients at community pharmacy level. The variable of interest that an epidemiologist wants to test is categorical.
What is the most appropriate test-statistics to be used in this case?
A. T-test
B. Chi-square test
C. One way ANOVAs test
D. The fisher-exact test
ANSWER: B
166. A researcher conducted a study to see the association between cigarette smoking during pregnancy and severe acute malnutrition status (SAM) among under five-year-old children. The reported odd ratio by the researcher was 9.62.
What is the most appropriate interpretation of odd ratio by the researcher?
A. There is no association between SAM and cigarette smoking
B. The odds of SAM is low among mothers who smoke cigarette
C. The odds of SAM is high among mothers who smoke cigarette
D. The odds of SAM is high among mothers who did not smoke cigarette
ANSWER: C

167. A head of a midwife in a referral hospital was coaching, assisting and trying to solve problems of her subordinates. Besides, she was influencing the staff members to work hard for the achievement of organizational goals.

What function of management most likely goes in line with this scenario?

- A. Leading
- B. Planning
- C. Organizing
- D. Controlling

ANSWER: A

168. A midwife implemented her operational plan derived from the five years strategic plan. In the meantime, she was continuously checking her achievement against intended goals and taking any corrective actions where necessary to make sure that she was on track.

What most appropriate management function did the midwife follow in this scenario?

- A. Controlling
- B. Organizing
- C. Planning
- D. Leading

ANSWER: A

169. A head of obstetrics ward has developed a spirit of trust, enthusiasm, and genuine involvement with his employees. He has also the capacity to understand the feelings of others in each department for the achievement of the organization's goal.

What is the most likely managerial skill of the manager in this scenario?

- A. Conceptual skill
- B. Technical skill
- C. Tactical skill
- D. Human skill

ANSWER: D

170. A chief executive director in a district hospital prepared a plan. In his plan, he stated more about what his organization intended to be in the future and how it would get there and he tried to show the best path to reach the organization's mission.

What is the most likely type of planning the director prepared in this scenario?

- A. Tactical planning
- B. Strategic planning
- C. Operational planning
- D. Contingency planning

ANSWER: B

171. A team of health care providers in a health facility were discussing about a preparation of strategic plan for their institution. The team leader came with a finding that his health care facility was equipped with sufficient material, financial and trained man power resources.

What strategic planning tool was presented by the team leader in this scenario?

- A. Threat
- B. Strength
- C. Weakness
- D. Opportunity

ANSWER: B

172. A group of midwives are supposed to be trained on shifting medical records to electronic database and are instructed on how they could integrate the data system. As a result, they are using the data for effective problem solving, and informed decision-making.

Which health system building blocks are most likely addressed in this scenario?

- A. Health Financing
- B. Human resource policies and procedures
- C. Management of medicine and health supplies
- D. Health Information and Monitoring or Evaluation

ANSWER: D

173. A senior midwife in a health care facility was stringently working on performance appraisal, training and development, discipline and compensation administration of his subordinates.

What most appropriate human resource management activity goes in line with this scenario?

- A. Selection
- B. Retaining
- C. Orientation
- D. Human resource planning

ANSWER: B

174. A manager at a certain referral hospital is eager to help staff members grow academically. At the same time, he advises them to become responsible and show their initiative while delivering services to their clients.

What leadership style most likely agrees with the above scenario?

- A. Democratic
- B. Dictatorial
- C. Autocratic
- D. Anarchic

ANSWER: A

175. A senior midwife provides poor mentorship and insufficient training for his graduate level midwives due to the hot weather condition. Moreover, conflicts and instabilities affect students' performance especially on their clinical decision making.

What are the most likely factors affecting clinical decision making in this scenario?

- A. Seasonal vibration
- B. Knowledge and skill
- C. Environmental factors
- D. Perception and Personality

ANSWER: C

176. A senior midwife generated health an indicator from HMIS data to decide the importance of the new project for reduction of maternal mortality. Then he has identified the level of political commitment, the allocation of resources, the degree of decentralization, and mechanisms for community participation.

What is the most likely indicator generated by the midwife in this case?

- (A) Mortality indicator
- (B) Population indicator
- (C) Health policies indicator
- (D) Health service coverage indicator

ANSWER: C

177. A midwife is working in a public health facility which provides services for about 25,000 catchment population. The health care facility mainly focuses on disease prevention and health promotion activities on top of the curative services.
What is the most likely category of the health care facility according to the new health tier system in Ethiopia?
A. Health Post
B. Primary hospital
C. Urban Health center
D. Rural health center
ANSWER: D
178. A health team conducting continuous and systematic review of a project activity aiming at checking the progress of the activities against the plan. In the meantime, corrective action has to be taken if any gap is detected during the process.
What is the most likely component of program monitoring and evaluation done in this case?
A. Evaluation
B. Monitoring
C. Coordination
D. Supportive supervision
ANSWER: B
179. In a referral hospital, the management established a system for controlling activities that would be carried out in the hospital, and they did monitor and evaluate periodically to see the achievements. What most likely component of controlling would be done at last by the management?
A. Taking corrective actions
B. Measuring actual performance
C. Comparing performance with the standards
D. Establishing standards to measure performance
ANSWER: A
180. A program manager is dealing with a combination of measurements and judgments that permit conclusions to be made about outcomes or benefits of the program or method.
What is the most appropriate type of evaluation addressed in this case?
A. Impact Evaluation
B. Process Evaluation
C. Formative evaluation
D. Summative Evaluation
ANSWER: D
181. A group of health workforce has been assigned for an outreach activity to figure out the health problems of a particular community in five kebeles. The team has randomly selected two kebeles and has the map.
What is the most likely next step that the team is expected to do?
A. Community Intervention
B. Problem Identification
C. Community survey
D. Prioritization
ANSWER: C

182. A midwife assigned to a health facility has provided health education session on contents of breast milk and feeding for postnatal women. Accordingly, he has emphasized that water is the first most abundant component of breast milk, and water should not be given to a baby during the first six months.

What is the most likely next most abundant component of breast milk in this case?

- A. Lactose
- B. Glucose
- C. Fructose
- D. Glycogen

ANSWER: A

183. Occupational health and safety department provided a half day orientation for new midwifery staff members regarding instrument processing for reusable medical and surgical supplies. Accordingly, the midwives were trained on the steps and procedures of instrument processing.

What is the most likely first step in this case scenario?

- A. Chemical disinfection
- B. Decontamination
- C. Autoclaving
- D. Cleaning

ANSWER: B

184. A 20-year-old female student visited a nearby health facility for abortion service. She was examined and was found to be at the 8-week of gestation. However, the staff on duty had a strong belief that termination of pregnancy was against his religion. He shouted at her not to terminate and rather to deliver it. The student returned to her home and went for a traditional healer to terminate her pregnancy.

What is the most likely predisposing behavioral factor observed in this case?

- A. Norms
- B. Values
- C. Attitude
- D. Knowledge

ANSWER: C

185. A 26-year-old male prisoner from the early correction facility was presented to a medical OPD with a complaint of rash, pruritus and skin lesions of two days duration. He said that he had been with many prisoners in a single cell.

What is the most likely diagnosis in this case?

- A. Scabies
- B. Cholera
- C. Typhus fever
- D. Relapsing fever

ANSWER: A

186. A junior midwife was following a 30-year-old laboring mother in a health centre without partograph chart. The labor poorly progressed and finally the contraction became irregular and weak. The fetal station was high and the cervix was not effaced as expected. He arranged a referral to a hospital but delayed due to lack of transport access. The mother became exhausted and eventually passed away on arrival at the referral site.

What is the most likely cause of death in this case?

- A. Obstructed labor
- B. Preeclampsia
- C. Infection
- D. Abortion

ANSWER: A

187. A 19-year-old couples met at a café out of their village during the evening. They seemed to be in a good mood whispering, hugging, cuddling and kissing each other. Since it was getting dark, they decided to spend the night together and booked a room. She was taking Depo-Provera injection regularly. Their HIV status was non-reactive, but they did not have condom because they were in a hurry.

What is the most likely health problem they might face?

- A. HIV/AIDS
- B. Sexual violence
- C. Teen age pregnancy
- D. Sexually transmitted infections

ANSWER: D

188. A group of investigators wanted to check the validity of cervical cancer screening test. The group has recruited 20 women for the screening test. According to the test result, five of them tested positive with the disease while eight of them were tested negative without the disease; and four were tested positive without the disease whereas three of them were tested negative with disease.

What is the sensitivity of the test in this case?

- A. 0.33
- B. 0.56
- C. 0.63
- D. 0.72

ANSWER: C

189. A regional health bureau received a report of malaria outbreak in a community. Experts conducted rigorous assessments and the final report showed an epidemic curve with successive series of peaks.

What type of outbreak most likely occurred in the community?

- A. Continuous common source
- B. Point source
- C. Propagated
- D. Mixed

ANSWER: C

190. An epidemiologist was invited by the regional health bureau to make a presentation on disease occurrence for the newly recruited staff members of the bureau. In his presentation, epidemiologist emphasized that an epidemic was spreading over several countries or continents, and affecting a large number of people.

What is the most likely type of disease occurrence based on the above case?

- A. Epidemic
- B. Outbreak
- C. Pandemic
- D. Endemic

ANSWER: C

191. A midwife was preparing an action plan for cervical cancer screening program. He held a discussion with woreda health bureau representatives and listed all necessary resources required for implementation.

What is the most likely component of action plan undertaken in this case?

- A. Input identification
- B. Problem identification
- C. Outcome identification
- D. Intervention identification

ANSWER: A

192. A Maternal unit staffs have planned to provide health education on personal hygiene for post natal mothers. As a teaching aid, they want to use a method which can speed up and enhance understanding, stimulate thinking and create opportunities for active learning.

What is the most preferable teaching aid they need to use?

- A. Audio
- B. Visual
- C. Written-word
- D. Direct experience

ANSWER: B

193. A school age child visited a health center with complaints of diarrhoea, vomiting and abdominal cramp. During history taking, the midwife learned that the child was exposed to contaminated water.

What is the most likely mode of transmission for the above scenario?

- A. Biological transmission
- B. Mechanical transmission
- C. Vehicle-borne transmission
- D. Direct horizontal transmission

ANSWER: C

194. A health care provider observed multiple cases with symptoms of abdominal pain from a nearby rural school. A result from laboratory investigation identified ascariasis egg in a stool sample of the majority of the children. An educational intervention on prevention of this communicable disease was delivered.

What is the most important prevention mechanism that is unique to this scenario?

- A. Encouraging students to wear shoes
- B. Educating to avoid eating raw meat
- C. Washing hands after using the latrine
- D. Avoiding eating soil contaminated food

ANSWER: D

195. A pregnant woman in her second trimester came to ANC clinic seeking advice on treating drinking water while she was out of town for a field work. The place where she was going had tap water source, but she was not sure about its safety. She wanted to use safe, readily available and effective water treatment technique.

What is most likely water treatment technique to be advised for the woman?

- A. Using iron or its compounds
- B. Using silver or its compounds
- C. Using iodine or its compounds
- D. Using chlorine or its compounds

ANSWER: D

196. More than 65 people who attended a cultural wedding ceremony visited a health center for a sudden onset of vomiting, watery diarrhoea, fever and abdominal cramp. The health care provider suspected staphylococcal food poisoning and treated with fluid and electrolyte replacement.

What is the most appropriate food borne disease control measure in the above scenario?

- A. Avoid eating canned foods from bulging containers.
- B. Reduce time interval from food preparation to consumption
- C. Improve proper hand washing practice before and after eating
- D. Avoid consuming raw and under-cooked eggs and egg products

ANSWER: B

197. A 30-day-old home delivered baby was referred from the nearby health center to a hospital. The neonate was not able to breastfeed and was under critical condition. After thorough investigations, the baby was diagnosed with neonatal sepsis and admitted to neonatal intensive care unit. Such like cases are being reported repeatedly from the hospital.

What is the most appropriate intervention needed in this scenario?

- A. Improve proximity and accessibility of hospitals
- B. Improve availability of staff in the health center
- C. Improve referral system among health facilities
- D. Improve awareness on institutional delivery

ANSWER: D

198. A 16-year-old high school student came to adolescent and youth friendly service unit to receive condom. He came to the unit at the 15-minute break and was in a hurry to return back to school. On the other hand, the midwife did not ask him whether he has time to discuss about other reproductive health issues. Rather she started to provide counselling on negative effect of early age sexual initiation on his health and future carrier.

Which principle of education is most likely overlooked in this case?

- A. Principles related to abilities
- B. Principles related to motivation
- C. Principle cultural appropriateness
- D. Principles related to nature of learning

ANSWER: B

199. A two-year-old child was brought to MCH clinic with a history of diarrhoea for the last three days. The child was restless and drunk fluid eagerly. His eyes were sunken and skin pinch went slowly on physical examination. The health care provider suspected the risk of developing micronutrient deficiency if appropriate prevention measure was not taken.

What is the most likely micronutrient deficiency to be prevented in this case?

- A. Cobalt
- B. Iodine
- C. Iron
- D. Zinc

ANSWER: D

200. A newly graduated midwife was assigned to provide education on family planning utilization for a group of women. During the session, she tended to use medical terms and English words in her speech. In the middle of the session, she observed that the women were not attending to her education. Rather, they were talking to each other about their personal issue.

Which principle of education was most likely overlooked in this case?

- A. Principles related to motivation
- B. Principle of cultural appropriateness
- C. Principles related to nature of learning
- D. Principles related to need and abilities

ANSWER: D