



Milki health science tutorial center



+251910690594, +251911058885

Tajaajiloota nuti kenninu



Barnoota dabalataa kennuu

Exit Exam'f/ COCf isin qopheessuu.

Leenjiwwan gaggabaaboo kennuu fi K.K.F.



Teessoo: Magaalaa Naqamtee, Gamoo OSSA, Darbii 3ffaa

[Back](#)

Time left 4:22:49

Question 46

Not yet answered

Marked out of 1.00

 [Flag question](#)

Members of rural community in a district asked a nurse about how to keep the safety of our water for drinking. A nurse wanted to instruct the community about the most cost effective and reliable method of disinfecting water at household level. What is the most likely water treatment technique instructed by the nurse?

- ☐ a. Chlorination of water
- ☐ b. Home Candle Filters
- ☐ c. Boiling of water
- ☐ d. Cloth filtration

[Previous page](#)[Next page](#)

Search...



[Back](#)

Time left 4:28:53

Question 24

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse has been attending a multi-gravid woman with first stage of labor in local health center. The woman was in the latent phase of labor and had persisted uterine contraction for 10 hour. The cervical dilatation remains less than 4cm and fetal heart beat was 140/min. The nurse referred the mother to a hospital. What is the most likely explanation for referral of this patient?

- ☐ a. PROM
- ☐ b. Prolonged labor
- ☐ c. Fetal distress
- ☐ d. Uterine rupture



Search...





Time left 4:23:57

Question 42

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse who is working as a ward head in a hospital developed annual plan with hospital staff teams. However, they failed to achieve the plan because of inability of mobilizing the human and material resources from the hospital. What is the most likely management function that causes low achievement of this ward?

- ☐ a. Directing
- ☐ b. Planning
- ☐ c. Controlling
- ☐ d. Organizing

[Previous page](#)[Next page](#)

Search...





Time left 4:22:25

Question 47

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 31-year-old daily laborer chews khat everyday despite repeated conflict with his wife due to economic problem of his substance use habit. In the last two years, his working hours were only in the morning and he spends the whole afternoon using khat to feel free from stressing symptoms. What is the most likely diagnosis of this person?

- ☐ a. Stimulant withdrawal
- ☐ b. Stimulant intoxication
- ☐ c. Unspecified stimulant related disorder
- ☐ d. Stimulant use disorder

[Previous page](#)[Next page](#)

Search...



**Question 39**

Not yet answered

Marked out of 1.00

 Flag question

A postpartum woman from a rural area was admitted to gynecologic ward with a diagnosis of vesico-vaginal fistula. A nursing diagnosis was made as "Disturbed body image related to foul smelling odor as manifested by woman's verbalization". What is the priority nursing intervention for the woman?

- ☐ a. Empower the woman with economical status
- ☐ b. Encourage the woman to take balance diet
- ☐ c. Provide information on self-care activities
- ☐ d. Teach the woman about positive self-talk

[Previous page](#)[Next page](#)

Search...





Time left 4:23:44

Question 43

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 54-year-old man was believed to be in good health when admitted for an elective surgery. Few hours after surgery, the nurse noted a sharp increase in his pulse rate and brief intense grand mal convulsion. His wife reported that he has 10 year history of excessive alcohol drinking. What is the priority nursing action for this patient?

- ☐ a. Consult the physician immediately
- ☐ b. Record findings in recording book
- ☐ c. Follow the patient closely
- ☐ d. Handover the patient properly to a nurse in next turn

[Previous page](#)[Next page](#)

Search...



**Question 48**

Not yet answered

Marked out of 1.00

 Flag question

A nurse who is works in scabies prevalent community wants to design a different means to prevent the disease in the community in collaboration with stakeholders. What is the most appropriate prevention strategy to be used by the nurse?

- ☐ a. Avoidance drinking of contaminated water
- ☐ b. Discharging the accumulated water from the area
- ☐ c. Provision of adequate water for personal hygiene
- ☐ d. Prevention swimming and crossing in the infected water

[Previous page](#)[Next page](#)**Quiz navigation**

Search...





Marked out of 1.00

 [Flag question](#)

A nurse is giving neonatal care for a baby immediately after delivery. The baby has severe chest indrawing, difficulty of breathing, cyanosed extremities and was unable to suck breast. On examination, the baby has normal grasping and moro reflex. What is the most priority nursing intervention for the baby?

- ☐ a. Put the baby in radiant warmer
- ☐ b. Clear airways
- ☐ c. Encourage breast feeding
- ☐ d. Give oxygen to the baby

[Previous page](#)[Next page](#)

Quiz navigation

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16



Search...



[Back](#)

Time left 4:27:33

Question 30

Not yet answered

Marked out of 1.00

 [Flag question](#)

The morning shift nurse had left award earlier by giving and recoding the 8 O'clock medication. The afternoon nurse checked the patient's card and found the medication was administered earlier than the schedule. The nurse become angry and wants to share the issue to the head nurse. What is the most appropriate reason to accuse the morning shift nurse?

- ☐ a. Breach of duty
- ☐ b. Omission of care
- ☐ c. Commission of care
- ☐ d. Negligence



Search...





Marked out of 1.00

 [Flag question](#)

A 68-year-old woman has come to a nurse with a manifestation of chills, rigor, fever, head ache, diarrhea, hallucinations, abdominal pain, and ache. The nurse suspected that the patient is with malaria. After observing repeated cases, the nurse wanted to design disease prevention strategy that can apply in the community. What is the most likely strategy to prevent the disease in this scenario?

- ☐ a. Isolation of an individual who develop the disease
- ☐ b. Avoid drinking of contaminated water
- ☐ c. Discharging the accumulated water from the area
- ☐ d. Treatment of an individual who develop the disease

[Previous page](#)[Next page](#)

Search...



[Back](#)

Time left 4:27:44

Question 29

Not yet answered

Marked out of 1.00

 [Flag question](#)

A neonate, who was diagnosed with neonatal sepsis, was admitted in a hospital. The nurse in charge examined the baby and the findings showed central cyanosis, chest in drawing, tachycardia and tachypnea. What is the most likely nursing diagnosis of this case?

- ☐ a. High risk for hypoventilation
- ☐ b. Ineffective airway clearance
- ☐ c. Ineffective breathing pattern
- ☐ d. Ineffective tissue perfusion

[Previous page](#)[Next page](#)

Search...





Time left 4:21:32

Question 50

Not yet answered

Marked out of 1.00

 [Flag question](#)

The nurse wants to test whether the mean age of adults who visited the youth friendly service is not different from 25 years. The nurse took sample of 196 adult patients from population. The mean age of sampled adults was 20 years and the given population standard deviation was six years. What is the most appropriate statistical test used for the given hypothesis?

- ☐ a. Standard normal Z- test
- ☐ b. Chi-square test
- ☐ c. Student t-test
- ☐ d. Paired t-test

[Previous page](#)[Next page](#)

Search...





Time left 4:23:22

Question 44

Not yet answered

Marked out of 1.00

 [Flag question](#)

A breast feeding mother came to postnatal clinic with the complaints of left side breast swelling and unable to feed her baby. On physical examination, the nurse identified that she has irregular mass and breast tenderness on her both breasts. What is the most likely nursing diagnosis for this client?

- ☐ a. Risk for imbalanced body temperature
- ☐ b. Risk for impaired skin integrity
- ☐ c. Risk for infection
- ☐ d. Risk for injury

[Previous page](#)[Next page](#)**Quiz navigation**

Search...





Time left 4:24:35

Question 40

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 26-year-old Gravid II Para I pregnant woman came to ANC clinic with complaints of headache, dyspnea, and fatigue for three days. Her pulse rate was 78/min, Blood pressure was 100/60 mmHg and fetal heart rate was 140/min. On physical examination, she had pale conjunctiva and 32 weeks gestational age pregnancy. Her hemoglobin concentration was 10g/dl. What is the most likely indicative characteristic for the woman's problem?

- ☐ a. Headache
- ☐ b. Pale conjunctiva
- ☐ c. Dyspnea
- ☐ d. Hgb 10 g/dl



Search...





Nursing (comprehensive)

[Back](#)

Time left 3:53:21

Question 51

Not yet answered

Marked out of 1.00

[Flag question](#)

A two-day-old male neonate was admitted to NICU for thermal care. On physical examination, the newborn has pink palms and soles, and yellow discoloration in the face and eyes. Based on IMNCI classification, what is the most likely diagnosis of the neonate?

- ☒ a. Severe jaundice
- ☐ b. Uncomplicated jaundice
- ☐ c. No jaundice
- ☐ d. Jaundice

[Clear my choice](#)[Previous page](#)[Next page](#)**Quiz navigation**



Time left 3:51:38

Question 53


Not yet answered

Marked out of 1.00

[Flag question](#)

A nurse working in an ICU always enters earlier to assess the order and revises cares for all patients. One day, the nurse gets wrong order and doubtful medication for her critically ill patient. Then, the nurse immediately consults the doctor who ordered the medication before she carries out the medication. What is the most likely reason that the nurse wants to consult the doctor?

- ☐ a. To prevent mal practices that risk the patients and to safeguard them
- ☐ b. Fears not to take accountability for any bad consequence
- ☐ c. Since he is newly assigned, he is not clear with ICU procedures
- ☐ d. Lacks confidence to work independently as professional nurse

[Previous page](#)[Next page](#)**Quiz navigation**

10:39 AM P
Question.AI

Network connection currently unavailable.

Solution 1

Solution 2

Solution 3

Solution 4

Question

A nurse working in an adult emergency OPD observed six individuals who developed hepatitis A disease. The incubation periods for each affected person were 29, 31, 24, 29, 30 and 25 days. Calculate the mode of hepatitis A incubation period in this Scenario? The most frequently appearing data.

- ☐ A. 28
- ☒ B. 27
- ☐ C. 31
- ☐ D. 29



Ask

[Back](#)

Time left 3:53:21

Question 51

Not yet answered

Marked out of 1.00

[Flag question](#)

A two-day-old male neonate was admitted to NICU for thermal care. On physical examination, the newborn has pink palms and soles, and yellow discoloration in the face and eyes. Based on IMNCI classification, what is the most likely diagnosis of the neonate?

- ☒ a. Severe jaundice
- ☐ b. Uncomplicated jaundice
- ☐ c. No jaundice
- ☐ d. Jaundice

[Clear my choice](#)[Previous page](#)[Next page](#)**Quiz navigation**

[Back](#)

Time left 3:53:21

Question 51

Not yet answered

Marked out of 1.00

[Flag question](#)

A two-day-old male neonate was admitted to NICU for thermal care. On physical examination, the newborn has pink palms and soles, and yellow discoloration in the face and eyes. Based on IMNCI classification, what is the most likely diagnosis of the neonate?

- ☒ a. Severe jaundice
- ☐ b. Uncomplicated jaundice
- ☐ c. No jaundice
- ☐ d. Jaundice

[Clear my choice](#)[Previous page](#)[Next page](#)**Quiz navigation**

[Back](#)

Time left 3:46:32

Question 58

Not yet answered

Marked out of 1.00

[Flag question](#)

A woman was delivering her baby at the 34th week of gestation. Immediately after birth, the baby became lethargic, does not cry, and has weak Moro reflex. What is the most likely diagnosis for this condition?

- ☐ a. Asphyxia
- ☐ b. Hypothermia
- ☐ c. Hypoxemia
- ☐ d. Hypoxia

[Previous page](#)[Next page](#)**Quiz navigation**

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36



[Back](#)

Time left 3:46:32

Question 58

Not yet answered

Marked out of 1.00

[Flag question](#)

A woman was delivering her baby at the 34th week of gestation. Immediately after birth, the baby became lethargic, does not cry, and has weak Moro reflex. What is the most likely diagnosis for this condition?

- ☐ a. Asphyxia
- ☐ b. Hypothermia
- ☐ c. Hypoxemia
- ☐ d. Hypoxia

[Previous page](#)[Next page](#)**Quiz navigation**

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36



A newly graduated female nurse planned to start professional services and applied to a private hospital. The human resource of the hospital asked her to bring the evidence that indicates legal permit that the government agency grants her to practice as a BSC nurse. What is the most likely requirement the nurse must fulfill?

- ☐ a. License evidence
- ☒ b. Certification evidence
- ☐ c. Accreditation evidence
- ☐ d. Registration evidence

[Clear my choice](#)

[Previous page](#)

[Next page](#)

Quiz navigation



Time left 4:05:02

Question 41

Not yet answered

Marked out of 1.00

Flag question

A primigravida pregnant woman was admitted to a gynecology ward with hyperemesis gravidarum. During admission, her blood pressure was 80/50mmHg and Pulse rate was 100/min. She had dry mouth and poor concentration. Laboratory investigation results show urine ketone 3+, urine proteins 2+ and her HCT is 35 percent. She has been managed with two bags of ringer lactate with two vials of 40 percent dextrose. What is the most appropriate sign that indicates her improvement?

- ☐ a. Ketone free urine
- ☐ b. Protein free urine
- ☐ c. Increase blood glucose
- ☐ d. Increased hematocrit level

[Previous page](#)[Next page](#)**Quiz navigation**



Time left 4:02:45

Question 42

Not yet answered

Marked out of 1.00

[Flag question](#)

An eleven-year-old child was brought to a clinic due to restlessness while playing and he does not follow instructions in class. He cannot read, listen and solve simple mathematics questions. He also loses learning materials, leaves class, runs, and climbs tree frequently. What is the most likely type of childhood behavioral disorders in this case?

- ☐ a. Mental retardation disorder
- ☒ b. Down's syndrome
- ☐ c. Attention deficit hyperactive disorder
- ☐ d. Childhood schizophrenia

[Clear my choice](#)[Previous page](#)[Next page](#)**Quiz navigation**

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27



[Back](#)**Question 53**

Not yet answered

Marked out of 1.00

 Flag question

A postnatal woman with cesarean section wound developed vomiting and bleeding in the recovery unit. Her blood pressure was 90/50mmHg, temperature was 36 degree centigrade, pulse rate was 120/min, and urine output was 500ml/24 hour. Her hematocrit level was 38 percent (normal range 33-42 percent). Which parameter indicates maternal wellbeing after four hour?

- ☐ a. Urine output of 20ml/hour
- ☐ b. Stable Vital sign
- ☐ c. Pink congestive
- ☐ d. Initiation of breast feeding



Search...



[Back](#)**Question 56**

Not yet answered

Marked out of 1.00

 Flag question

A mother is on anticonvulsant therapy with magnesium sulfate for severe pre-eclampsia. During the course her Pulse rate was 62/min, Diastolic blood pressure was 100mmHg, respiratory rate was 15/min, and urine output was 20 ml per hour over preceding four hours. On physical examination, her patellar reflex is absent. Which management is the most appropriate for this case?

- ☐ a. Change the drug to diazepam
- ☐ b. Providing calcium gluconate
- ☐ c. Resuscitating the mother
- ☐ d. Continue administering magnesium sulfate



Search...



[Back](#)

Time left 4:21:18

Question 51

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse observed medication administration error made by his friend working in medical ward. Due to this, the patient developed minor complication and treated. What is the first thing the nurse must do?

- ☐ a. Follow secretly and supervise
- ☐ b. Confront his friend
- ☐ c. Report to the nurse director
- ☐ d. Report to the legal judiciary body

[Previous page](#)[Next page](#)**Quiz navigation**

Search...





Time left 4:01:30

Question 43

Not yet answered

Marked out of 1.00

[Flag question](#)

A 16-year-old boy with brief psychotic disorder came to a clinic with contractions of muscles that result in abnormal movement, tongue protrusion and difficulty of breathing. Symptoms developed after taking two doses of prescribed haloperidol 5 mg orally. What is the best initial treatment for this boy?

- ☐ a. Switch to another antipsychotic
- ☐ b. Reduce dose of antipsychotic
- ☐ c. Discontinue the drug and refer
- ☐ d. Give 10 mg diazepam intravenously

[Previous page](#)[Next page](#)**Quiz navigation**

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36





Time left 4:04:06

Question 41

Not yet answered

Marked out of 1.00

[Flag question](#)

A primigravida pregnant woman was admitted to a gynecology ward with hyperemesis gravidarum. During admission, her blood pressure was 80/50mmHg and Pulse rate was 100/min. She had dry mouth and poor concentration. Laboratory investigation results show urine ketone 3+, urine proteins 2+ and her HCT is 35 percent. She has been managed with two bags of ringer lactate with two vials of 40 percent dextrose. What is the most appropriate sign that indicates her improvement?

- ☐ a. Ketone free urine
- ☐ b. Protein free urine
- ☒ c. Increase blood glucose
- ☐ d. Increased hematocrit level

[Clear my choice](#)[Previous page](#)[Next page](#)**Quiz navigation**

[Back](#)

Time left 4:00:31

Question 44

Not yet answered

Marked out of 1.00

[Flag question](#)

A nurse works in district X at which an enlarged thyroid gland, speech defects and deafness are common among children living in the district. What is the most likely cause of the problem in this scenario?

- ☐ a. Folic acid deficiency
- ☐ b. Iron deficiency
- ☐ c. Iodine deficiency
- ☐ d. Vitamin A deficiency

[Previous page](#)[Next page](#)**Quiz navigation**

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36



[Back](#)

Time left 3:55:11

Question 49

Not yet answered

Marked out of 1.00

[Flag question](#)

Gender office of the city administration organizes a workshop on improving gender equality. Based on the invitation, the CEO of the general hospital made a speech at the event representing the hospital. What is the most likely managerial role played by the CEO?

- ☐ a. Liaison
- ☐ b. Influencer
- ☐ c. Figurehead
- ☐ d. Monitor

[Previous page](#)[Next page](#)**Quiz navigation**

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36



MILKI HEALTH SCIENCE TUTORIAL CENTER



0910 69 0594
0911 05 8885

TAJAAJILA KENNINU

- ✓ **Barnoota Dabalataa Kennu**
- ✓ **Exit Exam/ COC'f isin qopheessuu**
- ✓ **Leenjiwwan gaggabaaboo kennuu fi kkf**



Teessoon: Magaalaa Naqamtee
Gamoo OSSA Darbii 3^{ffaa}

[Back](#)

Time left 3:56:51

Question 47

Not yet answered

Marked out of 1.00

[Flag question](#)

A breast feeding mother came to postnatal clinic with the complaints of left side breast swelling and unable to feed her baby. On physical examination, the nurse identified that she has irregular mass and breast tenderness on her both breasts. What is the most likely nursing diagnosis for this client?

- ☐ a. Risk for imbalanced body temperature
- ☐ b. Risk for impaired skin integrity
- ☒ c. Risk for infection
- ☐ d. Risk for injury

[Clear my choice](#)[Previous page](#)[Next page](#)**Quiz navigation**

Time left 3:50:20

Flag question

- ☐ a. Hypothermia
- ☐ b. Hypoxemia
- ☐ c. Asphyxia
- ☐ d. Hypoxia

Next page

Quiz navigation



Question 20

Not yet answered

Marked out of 1.00

 Flag question

In a district only 5 percent of households have latrine and intestinal parasite was the top ten diseases in the catchment. Zonal health administration initiated a project to increase household latrine coverage by 20 percent each year. The project provides essential materials for latrine construction. But a year later, the project assessment indicated that less than half of the target has been reached. What is the most likely reason for failure of the project?

- ☐ a. Availability of resource
- ☐ b. Complexity of the problem
- ☐ c. Interest and support of the authorities
- ☐ d. Low community acceptability

[Previous page](#)[Next page](#)

Search...



[Back](#)

Time left 3:50:07

Question 100

Not yet answered

Marked out of 1.00

 [Remove flag](#)

Gender office of the city administration organizes a workshop on improving gender equality. Based on the invitation, the CEO of the general hospital made a speech at the event representing the hospital. What is the most likely managerial role played by the CEO?

- ☐ a. Liaison
- ☐ b. Influencer
- ☐ c. Figurehead
- ☐ d. Monitor

[Previous page](#)[Finish attempt ...](#)

Search...





Time left 4:20:21

Question 54

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 42-year-old male patient was going to have amputation due to gangrene. If he is amputated, he will lose his properties which will be transferred to his brother according to cultural practices in the area. If the procedure is not done, the patient will die. The nurse in charge convinced the patient about the procedure. What principle is most likely applied in the case?

- ☐ a. Veracity
- ☐ b. Proportionality
- ☐ c. Compassion
- ☐ d. Paternalism

[Previous page](#)[Next page](#)

Search...



**Question 55**

Not yet answered

Marked out of 1.00

 Flag question

A mother brought her three-year-old child to an emergency clinic. The child was comatous and apenic on arrival. Also, no sign of palpable chest for ten seconds as well as dilated pupils, and absence of respiratory movements which are the typical findings. What is the most likely initial nursing intervention for the above case?

- ☐ a. Chest compression and gastric lavage
- ☐ b. NG tube feeding and chest compression
- ☐ c. Middle chest thrust and gastric lavage
- ☐ d. Mechanical ventilation and chest compression

[Previous page](#)[Next page](#)

Search...



**Question 52**

Not yet answered

Marked out of 1.00

 [Flag question](#)

A postpartum mother visited a hospital after three hours of home delivery with a complaint of bleeding for three hours. On arrival, she has slight vaginal bleeding, pale conjunctiva, and cold clammy skin. The uterus is soft and spongy on palpation. Her blood pressure was 90/60 mmHg, pulse rate was 82/min and temperature was 37.3 degree centigrade. What is the priority nursing intervention for this mother?

- ☐ a. Apply uterine tamponade
- ☐ b. Massage the uterus
- ☐ c. Give oxytocine IM
- ☐ d. Start IV fluid

[Previous page](#)[Next page](#)**Quiz navigation**

Search...



[Back](#)

Time left 3:51:39

Question 93

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 24-year-old Gravida II Para I woman came to labor ward with complaints of back pain and gash of fluid for four hours. On physical examination, the nurse identified cervical dilation four cm for 14 hours duration, station three, uterine contraction three/10min lasting for 18 second and ruptured membrane. What is the most likely sign that indicates prolonged labor for this client?

- ☐ a. Station
- ☐ b. Status of membrane
- ☐ c. Cervical dilation
- ☐ d. Uterine contraction



Search...



[Back](#)

Time left 3:54:06

Question 83

Not yet answered

Marked out of 1.00

 [Flag question](#)

A two-day-old male neonate was admitted to NICU for thermal care. On physical examination, the newborn has pink palms and soles, and yellow discoloration in the face and eyes. Based on IMNCI classification, what is the most likely diagnosis of the neonate?

- ☐ a. Uncomplicated jaundice
- ☐ b. Jaundice
- ☐ c. No jaundice
- ☐ d. Severe jaundice



Search...





Time left 3:50:50

Question 97

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 34th week pregnant woman was admitted to a maternity ward with hyper emesis gravidarum. At the time of admission, her blood pressure was 90/50 mmHg, Pulse rate was 100/min and she was vomiting frequently. She had dry mouth and poor concentration. Her lab investigation results show that 3+ ketone in urine, 2+ proteins and her HCT is 35 percent. What is the most appropriate management for this woman?

- ☐ a. Terminate the pregnancy
- ☐ b. Start RL with 40 percent dextrose
- ☐ c. Start NG tube feeding
- ☐ d. Put the woman on NPO

[Previous page](#)[Next page](#)

Search...



Flag question

- ☐ a. Suggest alternative sources of childbearing
- ☐ b. Reassure the couple for psychological upset
- ☐ c. Provide medication to resume fertility
- ☐ d. Advise to reduce alcohol consumption

Next page



[Back](#)

Time left 3:53:32

Question 85

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse working in ART clinic of a hospital wants to summarize CD4 T cell counts for 13 HIV positive patients who were treated with highly active antiretroviral therapy (HAART) for the last six months. Their CD4 T cell counts for the 13 subjects were 230, 205, 313, 207, 227, 245, 173, 58, 103, 181, 105, 301, and 169. What is the mean CD4 T cell count of HIV positive patients?

- ☐ a. 193.0
- ☐ b. 193.6
- ☐ c. 205.0
- ☐ d. 206.1



Search...





Not yet answered

Marked out of 1.00

 [Flag question](#)

A 30-year-old pregnant woman has come to an ANC clinic with complaints of headache, abdominal pain, vomiting and difficulty of breathing during sleeping for two weeks duration. Her blood pressure is 100/70mmHg, Pulse rate is 80/min and fetal heartbeats 140/min. On physical examination, she has markedly enlarged abdomen and gestational age of 34weeks. The nurse has suspected that the woman is at greater risk for amniotic fluid disorder. Which is the most pertinent characteristic that indicates the disorder?

- ☐ a. Difficulty of breathing
- ☐ b. Vomiting
- ☐ c. Enlarged abdomen
- ☐ d. Abdominal pain

Previous page

Next page



Search...





Not yet answered

Marked out of 1.00

 [Flag question](#)

A 30-year-old pregnant woman has come to an ANC clinic with complaints of headache, abdominal pain, vomiting and difficulty of breathing during sleeping for two weeks duration. Her blood pressure is 100/70mmHg, Pulse rate is 80/min and fetal heartbeats 140/min. On physical examination, she has markedly enlarged abdomen and gestational age of 34weeks. The nurse has suspected that the woman is at greater risk for amniotic fluid disorder. Which is the most pertinent characteristic that indicates the disorder?

- ☐ a. Difficulty of breathing
- ☐ b. Vomiting
- ☐ c. Enlarged abdomen
- ☐ d. Abdominal pain

Previous page

Next page



Search...



[Back](#)

Time left 3:53:16

Question 85

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse working in ART clinic of a hospital wants to summarize CD4 T cell counts for 13 HIV positive patients who were treated with highly active antiretroviral therapy (HAART) for the last six months. Their CD4 T cell counts for the 13 subjects were 230, 205, 313, 207, 227, 245, 173, 58, 103, 181, 105, 301, and 169. What is the mean CD4 T cell count of HIV positive patients?

- ☐ a. 193.0
- ☐ b. 193.6
- ☐ c. 205.0
- ☐ d. 206.1



Search...





Marked out of 1.00

 [Flag question](#)

A primigravida pregnant woman was admitted to a gynecology ward with hyperemesis gravidarum. During admission, her blood pressure was 80/50mmHg and Pulse rate was 100/min. She had dry mouth and poor concentration. Laboratory investigation results show urine ketone 3+, urine proteins 2+ and her HCT is 35 percent. She has been managed with two bags of ringer lactate with two vials of 40 percent dextrose. What is the most appropriate sign that indicates her improvement?

- ☐ a. Increase blood glucose
- ☐ b. Ketone free urine
- ☐ c. Protein free urine
- ☐ d. Increased hematocrit level

[Previous page](#)[Next page](#)

Quiz navigation



Search...





Time left 3:52:08

Question 91

Not yet answered

Marked out of 1.00

 [Flag question](#)

A two-month-old infant was brought to a hospital with a complaint of cough of two days duration. The infant is alert and can feed breast milk. The infant respiration is 50/min, pulse rate is 110/min and temperature is 38 degree centigrade. The nurse wants to classify this illness based on IMNCI protocol. What is the most likely diagnosis of the child?

- ☐ a. Severe pneumonia
- ☐ b. Cough/Cold
- ☐ c. Very severe pneumonia
- ☐ d. Pneumonia

[Previous page](#)[Next page](#)

Search...





Nursing (Comprehensive)

[Back](#)

Question 89

Not yet answered

Marked out of 1.00

 Flag question

A nurse works in district X at which an enlarged thyroid gland, speech defects and deafness are common among children living in the district. What is the most likely cause of the problem in this scenario?

- ☐ a. Vitamin A deficiency
- ☐ b. Iron deficiency
- ☐ c. Iodine deficiency
- ☐ d. Folic acid deficiency

[Previous page](#)[Next page](#)

[Back](#)

Time left 3:54:50

Question 79

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 32-year-old soldier saw terrorist attack on television and starts to experience feelings of panic. The panic disappeared within a few hours; however, for the next few nights, he had night mares with frequent thoughts about dying. Before 10 years, he survived ambush which killed three of his close friends in front of him. What is the most likely diagnosis for the above case?

- ☐ a. Adjustment disorder
- ☐ b. Post traumatic stress disorder
- ☐ c. Complicated grief
- ☐ d. Acute stress disorder



Search...



[Back](#)

Time left 3:51:51

Question 92

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse is preparing a care plan for a 45-year-old newly admitted patient who was relatively healthy one day ago. On assessment, the patient looks confused, disoriented, and irritable and could not recognize where he is. He frequently pulls out the infusion set. What is the priority nursing intervention for this patient?

- ☐ a. Restrain the patient
- ☐ b. Assist his orientation
- ☐ c. Assist his memory
- ☐ d. Control his behavior

[Previous page](#)[Next page](#)

Search...





Time left 3:54:39

Question 80

Not yet answered

Marked out of 1.00

 [Flag question](#)

A female nurse attending hospital reform training introduces herself as she is serving with skill and knowledge expected for the specific position she is assigned to. What is the most likely managerial level of a nurse expressed in this case?

- ☐ a. Top level
- ☐ b. Second level
- ☐ c. First level
- ☐ d. Middle level

[Previous page](#)[Next page](#)**Quiz navigation**

1

2

3

4

5

6

7

8



Search...





Time left 3:54:16

Question 82

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 16-year-old boy with brief psychotic disorder came to a clinic with contractions of muscles that result in abnormal movement, tongue protrusion and difficulty of breathing. Symptoms developed after taking two doses of prescribed haloperidol 5 mg orally. What is the best initial treatment for this boy?

- ☐ a. Reduce dose of antipsychotic
- ☐ b. Switch to another antipsychotic
- ☐ c. Discontinue the drug and refer
- ☐ d. Give 10 mg diazepam intravenously

[Previous page](#)[Next page](#)**Quiz navigation**

Search...



[Back](#)**Question 67**

Not yet answered

Marked out of 1.00

 Flag question

A 44-year-old man, who has a history of excessive alcohol drinking for five years, came with persistent depressed mood and marked loss of interest. He stopped drinking before two days. The nurse diagnosed the case as he has alcohol induced depressive disorder. What is the most important criterion to support this diagnosis?

- ☐ a. Symptoms developed during substance withdrawal
- ☐ b. Persistent depressed mood
- ☐ c. Clinically significant distress
- ☐ d. Drinking excessive alcohol for five years



Search...



[Back](#)**Question 88**

Not yet answered

Marked out of 1.00

 Flag question

A 26-year-old man came to a hospital and reported auditory hallucination of one day duration. The onset is reported to be two hours after drinking five bottles of beer. He has no previous history of mental illness and no insight of what happened to him. What is the most likely diagnosis for this case?

- ☐ a. Brief psychotic disorder
- ☐ b. Schizophreniform disorder
- ☐ c. Alcohol withdrawal syndrome
- ☐ d. Alcohol induced psychotic disorder

[Previous page](#)[Next page](#)

Search...





Time left 3:55:11

Question 77

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 32-year-old male patient came to the health center for voluntary counseling and testing. The nurse in charge nods his head while the patient is talking. However, the patient was not telling the details about his past experience. What question most likely encourages open discussion to get full information?

- ☐ a. What do you want to talk about?
- ☐ b. Can we discuss about your partners and friends?
- ☐ c. How many partners do you have?
- ☐ d. How have been things going?

[Previous page](#)[Next page](#)

Search...



[Back](#)

Time left 3:53:52

Question 84

Not yet answered

Marked out of 1.00

 [Flag question](#)

A Zonal health administration inspection team visits woreda health offices every quarter. One of a leader from a woreda explains his experience of leadership for the participants as he involves staffs in the decisions, shares responsibilities and creates chances for staffs to be innovative. What leadership style is most likely used by the woreda administrator?

- ☐ a. Transformational
- ☐ b. Authoritarian
- ☐ c. Laissez faire
- ☐ d. Democratic



Search...





Time left 3:58:57

Question 68

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 70-year-old patient came to an adult OPD with complaints of loss of body weight, fatigue and unable to remember things. On history taking, the nurse found that he routinely forgets taking meal and name of things. On physical examination, he became underweight. What is the most likely intervention for this patient?

- ☐ a. Participate in exercise and fitness programs
- ☐ b. Engage in recreational and educational program
- ☐ c. Provide psychological and food services
- ☐ d. Facilitate hearing aid and hearing-aid usage



Search...





Back

Time left 3:58:43

Question 69

Not yet answered

Marked out of 1.00

 [Flag question](#)

A pregnant woman came to a hospital at her 36th gestation with complaint of headache, anxiety, paler palm and weakness.

Laboratory investigation indicates that her hemoglobin level was 10 g/dL (Normal Hgb for late pregnancy=11.0-13.8 g/dl) and she was diagnosed with anemia. What patient's outcome criterion is the most important to prioritize nursing diagnosis?

- ☐ a. Free from anxiety
- ☐ b. Having Hgb greater than 13 g/dL
- ☐ c. Subside from pale discoloration
- ☐ d. Free from headache

Previous page

Next page



Search...



[Back](#)**Question 72**

Not yet answered

Marked out of 1.00

 Flag question

A pregnant woman at GA of 14-week was admitted to a hospital for severe and persistent nausea and vomiting. Physical examination shows that she had dry lip and mucous membrane, is acutely sick looking and warm to touch. Laboratory results show ketone +4 and urine output of 300ml in 24hours. What is the most appropriate plan of action for this mother?

- ☐ a. Administer anti-emetics immediately
- ☐ b. Providing total parenteral nutrition
- ☐ c. Administering fluid for the mother
- ☐ d. Advise the mother to take oral food

[Previous page](#)[Next page](#)

Search...



**Question 70**

Not yet answered

Marked out of 1.00

 Flag question

A woman, who has been using IUCD for six months, came to a clinic with complaints of lower abdominal pain and pain during sexual intercourse. On vaginal examination, uterine and cervical tenderness were identified. What is the priority nursing diagnosis for this woman?

- ☐ a. Risk for infection
- ☐ b. Risk for trauma
- ☐ c. Risk for injury
- ☐ d. Risk for bleeding

[Previous page](#)[Next page](#)**Quiz navigation**

1

2

3

4

5

6

7

8



Search...



[Back](#)

Time left 3:58:22

Question 71

Not yet answered

Marked out of 1.00

Flag question

A nurse works as surveillance team member of a national COVID-19 prevention and control. The team detected COVID-19 case on each day and reported immediately to the ministry of health. What is the most likely good feature of the surveillance done by the team?

- ☐ a. The team had strong laboratory services for accurate diagnosis
- ☐ b. The team had comprehensive action
- ☐ c. The team had timely reporting of cases
- ☐ d. The team used combined passive

[Back](#)

Time left 3:55:01

Question 78

Not yet answered

Marked out of 1.00

 [Flag question](#)

A five-year-old child was brought to a clinic with fever, cough, red eye, runny nose and general macula-papular skin rash. The child was admitted to pediatric ward and immediately switched to nasal oxygen. And vitamin "A" 200,000 IU was provided on admission. What is the most likely complication in this case?

- ☐ a. Seizure shock
- ☐ b. Croup
- ☐ c. Cerebral palsy Diarrhoea
- ☐ d. Pneumonia



Search...



[Back](#)**Question 65**

Not yet answered

Marked out of 1.00

 Flag question

A three-year-old child with complaints of change in social interaction and repetitive behavior was brought to a hospital. On assessment, the child loves to play alone, used multiple nonverbal behaviors, plays with the same toys and did not speak words appropriately. What is the most likely diagnosis of this child?

- ☐ a. Childhood depression disorder
- ☐ b. Autism spectrum disorders
- ☐ c. Down syndrome
- ☐ d. Attention deficit

[Previous page](#)[Next page](#)

Search...



[Back](#)

Time left 4:01:05

Question 58

Not yet answered

Marked out of 1.00

 [Flag question](#)

A six -year-old male child with bed wetting problem came with his mother. The child had no problem in the day time but has history of bed wetting four to five times in the morning per week. He did not have history of dysuria, urinary frequency and increased thirst. What is the most appropriate nursing intervention for this child?

- ☐ a. Encourage the family to use diaper until the child stops by him self
- ☐ b. Decrease the amount of fluid consumed during the day
- ☐ c. Advice the mother to scold the child not to repeat



Search...



**Question 64**

Not yet answered

Marked out of 1.00

 Flag question

A two-year-old baby was admitted to pediatric ward with a diagnosis of septic arthritis. The nurse planned for an intervention that would enhance wound healing. What is the priority intervention the nurse should give?

- ☐ a. Administer high protein diet
- ☐ b. Administer systemic antibiotic
- ☐ c. Instill antibiotics to the affected joint
- ☐ d. Debride of infected wound

[Previous page](#)[Next page](#)**Quiz navigation**

1

2

3

4

5

6

7

8



Search...





Not yet answered

Marked out of 1.00

 [Flag question](#)

A 25-year-old woman came to a hospital with a complaint of distressing dream of an attempted rape for the last six months. She had difficulty in falling and staying asleep, unable to have love feeling, irritable and outburst to tear. The nurse is considering a diagnosis of post-traumatic stress disorder. What criterion is necessarily lacking to support the diagnosis?

- ☐ a. Efforts to avoid activities, places, or people that remind the trauma
- ☐ b. Recurrent experiences of feeling detached from self
- ☐ c. Direct exposure to traumatic event or witnessing as it occurred to others
- ☐ d. Recurrent experiences of unreality of surroundings

[Previous page](#)[Next page](#)

Search...



**Question 72**

Not yet answered

Marked out of 1.00

 Flag question

A pregnant woman at GA of 14-week was admitted to a hospital for severe and persistent nausea and vomiting. Physical examination shows that she had dry lip and mucous membrane, is acutely sick looking and warm to touch. Laboratory results show ketone +4 and urine output of 300ml in 24hours. What is the most appropriate plan of action for this mother?

- ☐ a. Administer anti-emetics immediately
- ☐ b. Providing total parenteral nutrition
- ☐ c. Administering fluid for the mother
- ☐ d. Advise the mother to take oral food

[Previous page](#)[Next page](#)**Quiz navigation**

Search...



[Back](#)

Time left 3:56:14

Question 74

Not yet answered

Marked out of 1.00

 [Flag question](#)

In a health center, conflict arisen between OPD case team staffs on the issue of managing patients with COVID-19 infection. However a, head of the health center wants to resolve the conflict by considering assertiveness and cooperation between parties. What is the most likely conflict resolution strategy used by the head?

- ☐ a. Accommodation
- ☐ b. Collaboration
- ☐ c. Competition
- ☐ d. Avoidance



Search...





Nursing (Comprehensive)

[Back](#)

Time left 3:59:20

Question 66

Not yet answered

Marked out of 1.00

 [Flag question](#)

A two-year-old male child came to a hospital due to cough, fever, frequent vomiting, and uncontrolled jerky movement once at home. With physical examination, the child was irritable and has body temperature of 36.5 degree centigrade and was able to drink. What most likely danger sign does the nurse identify for this case?

- ☐ a. Fever
- ☐ b. Cough
- ☐ c. Convulsion
- ☐ d. vomiting everything



Search...





Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse working in woreda health office is planning to design substance use prevention plan as part of routine school health services. During preliminary assessment in the school, many young students use khat because of abundant availability of it in the surrounding areas. What is the most recommended effective measure of prevention of khat use in this school?

- ☐ a. Teach the students and the community around the school regarding its health impact
- ☐ b. Work with community police to clear school surrounding from Khat shops
- ☐ c. Discuss with the school administration to impose serious punishment on users
- ☐ d. Push policy makers to impose sanction by researching the health impact



Search...





Time left 4:00:50

Question 59

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 24-year-old pregnant woman came to an ANC clinic with complaints of headache and dizziness for two weeks. Her pulse rate is 70/min, gestational age 20weeks and fetal heart beat is 150/min. Laboratory investigation shows that Hgb 10g/dl and serum iron level less than 4 micro gram/dl (Normal range of serum iron = 65–75 micro gram/100mL). What is the most appropriate management for this woman?

- ☐ a. Diet therapy
- ☐ b. Blood transfusion
- ☐ c. Ferrous sulphate
- ☐ d. Fluid therapy



Search...





Time left 4:34:32

Question 4

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 26-year-old female patient was admitted to a medical ward with a diagnosis of DKA. The patient selected a treatment option that is best for her after being informed about all treatment options. After five days of treatment, the patient progressed well and was discharged. However, since she has no money to pay, she is kept in the ward till the hospital manager will come and decide. What is the most likely encountered disrespect to the patient?

- ☐ a. Detention in facility
- ☐ b. Abandonment of care
- ☐ c. Non consented care
- ☐ d. No dignified care

[Previous page](#)[Next page](#)

Search...



[Back](#)

Time left 4:34:40

Question 4

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 26-year-old female patient was admitted to a medical ward with a diagnosis of DKA. The patient selected a treatment option that is best for her after being informed about all treatment options. After five days of treatment, the patient progressed well and was discharged. However, since she has no money to pay, she is kept in the ward till the hospital manager will come and decide. What is the most likely encountered disrespect to the patient?

- ☐ a. Detention in facility
- ☐ b. Abandonment of care
- ☐ c. Non consented care



Search...



[Back](#)

Time left 4:34:55

Question 3

Not yet answered

Marked out of 1.00

 [Flag question](#)

A measles outbreak occurred in the woreda following mass population displacement because of conflict. Then, the head of woreda health office reported the outbreak situation for the Medias and other people outside the organization. What is the most likely managerial role the head played at the time?

- ☐ a. Spokesperson
- ☐ b. Liaison
- ☐ c. Disseminator
- ☐ d. Monitor

[Previous page](#)[Next page](#)

Search...





Time left 4:34:17

Question 5

Not yet answered

Marked out of 1.00

 [Flag question](#)

The nurse working in university students clinic meet a 19-year-old fresh student fulfilling the diagnostic criteria for adjustment disorder with depressed mood. He occasionally had difficulty of sleeping but function optimally. What is the most appropriate nursing intervention for this case?

- ☐ a. Reassure him symptoms will subside with time
- ☐ b. Tell him to come back if symptoms worsen
- ☐ c. Give medication to help him sleep well
- ☐ d. Reassure and advise to contact counseling unit



Search...





Time left 3:51:06

Question 96

Not yet answered

Marked out of 1.00

 [Flag question](#)

A researcher wants to conduct trends of malaria case for the last five years in low land district. The researcher collected confirmed malaria cases from health facilities around the district. The results were 145 cases in 2014; 276 cases in 2015; 356 case in 2016; 217 cases in 2017 and 467 cases in 2018. Finally, the researcher wanted to present these findings with graph. What is the most appropriate method of data presentation to be used by a researcher?

- ☐ a. Frequency polygon
- ☐ b. Scatter plot
- ☐ c. Line graph
- ☐ d. Histogram



Search...



[Back](#)

Time left 4:01:16

Question 57

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 20 weeks pregnant woman came with the complaint of vaginal bleeding for three hours. She had abdominal and back pain. On examination, her hemoglobin level was 9g/dl (Normal range 10.5-13.7 g/dl), blood pressure 90/50mmHg and pulse rate 110/min. The woman was diagnosed for inevitable abortion. What is the priority nursing intervention for this woman?

- ☐ a. Resuscitate with normal saline
- ☐ b. Reassure the woman
- ☐ c. Provide analgesia
- ☐ d. Prepare for blood transfusion



Search...



**Question 60**

Not yet answered

Marked out of 1.00

 Flag question

A pregnant woman who came to a health center was diagnosed with malaria. Inquiring from the patient just if oral medication would be suitable for her, the nurse who made the diagnosis gave the patient a prescription. What is the most likely patient right that the nurse breached?

- ☐ a. The right to get information
- ☐ b. The right to choose treatment
- ☐ c. The right to get respect from the nurse
- ☐ d. The right to control one's own life

[Previous page](#)[Next page](#)**Quiz navigation**[1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#)

Search...





Time left 3:59:56

Question 63

Not yet answered

Marked out of 1.00

 [Flag question](#)

A five-month-old patient was brought to a hospital with the manifestations of pearly white, curdlish material visible on the tongue, palate, and buccal mucosa for two weeks duration. The patient was treated for recurrent infections previously. What is the most likely recommended medication to be given to this patient?

- ☐ a. Acyclovir
- ☐ b. Amoxicillin
- ☐ c. Predinsolon
- ☐ d. Nystatin

[Previous page](#)[Next page](#)**Quiz navigation**

Search...



[Back](#)

Time left 3:56:01

Question 75

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 30-month-old child was brought to the health center with a complaint of cough of three weeks duration. The nurse identified that the father of the child is on initiation phase of TB treatment and the child also had low grade fever. What is the most appropriate initial step to do for this child?

- ☐ a. Put the child on anti-TB treatment
- ☐ b. Initiate INH as TB prophylaxis
- ☐ c. Refer urgently to the hospital
- ☐ d. Counsel on the diagnosis of TB exposure



Search...





Time left 4:35:24

Question 1

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse works on maternal care clinic in a city. One day a woman came to a clinic to get family planning method, but she did not know about the types and effects of the available methods. The nurse gave all necessary information about each family planning methods. Then, the woman used family planning method based on her choice. What is the most likely approach of health education used by the nurse in this scenario?

- ☐ a. Directive approach
- ☐ b. Informed decision making approach
- ☐ c. Persuasion approach
- ☐ d. Deliberate attempt approach



Search...



[Back](#)

Time left 4:35:07

Question 2

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 28th week pregnant mother was admitted to a maternity ward for sever malaria. She was complaining fever, chills, headache, joint pain and loss of appetite. On physical assessment, her blood pressure is 100/60 mmHg, pulse rate 90/min, respiratory rate 24/min and temperature is 38.8 degree centigrade. What is the priority nursing diagnosis for this scenario?

- ☐ a. Impaired nutritional intake
- ☐ b. Hyperthermia
- ☐ c. Fatigue
- ☐ d. Fluid volume deficit



Search...





Time left 4:35:44

Question 100

Not yet answered

Marked out of 1.00

 [Remove flag](#)

Gender office of the city administration organizes a workshop on improving gender equality. Based on the invitation, the CEO of the general hospital made a speech at the event representing the hospital. What is the most likely managerial role played by the CEO?

- ☐ a. Liaison
- ☐ b. Influencer
- ☐ c. Figurehead
- ☐ d. Monitor

[Previous page](#)[Finish attempt ...](#)**Quiz navigation**

Search...



[Back](#)

Time left 4:34:04

Question 6

Not yet answered

Marked out of 1.00

 [Flag question](#)

An infant who was born from a known HIV positive mother is brought to a pediatric OPD with the clinical features of cough, stridor and oral thrush. The HIV status of the infant was positive for antibody test and unknown for DNA/PCR test due to unavailability of the test. What is the appropriate management for the child according to IMNCI?

- ☐ a. Initiate ARV treatment
- ☐ b. Initiate PMTCT care
- ☐ c. Advise to stop breastfeeding
- ☐ d. Advise on HIV prevention



Search...





Nursing (Comprehensive)

[Back](#)

Question 95

Not yet answered

Marked out of 1.00

 Flag question

A nurse was making physical examination for a postnatal mother. The mother was restless, irritable, had pressured speech, and unusual behavior. She had also obsessional and delusional thinking. What is the most likely diagnosis for this woman?

- ☐ a. Postpartum blue
- ☐ b. Postpartum schizophrenia
- ☐ c. Postpartum Psychosis
- ☐ d. Postpartum depression

[Previous page](#)[Next page](#)

[Back](#)

Time left 4:31:23

Question 16

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse researcher studied a hypothesis which read as "Frequent consumption of fruits and vegetables is protective for the occurrence of stomach cancer". He recruited 198 subjects recently diagnosed with stomach cancer and 396 subjects free of stomach cancer. All subjects were asked about their fruit and vegetable consumption during the past one year. What is the most likely study design that the researcher used?

- ☐ a. Ecological
- ☐ b. Cohort
- ☐ c. Cross-sectional
- ☐ d. Case-control



Search...





Time left 4:32:36

Question 11

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 30-year-old gravid-II, para-I pregnant woman came to adult OPD with a complaint of loss of consciousness and generalized convulsions. During physical examination, her temperature was 40 degree centigrade, hematocrit 15 percent and Hyper parasitaemia (greater than 5 percent load). The nurse diagnosed malaria. What is the most likely malaria type which infects the woman?

- ☐ a. Plasmodium falciparum
- ☐ b. Plasmodium vivax
- ☐ c. Plasmodium ovale
- ☐ d. Plasmodium malariae

[Previous page](#)[Next page](#)

Search...





Time left 4:31:10

Question 17

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 30-year-old female patient was diagnosed with ovarian cancer and the oncologist decided the need for radiation therapy. The nurse in charge explained the risk of infertility secondary to radiation therapy. She refused radiation therapy and at the same time she had strong desire to have children. The procedure was done to save her life; she was cured of ovarian cancer and develop infertility What ethical principle most likely forced the health care team's action?

- ☐ a. Beneficence
- ☐ b. Justice
- ☐ c. Nonmaleficence
- ☐ d. Autonomy



Search...



[Back](#)**Question 61**

Not yet answered

Marked out of 1.00

 Flag question

A head of health center discussed the bottlenecks of health care insurance service utilization with community leaders who came from the catchment kebeles. During discussion, the community leaders urged the health center head to address frequent hospital supply stock-out issues. What is the most likely problem prioritization criterion addressed by the head?

- ☐ a. Feasibility
- ☐ b. Community concern
- ☐ c. Sustainability
- ☐ d. Magnitude



Search...





Marked out of 1.00

 [Flag question](#)

A four-year-old child was brought to pediatric emergency with a complaint of breathing difficulty. On examination, the nurse suspected foreign body aspiration. What is the most important clinical feature the nurse should check to confirm this diagnosis?

- ☐ a. Sign of infections
- ☐ b. Hemoptysis
- ☐ c. Gagging
- ☐ d. Atelectasis

[Previous page](#)[Next page](#)

Quiz navigation

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24



Search...





Not yet answered

Marked out of 1.00

 [Flag question](#)

A 45-year-old mother came to a hospital with the complaints of bearing down sensation, dragging discomfort and low back pain. She was admitted to a gynecology ward for a diagnosis of second-degree uterine prolapse. On pelvic examination, soft and reducible mass bulging into the vagina and distending through vaginal introitus. She is now scheduled for surgical procedure. What is the most appropriate nursing care for this mother at this time?

- ☐ a. Physical and psychological preparation
- ☐ b. Advise the mother about exercise
- ☐ c. Teach about patient self-care
- ☐ d. Prescribe and administer sedative drugs

[Previous page](#)

[Next page](#)



Search...



**Question 7**

Not yet answered

Marked out of 1.00

 [Flag question](#)

A group of patients with substance use disorder were admitted to substance rehabilitation center. Most of them deny their problem. The nurse is preparing a teaching plan for them to help their recovery. What is the most appropriate information to be included in the nursing plan?

- ☐ a. Focus on change in the patient's motivation probably has the best impact
- ☐ b. Communicate one can use substance for their stress relieving effect
- ☐ c. Communicate that substance use is a problem not related to individual weakness
- ☐ d. Persons with substance-related problems can recover without formal treatment



Search...





Not yet answered

Marked out of 1.00

 [Flag question](#)

A four-year-old child was presented to the health center with diarrhea of five days duration. On physical examination, the skin pinch goes back within two seconds, and the child was very eager to drink water and had WFH z-score greater than -1. What is the most appropriate management of this child according to IMNCI?

- ☐ a. Give ORS solution in each loss
- ☐ b. Give RL solution intravascular over three hours
- ☐ c. Give ReSoMAL solution in each loss
- ☐ d. Give ORS over 4-hours period

[Previous page](#)[Next page](#)

Quiz navigation



Search...





Question 17

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 26-year-old woman came to a clinic and complained about persistent feeling of sadness, hopelessness, sleep disturbance and loss of interest. The mental status examination revealed that she had difficulty to sustain attention. What are the core symptoms in the diagnosis of the above patient?

- ☐ a. Distractibility and/or sleep disturbance
- ☐ b. Depressed mood and/or loss of interest
- ☐ c. Hopelessness and/or distractibility
- ☐ d. Loss of interest and/or hopelessness

[Previous page](#)[Next page](#)**Quiz navigation**

Search...



[Back](#)

Time left 4:33:32

Question 8

Not yet answered

Marked out of 1.00

 [Flag question](#)

Staff members in one of the public hospitals of Ethiopia, raised salary increment claim to the managers. The managers tried to convince the staff to continue their duties but the staffmembers resisted. At the end, the managers organized a task force and discussed with the health staff and agreed to work their regular time in the hospital and part of their time in private clinics. Finally the conflict is resolved. What is the most likely conflict resolution strategy applied by the managers?

- ☐ a. Lose-lose
- ☐ b. Win-lose
- ☐ c. Resolution aftermath
- ☐ d. Win-win



Search...





MoEEE

MoE - Exit Exam

Time left 4:33:05

A 68-year-old woman was brought to hospital due to progressive deterioration of memory and language ability in the last two years. Her son reports that she couldn't stay at home, wandering on the street and often miss to return to home. During examination, she is alert, has memory problem and talks alone appearing as if discussing with someone. What is the most likely outcome of treating this patient?

- ☐ a. Peaceful death within a short period of time
- ☐ b. Behavioral control with delaying of cognitive deterioration
- ☐ c. Eventual partial return of pre morbid level of functioning
- ☐ d. Eventual full return to pre morbid level of functioning

[Previous page](#)[Next page](#)

Quiz navigation



Search...





Not yet answered

Marked out of 1.00

 Flag question

A four month postpartum woman came to family planning clinic upon return of her menstrual period. The nurse discussed about contraception options, their side effects and time of initiation. The woman decided to use implanon. What is the appropriate site to administer this contraception?

- ☐ a. On the deltoid muscle
- ☐ b. On the Gluteal muscle
- ☐ c. Outer side of the thigh
- ☐ d. On inner side of upper arm

[Previous page](#)[Next page](#)

Quiz navigation



Search...



[Back](#)**Question 21**

Not yet answered

Marked out of 1.00

 Flag question

A nurse working in ART clinic, heard about HIV positive patient who is in love with his neighbor during counseling on his status. The nurse counseled him to stop the relationship. The patient offended and shouted; "I want to infect her and many more so shut-up your mouth". What is the appropriate action to be taken by the nurse?

- ☐ a. Accuse him for endangering the health of the public
- ☐ b. Tell his status to management team
- ☐ c. Further communication and counseling with the patient
- ☐ d. Tell his HIV status to the emergency nurse



Search...



[Back](#)**Question 20**

Not yet answered

Marked out of 1.00

 Flag question

A head of the health center and the health extension workers discussed about service delivery of the health center by involving the catchment farmers. On their discussion, the farmers requested an insurance scheme, which addresses their health and members of their family. What is the most likely health insurance scheme to be recommended in this scenario?

- ☐ a. Community-Based Health Insurance
- ☐ b. Individual Health Insurance
- ☐ c. Social Health Insurance
- ☐ d. Mandatory Health Insurance



Search...



**Question 15**

Not yet answered

Marked out of 1.00

 [Flag question](#)

A newly appointed head of the health center had discussion with catchment kebele leaders to identify real health problems that occurred in the community. The head identified trachoma as major health problem following poor latrine utilization though they have constructed latrine. What would be the most appropriate good care delivery approach used by the head?

- ☐ a. Disease control programs
- ☐ b. Convectional care in referral hospitals
- ☐ c. People-centered primary care
- ☐ d. Conventional care in clinics

[Previous page](#)[Next page](#)**Quiz navigation**

Search...





Time left 4:27:57

Question 28

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse working in an ICU always enters earlier to assess the order and revises cares for all patients. One day, the nurse gets wrong order and doubtful medication for her critically ill patient. Then, the nurse immediately consults the doctor who ordered the medication before she carries out the medication. What is the most likely reason that the nurse wants to consult the doctor?

- ☐ a. Since he is newly assigned, he is not clear with ICU procedures
- ☐ b. To prevent mal practices that risk the patients and to safeguard them
- ☐ c. Fears not to take accountability for any bad consequence
- ☐ d. Lacks confidence to work independently as professional



Search...





Time left 4:26:33

Question 34

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nine-year-old child was brought to a pediatric OPD with complaints of nausea and tenderness. The nurse has suspected that the child may have appendicitis. What additional sign is needed to reach in this diagnosis?

- ☐ a. Positive barlow sign
- ☐ b. Positive kernig sign
- ☐ c. Positive brudzinski sign
- ☐ d. Positive rovsing sign

[Previous page](#)[Next page](#)**Quiz navigation**

Search...





Flag question

- ☐ a. Birth of infant with normal weight for gestational age
- ☐ b. Birth of infant with low weight for gestational age
- ☐ c. Birth of infant with large weight for gestational age
- ☐ d. Birth of infant who initiates breast feeding successfully





Nursing (Comprehensive)

[Back](#)

Time left 4:26:21

Question 35

Not yet answered

Marked out of 1.00

 [Flag question](#)

A researcher has planned to assess prevalence of anemia and associated factor among ANC attendant mothers in health center 'X'. The researcher wants to review the cards of the mothers which were kept in electronic data base. What is the most likely sampling technique to be used by the researcher?

- ☐ a. Cluster sampling
- ☐ b. Multistage sampling
- ☐ c. Stratified sampling
- ☐ d. Simple random sampling



Search...



MILKI HEALTH SCIENCE TUTORIAL CENTER



0910 69 0594
0911 05 8885

TAJAAJILA KENNINU

- ✓ **Barnoota Dabalataa Kennu**
- ✓ **Exit Exam/ COC'f isin qopheessuu**
- ✓ **Leenjiwwan gaggabaaboo kennuu fi kkf**



Teessoon: Magaalaa Naqamtee
Gamoo OSSA Darbii 3^{ffaa}