

A 25 year old female patient presented with barking cough with associated drooling of saliva. On examination the patient prefers to sit in tripod position. Lateral neck x-ray showed thumb sign. What is the most likely diagnosis?

- ☒ a. Epiglottitis ✓
- ☐ b. Retropharyngeal abscess
- ☐ c. Exudating tonsillitis
- ☐ d. Bacterial tracheitis

The correct answer is: Epiglottitis

What is the most important preventive measure after an ARF episode?

- ☒ a. Secondary prophylaxis with benzathine penicillin ✓
- ☐ b. Daily aspirin
- ☐ c. Valve replacement surgery
- ☐ d. Long-term corticosteroids

The correct answer is: Secondary prophylaxis with benzathine penicillin

A 26 year old primigravid lady at 36 weeks gestation comes to the office due to light headedness, and nausea at bed time. Her pregnancy has been uncomplicated and medical history is unremarkable. BP is 115/75 mmHg when sitting, 110/70 mmHg when standing and 80/60 mmHg when supine. Obstetric palpation shows uterus consistent in size with 36 weeks gestation. Which of the following is the most likely explanation for this patient's hypotension while supine?

- ☐ a. Reduced peripheral arterial resistance
- ☒ b. Decreased venous return ✓
- ☐ c. Increased hemodilution
- ☐ d. Increased parasympathetic activity

The correct answer is: Decreased venous return

The paroxysmal stage of pertussis is characterized by:

- ☐ a. Fever
- ☐ b. Asymptomatic
- ☒ c. Paroxysmal cough with whoop ✓
- ☐ d. Mild cough and coryza

The correct answer is: Paroxysmal cough with whoop

A 6-year-old boy presents with exertional fatigue and headaches. Blood pressure is 150/95 mmHg in the upper limbs and 90/60 mmHg in the lower limbs. Femoral pulses are weak and delayed. Echocardiography shows concentric left ventricular hypertrophy. Which congenital heart disease best explains these findings?

- ☐ a. Ebstein anomaly
- ☐ b. Ventricular septal defect
- ☐ c. Dilated cardiomyopathy
- ☒ d. Coarctation of the aorta ✓
- ☐ e. Atrioventricular septal defect

The correct answer is: Coarctation of the aorta

A 33 years old woman Para-II presents to the labor and delivery unit shortly after giving birth at home. She reports significant vaginal bleeding since delivery. On examination her uterus is found to be “boggy” and poorly contracted on abdominal palpation. Her vital signs are BP=90/60, HR=120, RR=22, and she appears pale and diaphoretic. What is the most likely cause of her condition and what will be your first response for this condition?

- ☐ a. Uterine rupture and shout for help
- ☒ b. Uterine atony and shout for help ✓
- ☐ c. Retained placental fragments and remove the fragment by yourself
- ☐ d. Uterine atony and manage by yourself

The correct answer is: Uterine atony and shout for help



A patient with a history of PUD presents with severe, sudden-onset epigastric pain. A chest X-ray shows "free air under the diaphragm." Which of the following is the most appropriate next step in management?

- ☒ a. Immediate upper GI endoscopy ✖
- ☐ b. Barium swallow to locate the perforation
- ☐ c. High-dose IV Proton Pump Inhibitor (PPI) and observation
- ☐ d. Emergent surgical consultation for laparotomy

The correct answer is: Emergent surgical consultation for laparotomy

A 23 yrs old male patient is on his third post operative day after appendectomy for the indication of acute simple appendicitis. he develops abdominal distention, nausea, vomiting and failure to pass faces and flatus. His vital signs are in normal range and bowel sounds are absent. What is the most likely cause of the patients compliant?

- ☐ a. Internal hernias
- ☐ b. Post op adhesion
- ☒ c. Paralytic ileus ✓
- ☐ d. Large bowel obstruction

The correct answer is: Paralytic ileus

A 32 yrs old young adult admitted to surgical ward with consideration of primary lung abscess. A duty physician requested culture and sensitivity test before starting empiric antibiotics. What could be the expected pathogen in this patient?

- ☐ a. There will not be growth
- ☐ b. Predominately aerobic
- ☒ c. Mixed aerobic and anaerobic ✖
- ☐ d. Purely anaerobic

The correct answer is: Purely anaerobic

A 25-year-old male presents to the emergency department after sustaining a penetrating chest injury from a stabbing incident. On physical examination, you find a 3x4 cm sucking wound located over his right anterior-lateral lower chest. The patient is in moderate respiratory distress, and you can hear air being drawn into the wound with each inspiration. What is the immediate management for this patient?

- ☒ a. Cover the entire wound with a sterile dressing ✗
- ☐ b. Administer IV fluids and wait for further imaging
- ☐ c. Chest tube insertion
- ☐ d. Apply a three-sided sterile dressing to the wound

The correct answer is: Apply a three-sided sterile dressing to the wound

Confirmed visceral leishmaniasis in an immunocompetent adult. The best management for this type of patient is?

- ☐ a. Pentamidine
- ☒ b. Sodium stibogluconate plus paromomycin ✓
- ☐ c. Oral miltefosine
- ☐ d. Paromomycin

The correct answer is: Sodium stibogluconate plus paromomycin

A patient has an AST:ALT ratio of 2.5:1. This pattern is most characteristic of:

- ☐ a. Chronic Hepatitis C
- ☒ b. Alcoholic Liver Disease ✓
- ☐ c. Nonalcoholic Fatty Liver Disease (NAFLD)
- ☐ d. Acute Viral Hepatitis

The correct answer is: Alcoholic Liver Disease

A 35-year-old man with a history of rheumatic heart disease presents with fever, weight loss, and new-onset murmur. Blood cultures grow *Streptococcus viridans*. What is the most likely source of infection?

- ☐ a. Gastrointestinal
- ☐ b. Dental procedure
- ☒ c. Skin infection ✖
- ☐ d. Urinary tract infection

The correct answer is: Dental procedure

Which sign is pathognomonic for measles?

- ☒ a. Koplik spots ✓
- ☐ b. Maculopapular rash
- ☐ c. Coryza
- ☐ d. Cough

The correct answer is: Koplik spots

One of the following is not an indication for surgical intervention in a patient with suspected lung abscess

- ☐ a. Abscess rupture/pyopneumothorax
- ☐ b. Inability to exclude a cavitating carcinoma
- ☒ c. An abscess less than 6 cm in diameter ✓
- ☐ d. Contralateral lung contamination

The correct answer is: An abscess less than 6 cm in diameter

Most common cause of peptic ulcer disease worldwide?

- ☐ a. NSAIDs
- ☐ b. Stress
- ☒ c. Helicobacter pylori ✓
- ☐ d. Smoking

The correct answer is: Helicobacter pylori

A term neonate has recurrent focal seizures on day 3. The baby was born via emergency C-section for fetal distress, and the mother had preeclampsia. What is the most likely diagnosis?

- ☐ a. Neonatal meningitis
- ☒ b. Hypoxic-ischemic encephalopathy (HIE) ✓
- ☐ c. Pyridoxine dependency
- ☐ d. Hypocalcemia

The correct answer is: Hypoxic-ischemic encephalopathy (HIE)

A 54 year old known cardiac patient with atrial fibrillation on warfarin presented with two week history of bleeding during sexual activity with offensive vaginal discharge. Cervix is irregular, tender, with contact bleeding upon digital vaginal examination. Her Hgb is 7g/dl, INR is 2. What is the most likely diagnosis?

- ☐ a. PID
- ☒ b. Cervical cancer ✓
- ☐ c. Cervical polyp
- ☐ d. Bleeding diathesis

The correct answer is: Cervical cancer

A 25 year old pregnant woman came to OPD for her second ANC visit at 20weeks of gestation. Her vaccination history showed that she took TT1 a month back and now you provided her second dose and you are discussing about her next dose. When is the appropriate time to provide her next dose of TT vaccine?

- ☒ a. After a month ✖
- ☐ b. After a year
- ☐ c. After 6 months
- ☐ d. After 3 months



The correct answer is: After 6 months

Which type renal failure is commonly seen in a patient having active bleeding with hypotension?

- ☐ a. Acute tubulal necrosis
- ☐ b. Interstitial renal injury
- ☒ c. Pre-renal azotemia ✓
- ☐ d. Post renal azotemia

The correct answer is: Pre-renal azotemia

A 23-year-old man from Kobo presents with 7 days of fever, abdominal pain, and diarrhea followed by constipation. Examination reveals relative bradycardia and faint salmon-colored macules on the trunk. Blood cultures grow a gram-negative bacillus. Which of the following is the most likely diagnosis?

- ☐ a. Malaria
- ☐ b. Enteric fever
- ☒ c. Typhus ✖
- ☐ d. Relapsing fever

The correct answer is: Enteric fever

A 6-year-old presents with migratory arthritis 2 weeks after sore throat and shows dramatic improvement with aspirin. What additional finding supports the diagnosis?

- ☐ a. Thrombocytopenia
- ☐ b. Prolonged PR interval
- ☒ c. Subcutaneous nodules only ✖
- ☐ d. Persistent arthritis

The correct answer is: Prolonged PR interval

A 16 year old home servant lady came with a complaint of being raped four months back and she had been amenorrhic since then. On physical examination the pertinent finding was that, there was 20 weeks sized gravid uterus and FHB was positive by fetoscope. Urine HCG is positive, Hgb is 13.5g/dl and no gross congenital anomaly seen on ultrasound. She was seeking the pregnancy to be terminated. What is most appropriate next step for this client?

- ☐ a. Terminate the pregnancy unconditionally
- ☐ b. Push her to keep the pregnancy
- ☒ c. She needs to bring legal proof, if rape confirmed, terminate the pregnancy ✗
- ☐ d. She needs to bring the family member and if they agree terminate the pregnancy



The correct answer is: Terminate the pregnancy unconditionally



Which type of renal cast is commonly seen patient with acute glomerulonephritis?

- ☐ a. White blood cell cast
- ☐ b. Dysmorphic RBC with RBC cast
- ☐ c. Lipid cast
- ☒ d. Granular cast ✗

The correct answer is: Dysmorphic RBC with RBC cast

A 28-week preterm neonate is born via emergency cesarean section for placental abruption. The delivery room temperature is 23°C (73°F), and the baby is not dried immediately. On admission to the NICU, the axillary temperature is 35.1°C (95.2°F). What is the MOST important immediate intervention?

- ☒ a. Place under a radiant warmer ✓
- ☐ b. Start broad-spectrum antibiotics
- ☐ c. Obtain a complete blood count
- ☐ d. Administer IV fluids

The correct answer is: Place under a radiant warmer

Following are features of necrotizing fasciitis type II except:

- ☐ a. It can give rise to toxic shock syndrome
- ☒ b. It is monomicrobial-mainly beta hemolytic streptococci ✗
- ☐ c. It can occur in young healthy individuals also
- ☐ d. Organism is Clostridium perfringens

The correct answer is: Organism is Clostridium perfringens

A 48-year-old man with long-standing hypertension presents for follow-up. His BP is persistently 158/96 mmHg despite adherence to a thiazide diuretic and ACE inhibitor. He has hypokalemia and metabolic alkalosis. Which is the most likely underlying cause?

- ☒ a. Pheochromocytoma ✖
- ☐ b. Primary hyperaldosteronism
- ☐ c. Coarctation of the aorta
- ☐ d. Renal artery stenosis

The correct answer is: Primary hyperaldosteronism

A 35-year-old male presents to the ER with severe chest pain after being stabbed. On examination, he is in respiratory distress and hypotensive. Chest tube was inserted and approximately 2000ml of blood is drained immediately. What is the most appropriate first step in management?

- ☐ a. Administer pain medication
- ☐ b. Insert a chest tube
- ☐ c. Perform a thoracotomy
- ☒ d. Cover the wound with a sterile dressing ❌

The correct answer is: Perform a thoracotomy

A 45-year-old known type 2 diabetic patient presented complaining of cough and fever of 3 days duration. The patient also has myalgia and fatigue. On physical examination, RR is 26, Bp 120/80, PR 70, T 37.4. He is alert. CBC shows mild leukocytosis. Renal function test is normal. CXR shows lobal consolidation. What is the right management for this patient?

- ☐ a. Amoxicillin
- ☐ b. Ceftazidime and vancomycin
- ☒ c. Amoxicillin-clavulinic acid and Azitromycin
- ☐ d. Ceftraxion and azithromycin



The correct answer is: Amoxicillin-clavulinic acid and Azitromycin



36 year old woman at 38weeks gestation presents with sudden onset of painless vaginal bleeding. On examination the fetal heart rate shows recurrent variable decelerations with stable maternal vital signs. An emergency bedside ultrasound shows fetal blood vessels crossing the internal cervical os. What is the most likely diagnosis and how do you manage?

- ☒ a. Vasa previa and emergency cesarean delivery ✓
- ☐ b. Placenta previa and emergency cesarean delivery
- ☐ c. Abruptio placenta and vaginal delivery
- ☐ d. Occult cord prolapse and emergency cesarean section delivery

The correct answer is: Vasa previa and emergency cesarean delivery



Which of the following distinguish IgA nephropathy from post infectious glomerulonephritis?

- ☒ a. Gross hematuria ✖
- ☐ b. RBC casts
- ☐ c. Normal C3 level
- ☐ d. Proteinuria

The correct answer is: Normal C3 level



A 40-year-old man presents with heart failure symptoms. Echocardiography shows dilated ventricles with reduced ejection fraction. What is the most likely diagnosis?

- ☒ a. Hypertrophic cardiomyopathy ✖
- ☐ b. Constrictive pericarditis
- ☐ c. Dilated cardiomyopathy
- ☐ d. Restrictive cardiomyopathy

The correct answer is: Dilated cardiomyopathy

Gravida III Para II mother comes to emergency OPD with seven hours of pushing down pain and six hours of gush of fluid per vaginum. She has stable vital sign. She has 4 moderate contraction over 10 minutes and fetal heart rate of 136bpm. Cervix is 7cm dilated, station +2 with compound presentation of fetal head in occipito-anterior position and fetal hand palpated on the right side of the head. How do you manage labor in this mother?

- ☒ a. Admission and labor augmentation ✗
- ☐ b. Admission and pantograph follow-up
- ☐ c. Manipulating the prolapsed hand to revert it before admission
- ☐ d. Emergency cesarean section preparation

The correct answer is: Admission and pantograph follow-up

A 12-year-old child presents with migratory polyarthrititis, fever, and a history of sore throat 3 weeks ago.

Eradication of which organism would decrease ARF?

- ☐ a. Group B streptococcus
- ☒ b. Group A β -hemolytic Streptococcus ✓
- ☐ c. Streptococcus pneumonia
- ☐ d. Staphylococcus aureus

The correct answer is: Group A β -hemolytic Streptococcus

A 38-year-old man presents with intermittent fever for 3 weeks, night sweats, fatigue, and diffuse joint pain. He works as a livestock farmer and reports regular consumption of unpasteurized milk. On examination, temperature is 38.8°C, and there is mild hepatosplenomegaly. Laboratory studies show mild anemia and elevated ESR. Which of the following is the most appropriate investigation to confirm the diagnosis?

- ☐ a. Rose Bengal test
- ☐ b. Chest X-ray
- ☒ c. Weil–Felix test ✗
- ☐ d. Blood culture

The correct answer is: Blood culture



A 7-year-old child with Down syndrome presents with neck pain and progressive difficulty walking after a minor fall. What condition should be suspected?

- ☒ a. Disc herniation ✖
- ☐ b. Atlanto-axial instability
- ☐ c. Cervical spine fracture
- ☐ d. Osteomyelitis

The correct answer is: Atlanto-axial instability

The Following facts are true in Ludwig's angina except:

- ☐ a. Putrid halitosis is commonly found
- ☐ b. It can give rise to mediastinitis also
- ☒ c. It is caused by Staphylococcus aureus ✓
- ☐ d. Diffuse swelling in the submental and submandibular region is common

The correct answer is: It is caused by Staphylococcus aureus

A 6-year-old child presents with vomiting and diarrhea for 3 days. On admission, urine output is 0.4 mL/kg/hr, serum creatinine has increased 2.2 times baseline, and FeNa is 0.4%. After fluid bolus, urine output improves significantly. What is the most appropriate diagnosis?

- ☐ a. Postrenal obstruction
- ☒ b. Prerenal acute kidney injury ✓
- ☐ c. Acute interstitial nephritis
- ☐ d. Acute tubular necrosis

The correct answer is: Prerenal acute kidney injury

A 25-year-old electrician is brought to the emergency department after accidentally touching a high-voltage wire. He has an entry wound on his hand and an exit wound on his foot. The skin at both sites appears charred, and he complains of muscle pain and weakness. Vital signs are stable. Which of the following is the most immediate concern in this patient?

- ☐ a. Cardiac arrhythmias due to electrical current passage
- ☒ b. Psychological trauma from the incidence ✖
- ☐ c. Delayed wound infection at the burn sites
- ☐ d. Superficial skin burns at the entry and exit sites

The correct answer is: Cardiac arrhythmias due to electrical current passage

A 17 year old nulligravid girl diagnosed with primary amenorrhea after she presented with a complaint of cyclic lower abdominal pain and failure to see menses. On physical examination there is bulging and bluish membrane on the introitus. What is the most likely diagnosis?

- ☐ a. Low vaginal septum
- ☐ b. High vaginal septum
- ☒ c. Imperforate hymen ✓
- ☐ d. Blind vagina

The correct answer is: Imperforate hymen

A 12-year-old child with a history of congenital heart disease is diagnosed with infective endocarditis. Despite appropriate intravenous antibiotics, he develops recurrent embolic strokes. Echocardiography shows large, mobile vegetations on the mitral valve. Which of the following is the most appropriate next step in management?

- ☐ a. Switch to oral antibiotics
- ☐ b. Add corticosteroids
- ☒ c. Urgent surgical intervention ✓
- ☐ d. Monitor inflammatory markers
- ☐ e. Continue antibiotics for 2 more weeks

The correct answer is: Urgent surgical intervention

A 40-year-old male with blunt abdominal trauma presents with hypotension, tachycardia, and a positive FAST (Focused Assessment with Sonography for Trauma) scan showing free fluid in the abdomen. Which of the following is the most appropriate next step in management?

- ☒ a. CT scan of the abdomen ✖
- ☐ b. Observation and repeat physical exam
- ☐ c. IV fluid resuscitation and blood transfusion
- ☐ d. Immediate exploratory laparotomy

The correct answer is: Immediate exploratory laparotomy

A 60-year-old woman presents with chest pain on exertion that is relieved by rest. ECG is normal at rest. What is the most likely diagnosis?

- ☐ a. Unstable angina
- ☐ b. Stable angina
- ☒ c. Pericarditis ✗
- ☐ d. MI

The correct answer is: Stable angina

A 28 year old woman is in labor with her first child. She has had a normal antenatal course. She is at 39 weeks gestation and has been pushing for two hours. The fetal heart rate is 140BPM and regular. The fetal head is visible at the introitus but no further progress despite adequate contraction of the uterus. On examination the umbilical cord is seen protruding from the vagina. Which of the following is the most appropriate next step of management?

- ☐ a. Cesarean section delivery
- ☐ b. Vaginal delivery
- ☐ c. Forceps delivery
- ☒ d. Vacuum delivery ✗

The correct answer is: Cesarean section delivery

Which heart valves are commonly affected in chronic rheumatic heart disease?

- ☐ a. Pulmonary and aortic valves
- ☒ b. Aortic and mitral valves ✓
- ☐ c. Tricuspid and mitral valves
- ☐ d. Tricuspid and pulmonary valves

The correct answer is: Aortic and mitral valves

The first-line pharmacological treatment for the prevention of variceal re-bleeding in a cirrhotic patient is:

- ☐ a. ACE Inhibitors
- ☐ b. Non-selective beta-blockers (e.g., Propranolol)
- ☒ c. Statins ✗
- ☐ d. Loop diuretics

The correct answer is: Non-selective beta-blockers (e.g., Propranolol)

A 36yr old male from Jara refugee camp presents with fever, severe headache, and a maculopapular rash that began on the trunk and spread outward, sparing the palms and soles. He lives in crowded conditions with poor hygiene. Which vector is most likely responsible?

- ☐ a. Tick
- ☒ b. Louse ✓
- ☐ c. Mosquito
- ☐ d. Mite

The correct answer is: Louse

A 25 years old para-III mother who delivered three days back vaginally had history of prolonged rupture of membrane of 25hrs duration. She developed Endometritis and treated with antibiotics and her neonate developed early onset neonatal sepsis with culture positive for group B streptococcus. What measures could have prevented this infections in in the mother and the neonate?

- ☐ a. Immediate cesarean delivery
- ☒ b. Intrapartum antibiotics treatment ✓
- ☐ c. Vaccination
- ☐ d. Augmentation of labor

The correct answer is: Intrapartum antibiotics treatment

Most appropriate test to confirm H. pylori eradication?

- ☐ a. Endoscopy
- ☒ b. Stool microscopy ✖
- ☐ c. Serology
- ☐ d. Urea breath test

The correct answer is: Urea breath test

Which of the following is the most sensitive indicator of cholestasis and biliary obstruction?

- ☒ a. Alkaline phosphatase (ALP) ✓
- ☐ b. Alanine aminotransferase (ALT)
- ☐ c. Aspartate aminotransferase (AST)
- ☐ d. Serum Albumin

The correct answer is: Alkaline phosphatase (ALP)

A 19-year-old college student presents with fever, neck stiffness, and altered consciousness. CSF analysis shows low glucose, high protein, and neutrophilic pleocytosis. Which organism is most likely responsible?

- ☐ a. *Neisseria meningitides*
- ☐ b. Enterovirus
- ☐ c. *Cryptococcus neoformans*
- ☒ d. *Mycobacterium tuberculosis* ✖

The correct answer is: *Neisseria meningitides*

A 30-year-old male presents with on-and-off abdominal pain and vomiting over the previous week. He has been managing to pass small amount of stools occasionally. He has been admitted to the hospital four times with similar symptoms after his laparotomy for a perforated appendix 5 years ago. What is the most likely diagnosis?

- ☐ a. Recurrence of malignancy
- ☒ b. Anastomotic leak ✖
- ☐ c. Primary colonic tumor
- ☐ d. Postoperative adhesions

The correct answer is: Postoperative adhesions

A 25-year-old man from who was working at Humera as a daily laborer presents with prolonged fever, weight loss, massive splenomegaly, and pancytopenia. Which test is the gold standard test for diagnosis?

- ☐ a. Blood culture
- ☒ b. Splenic aspiration ✓
- ☐ c. Bone marrow aspiration
- ☐ d. rK39 antibody test

The correct answer is: Splenic aspiration

At what age should an HIV-exposed infant undergo definitive PCR testing for HIV diagnosis?

- ☐ a. 6 months
- ☒ b. 4-6 weeks ✓
- ☐ c. At birth
- ☐ d. 18 months

The correct answer is: 4-6 weeks

Single most important etiological factor in papillary carcinoma is:

- ☐ a. Autoimmune thyroiditis
- ☐ b. Multiple endocrine neoplasia syndromes
- ☐ c. Childhood irradiation
- ☒ d. Endemic goiter ✖

The correct answer is: Childhood irradiation

Tension pneumothorax is one of the reasons for which of the following types of shock?

- ☒ a. Hypovolemic shock ✖
- ☐ b. Distributive shock
- ☐ c. Septic shock
- ☐ d. Obstructive shock

The correct answer is: Obstructive shock

A 30 years old male presented with perianal itching and wetness of 2 months duration. On physical examination you find 0.5 by 0.5 cm measuring small opening at the 7 o'clock of the anal verge with minimal serous discharge. Digital rectal examination illustrates a small mucosal defect in the anal canal but there is no mucosal tear or swelling. What is the diagnosis of this patient?

- ☐ a. Intersphincteric perianal abscess
- ☒ b. Perianal fistula ✓
- ☐ c. Anal fissure
- ☐ d. Internal hemorrhoid

The correct answer is: Perianal fistula

Which of the following electrolyte abnormality is commonly seen in patient with chronic kidney disease?

- ☒ a. Hypokalemia ✖
- ☐ b. Hypophosphatemia
- ☐ c. Hypocalcemia
- ☐ d. Hypernatremia

The correct answer is: Hypocalcemia

A 35b year old mother delivered an alive male neonate weighing 3200gram with APGAR score of 8 &9 in 1st and 5th minutes respectively. Immediately postpartum, she had stable vital sign with well contracted 20 weeks sized uterus, no active vaginal bleeding and is able to breast feed. When should this mother be discharged?

- ☐ a. 4 hours
- ☒ b. 24 hours ✓
- ☐ c. 12 hours
- ☐ d. 6 hours



The correct answer is: 24 hours

A 60 year old para III mother come to regular OPD with protruding mass per vagina of eight years duration. She usually replace the mass to urinate. She has also urinary dribbling and on your inspection of external genitalia you observe a pinkish mass with central opening. What additional physical examination you should to localize the site of defect?

- ☒ a. Speculum examination ✓
- ☐ b. Examination with Valsalva maneuver
- ☐ c. Pessary fitting test
- ☐ d. Bimanual pelvic examination

The correct answer is: Speculum examination

A preterm neonate (30 weeks) on day 5 develops abdominal distension, bilious vomiting, and bloody stools. Abdominal X-ray shows pneumatosis intestinalis. What is the diagnosis?

- ☐ a. Meconium ileus
- ☒ b. Hirschsprung disease ✖
- ☐ c. Malrotation with volvulus
- ☐ d. Necrotizing enterocolitis (NEC)

The correct answer is: Necrotizing enterocolitis (NEC)

A 1-year-old suddenly chokes while eating peanuts and becomes cyanotic and unable to cry. What is the IMMEDIATE management?

- ☒ a. Tracheostomy ✖
- ☐ b. Back blows and chest thrusts
- ☐ c. Chest X-ray
- ☐ d. Nebulization
- ☐ e. Rigid bronchoscopy

The correct answer is: Back blows and chest thrusts

Most common malignant tumor in submandibular salivary gland is:

- ☐ a. Adenoid cystic carcinoma
- ☐ b. Mucoepidermoid tumor high grade
- ☒ c. Pleomorphic adenocarcinoma ✖
- ☐ d. Acinic cell tumor

The correct answer is: Adenoid cystic carcinoma

A 25 year old commercial sex worker presented to OPD with the complaint of multiple lesions over her genitalia. On examination, there are multiple 1 to 2 cm pink tan cauliflower like nodules over the vulva and perineum. What is the most likely diagnosis of her presentation?

- ☐ a. Condylomata lata
- ☐ b. Condyloma acuminata/ genital warts
- ☐ c. Squamous cell carcinoma
- ☒ d. Lichen sclerosis ✖

The correct answer is: Condyloma acuminata/ genital warts

A 40 years old man presented with frequent vomiting of ingested matter, abdominal distention, and crampy abdominal pain of 8 hrs. duration. On examination he is hypotensive, tachycardic and has abdominal distention with no guarding or rigidity and tender, irreducible inguinal swelling. Which of the following is the best next step in management option for this patient?

- ☐ a. Fluid resuscitation, NG tube decompression and start conservative management for bowel obstruction once his blood pressure is corrected
- ☐ b. Fluid resuscitation, NG tube decompression, send plain abdominal X-ray and do rectal tube deflation
- ☐ c. Take the patient immediately to operation room for emergency laparotomy
- ☒ d. Fluid resuscitation, NG tube decompression and send abdominopelvic ultrasound and plain abdominal X-ray ✓

The correct answer is: Fluid resuscitation, NG tube decompression and send abdominopelvic ultrasound and plain abdominal X-ray

Which one of the following micronutrient has no public health significance

- ☐ a. Iodine
- ☐ b. Iron
- ☐ c. Zinc
- ☒ d. Thiamine ✓

The correct answer is: Thiamine

A 28 yrs old male sustained a falling down accident from a 3 m tall tree. Immediately after the incident he had lost his consciousness for 5 minutes but get conscious subsequently. Then after about 2 hours he becomes confused and lethargic. Neurologic examination shows dilated pupils on the right side with decreased power on the left side of the body. Ct-scan shows biconvex(lens-shaped) collection on the right temporoparietal area. What is the likely diagnosis of this patient?

- ☐ a. Acute subdural hematoma
- ☒ b. Acute epidural hematoma ✓
- ☐ c. Acute intracerebral hemorrhage
- ☐ d. Acute Subarachnoid hemorrhage

The correct answer is: Acute epidural hematoma

A 65 years old patient who was admitted to ICU for acute coronary syndrome develops right upper quadrant pain, jaundice, high grade fever. His blood pressure dropped to 80/40 mmhg. Bedside ultrasound shows dilated common bile duct and on laboratory examination there is elevation of serum alkaline phosphatase. What is the likely diagnosis of this patient?

- ☒ a. Acute cholangitis ✓
- ☐ b. Acalculous cholecystitis
- ☐ c. Acute cholecystitis
- ☐ d. Choledocholithiasis

The correct answer is: Acute cholangitis

A 9-year-old with severe malaria develops AKI. Urinalysis shows muddy brown granular casts and FeNa of 3%. What is the most likely underlying pathology?

- ☐ a. Bilateral ureteric obstruction
- ☒ b. Tubular epithelial cell necrosis ✓
- ☐ c. Renal vasoconstriction
- ☐ d. Glomerular immune complex deposition

The correct answer is: Tubular epithelial cell necrosis

A child with congestive heart failure has hepatomegaly, pulmonary edema, and poor feeding even at rest. According to Ross classification, this child is in:

- ☐ a. Class II
- ☐ b. Class I
- ☐ c. Class III
- ☒ d. Class IV ✓

The correct answer is: Class IV

A 34 year old woman at 38 weeks of gestation presents with headache, blurred vision, and swelling of the feet. On physical examination her BP=180/120 mmhg and urine protein is+3. Blood test shows elevated liver enzyme and thrombocytopenia. She is given Mgso4 and hydralazine and delivered by vaginally after stabilization. Two days later on post partal period, she develops confusion, agitation, and hallucination. She is mostly symptomatic at night. What is the most likely diagnosis?

- ☒ a. Eclampsia ✖
- ☐ b. Rebound hypertension
- ☐ c. Post-partum psychosis
- ☐ d. Magnesium toxicity

The correct answer is: Post-partum psychosis

A 31 year old woman who is HIV positive delivers a healthy baby. She wants to breast feed her baby. What is the best advice for her?

- ☐ a. Breast feed only if the baby tests negative for HIV at birth and at six weeks
- ☒ b. Breast feed exclusively for the first six months and then introduce complementary feeding but the baby should be on prophylaxis ✓
- ☐ c. Avoid breast feeding and use formula feeding or cow milk
- ☐ d. Breast feed exclusively for the first six months without prophylaxis

The correct answer is: Breast feed exclusively for the first six months and then introduce complementary feeding but the baby should be on prophylaxis

A 30 years old woman who is 28 weeks pregnant presents to routine ANC visit. She is asymptomatic and no chronic medical condition. She is screened for Hepatitis B and found to be positive for hepatitis B Surface antigen (HBsAg). Her serum hepatitis B virus (HBV) DNA level is 2000IU/ml. which of the following is the most appropriate management for this patient?

- ☒ a. Administer hepatitis B immune globulin (HBIG) HBV vaccine to the new born ✓
- ☐ b. Start tenofovir (TDF) prophylaxis
- ☐ c. Start long term anti-viral treatment
- ☐ d. Perform cesarean section delivery

The correct answer is: Administer hepatitis B immune globulin (HBIG) HBV vaccine to the new born

The second top cause of cancer in Ethiopian woman is cervical cancer, a cancer that can be both prevented and detected early. Ministry of health of Ethiopia has launched a program for HPV vaccination. Which age groups are being targeted by this vaccination program?

- ☐ a. 15----25 years old
- ☐ b. 9---14 years old
- ☒ c. One age cohort of 14 years old ✖
- ☐ d. all under age 9

The correct answer is: 9---14 years old

A 29-year-old woman presents with painless swelling on the right side of her neck for 2 months, associated with low-grade fever, night sweats, and weight loss. There is no cough or hemoptysis. Examination reveals matted, non-tender cervical lymph nodes. Chest X-ray is normal. Fine-needle aspiration cytology (FNAC) of the lymph node shows caseating granulomas, and GeneXpert MTB/RIF is positive. What is the most appropriate management?

- ☐ a. Surgical excision of lymph nodes
- ☐ b. Broad-spectrum antibiotics
- ☒ c. Standard anti-tuberculosis therapy for 6 months 
- ☐ d. Anti-TB therapy for 12 months

The correct answer is: Standard anti-tuberculosis therapy for 6 months

A patient experiences recurrent episodes of fever separated by afebrile periods. Peripheral blood smear during febrile episodes reveals spirochetes. What mechanism explains the recurrent nature of the illness?

- ☒ a. Endotoxin release ✖
- ☐ b. Intracellular persistence
- ☐ c. Biofilm formation
- ☐ d. Antigenic variation

The correct answer is: Antigenic variation

A 28-year-old man presents with chronic fever, weight loss, and generalized lymphadenopathy. He reports multiple unprotected sexual encounters. Which of the following is the most appropriate initial diagnostic test?

- ☒ a. HIV-antibody test ✓
- ☐ b. HIV viral load (RNA PCR)
- ☐ c. p24 antigen alone
- ☐ d. CD4 cell count

The correct answer is: HIV-antibody test

A livestock handler from Mersa develops a painless black eschar on the forearm with surrounding edema. He is stable with no systemic symptoms. Best management?

- ☐ a. Vancomycin
- ☐ b. Amphotericin B
- ☒ c. Ciprofloxacin or doxycycline ✓
- ☐ d. Observation only

The correct answer is: Ciprofloxacin or doxycycline

Which of the following types of nephrotic syndrome had the worst prognosis?

- ☐ a. Focal segmental glomerulosclerosis
- ☐ b. C3 glomerulopathy
- ☒ c. Minimal change disease ✖
- ☐ d. Membranoproliferative glomerulonephritis
- ☐ e. Membranous nephropathy

The correct answer is: Focal segmental glomerulosclerosis

A child with Tetralogy of Fallot suddenly develops deep cyanosis, irritability, and squatting behavior. What is the immediate management?

- ☒ a. IV morphine and knee-chest position ✓
- ☐ b. IV furosemide
- ☐ c. IV antibiotics
- ☐ d. Emergency intubation

The correct answer is: IV morphine and knee-chest position

A 34-year-old known RVI patient on HAART with poor adherence and with baseline CD4 count of 48 cells/ μ l presents with progressive headache, fever, and confusion for 10 days. He also has new-onset right-sided weakness and one episode of generalized seizure. He has a history of chronic weight loss and recurrent oral candidiasis. On examination, he is disoriented with right hemiparesis. Fundoscopy is normal. What is the most likely diagnosis for this patient?

- ☐ a. Brain abscess
- ☒ b. CNS toxoplasmosis ✓
- ☐ c. Pyogenic meningitis
- ☐ d. Disseminated tuberculosis

The correct answer is: CNS toxoplasmosis



A 28-year-old man returns from Harra presented with high-grade fever, jaundice, and confusion. Peripheral blood smear shows *Plasmodium falciparum* with 12% parasitemia. Blood glucose is 48 mg/dL. What is the most appropriate initial management?

- ☒ a. IV artesunate ✓
- ☐ b. Oral chloroquine
- ☐ c. IV quinine
- ☐ d. Oral artemether-lumefantrine

The correct answer is: IV artesunate

A 37 years old woman who is HIV positive has a recent viral load of 1200 copies/ml and is in labor. She has been on HAART since the 2nd trimester. What is the best mode of delivery for her?

- ☐ a. Forceps delivery
- ☐ b. Vaginal delivery
- ☒ c. Cesarean delivery ✓
- ☐ d. Vacuum extraction

The correct answer is: Cesarean delivery

A 12 year old female student comes to the clinic with her mother to ask about the best preventive method for cervical cancer after hearing about different prevention options on social media. What should be your response?

- ☒ a. HPV vaccine ✓
- ☐ b. Avoiding Multiple Sexual Partner
- ☐ c. Avoid sexual intercourse
- ☐ d. Regular pap smear

The correct answer is: HPV vaccine

Which of the following is NOT a feature of glomerular hematuria?

- ☒ a. Blood clots ✓
- ☐ b. Dysmorphic RBCs
- ☐ c. RBC casts
- ☐ d. Significant proteinuria (>500 mg/day)

The correct answer is: Blood clots

Adefrs an 18 yrs old tall, slim adolescent who used to smoke 12 packs per year for the past two years presented to emergency department with the complaint of sudden onset left side pleuritic chest pain with accompanying dyspnea on exertion for the past 8 hrs. A day back he flew from Dubai to Addis. Physical finding is consistent with decreased air entry and hyperresonance on the affected side. What could be the likely cause for Adefrs's presentation?

- ☐ a. Spontaneous pneumothorax
- ☒ b. Lung cancer ✗
- ☐ c. Massive hemothorax
- ☐ d. Pulmonary embolism

The correct answer is: Spontaneous pneumothorax

A 25 years old known RVI patient who has had laparotomy for appendiceal abscess 9 days back presented with discharge from the wound site for the past 2 days. On physical examination there is a small separation of the wound edges with serous discharge coming out of the wounds but the underlying fascia is intact. Which of the following correctly matches the diagnosis and the management of the above patient?

- ☐ a. Deep surgical site infection – wound care and antibiotics
- ☐ b. Evisceration – surgical intervention
- ☒ c. Anastomotic leak – laparotomy ✓
- ☐ d. Wound dehiscence - wound care and subsequent closer

The correct answer is: Anastomotic leak – laparotomy

One of the following is ODD

- ☐ a. Pulselessness
- ☐ b. Painless
- ☐ c. Paralysis
- ☒ d. Poikilothermic ✖

The correct answer is: Painless

A 30 year old known type I diabetic woman is planning to conceive. She has no history of diabetic complication and has good glycemic control. Which of the following is the most appropriate time for screening diabetic retinopathy)

- ☒ a. Before pregnancy ✓
- ☐ b. During the first trimester
- ☐ c. During the third trimester
- ☐ d. During second trimester

The correct answer is: Before pregnancy

Characteristic feature of septic shock include following except:

- ☐ a. Hypotension
- ☐ b. Tachycardia
- ☐ c. Oliguria
- ☒ d. Alkalosis ✓

The correct answer is: Alkalosis

A 37 year old primigravida woman visited you at your office for seeking advice about her pregnancy at gestational age of ten weeks. Otherwise she has no chronic medical condition or any complaint. Which of the following best explains the client's risk?

- ☐ a. She is at low risk of giving birth by cesarean section
- ☐ b. She is at low risk of having an increased risk of preterm labor
- ☒ c. She is at high risk of having congenitally anomalous fetus ✓
- ☐ d. She is at low risk of having increased miscarriage

The correct answer is: She is at high risk of having congenitally anomalous fetus



A 35 year old woman has an IUCD device inserted for contraception. Two weeks later, she develops fever, lower abdominal pain, and vaginal discharge. On pelvic examination, there is cervical motion tenderness. A pregnancy test is negative. Which of the following is the most likely complication of the IUCD insertion and what do you do for the IUCD?

- ☐ a. PID and remove IUCD
- ☒ b. PID and leave IUCD inside with PID treatment 
- ☐ c. Ovarian torsion and remove IUCD
- ☐ d. Uterine perforation and remove IUCD

The correct answer is: PID and leave IUCD inside with PID treatment

A 29 year old gravida II para I mother came to ANC at 21 weeks of gestation. Her blood group is O-negative and she didn't take Anti-D during her last delivery. Indirect coombs test is negative. What is the next step in management of this mother?

- ☐ a. Give Anti-D during this pregnancy at 28 weeks and within 72 hrs post-delivery if neonatal blood group becomes positive
- ☐ b. Give Anti-D during this pregnancy at 28 weeks and within 72 hrs post-delivery irrespective of the neonatal blood group and Rh type
- ☒ c. Determine antibody titer ✗
- ☐ d. Give Anti-D within 72 hrs of post delivery

The correct answer is: Give Anti-D during this pregnancy at 28 weeks and within 72 hrs post-delivery if neonatal blood group becomes positive

A patient with a brain abscess secondary to chronic sinusitis is started on empiric therapy. Which regimen is most appropriate?

- ☐ a. Acyclovir + cefepime
- ☒ b. Ampicillin + gentamicin ✖
- ☐ c. Ceftriaxone + metronidazole
- ☐ d. Vancomycin alone

The correct answer is: Ceftriaxone + metronidazole



A 2-year-old child presents with sudden onset cyanosis. Oxygen saturation does not improve with 100% oxygen. Echocardiography shows parallel great vessels. What is the most urgent lifesaving intervention?

- ☐ a. Immediate VSD closure
- ☐ b. IV diuretics
- ☒ c. Beta-blocker therapy ✖
- ☐ d. Prostaglandin E1 infusion



The correct answer is: Prostaglandin E1 infusion



A 4-week-old first-born male presents with progressive projectile non-bilious vomiting and weight loss. On physical examination, he has sunken eye ball, irritable and on abdomen he has palpable olive-shaped mass. What is the MOST important initial management step?

- ☐ a. Observation
- ☐ b. Antibiotics
- ☐ c. Immediate surgery
- ☒ d. IV fluids and electrolyte correction ✓
- ☐ e. Oral feeding

The correct answer is: IV fluids and electrolyte correction



A 26 years old primigravida lady whose GA is 38 weeks plus 5 days admitted with the diagnosis of latent first stage of labor (cervix 3cmdilated,80% effaced, station -1, membrane intact and clinically adequate pelvis). After 8 hrs of stay she was reevaluated and the pelvic assessment showed cervix 3cm and 80% effaced with the same station. ARM was done it showed G-III Meconium stained amniotic fluid. What is the most appropriate step in managing this patient?

- ☒ a. Augmentation with oxytocin ✗
- ☐ b. Instrumental assisted delivery
- ☐ c. Prepare her for emergency cesarean delivery
- ☐ d. Give her additional 4 hrs

The correct answer is: Prepare her for emergency cesarean delivery

A newborn presents with cyanosis, single loud S2, and chest X-ray shows an egg-on-string appearance. Which physiologic shunt is essential for survival?

- ☐ a. Ventricular level shunt
- ☐ b. Atrial level shunt
- ☒ c. Any intracardiac shunt ✓
- ☐ d. Patent ductus arteriosus

The correct answer is: Any intracardiac shunt

A 35 year old stage two hypertension presents for contraceptive counseling. She has no history of other medical problems. She is married and sexually active with her husband, but she doesn't want to have more children. She is currently using condom but didn't satisfied with it. What is the least ideal contraceptive option for this patient?

- ☒ a. Combined oral contraceptive ✓
- ☐ b. Depo medroxyprogesterone acetate injection
- ☐ c. IUCD
- ☐ d. Implant

The correct answer is: Combined oral contraceptive

newborn has delayed passage of meconium stools and barium enema radiograph shows dilated proximal colon and small obstructed distal colon. What should be the next gold standard diagnostic test?

- ☐ a. abdominal CT-scan
- ☒ b. Chromosome analysis ✖
- ☐ c. stool studies
- ☐ d. rectal suction biopsies

The correct answer is: rectal suction biopsies

A 30-year-old male is brought to the emergency department after a fall from a height. He is complaining of severe chest pain and shortness of breath. On examination, you notice decreased breath sounds on the left side and a shift of the trachea to the right and hypotension. Chest X-ray confirms a large radiolucent area in the left thoracic cavity. What is the most likely diagnosis?

- ☒ a. Tension pneumothorax ✓
- ☐ b. Pneumothorax
- ☐ c. Hemothorax
- ☐ d. Pulmonary contusion

The correct answer is: Tension pneumothorax

A 58-year-old man presents with progressive shortness of breath, orthopnea, and bilateral leg swelling. On examination, BP is 150/95 mmHg, bibasal crackles are present, and an S3 gallop is heard. Chest X-ray shows cardiomegaly and pulmonary congestion. What is the most likely underlying mechanism?

- ☐ a. Increased venous return
- ☐ b. Reduced cardiac output due to systolic dysfunction
- ☐ c. Reduced preload
- ☒ d. Increased heart rate ✗

The correct answer is: Reduced cardiac output due to systolic dysfunction

A 26 year old Para I Mother came to OPD for preconceptual care. On her previous delivery she gave birth of a baby with spinal bifida. When and what preventive therapy should started in this client to prevent recurrence of the event?

- ☐ a. one month before pregnancy and iron tablet
- ☐ b. one to three months before conception & the first three months of pregnancy and high dose iron tablet
- ☐ c. First three months of pregnancy and high dose folic acid
- ☒ d. one to three months before conception & the first three months of pregnancy and high dose folic acid ✓

The correct answer is: one to three months before conception & the first three months of pregnancy and high dose folic acid

A 60-year-old female presents with a history of blunt trauma to the abdomen. She has persistent hypotension despite IV fluids and blood transfusion. A CT scan shows a large retroperitoneal hematoma. What is the most likely site of injury?

- ☐ a. Spleen
- ☒ b. Colon ✖
- ☐ c. Pancreas
- ☐ d. Kidneys

The correct answer is: Kidneys

Mintewab is a 65 yrs old known cardiac patient on follow up for the past 10 months currently presented with severe crampy postprandial abdominal pain of 6hrs duration. The pain aggravated by taking a meal. On physical examination she has severe pain out of proportion with the physical findings. CT angiography result shows huge thrombus at the bifurcation of superior mesenteric artery. What is the likely cause?

- ☒ a. Acute mesenteric ischemia ✓
- ☐ b. Chronic mesenteric ischemia
- ☐ c. Portal vein thrombosis
- ☐ d. Diverticulosis

The correct answer is: Acute mesenteric ischemia

An 8-month-old infant presents with episodic crying, drawing up of legs, vomiting, and red currant jelly stools. What is the MOST likely diagnosis?

- ☒ a. Intussusception ✓
- ☐ b. Hirschsprung disease
- ☐ c. NEC
- ☐ d. Appendicitis
- ☐ e. Volvulus

The correct answer is: Intussusception

A patient with chronic heart failure is started on an ACE inhibitor. What is the main benefit of this drug in heart failure?

- ☒ a. Reduce ventricular remodeling ✓
- ☐ b. Increase myocardial contractility
- ☐ c. Reduce heart rate
- ☐ d. Increase preload

The correct answer is: Reduce ventricular remodeling

Which one of the following statements is incorrect?

- ☐ a. Starting a patient on alpha adrenergic antagonist will raise PSA value by 50% in a patient with BOO
- ☐ b. Combining adrenergic antagonist with 5 alpha reductase inhibitors have synergistic effect
- ☒ c. Bladder outlet obstruction is exclusively for males ✓
- ☐ d. Urethral catheterization is contraindicated unless urethral injury is ruled out

The correct answer is: Bladder outlet obstruction is exclusively for males

A 12-year-old presents with fever, pallor, petechiae, oliguria, and history of bloody diarrhea one week ago. Labs show anemia, thrombocytopenia, and elevated creatinine. Which lab abnormality is MOST expected?

- ☒ a. Prolonged PT and aPTT ✖
- ☐ b. Elevated anti-streptolysin O
- ☐ c. Schistocytes on peripheral smear
- ☐ d. Low complement levels

The correct answer is: Schistocytes on peripheral smear

A 60 years old male presented with urinary frequency, urgency, dysuria, hesitancy, and dribbling of 6 months duration. He developed inability to void over the past 24 hrs and severe suprapubic pain. Physical examination shows stable vital signs, suprapubic tender swelling and enlarged prostate with smooth borders and palpable median sulcus. What is the best next appropriate management of this patient?

- ☒ a. Initiate Alpha blocker agents ✖
- ☐ b. Transurethral catheterization
- ☐ c. Give analgesics and follow conservatively for 24 hours
- ☐ d. Emergency transurethral resection of prostate

The correct answer is: Transurethral catheterization



A 24 year old commercial sex worker complaining of profuse, whitish vaginal discharge with fishy odor during intercourse. Microscopic examination revealed clue cells. What is the most likely diagnosis in this patient and how do you treat her?

- ☐ a. Syphilis and B. Penicillin IM
- ☐ b. Gonorrhea and ceftriaxone IM injection
- ☐ c. Trichomonas Vaginalis and Metronidazole PO
- ☒ d. Bacterial vaginosis and metronidazole PO ✓

The correct answer is: Bacterial vaginosis and metronidazole PO

A primigravida lady at gestational age of 25 weeks presented with crampy lower abdominal pain of five hours duration. She has associated passage of watery fluid per vagina. Otherwise no history of vaginal bleeding/passage of tissue. On physical examination vital signs are stable, no abdominal tenderness, cervix is closed and watery fluid per vaginum is noted. On ultrasound examination cardiac activity is negative, amniotic fluid is minimal and biparital diameter is 14 weeks. What is the likely diagnosis?

- ☐ a. Missed abortion
- ☐ b. Threatened abortion
- ☒ c. Inevitable abortion ✖
- ☐ d. Incomplete abortion

The correct answer is: Missed abortion



What should be done for the above patient?

- ☐ a. Do abscess drainage
- ☒ b. Start IV antibiotics ✖
- ☐ c. Do emergency laryngoscopy in ICU
- ☐ d. Start IV dexamethasone

The correct answer is: Do emergency laryngoscopy in ICU

Which of the following is not an indication for amputation?

- ☐ a. Dead-loss limb
- ☐ b. All
- ☒ c. None ✓
- ☐ d. Dead limb
- ☐ e. Deadly limb

The correct answer is: None

One of the following is not the common sign of Graves' disease:

- ☐ a. Pretibial myxedema
- ☐ b. Cardiac Failure
- ☒ c. Proximal myopathy ✓
- ☐ d. Proptosis

The correct answer is: Proximal myopathy

A term neonate fails to pass meconium beyond 48 hours of life and develops abdominal distension. Rectal biopsy shows absence of ganglion cells. What is the diagnosis?

- ☒ a. Meconium plug syndrome ✖
- ☐ b. Hirschsprung disease
- ☐ c. Cystic fibrosis

The correct answer is: Hirschsprung disease

While you are evaluating a suspected sexual abused girl, you are worried about the possibility of pregnancy as the incident happened four days back and her LMP was 20 days back. What is the best option of contraception for this girl to prevent pregnancy?

- ☒ a. Depo-Provera ✖
- ☐ b. Post pill/levonorgestrel
- ☐ c. IUCD
- ☐ d. Implanon

The correct answer is: IUCD

Which complication of portal hypertension is defined by an absolute neutrophil count (ANC) ≥ 250 cells/mm³ in ascitic fluid?

- ☐ a. Hepatorenal Syndrome
- ☒ b. Spontaneous Bacterial Peritonitis (SBP) ✓
- ☐ c. Budd-Chiari Syndrome
- ☐ d. Hepatic Hydrothorax

The correct answer is: Spontaneous Bacterial Peritonitis (SBP)

A 2 year and 3mo old male toddler presented to WCSH pediatric OPD by his mother with a complain of decrease spontaneous movements on his left upper and lower extremities which is more severe on the arm than the leg .He started right hand preference around the age of 15 month and walk around 18monhth of age.On physical examination the child walks on tiptoe and circumductive gait is apparent. What is the most likely diagnosis

- ☐ a. Spastic hemiplegia
- ☐ b. Spastic diplegia
- ☒ c. Spastic monoplegia ✖
- ☐ d. Extrapyramidial CP

The correct answer is: Spastic hemiplegia

A 25-year-old university student presents complaining of productive cough of 2 weeks duration. The patient also has fever loss of appetite and weight loss. On examination, crepitation over right posterior chest. His HIV sero status is negative. Sputum acid fast examination is positive for acid fast bacilli. What type finding is expected in this patient's chest X-ray?

- ☐ a. Multilobar consolidation
- ☐ b. Cavitory lesion on the upper lobe
- ☐ c. Cavitory lesion on the middle and lower lobes
- ☒ d. Interstitial infiltration ✖

The correct answer is: Cavitory lesion on the upper lobe

