

6. For a fixed number of ideal stages in a distillation column, as the reflux ratio is increased, the difference in composition between the top and bottom product streams
- A. increases
 - B. decreases
 - C. remains unaffected
 - D. passes through a maximum

Answer: Option A

Explanation:

No answer description available for this question. [Let us discuss.](#)

[View Answer](#)

[Discuss in Forum](#)

[Workspace](#)

[Report](#)

27. According to Chilton-Colburn analogy for mass transfer, $N_{St}.N_{Sc}^{2/3}$ is equal to
- A. f
 - B. $f/2$
 - C. $2f$
 - D. $1/f$

Answer: Option B

Explanation:

No answer description available for this question. [Let us discuss.](#)

[View Answer](#)

[Discuss in Forum](#)

[Workspace](#)

[Report](#)

28. A pure drug is administered as a sphere and as a cube. The amount of drug is the same in the two tablets. Assuming that the shape and size do not influence the mass transfer, the ratio of rate of dissolution in water at $t = 0$ for the cubic to spherical tablet is
- A. 0.54
 - B. 1.04
 - C. 1.24

D. 1.94

Answer: Option **C**

Explanation:

No answer description available for this question. [Let us discuss.](#)

[View Answer](#)

[Discuss in Forum](#)

[Workspace](#)

[Report](#)

29. Minimum reflux ratio in a distillation column results in

- A.** optimum number of trays.
- B.** minimum reboiler size.
- C.** maximum condenser size.
- D.** minimum number of trays.

Answer: Option **C**

Explanation:

No answer description available for this question. [Let us discuss.](#)

[View Answer](#)

[Discuss in Forum](#)

[Workspace](#)

[Report](#)

30. Flooding in a distillation column is detected by a sharp

- A.** increase in Murphree plate efficiency.
- B.** decrease in pressure drop.
- C.** decrease in liquid hold up in the column.
- D.** increase in pressure drop.

Answer: Option **D**

Expla