

Public Health COC Exam

Part One

Direction: Each of the following questions is followed by four possible alternative answers. Read each question carefully and blacken the letter of your best choice on the separate answer sheet provided.

1. A 21 yrs old male pt. came with complaint of productive cough of one month duration. He also has low grade fever, night sweats, weight loss & loss of appetite. On examination he is chronically sick looking with vital signs in the normal range. He has no remarkable syndromes. Laboratory investigation of Sputum microscopy showed mycobacterium bacilli. What is the best next step of the management of the pt?

A) _____

2. A 40yr old known RVI pt. who has been on ART for the last three yrs came with a complaint of whitish coating over his tongue with difficulty & pain during swallowing of two wks duration. His Vital signs are in a normal range. On examination there is cheese coating over the tongue, extending to the pharynx, which bleeds while scraped using a spatula.

What is the WHO clinical stage of the pt?

A) Stage I B) Stage II C) Stage III D) Stage IV

3. A 20 yr old male came with complaints of on and off type high grade fever of five days duration. He also has history of headache and arthralgia. He is acutely sick looking & his cloth is infested with body lice. His PR is 116/min, RR are 20/min, Temp. is 40.5 ° C and BP is 100/70 mmHg. Blood film shows multiple **spirochetes**.

What is the drug of choice for the management of this pt.

A) Doxycycline C) Erythromycin
B) Ciprofloxacin D) Penicillin

4. A 30 yr old male pt. came to a clinic after he sustained a nail prick injury at his leg while working at a construction site a day back. Upon evaluation the wound is deep & clean. What is the most appropriate next step in the management of this pt?

A) PO antibiotics C) Wound debridement
B) Tetanus anti-toxin D) Tetanus immunoglobulin

5. A 34 yr old male pt came with a complaint of burning of epigastric pain of two wks duration. He was repeatedly treated with anti acid & omeprazole for the last three yrs. He is chronically sick looking with vital sign in the normal range. On examination he has pale conjunctiva & non icteric and there is mild epigastric tenderness. Laboratory investigation results show hemoglobin level of 8.4mg/dl and negative H.Pylori test. What is the most likely anticipated complication in this pt?

A) Gastric outlet obstruction C) Perforated peptic ulcer
B) Gastrointestinal bleeding D) Gastric lymphoma

6. A 26 yrs old male presented with complaints of bloody diarrhea & fever of 12 hrs duration. He reported that a day back he had eaten raw vegetables at a local restaurant. His PR is 116/min, Temp 38 ° C , BP 90/60mmHg and RR 20/min. He has no remarkable systemic findings on examination. What is the most likely Dx of this pt?

A) Shigellosis C) Taeniasis
B) Giardiasis D) Cholera

7. A 54 yr old male came with a complaint of abdominal distention of one month duration. His PR is 100/min, RR 20/min, BP 100/70 mmHg and temp is 36.8°C. On examination he has greatly distended abdomen with positive signs of fluid collection, gynecomastia & erythematous palms. On investigation HBSAg is Reactive. What is the most likely Dx of this pt?

A) Chronic liver disease 2° to Autoimmune hepatitis
B) Chronic liver disease 2° to Chronic Viral hepatitis
C) Chronic liver disease 2° to drug induced hepatitis

D) Chronic liver disease 2° to alcoholic hepatitis

8. A 24 yr old male came with a complaint of cough of seven days duration. He also has high grade fever, chest pain & fast breathing. Otherwise, he has no history of hospital admission of Rx for similar condition. His PR is 108/min, RR 26/min and temp. 38.1°C. On examination there is dullness and bronchial breath sound over over the left middle 1/3rd of the posterior chest. Chest X-Ray shows that there is a homogeneous opacity over the middle third of the right chest. What is the next best step to the management of this pt?

A) Clarithromycine

C) Ceftriaxone

B) Augmentin

D) Cefalexine

9. A 28 yr old male presented with complaints of shortness of breath & dry cough of two wks duration. He also has low grade fever & decrease appetite. He is chronically sick looking with vital signs in the normal range. On respiratory examination, there is dullness and abscess air entry over the posterior lower 1/3rd of the left chest. What is the most likely explanation for this finding?

A) Consolidation

C) Bronchieotasis

B) Lung abscess

D) Pleural effusion

10. A 29 yr old female who is known asthmatic pt. on intermittent Salbutamol puff for the last two yrs came for routine follow up. She states that she has been using her Salbutamol puff more than three times a day to get relieved to the last one month. What is most appropriate step to the management of this pt?

A) Oral corticosteroid

C) Low dose inhaled corticosteroid

B) Long acting B-agonist (LABA)

D) High dose inhaled corticosteroid

11. A 27 yr old female came with a complaint of fever of five days duration. She has history of IV drug use for the last one year. Her BP is 120/78 mmHg, PR is 120/min, RR 24/min & temp. is 39.2°C. On cardiovascular examination, there is systemic murmur at the apex. Laboratory investigation result shows Hgb level of 9.5 mg/dl & positive blood culture for **staphylococcus aureus**. What is the most likely Dx of this Pt.?

A) Infective endocarditis

C) Acute Pericarditis

B) Acute rheumatic fever

D) Myocarditis

12. A 66 yr old male came to a clinic for BP check up. Two months back his BP was 144/92 mmHg & he was advised to have salt free diet & to start follow up. Now his BP is 158/99 mmHg, PR 74/min, RR 14/min & temp. is 36.7°C. There is no remarkable systemic finding on examination. His serum creatinine is 0.9 mg/dl (NR Serum Cr.= 0.6-1.2 mg/dl). What is the next step in management of this Pt.?

A) Start single PO anti hypertensive drug
measurement

C) Appoint her after two days for repeat

B) Start combined PO anti hypertensive drug
him home

D) Advise on life style modification & send

13. A 28 yr male came to an Em OPD with a complaint of shortness of breath of six days duration. He also had pleuritic type of chest pain which improves with leaning forward & worst with supine position, high grade fever & dry cough of four days duration. He is mostly sick looking with BP of 70/50 mmHg, PR of 112/min, RR of 32/min, temp. of 38.7°C. Distended neck veins & distal heart are the findings on cardiovascular examination. What is the most likely Dx of the Pt.?

A) Severe Pneumonia

C) Pulmonary embolism

B) Carotid stenosis

D) Spontaneous pneumothorax

14. A 36 yr old male came with complaint of watery diarrhea & vomiting of two days duration. He was being treated with oral antibiotic & ORS but has no improvement. Three days back he started to complain of decreased urine output & headache. He is lethargic with BP of 140/97 mmHg, PR 122/min, RR of 24/min & temp. of 36.8°C. On investigation serum creatinine level is 5.4 mg/dl & stool examination shows trophozoite of Giardiasis (NR Serum Cr.= 0.6-1.2 mg/dl). What is the most likely cause of the pts. Current condition?

A) Oral antibiotics

C) Sepsis of GI focus

B) GI fluid loss

D) Obstructive uropathy

15. A 35 yr old woman came with complaint of right side flank pain of one day duration. She also has high grade fever & repeated vomiting. She is acutely sick looking with BP 90/60 mmHg, PR 124/min, RR of 22/min & temp. of 39.2°C. On examination there is right side costovertebral angle tenderness. Laboratory investigation result shows WBC count 19,000/mm³ & 12 WBC/HPF on urinalysis. What is the next best step in the management of this pt?

A) Abdominal Ultrasound

C) Start PO antibiotics

B) Urine analysis

D) Start IV antibiotics

16. A 40 yrs old male pt. presented with complaint of pain & swelling of the first toe of his right foot which began at night time four days ago. The pain & swelling was progressively worsening. He had similar condition a year back. On musculoskeletal examination joints of toe affected & toe swelling, warm to touch & extremely tender. What is the most likely clinical Dx?

A) Rheumatoid arthritis

C) Septic arthritis

B) Acute gouty arthritis

D) Osteoarthritis

17. A 19 yr old male pt presented with a complaint of easy fatigability associated with occasional fever for the past three weeks. He also had repeated episodes of epistaxis in the past three days. He is acutely sick looking, in pain. With BP of 100/70 mmHg, PR of 124/min, RR of 24/min and temp. of 37.8°C. On examination he has pale conjunctiva, CBC results revealed WBC count of 40,000/mm³, Hgb level of 10 mg/dl & platelet count 60,000/ml. What is the most likely clinical Dx?

A) Severe sepsis

C) Acute osteomyelitis

B) Hemolytic anemia

D) Hematologic malignancy

18. A 40yr old female pt. was admitted to a medical ward with a diagnosis of severe community acquired pneumonia five days back. She was started on prompt treatment with IV antibiotics but her vital signs fail to improve. On the sixth day of admission, she developed confusion. Her BP is 80/40 mmHg, PR is 136/min, feeble RR are 26/min & temp is 38.8°C. What is the most appropriate time to start vasopressor medication for the pt?

A) After the fever subsides

C) Immediately after the first record of BP

B) After 24 hrs of IV antibiotics

D) When BP fails to improve after adequate IV

fluid

19. A 22 yr old female student tried to commit suicide & ingested a chemical. She brought to emergency unit within half an hour. The mother claimed the chemical was "malathion". At presentation she had one episode of vomiting. Otherwise she is with vital signs in the normal range. What is the next best step in the management of this pt?

A) Decontaminate with charcoal

C) Give her IV atropine

B) Start IV fluid with NS

D) Give IV antibiotics

20. A 45 yr old male pt came with complaint of untreatable vomiting of ingested matter which was intermittently blood tinged of four months duration. He has lost significant amount of weight since the onset of his problem. He looks emaciated, with BP of 75/60 mmHg & PR 100/min. Plain abdominal X-Ray result shows enlarged gastric bubble & dilated proximal duodenum with paucity of air in the small bowel. What is the most likely electrolyte abnormality that this pt. could have?

A) Hypokalemia & Hypochloremia

C) Hypernatremia & Hypermagnesemia

B) Hyperkalemia & Hyperchloremia

D) Hypercalcemia & Hyperphosphatemia

21. A 38 yrs old female came with a complaint of right breast lump of one year duration which was gradually increasing in size. She also has an associated irregularity on the covering skin with nipple inverted. On examination there is a 1x4 cm sized firm & irregular mass on the right breast with pus orange appearance of the covering skin & a 2x2 cm sized right axillary lymphadenopathy. What is the most likely Dx of this

pt?

- A) Chronic breast abscess
- B) Breast TB
- C) Breast cancer
- D)

22. A 32 yr old male came to the OPD after he sustained a cut injury to his hand while working on a construction at a work place. He stated that he was vaccinated for tetanus twice about eight years back. What is the appropriate management for the pt?
- A) Give him only tetanus immunoglobulin (TIG)
 - B) Give him only tetanus anti toxoid (TAT)
 - C) Give him both TAT & TIG
 - D) Clean wound & send home
23. A 20 yrs old male came to emergency OPD with complaint of abdominal pain, vomiting of ingested water & loss of appetite associated with low grade fever of two days duration. He was holding his right side abdomen & bending down while coming to the OPD. On examination there is direct tenderness at the right lower abdomen. What is the most likely Dx of the pt?
- A) Peptic ulcer disease
 - B) Acute cholecystitis
 - C) Acute appendicitis
 - D) Acute pancreatitis
24. A 54 yr old female came with a complaint of abdominal pain of three days duration. She also has abdominal distention, failure to pass stool & intermittent vomiting of ingested water. Her BP is 80/60 mmHg, PR is feeble. There is distended abdomen with asymmetric shape & hyper tympanic on examination. What is the first line of management of this pt?
- A) Insert an NG tube & observe
 - B) Prepare her for emergency surgery
 - C) Resuscitate & insert a rectal tube
 - D) Start her on broad spectrum antibiotics
25. A 40 yr old male came with a complaint of bleeding just after defecation of three weeks duration. He also has excruciating pain during defecation and bleeding is bright red. He was having constipation two weeks prior to the onset of his current symptoms. On examination, there is a linear mucosal defect with a severe tenderness and raised border on the posterior wall of the rectum. What is the most likely Dx of this pt?
- A) Hemorrhoid
 - B) Anal fissure
 - C) Fistula in-ano
 - D) Ano-rectal cancer
26. A 68 yrs old male presented with complaints of failure to pass urine and lower abdominal pain of 10 hours duration. He has been to a nearby HC for the same complaint and the nurse tried inserting NG tube to relieve the retention but failed. On examination bladder is palpable and tender, and on digital rectal examination the prostate is enlarged and firm with movable rectal mucosa. What is the most appropriate initial step in the management of this pt?
- A) Condom catheter
 - B) Suprapubic catheter
 - C) Aspiration with needle
 - D) Trans urethral catheter
27. An 18 yr old student came to a HC with complaint of pain in the groin and lower abdominal area of six hours duration. His BP is 110/70mmHg, PR is 110/min, temp is 37.4°C and RR are 22/min. On genitourinary examination there is swollen, tender and high riding testis. Laboratory investigation results shows WBC of 10000/mm³ and urine analysis is non-revealing. What is the most Dx for the pt?
- A) Varicocele
 - B) Testicular torsion
 - C) Fournier's gangrene
 - D) Epididymo-orchitis
28. A 10yr old male presented to a HC with a complaint of severe pain over the right forearm after he sustained falling down accident from a two meter height. On musculoskeletal examination, the right forearm is swollen and tender with palpable deformity. Right forearm X-ray shows that there is a transverse fracture over the distal 1/3 of the ulnar bone, What is the appropriate initial step in the mgt of the pt?
- A) Apply posterior slab
 - B) Apply circular cast
 - C) Apply arm sling
 - D) Apply traction
29. A 12 yrs old male presented to a hospital with a complaint of severe pain over the right side of his pelvic which is not relieved by strong analgesics of five months duration. Pelvic x-ray shows that there is skin

appearing reaction. What is the most likely Dx?

- A) Paget's sarcoma
- B) Chondrosarcoma
- C)
- D)

30. A 35 yrs old male came with complaint of fast breathing and chest tightness after he fell down from the second floor to a stony ground on his chest. His RR is 35/min, and oxygen saturation is 70%. There is shifting of trachea to the left side and absent air entry on the right lung field with hyper resonant note on examination. What is the next step in the management of this pt?

- A) Refer him directly to hospital
- B) Put him on oxygen & order chest x-ray
- C) Insert a chest tube & put him on oxygen
- D) Do needle thoracentesis at 2nd intercostals space

31. A 4 year old male child was presented immediately after he sustained electrical burn injury to his arm. On musculoskeletal examination there is circumferential eschar over his forearm with a 3x4 cm wound over his forearm. He was resuscitated at the hospital. What is the most common complication this pt can develop?

- A) Arrhythmia
- B) Acute renal failure
- C) wound contracture
- D) Compartment syndrome

32. A 50 yr old male came with a complaint of left leg swelling of six months duration which worsens up on standing & long distance walking & relieved during sleeping. On examination there are superficial coiled, tortuous vessels seen on the medial surface of his left leg extending to his thigh. What is the appropriate initiated management for this pt?

- A) Encourage him to stay in supine position for longer hours
- B) Advise him to use elastic bandage on daytime
- C) Start him an anti coagulant medication
- D) Refer him to a specialist directly

33. An 18 month old child brought with a complaint of intractable vomiting of ingested matter & discomfort in his neck since two days ago. Chest x-ray showed that there is a circular opaque lesion behind the trachea. What is the most likely Dx of this child?

- A) Foreign body aspiration
- B) Foreign body ingestion
- C) Esophageal stenosis
- D) Esophageal tear

34. A three week infant born at 32 weeks of gestation & weighing 1750g, was brought to OPD with complaint of abdominal distension, feeding intolerance, and vomiting. On examination, the infant is lethargic with grossly distended & tender abdomen. Laboratory investigation shows leucopenia & thrombocytopenia. Plain abdomen x-ray shows air in the bowel wall. What is the most appropriate immediate step in the management of the infant?

- A) NPO with IV fluids
- B) Systemic antibiotic
- C) Surgical consultation
- D) Nasogastric decompression

35. A four day infant, delivered at home, was presented with complaints of irritability & difficulty in opening of the mouth associated with hunger & crying. Physical examination revealed tachypnea, tachycardia, fluctuating temperature & severe spasms with stimulation. What measure could have prevented this condition from occurring?

- A) Active immunization of the neonate with DPT
- B) Maternal immunization with tetanus toxoid
- C) Human tetanus immunoglobulin
- D) Prophylactic antibiotics at birth

36. A neonate born to a diabetic mother, was brought to the clinic with irritability & jitteriness. It is large, plethoric & tremulous. Physical examination reveals tachycardia, tachypnea & hypothermia. What is the most likely explanation for this condition?

- A) Hyperglycemia
- B) Hypercalcemia
- C) Hypoglycemia
- D) Polycythemia

37. A one year old infant was brought to the clinic with complaint of low grade fever and runny nose followed by paroxysm of whooping cough of two weeks duration. He has shortness of breath & becomes restless during the episodes of cough, and vomits following the paroxysms of cough. The mother says that the infant has not been vaccinated. Which vaccine could have prevented this condition?
- A) BCG B) OPV C) DPT D) Vitamin A
38. A mother brought her two yr old child with complaints of coryza, cough, conjunctivitis, and fever. The mother states that the child has a rash that began from the face & spread caudally. The immunization record is not up-to-date. Physical examination is remarkable for a morbilliform rash over the body including the palms, and tiny grayish white dots on the buccal mucosa next to the third molar. What is the most likely Dx?
- A) Rubella B) Mumps C) Measles D) Varicella.
39. A five yr old boy was brought to HC after having watery diarrhea of one day duration. He also has abdominal pain & vomiting of ingested matter, He drinks water eagerly. On examination, the child is irritable & has sunken eye balls & skin pinch goes back slowly. What is the most appropriate initial management for the pt?
- A) Give fluid & food to treat diarrhea at home C) Give fluid for severe dehydration
B) Give fluid & food for some dehydration D) Give antibiotics for the infection
40. A four yr old girl was brought to a HC with complaints of cough, fever, & shortness of breath of two weeks duration. She has weight loss & loss of appetite. She was given Amoxicillin syrup a week back for the same complaint but she has no significant improvement. Her father has been coughing for the past six months. She is fully vaccinated. Her vital signs & anthropometry are in the normal range. Her ESR is 80 mm/hr. What is the most likely diagnosis?
- A) Bronchitis C) Presumptive tuberculosis
B) Pneumonia D) Hyperactive airway disease
41. An eight yr old child came to an OPD complaining of intermittent fever of one day duration. He also has repeated vomiting of ingested matter, one episode of generalized abnormal body movement & reddish discoloration of urine. On examination, his temp. is 39.6°C, GCS is 14/15 has pale conjunctiva with icteric sclera. A laboratory investigation result shows Hematocrite level of 28%, RBS of 45mg/dl & serum creatinine level of 1.1 mg/dl. (NR Serum Cr.= 0.6-1.2 mg/dl). What is the most prevalent severity sign in this child?
- A) Jaundice C) Renal failure
B) Convulsion D) Haemoglobinuria
42. A five yr old child was presented with a complaint of foul smelling diarrhea associated with crampy abdominal pain, bloating of three weeks duration. Stool examination shows a trophozoite stage of a parasite. What is the most appropriate first line drug for this case?
- A) Tinidazole C) Albendazole
B) Praziquantel D) Metronidazole
43. A seven yr old child has easy fatigability, dyspnea, & body swelling. On examination, the child is in cardio-respiratory distress with tachypnea and tachycardia, & he has paper white conjunctiva & palms, a gallop rhythm, hepatomegaly, & pitting pedal edema. Management for heart failure was initiated at the emergency. What is the most likely precipitant of the heart failure in the child?
- A) Rheumatic fever C) Myocarditis
B) Severe anemia D) Arrhythmia
44. A seven yr old child came with a referral from another Hospital. She has been on Frusemide & prednisolone for acute rheumatic fever for the past two weeks. What is the most appropriate next step in the pt's care?
- A) Follow up with weekly Benzathine penicillin C) Follow up with monthly Benzathine penicillin
B) B.Penicillin for subsequent 10 days D) Follow up with single dose of Benzathine

pencillin

45. A four yr old child was presented with complaints of barking cough, hoarseness of voice & inspiratory stridor. He was fully vaccinated. On physical examination. The child is cyanotic & in respiratory distress, his temp. is 38.5 °c , has stridor at rest & pharynx was cryptematous. What is the initial Rx for this child?
- A) Dexamethazone
B) Use of coal mist
C) Air way management
D) Nebulized racemic Epinephrine
46. A 10 month old boy was brought to an emergency OPD after having abnormal body movement which is characterized by flexion & extension of extremities, drooling of saliva & up rolling of the eyes that lasted for approximately three minutes. He was well except that he had fever preceding the abnormal body movement. His body temp is 39°C . Laboratory investigations are non-revealing. What is most appropriate next step in the management of this child?
- A) Diazepam
B) Antipyrotics
C) Antiepileptic Drugs
D) Reassure & send home
47. A 12 month old infant, who was on formula feeding since one month, was brought to the OPD for routine checkup. The mother reported that she has been feeding her child goat's milk. Vital signs were in the normal range. On CBC test, Hgb is 9.8g/l with MCV of 114.5fl, & Neutrophils are large & have hyper-segmented nuclei. What is the most likely explanation for this finding?
- A) β - thalassemia
B) Iron deficiency anemia
C) Sideroblastic anemia
D) Folate deficiency anemia
48. A four yr old child was presented with complaint of vomiting & diarrhea of three days duration. On physical examination, vital signs are in the normal range, weight for height is 60%, & there is bilateral pitting edema. What is the most likely Dx?
- A) Marasmus
B) Kwashiorkor
C) Moderate malnutrition
D) Marasmic-Kwashiorkor
49. A 5 yr old male child who was admitted in the ward & being treated for severe acute malnutrition (marasmus) for the past one week, currently developed profuse watery diarrhea of 12 hours duration. On physical he is lethargic with PR of 120/min, weak hair broken eye ball & skin pinch goes back slowly. What is the most appropriate next step in the management of this child?
- A) ResoMal 10 ml/kg over the first hour
B) RL IV 20ml/kg over 30 minute
C) RL IV 15ml/kg over the 3rd hour
D) RL 3ml/kg in 30 minute for b two hours.
50. A 15 yr old child came to an OPD for regular checkup. He reported that he has frequent micturition which is more at night, excessive thirst
51. A seven year old girl who is known diabetic pt. & is on Insulin, presented with complaints of abdominal pain & vomiting of two days duration. She also has increased frequency of micturition & thirst. She is confused & disoriented to time & place but her vital signs are in the normal range. Treatment was initiated immediately & after two hours her RBS becomes 170 mg/dl, & venous PH becomes 7.1 What is the most appropriate next step in the management of the pt.
- A) Add 5% Dextrose to fluid
B) Increase same dose of Insulin
C) Change IV fluids to oral
D) Discontinue the Insulin
52. A 32 yr old G-II mother came to the ANC clinic for first visit with complaints of severe nausea & persistent vomiting. She is amenorrhic for the past 12 weeks. She believes termination of the pregnancy since she is not tolerating these symptoms. What is the most appropriate advice for the mother?
- A) Advise her that this condition is treatable
B) Advise her to use ginger and spicy foods
C) Advise her on complications of abortion
D) Advise her on the line of termination
53. A 24 yr old G-III, P-I, A-I mother, Whose LNMP is not known but claims to be amenorrhic for the past one & half months, came to the HC with complaints of vaginal bleeding & severe lower abdominal pain of two days duration. She has history of recurrent STI treatment since the age of 18 yrs. Her BP is 90/50 mmHg, PR is 120/min, and tem. Is 36.2°C . On abdominal examination, there is left lower quadrant

tenderness & on pelvic examination cervix is closed & there is blood on examining finger. Urine HCG test was positive two days before her presentation. What is the most appropriate next step in the management of her condition?

- A) Abdominal Ultrasound
- B) Repeat urine HCG
- c) Serum Beta HCG
- D) Pelvic Ultrasound

54. A 28 year old G-III, P-I, A-I (Induced abortion) mother came to a HC for the first ANC visit. Her LNMP is not known but she claimed to be amenorrheic for the past 12 weeks. She is a known RVI Pt. on ART (TDF + 3CT + EPV) for the last five yrs. Her vital signs are in the normal range. Routine ANC investigations results are normal. What is the most appropriate management of the pt?

- A) Continue the same ART regimen
- B) Change the ART regimen
- C) Refer her to hospital
- D) Repeat the HIV test

55. A 32 yr old G-IV, P-III mother presented to an emergency OPD at 38 weeks of gestation with a complaint of bright red & painless vaginal bleeding of two days duration. She has no other complaint & her ANC course was uneventful. Her vital signs are in a normal range. Obstetric examination revealed a 38 wks sized uterus with a FHB of 144/min and no active vaginal bleeding. Ultrasound showed a singleton pregnancy with reassuring biophysical profile and placenta implanted in the lower uterine segment covering the internal cervical os. What is the next step in the management of the mother?

- A) Reassure her & send her home
- B) Admit her & start corticosteroids
- C) Admit her & prepare for elective CS
- D) Start induction of labor with oxytocin

56. A 26 yrs old G-III, P-II mother presented with a pushing down pain of six hours duration on a gestational stage of 39 wks. Her BP was 120/80 mmHg, PR is 85/min and temp is 36.6°C. On obstetric examination FHB is in the range of 140 to 150/min. Cervix is dilated at 5 cm & the cervix is at station -3. Up on artificial rupture of membrane, there was G-I meconium signal liquor and fetal bradycardia in the range of 70 to 80/min is noted for 3 minutes without recovery. What is the most likely Dx?

- A) Cord Prolapse
- B) Uterine Rupture
- C) Maternal Hypotension
- D) Meconium stained liquor prolapse

57. A 23 yrs old G-I woman presented to the labor ward at active first stage of labor at 40 weeks of gestation. She had no complicated prenatal course. Her BP is 100/60 mmHg. On examination fundal height is 40 cm with FHB of 156/min, and membranes are intact. After one hour of stay in the labor ward the FHB tracing becomes in the range of 160-166/min for two minutes. What is the most appropriate next step in the pt's care?

- A) Perform Amniotomy
- B) Perform cesarean delivery
- C) Put her in lateral position
- D) Put the woman in knee-chest position

58. A 24 yr old G-I woman who is at gestational age of 40 wks was started on induction of labor for gestational hypertension. She was given oxytocin for 12 hours & reached a cervical dilation of 6 cm. She underwent cesarean delivery for an indication of non-reassuring fetal heart rate pattern. After delivery of the placenta, profuse bleeding was noted from the uterus, reaching 1500 ml. What is the most likely cause for the pt's problem?

- A) Cervical laceration
- B) Uterine inversion
- c) Retained tissue
- D) Uterine Atony

59. A 59 ye old P-II mother came a clinic with a complaints of increased abdominal girth associated with bloating, early satiety and urinary frequency of six months duration. She stated that she used to have irregular menstrual cycle & her age at menarche was at 10 yrs. On pelvic examination, there is a solid, fixed and irregular left adnexal mass. Routine laboratory investigation results are unremarkable. What is the primary factor determining the prognosis of the pt?

- A) Duration of the illness
- B) Parity of the pt.
- C) Stage of the disease
- D) Family history

60. A 23 yr old single woman came to for a clinic for consultation about her reproductive health status. She had three sexual partners. Her age at first sexual intercourse was 11 yrs. Her past gynecologic, medical and surgical histories are unremarkable. All the routine tests including HIV and PAP smear results are normal. What is the next recommended PAP smear screening schedule for the woman?
- A) Every one year
 - B) Every three years
 - C) Every five years
 - D) Every 10 years
61. A 47 yrs old P-VI mother presented to a HC with a complaint of pelvic pressure and feeling of incomplete emptying of the bladder associated with low back pain of four years duration. Her past gynecologic, obstetric, medical and surgical histories are unremarkable. On pelvic examination, there is relaxed and open genital hiatus with a thin-walled, smooth, and reducible bulged mass, the posterior and anterior vaginal walls are also bulged. What is the most likely Dx?
- A) Pedunculated myoma
 - B) Uterovaginal prolapsed
 - C) Gardner's cyst
 - D) Lipoma of vagina
62. A five yr old child was brought a pediatric OPD with a complaint of ear pain and low grade intermittent fever of four days duration. On physical examination, temp is 37.9 °c but other vital signs were in the normal range, Otoscopic examination revealed a bulged and hyperemic membrane, and minimal middle ear effusion. What is the next step in the management of the pt?
- A) Treat with oral antibiotics
 - B) Send home with reassurance
 - C) Advice to ear wash twice daily
 - D) Admit and treat with IV antibiotics

Part Two

Direction: Each of the following questions is followed by four possible alternative answers. Read each question carefully and blacken the letter of your best choice on the separate answer sheet provided.

63. A four yr old girl has visited different health facilities with a complaint of frequent tonsillitis and was treated with oral antibiotics. Her parents noted that she has restless sleep, frequent choking episodes while sleep, and a loud snoring. Physical examination revealed her vital signs are in the normal range and tonsils are enlarged with no erythema and exudates. Otoscopic examination is normal. For which condition is the child at great risk?

- A) Respiratory failure
- B) Tonsillar neoplasm
- C) Vocal cord paralysis
- D) Obstructive sleep apnea

64. A 38 yr old male presented with diffuse submandibular swelling associated with fever and throat pain of one week duration. Incision and drainage was done for the Dx of submandibular abscess and an antibiotic is to be started, What type of antibiotics should be considered?

- A) Active against facultative anaerobes
- B) Active against gram positive aerobes
- C) Active against gram negative aerobes
- D) Active against aerobes & anaerobes

65. A 25 yr old man came complaining of plaque accumulation on his lower arch and bleeding with slight touch to his gingival area. On examination, there was plaque which bleeds with manipulation for examination. What is the effective long term control method for this case?

- A) Scaling
- B) Flossing

66. A 28-year-old male presented with post-operative pain after extraction of right lower side molar tooth was done five days back. The pain was dull, radiating to ear that not responding to analgesics. He claimed that he failed to follow the post operation instructions. On examination, there was foul smelling food debris on extraction site. What is the most likely Dx of the pt?

- A) Osteomyelitis
- B) Osteoradionecrosis
- C) Alveolar osteitis
- D) Garre osteomyelitis

67. A 30-year-old woman came with a complaint of a short, sharp and shock type of tooth pain which gets exacerbated during tooth brushing of two weeks duration on examination she had gingival recession but no dental caries.

What is the most likely diagnosis for this case?

- A) Acute periodontal pain
- C) irreversible pulpitis

B) Dentinal sensitivity

D) reversible pulpitis

68. A patient with avulsed primary central incisors after he sustained an accident an hour back came to the hospital. He has put the tooth in a clean plastic bag and brought it with him. What is the most effective management for the patient?

A) Replacement and fixation

C) ordering soft diet only

B) Reassure and send home

D) give iv antibiotics

69. A 46-year-old man came to the clinic for check up. On mental status examination he has poor eye contact, and depressed mood with restricted and mood congruent affect, his speech is low in volume and tone and his thoughts are irrational. He has suicidal ideation and plan, with a previous history of suicidal attempt, What is the suicide risk level of this patient?

A) Low risk

C) minimal risk

B) High risk

D) moderate risk

70. A 32-year-old female patient presented with hallucination, delusion and disorganized speech for the last seven months. She has stopped working since one year back. What is the most likely diagnosis?

A) Brief psychotic disorder

C) delusional disorder

B) Schizoaffective disorder

D) schizophrenia

71. A 32-year-old man came with a complaint of pain less mass on the right upper eyelid of three month duration, ocular finding of the right upper lid revealed a non-tender localized nodule under the skin.

What is the most likely clinical diagnosis of the patient?

A) Chidazion

C) sebaceous cyst

B) Hondeolum

D) epithelial inclusion cyst

72. A 34-year-old male patient came with a complaints pain and blurring of vision on his left eye of three days duration after he sustained a falling down accident. He also has headache fever and malaise. Findings on ocular examination were marked eyelid swelling. Restricted ocular motility. Visual reduction and proptosis. There is leukocytosis on CBC.

What is the most appropriate initial step in the management of the patient?

A) Admit and start him on the IV antibiotics

B) Send him home with oral antibiotics

C) Admit and start him on eye drops

D) Refer him to ophthalmologist

73. A 35-year-old female presented with sudden onset left eye pain and blurring of vision of three hours duration which is associated with severe headache and photophobia. She has no history trauma or similar symptoms. Ocular finding in the left eye shows visual acuity of hand motion. Injected conjunctiva, cloudy cornea and irregular poorly reactive pupil.

What is the most likely diagnosis of the patient?

A) Anterior uveitis

C) angle closure glaucoma

B) Herpetic keratitis

D) herpetic keratoconjunctivitis

74. A four-year-old child was brought by her parents with a complaint of whitish central spot on both eyes of one week duration. On history she was not vaccinated fully. She had history of maculopopular rash with runny nose and cough about two weeks back. On ocular examination. She has central corneal ulcer bilaterally.

What is the most appropriate initial management for the child?

- A) Vitamin A supplementation
- B) Topical antibiotics
- C) investigation
- D) referral

75. A 50-year-old male patient with parkinson's disease came with a complaint of skin lesion of three months duration, on examination, there are erythematous plaque lesions over the scalp. Eye brows and posterior auricular area bilaterally with yellowish greasy scaling

What is the most likely diagnosis?

- A) Dandruff
- B) Psoriasis
- C) lichen planus
- D) Seborrheic dermatitis

76. A 15 year old girl came with asymptomatic skin lesion of one year duration. Upon evaluation there are open and closed comedos with scattered papules over the face, mostly on the forehead.

What is the most appropriate Rx for this stage of the disease?

- A) Oral Dapsone
- B) Topical retinoid
- C) Photodynamic therapy
- D) Oral contraceptive pills

77. A known RVI pt. who has been on HAART for the past six months, came with a complaint of fever, chills and generalized reddening of the skin of the skin five days duration. His temp. is 39.3°C, BP is 90/60 mmHg, PR is 126/min and RR is 27/min. On examination, the skin appears glazed, eroded with areas of coalescing erythema and Nikolsky sign was positive. Which drug is the most likely cause of the pt's condition?

- A) Zidovudine
- B) Nevirapine
- C) Tenofovir
- D) Indinavir

78. A 23 year old housewife came with a complaint of pruritic skin lesion of three weeks duration. She stated that her symptom started a month after her husband came back from field work. On examination, there is skin colored and excoriated papules over the trunk and extremities mainly involving the axilla, nipples and the finger web spaces. What is the most likely Dx of this pt?

- A) Scabies
- B) Syphilis
- C) Papular Urticaria
- D) Pruritic papular eruption

79. A number of pts are waiting for their turn in line to be seen by a health professional. When the health professional came one of the pts who know him very well, asked him to be seen first. But the health professional refused and politely convinced him to stay in the queue with the other pts. Which ethical principle does the health professional implement?

- A) Justice
- B) Autonomy
- C) Beneficence
- D) Non-maleficence

80. A 35 yr old man came to a hospital with complaint of recurrent headache of two

weeks duration. During evaluation, the physician did not listen to the pt. but keep on asking him leading questions and interrupting him repeatedly. Which quality of compassionate care is violated in this care?

- A) Respect
- B) Empathy
- C) Motivation
- D) Commitment

81. A health care professional, who has been working in emergency department for the past two years with a strong sense of motivation, started to come late to work. All his co-workers witnessed his gradual behavioral change, they further explained that he isolates himself and becomes uncompassionate to the pts too. What is the most likely cause of his behavioral change?

- A) Burnout
- B) Compassion fatigue
- C) The wider health care context
- D) Unbalanced focus between biomedical

models

82. A pt came to a HC for treatment. On evaluation, the health care provider cannot speak the language of the pt which made it difficult to communicate. There is also wide cultural difference which made the case very strong to the health professional. What is the most appropriate principle to be followed in managing the pt?

- A) Recognize diversity
- B) Uphold responsibility
- C) Value work place culture
- D) Challenge dignity barrier

83. A 28 yr old depressed male pt came for a psychiatric evaluation. While interviewing, the psychiatrist used a statement like "I understand how difficult it is being unable to sleep for three days". Then, the pt continued to tell his problems without reservations. What is the most likely reason of using empathic statement during the interview?

- A) To help the pt to disclose himself
- B) To help the provider to lead the interview
- C) To help the pt to feel being understood
- D) To help the provider to get more information

84. A health professional, who is working in a referral hospital, also works in a private diagnostic center which he owns. This has resulted in a conflict of interest between the hospital and him. What is the best recommendation to reduce the conflict in this case?

- A) Disclose his affiliation
- B) Use external monitoring
- C) Discontinue his affiliation
- D) Continue his affiliation secretly

85. A health professional uses an open-ended question at first and close-ended question in the middle of an interview to help him reach at diagnoses. Later, he talks about Dx and alternative treatment. At the end, he developed a treatment plan collaboratively with the pt. and agreed to divide responsibility for the next session. What is the evidence that understanding is established in the case?

- A) Two way communications is implemented

- B) Interview technique of the health professional
- C) Dx and Rx alternative is communicated
- D) Decision on the treatment plan is made collaboratively

86. A physician who has examining a 48 yr old female pt, reached at a Dx with a poor prognosis. What is the most likely issue that should be considered in disclosing the Dx?

- A) Worthiness of telling the news
- C) Psychological issues
- B) Availability of professional support
- D) One's ability to accept the pt's reaction

87. A public health officer, who is interviewing pts, uses a separate folder to record cocky pt's history and all records related to his pts. He was a secured cabinet with a lock to put the records of the pt's and to maintain confidentiality. What is the primary reason that the health professional tries to maintain confidentiality?

- A) To show respect
- C) To prevent harm
- B) To promote trust
- D) To promote autonomy

88. In an outpatient clinic with a very crowded waiting area, a health professional was giving seat and telling pts to wait until he calls their name. One pt stood up and insulted him but he politely asked him why he is reacted that way and listened to his complaint. What professional moral value explains the health professional's judgment?

- A) Giving attention to pt is good
- C) Considering what pts say is helpful
- B) Pts have the right to be aggressive
- D) Treating pt with consideration is good

89. The number of cancer pts & related deaths is increasing worldwide. Following this, country leaders are organizing pharmaceutical research institutes to produce effective treatment for this disease. What is the most appropriate research type for this scenario?

- A) Basic
- C) Exploratory
- B) Applied
- D) Explanatory

90. A researcher has traced published & unpublished literatures related to his research area from national and international research outputs. Finally, he has got many related literatures and uses them accordingly. What is the most likely violated literature review criterion in this sensrio?

- A) Comprehensive
- C) Relevance
- B) Up-to-date
- D) Balance

91. Investigators wanted to assess the effect of school based iron supplementation on cognitive development. They randomly assigned participants in to two groups, those who will receive and will not receive iron supplementation. The supplementation will provided on a weekly basis for one year. What is the most appropriate study design for the above scenario?

- A) Prospective Cohort
- C) Experimental
- B) Cross-sectional
- D) Case control

92. A research institution wants to identify the type of cardiovascular morbidity in a community. The institution grouped the hospitals based on their rank as primary, secondary &

tertiary level. The hospitals were randomly selected from each group to conduct the study. What is the most appropriate sampling method used in this scenario?

- A) Sample random sampling
- B) Cluster random sampling
- C) stratified random sampling
- D) systematic random sampling

93. A hospital manager wishes to know what proportions of pts admitted to the orthopedic department has died. A 95% confidence interval is desired with a width intervals of 0.09 of the population proportion are estimated as 0.20 from a previous study. What is the sample size needed for the above study?

- A) 10
- B) 87
- C) 171
- D) 174

94. An epidemiologist has planned to estimate the prevalence of sudden infant death syndrome (SIDS) among infants admitted to the ICU of a hospital. What is the most appropriate measurement tool to be used?

- A) Observational checklist
- B) Document review checklist
- C) self administered questionnaire
- D) face to face interview questionnaire

95. The Ethiopian medical journal reported a study of 13 HIV positive pts, who were treated with HAART for at least six months. The CD4 T-cell counts as baseline were as listed below. 130,168,205,313,207,227,245,173,103,181,105,301,169. What is the Inter Quartile Range (IQR) for this data set?

- A) 87
- B) 149
- C) 210
- D) 245

96. Hemoglobin A1C is a serum measure often used to monitor compliance among diabetic pts. A sample of five values of Hgb A1C obtained from a single diabetic pt. are 8.5%, 9.3%, 7.9%, 9.2% and 10.3%. What is the most appropriate measure of variation for this sample?

- A) Standard deviation
- B) Inter quartile range
- C) variance
- D) range

97. Cardiovascular risk factors such as diastolic blood pressure (mmHg) and HDL cholesterol levels (mg/dl) were measured among children in every 3 year intervals. It was noted that the coefficient of variant (CV) was 7.1% for blood pressure and 10.4% for HDL cholesterol. What is the correct statement about the above scenario?

- A) The variability in cholesterol level measurements and BP cannot be compared
- B) There is high variability in cholesterol level measurements than blood pressure
- C) There is high variability in blood pressure measurements than cholesterol level
- D) There is consistence of measurements in cholesterol level than blood pressure.

98. A study was conducted among 18male and 11 female hypertensive pts to measure tire serum creatinine concentration level. The male pts had a mean level of 1.7 mg/dl with a standard error of the mean of 0.4, while the female pts had a mean of 3.4 mg/dl with standard error of the mean of 1.03. The critical values for the distribution are 2.052 at degree of freedom. What is the 95% confidence interval of mean difference?

- A) -1.10, 4.50
- B) 0.89, 2.51
- C) -1.23, 4.63
- D) 0.85, 2.55

99. A group of researchers planned to conduct a study on Alzheimer's pts. The researchers are interested in comparing the rate of obesity between sexes of pts. From the total 20 pts included, 10 of them are males. What is the most appropriate statistics for this scenario?

- A) T-test
- B) Z-test
- C) χ^2 -test
- D) F-test

100. A group of researchers want to examine the association between ANC service utilization and birth

outcome. They have included 250 pregnant women who have ANC follow up and do not have. 10 among ANC attendants and 200 among non-ANC attendants delivered neonates with low birth weight. What is the value of relative risk for this scenario?

- A) 0.06 B) 1.10 C) 10.00 D) 0.25

101. A national wise case control study was conducted to assess the association of life skill training and academic dismissal among fresh female university students. Odds ratio of 0.4 was reported for students who took life skill training compared to students who did not take the training. What is the most appropriate interpretation of this finding?

- A) The odds of dismissal is 0.4 times less likely in those who took life skill training than those who did not
B) The odds of dismissal is 0.6 times less likely in those who took life skill training than those who did not
C) The odds of dismissal is 40% more likely in those who took life skill training than those who did not
D) The odds of dismissal is 60% more likely in those who took life skill training than those who did not

102. A manager of a HC engaged with the staffs during a sanitation campaign in the compound. The manager showed the staff who it can be done, and also supported and encouraged them to finish the sanitation campaign effectively and efficiently. What management function is most likely applied?

- A) Leading B) Planning C) Evaluation D) Organizing

103. A medical officer is performing a quarterly supervision in the laboratory unit of a health center. He assessed the available materials and their functionality using an inventory checklist. What managerial skill is being implemented?

- A) Conceptual B) Technical C) Communication D) Human relation

104. A manager in a primary hospital assigned the health work forces of the hospital based on their expertise. He also allocated the resources based on the prioritized problems of the hospital. What managerial principle is applied?

- A) Order B) Equity C) Remunerate D) Unit of direction

105. A health center's manager plans to improve the utilization of institutional delivery until the end of 2012 E.C by assigning committed midwives and creating awareness in the community.

What is the most appropriate principle of management to be applied?

- A) Converge of work C) management of objective

- B) Learning from experience D) management of exception

44. A performance management team of a primary hospital selected twelve indicators of health services to assess the monthly performance of a hospital out of the twelve service indicators, three indicators showed very low performance when compared with the intended plan.

Who is the primary responsible decision maker to solve the problem?

- A) Zonal health office C) primary health care unit
B) Regional health bureau D) federal ministry health
45. A manager of a primary hospital faced shortage of inpatient rooms in the hospital. He then asked the regional health bureau to provide financial supports however the regional health bureau could not provide the support. Thus the manager mobilized stakeholders and the society to make financial aid to the hospital to build inpatient room and it was successful.
Which principle Of primary health care is most likely applied?
A) Equity C) community participation
B) Decentralization D) inter-sectoral collaboration
46. The ministry of health has been working on community awareness creation and publicizes the important concepts and practices of self- responsibility in health and health related issues as most of the country's health problems are preventable.
What policy emphasis is most appropriate in this case?
A) Development of environmental health
B) Promotion of occupational health and safety
C) Development of an appropriate health service management system
D) Control of communicable diseases, epidemic and diseases related poor living conditions
47. A manager of a primary hospital conducted an assessment on the financial sources for the health care services provided in the hospital. And it was found that about 37% of the patients pay from their pocket to get services.
What is the most likely conclusion about the financial source of the hospital?
A) Private contribution
B) Federal and regional governments
C) Non-governmental organizations contribution
D) Grants and loans from bilateral and multilateral donors
48. A Non-governmental organization has been implementing the integrated management of neonatal and childhood illnesses in a country eight years and phased out before one year. As a program evaluator. A health officer want to asses the awareness, attitude and behavior of mothers towards the treatment of neonatal and childhood illnesses.
What type of evaluation is the most appropriate?
A) Impact evaluation C) formative evaluation
B) process evaluation D) summative evaluation
49. A country located at seas bore was hit by an earthquake followed by a massive land slide. The disaster caused loss of life, property damage and environmental degradation. What is the most appropriate category of the category?
A) Biological disaster C) Geographical disaster
B) Complex disaster D) Hydrological disaster
50. In a metropolitan city of a country, gun violence disaster became a major public health problem. To address the problem, the responsible bodies established a team to solve this public health problem. What is the most appropriate initial management to be taken by the team?
A) Investigation of the cause of the violence C) Integrated approach to stop the gun violence

- B) Provision of support for the victim families D) Public awareness to cope with consequence of the violence
51. A hospital's waste disposal team planned to dispose a mixed healthcare waste, which was mostly composed of contaminated bandage, gloves, paper and card boards. What will be the most appropriate waste management technique?
- A) Separating & disposing C) Incinerating all & land filling the ash
B) Disinfecting all and land filling D) Controlling burning and burial of the ash.
52. Both developing and developed countries in the world use chlorine as the main water disinfectant compared to other available resources though it has known side effects. What is the most likely reason?
- A) Chlorine is cheap and available in all parts of the world
B) Chlorine can kill all type of microorganisms in the water
C) Chlorine is safe to use and easy to apply in all types of water
D) Chlorine protect the water from contamination after treatment
53. A public health expert received a report of contamination of alike by pesticides and mercury from industries in the locality consequently, small aquatic animals will likely consume these chemicals which will be eaten by fishes, later, the fish will be used as food by people. What is the mostly likely fate of the chemicals?
- A) Bioaccumulate

B) Excreted by the animal

C) Disintegrate by the metabolism

D) Accumulate as a sediment in the water
54. A 35-year-old patient visited a hospital seeking medical treatment for his eye. During examination, he reported that he is a carpenter and also works as a freshman at a fast flowing large river which is three kilometers away from his home in his spare time. What is the most likely disease he is at high risk of acquiring?
- A) filariasis B) leishmaniasis C) onchocerciasis D) trypanosomiasis
55. A farmer working at the edge of a river experienced repeated insect bite by insects coming from the river. Two weeks later he became febrile and started developing papules which then changed to ulcer. What is the most likely vector transmitting this disease?
- A) Black fly B) sand fly C) flesh fly D) fruit fly
56. A health officer assessed the exposure of people to the special radiation in a city. He found out that people working at supermarket and shop checkouts use special red or green light emitting devices for detecting barcodes. Similarly, night clubs & theaters use special colored lights at the stages. What is the most likely radiation type they are exposed to?
- A) Laser radiation C) Microwave radiation
B) Infrared radiation D) Radiofrequency radiation
57. A health officer, who was assigned to assess the cause of death of children in a community, found that an intermediate determinant of child mortality was the main factor which leads to death in the community. What is the most likely cause of death in the community?
- A) Child care practices C) Lower respiratory tract infections
B) Source of water supply D) Occupation, literacy and social position

58. A researcher used valid, specific, sensitive & reliable indicator to measure the reproductive health related problems in a community. Which indicator selection criteria best describes this case?
- A) Usefulness
B) Understandable
C) Representativeness
D) Scientifically robustness
59. A 17 yr old primigravid lady presented for antenatal check up to a nearby HC. During examination, the midwife saw that all parts of her external genitalia was removed traditionally. Which is the most likely anticipated complication?
- A) Infection
B) Preterm birth
C) Prolonged labor
D) Low birth weight
60. The national health survey institute conducted screening for asbestosis disease among asbestos factory workers. The institute assess the exposure asbestos dust and prevalence disease in comparison with a control group. What is the most appropriate type of screening for this case?
- A) Mass screening
B) Targeted screening
C) Multiphase screening
D) Opportunistic screening
61. A team of epidemiologists at a specialized hospital screened 1000 pts known to have a chronic disease and 1000 controls without the disease with a new test. The sensitivity and specificity of the test, classified as high, calculated by the team was 94.6% and 98.6%. what is the most appropriate interpretation of the above result"
- A) The test misclassifies most people who do have diseases as free
B) The test will correctly identify most people who are disease free
C) The test misclassifies most people who do not have disease as positive
D) The test will correctly identifies people with the diseases but not without
62. A public health surveillance department has included twenty diseases under the routine surveillance program. From this disease list, poliomyelitis, dracunculiasis and NNT are placed in one category. What is the most likely criterion to select these diseases for national surveillance?
- A) They are international concerns
B) They are targeted for eradication
C) They have high pandemic potential
D) They have significant public health importance
63. An outbreak investigation team received a report of suspected malaria cases from a local health extension worker. The report stated that there were 15 new cases in the area within three days alone. What is the most important information required to decide as an outbreak?
- A) Bed net utilization
B) Status of the cases
C) Size of the population
D) Source of the infection
64. An organization develops an action plan to tackle obesity through community based intervention program in a district. In the proposed action plan, nutritional education and school physical activity were given much emphasis during the planning. What is the prevention interventions planned?
- A) Primary prevention
B) Secondary prevention
C) Tertiary prevention
D) Primordial prevention
65. Various scholars from different disciplines were invited to educate the youth of a community to increase their awareness on HIV/AIDS from social, psychological and health prospective. What teaching method is best to be applied in this case?
- A) Panel discussion
B) Forum
C) Meeting
D)
66. A boarding school spends huge amount of money to manage the school's waste which is mainly composed of food leftovers., dung from animal breeding and feces from the student's toilet. A team was assigned to find a cost/resource recovery waste management strategy. Which strategy is the most appropriate?
- A) Recovering energy from incinerating of the waste

- B) Recovering valuable items for reuse from the waste
 - C) Recovering biogas from the waste using anaerobic digestion
 - D) Recovering domestic fuel by recovering the dung from the waste
67. Students who are a university's cafeterias are often diagnosed with typhoid fever despite proper cooking and storage. Hence, the university's administration seeks an advice to control this disease. What is the most appropriate intervention for the case?
- A) Testing of food handlers
 - B) Checking the quality of the water
 - C) identifying the source of the raw food
 - D) Identifying the presence of rodents in the kitchen
68. One of the best malaria control strategy which was proved to reduce the prevalence of the disease significantly was indoor residual spray with DDT. However, despite the advantages it is completely banned in most countries since 1972. What is the most likely explanation for hanging?
- A) The toxicity level of the DDT to human is very high
 - B) DDT is expensive to implement at large scale
 - C) DDT is persistent in the environment
 - D) Mosquitoes became resistant to DDT
69. A medical center in a big city is under re-innovation to upgrade its service. For this reason, there is movement of various vehicles which suspends the dust in the city. Consequently the health workers, patients and visitors experiences respiratory problem to the silica in the dust. What is the costly effective preventive measure?
- A) Using breathing masks
 - B) Spraying water regularly
 - C) Shading the construction site
 - D) Moving to other site till the construction is done
70. A mother of three children came to a health facility to get family planning service. She reported that when her husband gets drunk, he tries to force her sexually without her will because he needs more children, but she has no desire. She always stressed around him yet she want to stay in the marriage. What is the most appropriate advice to this mother?
- A) Community health prevention
 - B) Discusses with patent of the husband
 - C) Promote her to use contraceptive
 - D) Link her to police station and lawyer
71. A 15 yr old female was kidnapped by a group of individuals while she was collecting firewood in rural area. After six months, a messenger came to the girl's parents and request them an engagement & marriage. Her parents accepted the messenger. The worda reproductive health program officer plans an intervention to decrease such practice in the community. What is the most appropriate intervention to be done by the officer?
- A) Empowering women and raising their status
 - B) Promote access to emergency contraception
 - C) Reducing poverty and alcohol consumption in the community
 - D) Establish protocols that indicate appropriate care and referral for victims of violence
72. Based on a recently report epidemic of watery diarrhea disease from the regional health bureau, a health officer planned to work with the community and take an appropriate community based intervention on the existing health problem. What is the most appropriate next step in the intervention?
- A) Community empowerment
 - B) Community development
 - C) Community organization
 - D) Community mobilization
73. A worda health facility prepared a sample poster on female genital mutilation and planned to distribute it to the community. What is the most appropriate next step to be done by the facility?
- A) Community mobilization
 - B) Audience segmentation
 - C) Need assessment
 - D) Pre-testing

73. A health professional is assigned in a nutrition intervention project, which is responsible to address nutritional deficiencies using nutrition specific interventions. What will be the area of focus of this intervention?
- A) Inadequate food intake
 - B) Inadequate access to food
 - C) Inadequate access to health service
 - D) Inadequate care of mothers and children
74. A 20 month old female child was presented with watery diarrhea of two days duration. On anthropometric assessment, her weight, height and MUAC was found to be 7.6 kg, 75cm and 10.7cm respectively. She has no edema and passed the appetite test. Which supplementation is contraindicated for the child?
- A) Iron
 - B) Zinc
 - C) Vitamin A
 - D) De-worming medication