



Prophylaxis against infective endocarditis (IE) is not reasonable for which of the following patients before dental procedures that involve manipulation of gingival tissue, manipulation of the periapical region of teeth, or perforation of the oral mucosa?

Select one:

- ☐ a. Patients with Rheumatic Severe Mitral Stenosis
- ☐ b. Patients with previous IE
- ☐ c. Patients with prosthetic cardiac valves
- ☐ d. Patients with unrepaired cyanotic congenital heart disease

A 75-year-old woman with diabetes and hypertension awakens with left-side body weakness and complains of double vision. On examination, she has a dense left hemiparesis. The right eye cannot move medially, she has ptosis, and the right pupil is dilated. Where is the most likely site of the lesion?

Select one:

- ☐ a. Left midbrain
- ☐ b. Left pons
- ☐ c. Right midbrain
- ☐ d. Right pons

Question **1**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **2**

Not yet answered

Marked out of 1.00

🚩 Flag question

Yuzpe method is a type of:

Select one:

- ☐ a. Post-coital IUCD
- ☐ b. Minilap sterilization
- ☒ c. Post-coital hormonal contraception
- ☐ d. Male contraceptive method

[Clear my choice](#)

Question **3**

Not yet answered

Marked out of 1.00

🚩 Flag question

An 80- year- old man presents with a basal ganglia hemorrhage. The most likely cause in a patient this age with a hemorrhage in this location is which of the following?

Select one:

- ☐ a. Hemorrhagic cerebral metastasis
- ☐ b. Arteriovenous malformation
- ☐ c. Hypertension
- ☐ d. Cerebral amyloid angiopathy

Question **4**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which of the enumerated hormones are the least active in the catabolic phase?

Select one:

- ☐ a. Estrogens
- ☐ b. Glucagon
- ☐ c. Insuline
- ☐ d. Aldosterone

Question **5**

Not yet answered

Marked out of 1.00

🚩

The following changes are expected in a patient with severe vomiting due to gastric outlet obstruction, except

Select one:

- ☐ a. Acidosis
- ☐ b. Hypochloremia
- ☐ c. Hyponatremia
- ☐ d. Hypokalemia

Question **6**

Not yet answered

Marked out of 1.00

🚩 Flag question

The maintenance fluid requirement for a patient weighing 24 Kg is

Select one:

- ☐ a. 1000ml/24hrs.
- ☐ b. 4000ml/24 hrs.
- ☐ c. 2000 ml/24hrs.
- ☒ d. 1580 ml/24hrs.

[Clear my choice](#)

Question **7**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question 8

Not yet answered

Marked out of 1.00

🚩 Flag question

A 49-year-old male comes to medical OPD for an evaluation as part of his medical check-up. He reports feeling well. He denies previous history of diabetes or coronary artery disease. Physical examination is unremarkable except for his BP, which was measured at 165/95 mmHg. His BP re-checked after 5 min was 160/95 mmHg. What would your recommendation be to this patient?

Select one:

- ☐ a. Initiate drug therapy
- ☐ b. Come back for a follow up after 6 months
- ☐ c. Reassurance
- ☐ d. Come back for a follow up in 3 months

Question 9

Not yet answered

Marked out of 1.00

🚩 Flag question

A 20 year old patient presented with perianal swelling and pain of two days duration. He had a 2x3 cm tender mass in the perianal region. He was diagnosed with a perianal abscess. He had no comorbidities. What is the best treatment strategy?

Select one:

- ☐ a. I + ACY- D under regional anesthesia
- ☐ b. giving PO antibiotics only
- ☐ c. giving IV antibiotics only
- ☐ d. I + ACY- D under local anesthesia

Which of the following therapies are indicated in asymptomatic Hypertrophic cardiomyopathy (HCM) patients with no outflow obstruction?

Select one:

- ☐ a. No need of treatment in Asymptomatic patients
- ☐ b. Amiodarone to reduce risk of sudden death
- ☐ c. Beta blocker to reduce atrial fibrillation risk
- ☐ d. Calcium channel blocker as positive Inotropic agents to improve cardiac performance

Question **11**

Not yet answered

Marked out of 1.00

🚩 Flag question

The minimum caloric requirement for an adult male in bed without fever or unusual losses is

Select one:

- ☐ a. 15 - 20 cal/kgbody Wt.
- ☐ b. 50 - 60 cal/kg body Wt.
- ☐ c. 35 - 50 cal/kg body Wt.
- ☐ d. 25 - 30 cal/kg body Wt.

Question **10**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **12**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which statement is correct in regard to choosing ART for PMTCT?

Select one:

- ☐ a. The best prevention of HIV transmission is achieved with drug combinations that lower the viral load during pregnancy.
- ☐ b. Whenever possible ART should be given only during the last week before delivery to avoid abnormalities in the developing fetus.
- ☐ c. No more than two drugs should be used because drug toxicity is too great when three or more drugs are used for pregnant women.
- ☐ d. The ARV drugs that are used should be those that can be administered by injections, as they result in the highest drug levels in the blood.

Question **13**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 64-year-old man with coronary artery disease and atrial fibrillation is referred for evaluation of fevers, arthralgias, pleuritis, and malar rash. The symptoms have developed over the past 6 months. The pleuritis has responded to steroid therapy, but prednisone has been unable to be tapered off due to recurrence of symptoms at daily steroid doses +ADw-15 mg of prednisone. His medications include aspirin, procainamide, lovastatin, prednisone, and carvedilol. You suspect drug-induced lupus due to procainamide. Antibodies directed against which of the following proteins is most likely to be positive?

Select one:

- ☐ a. Cardiolipin
- ☐ b. Histone
- ☐ c. Double-strand DNA
- ☐ d. Ribonucleoprotein (RNP)

Causes of lactic acidosis include all except

Select one:

- ☐ a. Infections
- ☐ b. Hypoxia
- ☐ c. Circulatory failure
- ☐ d. Acute Myocardial infarction

Question **14**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 40-year-old woman is brought to the emergency OPD after a generalized tonic-clonic seizure lasting for longer than 5 minutes without regaining her consciousness. Which of the following is the best initial treatment for this patient?

Select one:

- ☐ a. Phenobarbital 20 mg/kg IV
- ☐ b. Diazepam 10 mg IV
- ☐ c. Valproate 25 mg/kg IV
- ☐ d. Phenytoin 20 mg/kg IV

Question **15**

Not yet answered

Marked out of 1.00

🚩 Flag question

A single most practical method of assessing adequacy of tissue perfusion in a surgical patient is

Select one:

- ☐ a. Pulse rate
- ☐ b. Blood pressure
- ☐ c. daily weight
- ☐ d. Urine out put

Question **16**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which of the following is true regarding spinal epidural abscess?

Select one:

- ☐ a. Fever, back pain and local spine tenderness are usually first manifestations
- ☐ b. Cerebrospinal fluid examination is helpful to identify the specific etiology
- ☐ c. The treatment includes broad spectrum antibiotic therapy for 2 weeks
- ☐ d. Spine CT scan is the investigation of choice

An old man was evaluated a month after he has developed bilateral Lower limb weakness accompanied by difficulty controlling his bowel and bladder. He also claims that he feels less sensation below his mid-chest. On examination, he was found to have Normotonic and Normoreflexic Paraparesis but with bilateral upgoing plantar responses. There was a Sensory level at about Xyphisternum. On investigation, he had Prostate enlargement with high PSA (400ng/ml). If the patient is able to afford only for a single Imaging, which of the following levels do you prefer?

Select one:

- ☐ a. Lumbosacral Spine MRI
- ☐ b. Brain MRI
- ☐ c. Cervical Spine MRI
- ☐ d. Thoracic Spine MRI

Question **17**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **18**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **19**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which of the following pubertal event would occur even in the absence of ovarian estrogen production

Select one:

- ☐ a. Pubarche
- ☐ b. Vaginal cornification
- ☐ c. Menarche
- ☐ d. Thelarche

Question **20**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which of the following is correct regarding the first dose and route of aspirin use in suspected Acute Coronary Syndrome?

Select one:

- ☐ a. 162 or 325 mg enteric coated
- ☐ b. 162 or 325 mg non-enteric coated chewed
- ☐ c. More than 325
- ☐ d. 81 mg orally

Question **21**

Not yet answered

Marked out of 1.00

🚩 Flag question

A student is nervous for a big exam and is breathing rapidly, what do you expect out of the followings?

Select one:

- ☐ a. Metabolic Acidosis
- ☐ b. Respiratory Acidosis
- ☐ c. Metabolic Alkalosis
- ☒ d. Respiratory Alkalosis

[Clear my choice](#)

Question **22**

Not yet answered

Marked out of 1.00

🚩 Flag question

Mister +IBg-Z+IBk- is a 60 year old merchant from Hawassa who presented with the complaint of cough of 02 weeks productive of mucoid sputum with no other significant complaint but had habit of cigarette smoking 10 pack-years. On physical examination: V/S+AD0-stable+ADs- RS+AD0-right posterior lower 2/3rd absent air entry with stony dullness on percussion. For the above complaint and findings he was investigated with CBC+AD0-normal, OFTs+AD0-normal, CXR+AD0-right hemi-thorax opacity, pleural fluid analysis shows +AD0-LDH+AD4-2/3 upper serum limit. What is the likely diagnosis and next investigation modality?

Select one:

- ☐ a. Lung ca and pleural fluid cytology
- ☐ b. Heart failure and echocardiography
- ☐ c. Lung ca and pleural biopsy
- ☐ d. CKD and renal function test

Question **23**

Not yet answered

Marked out of 1.00

🚩 Flag question

All of the following statements about abruption placenta are false

Select one:

- ☐ a. patients often present with painful vaginal bleeding
- ☐ b. ultrasonography can effectively establish the diagnosis
- ☐ c. fetal distress is a common occurrence
- ☐ d. maternal hypertension is the most often associated etiologic factor

Question **24**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which one of the following is not true about pre-transfusion testing and transfusion?

Select one:

- ☐ a. Testing for RBC blood group antigens
- ☐ b. Cross matching between the donor+IBk-s sera and the recipient+IBk-s RBCs
- ☐ c. Group O may be given to A, AB, B, or O patients.
- ☐ d. Serologic compatibility for the recipient+IBk-s and donor+IBk-s groups

A 23-year-old woman was diagnosed with systemic lupus erythematosus based upon the presence of polyarthritis, malar rash with photosensitivity, and oral ulcerations. Antibodies to double-stranded DNA, Smith protein, and antinuclear antibodies were present in high titers. A urinalysis is normal. The patient is requesting treatment for the joint symptoms as she feels they limit her activities of daily living. What is the best choice for initial therapy in this individual?

Select one:

- ☐ a. Methotrexate, 15 mg weekly
- ☐ b. Physical therapy only
- ☐ c. Hydroxychloroquine, 200+IBM-400 mg daily
- ☐ d. Prednisone, 1 mg/kg daily

A 23-year-old woman was diagnosed with systemic lupus erythematosus based upon the presence of polyarthritis, malar rash with photosensitivity, and oral ulcerations. Antibodies to double-stranded DNA, Smith protein, and antinuclear antibodies were present in high titers. A urinalysis is normal. The patient is requesting treatment for the joint symptoms as she feels they limit her activities of daily living. What is the best choice for initial therapy in this individual?

Select one:

- ☐ a. Methotrexate, 15 mg weekly
- ☐ b. Physical therapy only
- ☐ c. Hydroxychloroquine, 200+IBM-400 mg daily
- ☐ d. Prednisone, 1 mg/kg daily

Question **25**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **25**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 23-year-old woman was diagnosed with systemic lupus erythematosus based upon the presence of polyarthritides, malar rash with photosensitivity, and oral ulcerations. Antibodies to double-stranded DNA, Smith protein, and antinuclear antibodies were present in high titers. A urinalysis is normal. The patient is requesting treatment for the joint symptoms as she feels they limit her activities of daily living. What is the best choice for initial therapy in this individual?

Select one:

- ☐ a. Methotrexate, 15 mg weekly
- ☐ b. Physical therapy only
- ☐ c. Hydroxychloroquine, 200+IBM-400 mg daily
- ☐ d. Prednisone, 1 mg/kg daily

Which one of the following is not true about the umbilical and paraumbilical hernias?

Select one:

- ☐ a. Umbilical hernia is common in children.
- ☐ b. Congenital hernias usually close spontaneously.
- ☐ c. Umbilical hernias should be repaired before 5 years of age.
- ☐ d. Surgical repair is necessary in adults.

Question **25**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **26**

Not yet answered

Marked out of 1.00

🚩 Flag question

After severe trauma patients are invariably losing weight which of the enumerated factors is mainly responsible for this weight loss?

Select one:

- ☐ a. Utilisation of glycogen
- ☐ b. Suppression of insulin production
- ☐ c. Utilisation of Body proteins
- ☐ d. Loss of intracellular water

Question **27**

Not yet answered

Marked out of 1.00

🚩 Flag question

A year-old man without a history of known medical problems presents with a 2-day history of a progressively worsening headache associated with fever and neck stiffness. His neurologic examination is notable for a Kernig sign. LP reveals a protein of 55 mg/dL, glucose of 50 mg/dL (serum glucose of 100 mg/dL), 250 white blood cells/+A7w-L with 60+ACU- lymphocytes, and cultures/Gram stain pending. These CSF studies are most consistent with which of the following?

Select one:

- ☐ a. Subarachnoid hemorrhage
- ☐ b. Bacterial meningitis
- ☐ c. Cannot tell without knowing the result of the cultures and Gram stain
- ☐ d. Viral meningitis

Question **28**

Not yet answered

Marked out of 1.00

🚩 Flag question

The most important step in the treatment of both septic shock and hemorrhagic shock is

Select one:

- ☐ a. intravenous fluids
- ☐ b. Foley catheter
- ☐ c. antibiotics
- ☐ d. Swan-Ganz catheter

Question **29**

Not yet answered

Marked out of 1.00

🚩 Flag question

The most common cause of surgical wound infection is

Select one:

- ☐ a. wound dressings removed too early
- ☐ b. contamination during the operation
- ☐ c. trauma due to over retraction
- ☐ d. inadequate skin preparation

Question **30**

Not yet answered

Marked out of 1.00

🚩 Flag question

Which of the following is not true of +ACI-banked+ACI-blood ?

Select one:

- ☐ a. Plasma lactate falls.
- ☐ b. Plasma glucose is lower.
- ☐ c. It has a lower pH.
- ☐ d. The red cell sodium is high.

Question **31**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **32**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 32-year-old woman was admitted to labor ward for induction of labor. Her gestational age from last menstrual period was 42 weeks. What is the most likely cause of her prolonged pregnancy?

Select one:

- ☐ a. Wrong date
- ☐ b. Lack of ANC
- ☐ c. Previous history of similar condition
- ☐ d. Female fetus

Question **33**

Not yet answered

Marked out of 1.00

🚩 Flag question

The common denominator in the various forms of shock is

Select one:

- ☐ a. circulating blood volume less than the size of the intravascular compartment.
- ☐ b. reduction of plasma volume.
- ☐ c. high hematocrit values.
- ☐ d. Blood loss

Question **34**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 78-year-old right hand male with diabetes and hypertension presents with sudden onset of right hemiparesis affecting mainly the face and arm. He seems to be able to understand and follow commands, however, cannot verbalize, and appears frustrated when asked questions because he cannot answer. Where is the location of the vascular occlusion?

Select one:

- ☐ a. Left posterior cerebral artery
- ☐ b. Right anterior cerebral artery
- ☐ c. Left anterior cerebral artery
- ☐ d. Left middle cerebral artery

Question **35**

Not yet answered

Marked out of 1.00

🚩 Flag question

The most likely cause of abnormal uterine bleeding in 13 years old girl is:

Select one:

- ☐ a. Anovulation
- ☐ b. Systemic bleeding diatheses.
- ☐ c. Ectopic pregnancy.
- ☐ d. Trauma

Question **36**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 45 year-old house wife female patient from Tula came to medical EOPD with the complaint of chest tightness, dyspnea, wheezing, unable to complete a sentence of 3 hours duration. She is known asthmatic patient with poor adherence and poor control. On physical examination she had O₂ saturation of 70+ACU- with room air, tachycardic, tachypenic, agitated. CBC+AD0-normal, CXR+AD0-not remarkable. All of the following can be done as part of management approach of acute severe asthma except?

Select one:

- ☐ a. High dose of SABA
- ☐ b. Magnesium Sulfate
- ☐ c. Antibiotics
- ☐ d. Supplementation of high concentration oxygen

Question **37**

Not yet answered

Marked out of 1.00

🚩 Flag question

A patient presents with 3 weeks of pain in the lower back. All the following are risk factors for serious causes of spine pathology except

Select one:

- ☐ a. duration of pain more than 2 weeks
- ☐ b. age more than 50 years
- ☐ c. urinary incontinence
- ☐ d. bed rest without relief

Question **38**

Not yet answered

Marked out of 1.00

🚩 Flag question

A 40 years old male patient came to the clinic for an evaluation of acute onset severe Left retro-orbital pain which was so excruciating that he moves around during the attack. He was having repetitive attacks during each consecutive evening, each attack staying only for a few minutes. During each attack, he has also noticed Left eye ptosis +ACY- lacrimation, Left nasal congestion +ACY- rhinorrhoea, and occasional Left sided photophobia. He had similar attacks a year back which after about month duration of sickness. Which primary headache do you think is this patient suffering from?

Select one:

- ☐ a. Tension Type Headache
- ☐ b. Migraine Headache
- ☐ c. Cluster Headache
- ☐ d. Primary Cough Headache

Question **39**

Not yet answered

Marked out of 1.00

🚩 Flag question

Hypokalemia is likely to occur in all EXCEPT

Select one:

- ☐ a. Conn+IBk-s syndrome / Hyperaldosterinism
- ☐ b. Alkalosis
- ☐ c. Paralytic ileus
- ☐ d. Acidosis

Mister +IBg-X+IBk- is 50 year old prisoner presented with the one month history of cough productive of whitish sputum, progressive dyspnea +ACY- right sided pleuritic chest pain. Had history of passive smoking from his cell-mates otherwise no history of active smoking. On examination , V/S+AD0-tachypenic , tachycardic with SPO2 of 85+ACU- on room air +ACY- 95+ACU- with 5Ls of intranasal oxygen +ADs- decreased air entry dullness on percussion over the right lower 2/3rd of posterior lung field. He underwent thorough investigation and result turns out to be small cell lung ca and the treating physician planned therapeutic thoracocentesis and pleurodesis, but after a gush of fluid measuring about 2 Ls was drained his situation got sudden deterioration with frothy sputum production. What is the likely cause of the sudden derangement?

Select one:

- ☐ a. Tension pneumothorax
- ☐ b. None
- ☐ c. SCLC
- ☐ d. Air embolism
- ☐ e. Re-expansion pulmonary edema

Question **40**

Not yet answered

Marked out of 1.00

 Flag question

Question **41**

Not yet answered

Marked out of 1.00

Flag question

A 24-year-old woman enters labor at 42 weeks in her first pregnancy. She is found to be 3 cm dilated and completely effaced with the vertex at zero station. Contractions are occurring every 5 minutes, lasting 45-60 seconds. The fetal heart rate is 120 beats/minute. What is your plan of management?

Select one:

- ☐ a. Rupture the amniotic sac and place an internal scalp electrode to confirm fetal heart rate. If the rate of 120 minute is confirmed, plan an immediate scalp pH to rule out fetal distress.
- ☐ b. Send her home to wait the active phase of labor. She should return when the contractions are every 3 minutes or when the bag of water breaks.
- ☐ c. Administer oxytocin to augment contractions. The frequency and duration are insufficient to expect a normally progressive labor
- ☐ d. Admit her to the hospital, and observe her labor which is normally progressive at this point.

Question **42**

Not yet answered

Marked out of 1.00

Flag question

A 30-year-old female patient with known Rheumatic Severe Mitral stenosis and Mild Mitral Regurgitation presented to OPD with palpitation and easy fatigability with resting ECG suggesting Atrial fibrillation with heart rate of 120/min. What is your recommendation regarding anticoagulation in this patient?

Select one:

- ☐ a. No need of anticoagulation since CHA2DS2+IBM-VASc score is less than 2
- ☐ b. Control the rate and rhythm with Metoprolol and then no need for anticoagulation if the rate and rhythm is controlled
- ☐ c. Start Rivaroxaban since it reduces stroke risks significantly
- ☐ d. Start Warfarin, since it is first line anticoagulant in such a patient

Question **43**

Not yet answered

Marked out of 1.00

Flag question

What is the classical digital rectal examination finding of an anal fissure

Select one:

- ☐ a. A tear in the anoderm at 6 O clock
- ☐ b. An opening at the perianal area
- ☐ c. A tear in the anoderm at 9 O clock
- ☐ d. A tear at the dentate line

Question 44

Not yet answered

Marked out of 1.00

Flag question

A 64-year-old man with congestive heart failure presents to the emergency room complaining of acute onset of severe pain in his right foot. The pain began during the night and awoke him from a deep sleep. He reports the pain to be so severe that he could not wear a shoe or sock to the hospital. His current medications are furosemide, 40 mg twice daily, carvedilol, 6.25 mg twice daily, candesartan, 8 mg once daily, and aspirin, 325 mg once daily. On examination, he is febrile to 38.5+ALA-C. The first toe of the right foot is erythematous and exquisitely tender to touch. There is significant swelling and effusion of the first metatarsophalangeal joint on the right. No other joints are affected. Which of the following findings would be expected on arthrocentesis?

Select one:

- ☐ a. Presence of weakly positively birefringent rhomboidal crystals under polarized light microscopy
- ☐ b. Glucose level of +ADw-25 mg/dL
- ☐ c. Presence of strongly negatively birefringent needles haped crystals under polarized light microscopy
- ☐ d. Positive Gram stain

Question 45

Not yet answered

Marked out of 1.00

Flag question

Thirty minutes after an uneventful cholecystectomy, a 30-year-old woman is noted to be restless and cyanotic. Blood pressure is 120/70, pulse is 110 and respiration is 12/minute and shallow. The most appropriate procedure at this point is to

Select one:

- ☐ a. Start antibiotic therapy
- ☐ b. Sedate
- ☐ c. Start heparin therapy
- ☐ d. Ventilate obtain a lung scan

Which is not the function of the thyroid hormones?

Select one:

- ☐ a. Thyroid hormones decrease bone and protein turnover and the speed of muscle contraction and relaxation.
- ☐ b. They also increase gastrointestinal motility, leading to diarrhea in hyperthyroidism and constipation in hypothyroidism.
- ☐ c. Thyroid hormones are responsible for maintaining the normal hypoxic and hypercapnic drive in the respiratory center of the brain.
- ☐ d. They are important for fetal brain development and skeletal maturation.

Question **46**

Not yet answered

Marked out of 1.00

Flag question

For the patient mentioned on number 5, how would you like to approach the raised blood pressure?

Select one:

- ☐ a. Calcium channel blockers
- ☐ b. Beta blocker
- ☐ c. No need of anti-hypertensive
- ☐ d. Steroid initiation

Question **47**

Not yet answered

Marked out of 1.00

Flag question

Which site of urolithiasis is associated with lower urinary tract symptoms?

Select one:

- ☐ a. Caliceal stone
- ☐ b. Pelviureteric junction stone
- ☐ c. Proximal ureteric stone
- ☐ d. Vesicoureteric junction stone

Question **48**

Not yet answered

Marked out of 1.00

Flag question

Question **49**

Not yet answered

Marked out of 1.00

Flag question

For patients with Acute Coronary Syndrome (ACS) treated medically, what is the recommended duration of double antiplatelet therapy (DAPT)?

Select one:

- ☐ a. 1 month
- ☐ b. 1 year
- ☐ c. 6 months
- ☐ d. Forever, unless at high risk of bleeding

Question **50**

Not yet answered

Marked out of 1.00

Flag question

A 40-year-old female is admitted to the hospital. On examination she is found to have pulmonary edema. Her past history is only significant for hypertension. She is on a medical regimen consisting of a Hydrochlorothiazide, Amlodipine, Enalapril, and Carvedilol. Her family also gave a history of recent uncontrolled high BP and one other episode of pulmonary edema 2 months ago. Her BP was 220/110 mmHg. The rest of her examination is unremarkable. What is the initial diagnosis?

Select one:

- ☐ a. Reno-vascular hypertension
- ☐ b. Primary hyperaldosteronism
- ☐ c. Pseudo hypertension
- ☐ d. Coarctation of aorta

Choose the one which is not first line TB drug?

Select one:

- ☐ a. Capreomycin
- ☐ b. Rifapentine
- ☐ c. Ethambutol
- ☐ d. Isoniazid

Question **51**

Not yet answered

Marked out of 1.00

 Flag question

A patient with end-stage renal disease (ESRD) on hemodialysis has moderate pericardial effusion and referred for cardiac evaluation. What is your recommendation?

Select one:

- ☒ a. Intensify Hemodialysis
- ☐ b. Start NSAID with Prednisolone short course
- ☐ c. Refer to cardiologist to Perform pericardiocentesis.
- ☐ d. Switch to peritoneal dialysis

[Clear my choice](#)

Question **52**

Not yet answered

Marked out of 1.00

 Flag question

Which of the following perianal condition is painful?

Select one:

- ☐ a. Anal condyloma
- ☐ b. Internal hemorrhoids
- ☐ c. Anal fissure
- ☐ d. Perianal fistula

Question **53**

Not yet answered

Marked out of 1.00

 Flag question

Question **54**

Not yet answered

Marked out of 1.00

Flag question

A 45-year-old female patient is evaluated for progressive dyspnea and dry cough of 3 years duration with history of TB treatment although there was no improvement from the symptoms. On investigation: HRCT shows sub pleural, ground glass reticular opacity and lung biopsy result is consistent with cellular NSIP. No identified connective tissue disease or drug exposure was identified. What is the recommended initial management approach?

Select one:

- ☐ a. Steroids
- ☐ b. 2nd line anti-TB
- ☐ c. Cyclophosphamide
- ☐ d. MMF

Question **55**

Not yet answered

Marked out of 1.00

Flag question

Main mechanism of action of progesterone only contraceptives is/are

Select one:

- ☐ a. Inhibition of tubal motility
- ☐ b. Inhibiting ovulation through higher centers
- ☐ c. Thickening cervical mucus
- ☐ d. Alteration of endometrial tissue

One of the following statement is not true regarding obstructive Jaundice.

Select one:

- ☐ a. Presence of pain is suggestive of stone obstruction rather than tumour.
- ☐ b. Patient with obstructive jaundice secondary to carcinoma are older compared to those due to stone.
- ☐ c. Gall bladder is never palpable in stone disease.
- ☐ d. Abdominal ultrasound is the most useful means of investigation for obstructive jaundice

Question **56**

Not yet answered

Marked out of 1.00

Flag question

A pregnant woman with fibroid uterus develops acute pain in abdomen with low grade fever and leukocytosis at 26 weeks. What is the most likely diagnosis?

Select one:

- ☐ a. Degeneration of fibroid
- ☐ b. Preterm labor
- ☐ c. Torsion of fibroid
- ☐ d. Infection of fibroid

Question **57**

Not yet answered

Marked out of 1.00

Flag question

Choose best syndromic management for an individual comes with urethral discharge.

Select one:

- ☐ a. Ceftriaxone 250 mg IM stat plus azithromycin 1 gm po stat
- ☐ b. B. penicillin 2.4 million units IM stat
- ☐ c. Acyclovir 400 mg po TID for 07 days plus azithromycin 1gm po stat
- ☐ d. Ceftriaxone 250 mg IM stat

Question **58**

Not yet answered

Marked out of 1.00

Flag question

Severe chills, fever, and chest pain in a patient after transfusion of 75cc of blood usually indicates

Select one:

- ☐ a. pyrexial reaction
- ☐ b. hemolytic transfusion reaction
- ☐ c. citrate toxicity
- ☐ d. blood contamination

Question **59**

Not yet answered

Marked out of 1.00

Flag question

35 year-old woman is now arriving in the emergency room after an automobile accident that caused injury to several of her organ system .To improve her respiratory status. The woman is placed on a respirator with the addition of positive end-expiratory pressure /PEEP/. She suddenly develops restlessness, hypotension and more profound hypoxemia. The most likely diagnosis is

Select one:

- ☐ a. decreaded venous return
- ☐ b. myocardial infacrction
- ☐ c. plumunary embolism
- ☐ d. tension pneumothorax

Question **60**

Not yet answered

Marked out of 1.00

Flag question

A victim of road traffic accident is brought to the emergency room unconscious and with a blood pressure of 80/60mmHg. The first step in management is

Select one:

- ☐ a. skull X-ray
- ☐ b. IV Ringer's lactate
- ☐ c. neurologic evaluation
- ☐ d. establish adequate airway

Question **61**

Not yet answered

Marked out of 1.00

Flag question

Which of the following is Not an indication for corticosteroids in pericarditis?

Select one:

- ☐ a. Recurrence despite colchicine
- ☐ b. Unresponsive to NSAID and colchicine
- ☐ c. Autoimmune etiology
- ☐ d. Acute Pericarditis in combination with NSAID
- ☐ e. None of the above

Question **62**

Not yet answered

Marked out of 1.00

Flag question

In normal wound healing, the predominant cell in the wound by the end of the first week is the

Select one:

- ☐ a. lymphocyte
- ☐ b. macrophages
- ☐ c. neutrophil
- ☐ d. fibroblast

Question **63**

Not yet answered

Marked out of 1.00

Flag question

The PMTCT clinic team at your hospital decides to use an out-patient testing strategy for pregnant women at the antenatal clinic. Which of the following accurately describes what will happen under this strategy?

Select one:

- ☐ a. All pregnant women will be tested for HIV whether or not they want to be tested.
- ☐ b. All pregnant women in the clinic will have to ask for an HIV test in order to be tested.
- ☐ c. All women will be tested a second time if their first test result is positive.
- ☐ d. All pregnant women are tested for HIV unless they refuse to be tested.

Question **64**

Not yet answered

Marked out of 1.00

Flag question

Which one is not true about epigastric hernias?

Select one:

- ☐ a. They are usually protrusions of properitoneal fat.
- ☐ b. They appear through a defect in the linea alba.
- ☐ c. They have peritoneal sacs.
- ☐ d. There may be multiple hernias.

Question **65**

Not yet answered

Marked out of 1.00

 Flag question

Which one is not true about epigastric hernias?

Select one:

- ☐ a. They are usually protrusions of properitoneal fat.
- ☐ b. They appear through a defect in the linea alba.
- ☐ c. They have peritoneal sacs.
- ☐ d. There may be multiple hernias.

Question **65**

Not yet answered

Marked out of 1.00

 Flag question

A 40 year old alcoholic male being treated for tuberculosis, but he has not been compliant with his medications. He complains of increasing weakness and fatigue. He appears to have lost weight, and his BP is 80/50mmHg. There is increased pigmentation over the elbows, cardiac exam is normal. In the advanced stage of this disease, the most likely electrolyte abnormalities will be

Select one:

- ☐ a. Low serum K⁺
- ☐ b. High serum Na⁺
- ☐ c. Low serum Na⁺
- ☐ d. High serum Na⁺ and low serum K⁺

Question **66**

Not yet answered

Marked out of 1.00

 Flag question

A 30 year old male patient who was doing well after chest tube insertion and intubation with ETT +ACV- put on mechanical ventilator for the diagnosis of acute respiratory failure 20 to? Plus massive pleural effusion, but on the 4th day of admission the ventilator begins to alarm for low exhaled tidal volume. Upon examination, increased bubbling was noted in the underwater seal chamber of the chest tube drainage. Which one of the following problems do these findings suggest?

Select one:

- ☐ a. Air trapping
- ☐ b. Bronchopleural fistula
- ☐ c. Asynchrony with ventilator
- ☐ d. Chest tube obstruction

Question **67**

Not yet answered

Marked out of 1.00

Flag question

Which one is false about IRIS?

Select one:

- ☐ a. Frequently seen in the setting of tuberculosis
- ☐ b. Continuation of ART is recommended
- ☐ c. Glucocorticoids are always required to control it
- ☐ d. Most common in patients with low CD4 count

Question **68**

Not yet answered

Marked out of 1.00

Flag question

Gastrointestinal bleeding includes the following manifestations except:

Select one:

- ☐ a. melena
- ☐ b. hematemesis
- ☐ c. Haematochezia
- ☐ d. hemoptysis

Question **69**

Not yet answered

Marked out of 1.00

Flag question

The normal response to surgery includes all except

Select one:

- ☐ a. Fever.
- ☐ b. Renal retention of sodium and calcium.
- ☐ c. Rise in serum sodium.
- ☐ d. Antidiuresis for 12 - 24 hours.

Question **70**

Answer saved

Marked out of 1.00

🚩 Flag question

A 49-year-old woman came with a chief complaint of irregular menses for the last 5 months. Her menses was regular before this condition and no history of recent contraceptive use. Pelvic ultrasonography revealed unremarkable result.

Select one:

- ☐ a. Cervical cancer
- ☐ b. What is the most likely cause of her problem?
- ☐ c. Ovulation dysfunction
- ☐ d. Submucosal myoma
- ☐ e. Uterine polyp

Question **71**

Not yet answered

Marked out of 1.00

🚩 Flag question

Question **72**

Not yet answered

Marked out of 1.00

Flag question

A 30-year-old pregnant woman at gestational age of 38 weeks was admitted to labor and delivery unit. An intern noted non-reassuring fetal rate and vaginal bleeding following gush of fluid per vaginal. What is the most likely cause of this problem?

Select one:

- ☐ a. Placenta previa
- ☐ b. Vasa previa
- ☐ c. Abruptio placenta
- ☐ d. Uterine rupture

Question **73**

Not yet answered

Marked out of 1.00

Flag question

A 45-year-old male patient who was diagnosed to have internal hemorrhoids, presented with perianal swelling and intermittent rectal bleeding of 2 years duration. He has to reduce the swelling manually after defecation. What is the grade of his hemorrhoids?

Select one:

- ☐ a. Grade IV
- ☐ b. Grade I
- ☐ c. Grade III
- ☐ d. Grade II

Question **74**

Not yet answered

Marked out of 1.00

Flag question

Which of the following is red flag sign in the diagnosis of Parkinson's disease?

Select one:

- ☐ a. freezing gait
- ☐ b. recurrent fall in the first year of diagnosis
- ☐ c. Bradykinesia
- ☐ d. resting tremor

Question **75**

Not yet answered

Marked out of 1.00

 Flag question

A 7-year-old girl was brought to Gynecology OPD by her mother for vaginal bleeding. Which of the following is the most common source of abnormal bleeding in this girl?

Select one:

- ☐ a. Vagina
- ☐ b. Uterus
- ☐ c. Urethra
- ☐ d. Ovary

Question **76**

Not yet answered

Marked out of 1.00

 Flag question

Regarding potassium administration, all are true except

Select one:

- ☐ a. It should not be given in the early postoperative hours
- ☐ b. The iv fluid with added potassium should be properly labeled and run slowly
- ☐ c. It can be given through iv injection
- ☐ d. Preferably the patient should be on continuous ECG monitor

A 25 year old sexually active male presented to medical OPD with a complaint of non-pruritic, discrete macules distributed on his trunk and extremities, involve his palms and soles. Two months before the above complaint, he had single painless indurated chancre which healed by itself. Choose wrong statement for the above case

Select one:

- ☐ A. Has a chance to develop meningitis
- ☐ B. The above symptoms can resolve spontaneously.
- ☐ C. He is non-infectious to other individuals at this stage
- ☐ D. He has a chance to develop hair loss

Question **77**

Not yet answered

Marked out of 1.00

Flag question

56 years old woman has come to you with the complaints of hot flushes irritability, joint pains with lack of sleep. Most appropriate treatment would be

Select one:

- ☐ a. estrogen
- ☐ b. Combined estrogen, progesterone preparations
- ☐ c. Selective estrogen receptor modulators (SERMS)
- ☐ d. Vitamins

Question **78**

Not yet answered

Marked out of 1.00

Flag question

Question **79**

Not yet answered

Marked out of 1.00

Flag question

A 60-year-old man with a history of coronary artery disease develops sudden onset of paralysis of all four limbs. On examination his eyes are open, he appears alert, and he can consistently respond to complex questions and commands by blinking his eyes, but otherwise has minimal facial movement and no movement of the limbs. Which of the following best describes this condition?

Select one:

- ☐ a. Coma
- ☐ b. Locked-in syndrome
- ☐ c. Brain death

Question **80**

Not yet answered

Marked out of 1.00

Flag question

Post-operative complications in a nutritionally depleted patient includes all except

Select one:

- ☐ a. Wound dehiscence
- ☐ b. Pulmonary infection
- ☐ c. Prolonged paralytic ileus
- ☐ d. Wound infection

Question **81**

Not yet answered

Marked out of 1.00

Flag question

A 60-year female patient presented with a sudden onset of right upper quadrant pain accompanied by rigor, fever and vomiting. On examination the patient was in agony with blood pressure of 100/60 mm Hg, pulse rate of 96 beats per minute. She had mild icterus of the sclera and right upper quadrant tenderness. The most likely diagnosis of this case would be:-

Select one:

- ☐ a. Perforated peptic ulcer disease
- ☐ b. Acute Cholecystitis
- ☐ c. Acute Pancreatitis
- ☐ d. Ascending cholangitis

Question **82**

Not yet answered

Marked out of 1.00

Flag question

A-25 years old known stage IV RVI patient presented with headache of one week duration, associated with this she also complain neck pain and vomiting of ingested matter. Up on physical exam she was conscious and oriented, has right pronator drift and positive neck rigidity. Which one of the following is the next best step in the management of the above case?

Select one:

- ☐ a. Do brain MRI
- ☐ b. Send Blood culture
- ☐ c. LP and CSF analysis
- ☐ d. Start empiric antifungal

Question **83**

Not yet answered

Marked out of 1.00

Flag question

A 23-year-old woman from rural area presents with shortness of breath and has Mitral stenosis with MVA of 1.1cm² on Echocardiography. What will you do about rheumatic fever prophylaxis?

Select one:

- ☐ a. Prophylaxis for the next 10 years
- ☐ b. No prophylaxis as she is +AD4-21 years old
- ☐ c. Prophylaxis till she turns 40 years or life long
- ☐ d. Prophylaxis for the next 5 years

Question **84**

Not yet answered

Marked out of 1.00

Flag question

One of the following is inappropriate for replacement of isotonic dehydration

Select one:

- ☐ a. 5+ACU- Dextrose in saline
- ☐ b. Ringer's solution
- ☐ c. 5+ACU- Dextrose in water
- ☐ d. normal saline

W /ro +IBg-AB+IBk- is a 40 year old known asthmatic patient for the last 15 years with poor adherence to her controllers arrives at medical EOPD with complaints of dyspnea , loud expiratory wheeze with RR+AD0-40 ,SpO2 of 70+ACU- , BP+AD0- 160/95. Three hours after asthma exacerbation management initiation her wheeze disappeared and chest wall movement was decreased. What is the next possible prior intervention?

Select one:

- ☐ a. Prepare for intubation
- ☐ b. Beta blocker initiation
- ☐ c. Reduce dose of salbutamol
- ☐ d. No need of other intervention

Question **85**

Answer saved

Marked out of 1.00

🚩 Flag question

All the following statements are true Except

Select one:

- ☐ a. wound healing is accelerated in an environment of low tissue oxygen
- ☐ b. a dehiscd wound that is re-sutured gains strength faster than a primary wound.
- ☐ c. synthesis of new collagen is blocked during periods of deficiency of ascorbic acid
- ☐ d. wounds heal faster at 390C(102.2F) than at 370C (98.7F)

Question **86**

Not yet answered

Marked out of 1.00

🚩 Flag question

The maintenance fluid requirement for a patient weighing 24 Kg is

Select one:

- ☒ a. 1580 ml/24hrs.
- ☐ b. 1000ml/24hrs.
- ☐ c. 4000ml/24 hrs.
- ☐ d. 2000 ml/24hrs.

[Clear my choice](#)

Question **87**

Not yet answered

Marked out of 1.00

 Flag question

Which of the following statement is not true about prophylactic antibiotic in surgery?

Select one:

- ☐ a. Prolonged prophylaxis has no advantage over short course(24 hours)
- ☐ b. Antibiotics are best given parenterally before the time of induction of anesthesia
- ☐ c. Second generation cephalosporins are not recommended
- ☐ d. Prophylaxis is indicated for clean wounds where the consequences for infection are grave

Question **88**

Not yet answered

Marked out of 1.00

 Flag question

The first step in treatment of cardiac arrest is

Select one:

- ☐ a. intracardial adrenaline
- ☐ b. defibrillation
- ☐ c. closed cardiac massage
- ☐ d. ventilating the patient

Question **89**

Not yet answered

Marked out of 1.00

Flag question

A 36-year-old female came to referral clinic for follow-up after diagnosed with Peripartal cardiomyopathy during her first pregnancy 1 year back. Currently she is maintained on Furosemide, Metoprolol, Enalapril and Spironolactone. Her LV function by echocardiogram is now 40+ACU-, which has improved from 20+ACU-. She wanted to be pregnant and asked you about her second pregnancy. What do you recommend?

Select one:

- ☐ a. Counsel against pregnancy
- ☐ b. To go ahead with pregnancy
- ☐ c. Recommend discontinuation of Enalapril and then proceed with pregnancy
- ☐ d. Recommend reassessing LV function in 6 months and if normal, then proceed

Question **90**

Not yet answered

Marked out of 1.00

Flag question

Question 91

Not yet answered

Marked out of 1.00

 Flag question

A young man presented with acute onset ascending weakness initially affecting his lower Limbs followed by involvement of his Upper Limbs and then his bilateral lower face in 5 days. He had a 2 days history of watery diarrhoea about a week prior which subsided on its own. On examination, he had flaccid quadriparesis with absent DTRs and Facial diparesis. Nerve Conduction Study (NCS) was done and it suggested an AIDP variant of GBS under appropriate clinical setting. PIHCT was NR. LP was done and sent for CSF analysis. What do you expect from the CSF analysis?

Select one:

- ☐ a. Low Cell count +ACY- Low Protein
- ☐ b. High Cell count +ACY- Low protein
- ☐ c. Low Cell count +ACY- High Protein
- ☐ d. High Cell count +ACY- High Protein

Question 92

Not yet answered

Marked out of 1.00

 Flag question

Which is the first and most important activator of HYPERGLYCEMIA in trauma and surgery?

Select one:

- ☐ a. Cortisone
- ☐ b. Glucagone
- ☐ c. Epinephrine
- ☐ d. Testosterone

Question **93**

Answer saved

Marked out of 1.00

Flag question

A 26 years married woman with IUCD came with a missed thread of the loop. What would be an acceptable modality of management?

Select one:

- ☐ a. remove the IUCD and allow the pregnancy to continue
- ☒ b. perform ultrasound to locate the loop and exclude possibility of pregnancy
- ☐ c. perform a laparoscopy to locate and remove the IUCD
- ☐ d. perform a D+ACY-C to remove the IUCD

[Clear my choice](#)

Question **94**

Not yet answered

Marked out of 1.00

Flag question

A 25-year-old woman experiences episodic spells in which she suddenly stops what she was previously doing and stares for about a minute, sometimes picking at her nose and her shirt. She cannot recall what happens during the spell itself. She says, however, that she knows when a spell is going to happen because she experiences a warm sensation in her epigastric region, followed by a sensation of fear and a rapid recollection of episodes of past life experiences along with palpitations. She may have postictal confusion. What is the most likely diagnosis?

Select one:

- ☐ a. Absence seizures
- ☐ b. Focal impaired awareness seizures
- ☐ c. Psychogenic nonepileptic Seizures
- ☐ d. Generalized Anxiety Disorder

In an otherwise healthy patient, a deficiency of which of the following would have the LEAST effect on wound healing?

Select one:

- ☐ a. Vitamin A
- ☐ b. Iron
- ☐ c. Alpha -ketoglutarate
- ☐ d. Vitamin C

Question **95**

Not yet answered

Marked out of 1.00

Flag question

Important components of collagen synthesis and wound strength and contraction include all the following except.

Select one:

- ☐ a. Fibroblasts
- ☐ b. Normal PO₂
- ☐ c. Myofibroblasts
- ☐ d. Vitamin C

Question **96**

Not yet answered

Marked out of 1.00

Flag question

Which of the following is true regarding urolithiasis?

Select one:

- ☒ a. Stone recurrence is common after treatment.
- ☐ b. It can cause gall stone ileus.
- ☐ c. Life time prevalence of urolithiasis is lower than 1+ACU-.
- ☐ d. Females have higher risk compared to males.

[Clear my choice](#)

Question **97**

Not yet answered

Marked out of 1.00

Flag question

Which one is against the principle of wound care

Select one:

- ☐ a. Routine antibiotic coverage
- ☐ b. Incised wound can be closed by primary suture of the skin
- ☐ c. Delayed closure of infected wounds
- ☐ d. Meticulous debridement

Question **98**

Not yet answered

Marked out of 1.00

Flag question

A 25-year-old nulligravid lady presented to Gynecology OPD with a complaint of heavy menstrual bleeding of 2 years duration. On physical examination: pale conjunctiva and 16 weeks sized abdominopelvic mass that moves with cervix on bimanual exam. What is the most likely cause of her problem?

Select one:

- ☐ a. Fibroid
- ☐ b. Endometrial hyperplasia
- ☐ c. Adenomyosis
- ☐ d. Granulosa cell tumor

Question **99**

Not yet answered

Marked out of 1.00

Flag question

Which of the following is the formula for Minute Volume calculation?

Select one:

- ☐ a. FEV_1/FVC
- ☐ b. $FVC \times RR$
- ☐ c. $CO \times RR$
- ☐ d. Tidal volume $\times RR$

Question **100**

Not yet answered

Marked out of 1.00

Flag question