

1. A 26 year-old female presented with two months of excessive menses and gum bleeding. Physical examination revealed pale conjunctiva, petechiae and ecchymosis. Which of the following is the most likely diagnosis?

- A. Hemophilia ✗
- ☒ B. Idiopathic thrombocytopenic purpura ✓
- C. Scurvy ✗
- D. Vitamin K deficiency ✗

2. A 21 year-old woman presented with fatigue and light headedness. Physical examination showed marked conjunctival and palmar pallor. Which of the following physical findings is most likely to be seen in this patient?

- A. Apical diastolic murmur
- ☒ B. Ejection systolic murmur
- C. Narrow pulse pressure
- D. Rumbling mid diastolic murmur

Anemia - Ejection systolic murmur (for murmur)
Tachycardia

3. A health officer was evaluating an anemic patient. Laboratory test revealed: Hgb = 7 g/dl, MCV = 77 fl, WBC = 5400/ μ l, Platelet = 650,000/ μ l. Which of the following is the most likely diagnosis? 50 - 88

- A. Acute myelogenous leukemia
- B. Aplastic anemia
- ☒ C. Iron-deficiency anemia
- D. Megaloblastic anemia

Anemia of I. Defe. → microcytic
Hypochromic
Also elevated vit. B12

4. A 45 year-old woman presented with fatigue and orthostatic dizziness of two months duration. She has conjunctival pallor. Complete blood count revealed: Hgb = 8 g/dl, WBC = 4200/ μ l, and platelet = 170,000/ μ l. Which of the following co-morbidities is LEAST likely?

- A. Chronic kidney disease ✓
- ☒ B. Chronic obstructive lung disease
- C. Colorectal cancer ✓
- D. Disseminated tuberculosis ✓

Anemia
150 - 400,000

5. A 57 year-old man is admitted to the Medical Ward with hemorrhagic stroke. After a week the patient suddenly developed pleuritic chest pain and shortness of breath. Physical examination revealed signs of respiratory distress and a swelling involving his right calf and thigh. What is the most likely cause for his deterioration?

- A. Acute myocardial infarction
- B. Aspiration pneumonia
- C. Fat embolism
- ☒ D. Pulmonary embolism

DVT

1. A 26 year-old female presented with two months of excessive menses and gum bleeding. Physical examination revealed pale conjunctiva, petechiae and ecchymosis. Which of the following is the most likely diagnosis?

- A. Hemophilia ✗
- ☒ B. Idiopathic thrombocytopenic purpura ✓
- C. Scurvy ✗
- D. Vitamin K deficiency ✗

2. A 21 year-old woman presented with fatigue and light headedness. Physical examination showed marked conjunctival and palmar pallor. Which of the following physical findings is most likely to be seen in this patient?

- A. Apical diastolic murmur
- ☒ B. Ejection systolic murmur
- C. Narrow pulse pressure
- D. Rumbling mid diastolic murmur

Anemia - Ejection systolic murmur (for murmur)
Tachycardia

3. A health officer was evaluating an anemic patient. Laboratory test revealed: Hgb = 7 g/dl, MCV = 77 fl, WBC = 5400/ μ l, Platelet = 650,000/ μ l. Which of the following is the most likely diagnosis? 50 - 88

- A. Acute myelogenous leukemia
- B. Aplastic anemia
- ☒ C. Iron-deficiency anemia
- D. Megaloblastic anemia

Anemia of I. Defe. → microcytic
Hypochromic
Also elevated vit. B12

4. A 45 year-old woman presented with fatigue and orthostatic dizziness of two months duration. She has conjunctival pallor. Complete blood count revealed: Hgb = 8 g/dl, WBC = 4200/ μ l, and platelet = 170,000/ μ l. Which of the following co-morbidities is LEAST likely?

- A. Chronic kidney disease ✓
- ☒ B. Chronic obstructive lung disease
- C. Colorectal cancer ✓
- D. Disseminated tuberculosis ✓

Anemia
150 - 400,000

5. A 57 year-old man is admitted to the Medical Ward with hemorrhagic stroke. After a week the patient suddenly developed pleuritic chest pain and shortness of breath. Physical examination revealed signs of respiratory distress and a swelling involving his right calf and thigh. What is the most likely cause for his deterioration?

- A. Acute myocardial infarction
- B. Aspiration pneumonia
- C. Fat embolism
- ☒ D. Pulmonary embolism

DVT

6. A 26 year-old man with rheumatic heart disease came for follow-up. He has active precordium and rumbling mid-diastolic murmur. What valvular lesion is most likely in this patient? *MS*

- A. Aortic regurgitation
- B. Mitral regurgitation
- ☒ C. Mitral stenosis
- D. Tricuspid regurgitation

7. A 65 year-old male with type 2 diabetes mellitus presents to the Emergency Department with a squeezing, retrosternal chest pain of one hour duration. He has vague discomfort in his right arm and shoulder. The physical examination is notable for diaphoresis and mild tachycardia. Which of the following is the most likely diagnosis?

- ☒ A. Acute coronary syndrome
- B. Acute pericarditis
- C. Aortic dissection
- D. Pulmonary embolism

8. Which of the following is a major criteria for the diagnosis of rheumatic fever?

- A. Elevated ESR
- B. Fever
- C. Prolonged PR segment
- ☒ D. Polyarthralgia

9. A 29 year-old woman came to the OPD with shortness of breath and body swelling of three weeks. On physical examination she has raised JVP, accentuated P2 and grade IV diastolic murmur at the apex. Which of the following drug delays disease progression?

- A. Frusemide
- B. Penicillin
- ☒ C. Enalapril
- D. Digoxin

10. A 45 year-old male with heart failure is comfortable at rest. He develops dyspnea and fatigue to walk to a nearby shop and is forced to rest for five minutes before symptoms improved. What is the stage and functional status of this patient?

- A. Stage A, Class II
- B. Stage B, Class III
- ☒ C. Stage C, Class IV
- D. Stage C, Class III

11. A 50 year-old man was found to have a BP=170/100. He has a history of asthma. Which one is the most appropriate drug of choice?

- A. Atenolol
- B. Enalapril
- ☒ C. Hydralazine
- D. Furosemide

12. A 40 year-old man presented with fever, shaking chills of three days duration. He also has non-productive cough and right-sided pleuritic chest pain. He has no significant past medical history. Physical examination revealed crepitation on the posterior right lower chest. Which one is the best initial investigation to confirm the diagnosis?
- ☒ A. Chest X-ray
 - B. Culture of sputum
 - C. Gram stain of sputum
 - D. White blood cell count
13. A 25 year-old woman who was diagnosed with pulmonary tuberculosis and started anti-tuberculosis treatment two weeks ago is found to be HIV positive on screening. On physical examination, she has pale conjunctiva and laboratory investigation shows CD4 count of 100cells/ μ L. Which one is the most appropriate first line antiretroviral regimen for this patient?
- A. ABC/3TC/NVP
 - ☒ B. TDF/3TC/EFV *1st (regimen)*
 - C. AZT/3TC/EFV *2nd (regimen)*
 - D. D4T/3TC/NVP
14. A 24 year-old man has generalized lymphadenopathy, aphthous ulcers, and gray-white plaques around his scrotum. Dark field microscopic examination demonstrates spirochetes. Which of the following is the treatment of choice?
- A. Ceftriaxone
 - B. Erythromycin
 - ☒ C. Penicillin
 - D. Tetracycline
15. A 30 year-old prisoner presents with intermittent fever and headache of two days duration. He claims that three of his roommates had similar symptoms and were treated in the prison. Blood film examination showed a spirochete. Which one of the following is the most appropriate antibiotic choice for this patient?
- A. Ceftriaxone
 - B. Ciprofloxacin
 - ☒ C. Procaine penicillin
 - D. Vancomycin
16. A 20 year-old male patient presents with abdominal swelling of three weeks duration. She lives in a rural village where the water source used is river water. Ultrasound examination reveals peri-portal fibrosis. Which one of the following is most likely cause for the patient's condition?
- A. River blindness
 - B. Ascariasis
 - C. Leishmaniasis
 - ☒ D. Schistosomiasis
17. A 22 year-old HIV positive patient who presented with shortness of breath was suspected to have Pneumocystis jirovecii pneumonia but he has no signs of respiratory distress and his CD4 count is 150/ μ L. Which of the following drugs is recommended for this patient?
- ☒ A. Co-trimoxazole
 - B. Fluconazole
 - C. Prednisolone
 - D. Pyrimethamine

23. A 30 year-old man presented with epistaxis, easy fatigability and intermittent fever for one month. Physical examination revealed hepatosplenomegaly. Which of the following is the best investigation to confirm the diagnosis?

- A. Blood culture
- ☒ B. Bone marrow aspiration
- C. Stool examination
- D. Ultrasound of abdomen

24. A 60 year-old man presented with sudden onset of right sided weakness on both upper and lower extremities before a day. The weakness is maximal at onset and during sleep. He is hypertensive but does not take medications. What is the most likely diagnosis?

- A. Hemorrhagic stroke
- ☒ B. Ischemic stroke
- C. Subarachnoid hemorrhage
- D. Transient ischemic attack

25. Which of the following is the most common cause of hemorrhagic stroke in Ethiopia?

- A. Aneurysm
- B. Anticoagulation
- C. Brain tumor
- ☒ D. Hypertension

26. A 24 year-old HIV positive woman presents with fever and skin rash over the left chest two days ago. The rash is vesicular and follows dermatomes (T4-5). She is confused and meningeal signs are positive. Which of the following is the most likely diagnosis?

- A. Fungal meningitis
- B. Pyogenic meningitis
- C. Tuberculous meningitis
- ☒ D. Viral Meningoencephalitis - *was Herpes Zoster*

27. A 30 year-old male presented with headache, fever and progressive weakness on the left side of his body two weeks ago. He has right ear discharge since two months. On physical examination BP=140/80 mmHg, PR=102/min, T=39°C. He is mentally alert and power on the left upper and lower extremities is 3/5. What is the most likely diagnosis?

- ☒ A. Brain abscess
- B. Brain tumor
- C. Ischemic stroke
- D. Subdural hematoma

18. A 60 year-old male had cough for five years with clear sputum production and shortness of breath. He has smoked a pack of cigarettes per day for 20 years. Laboratory examination reveals haemoglobin of 19 g/dl. X-ray of the chest shows hyperinflation without infiltrates. Which of the following is the most likely diagnosis?

- A. Bronchial asthma
- ☒ B. Chronic obstructive pulmonary disease
- C. Interstitial lung disease
- D. Pulmonary Tuberculosis

19. A 40 year-old alcoholic develops productive cough and fever. Chest x-ray shows an air-fluid level in the superior segment of the right lower lobe. Which of the following is the most appropriate pharmacotherapy?

- A. Ceftazidime and ciprofloxacin
- B. Ceftriaxone and gentamycin
- ☒ C. Metronidazole and crystalline penicillin (COPD + typhoid)
- D. Vancomycin and chloramphenicol

20. A 25 year-old man presents with abdominal pain, headache and fever of five days duration. The Health officer suspected the diagnosis of typhoid fever. Which of the following investigations confirms the diagnosis?

- A. Stool examination
- B. Blood culture
- C. Weil-Felix titer
- ☒ D. Widal test (typhoid)

21. A 27 year-old man complains of fever, rigors, productive cough with rusty-coloured sputum and pleuritic chest pain. Chest examination revealed increased tactile fremitus and dullness to percussion in the right lower lung field. What is the most likely diagnosis?

- ☒ A. Lobar pneumonia (S - AU)
- B. Bronchopneumonia
- C. Lung abscess
- D. Pleural effusion

22. A 35 year-old woman presented with trismus for a day. She also has shoulder and back pain. She reported that she sustained a puncture over the left foot a week ago. She is fully conscious. Which of the following is the most appropriate pharmacotherapy?

- A. Carbamazepine
- ☒ B. Diazepam
- C. Phenobarbital
- D. Phenytoin

28. A 60 year-old alcoholic patient presented with fever, headache and neck pain of two days duration. On examination body $T=39^{\circ}\text{C}$ and has neck stiffness. Bacterial meningitis was suspected but lumbar puncture was deferred because he has papilledema. Which of the following is the best empiric therapy for this patient in addition to Ampicillin?
- A. Vancomycin and Chloramphenicol
 - ☒ B. Ceftriaxone and Vancomycin
 - C. Ceftriaxone and Chloramphenicol
 - D. Cloxacillin and Ciprofloxacin
29. A 36-year-old male presented with fatigue and tea-colored urine for five days. Physical examination revealed jaundice and tender Hepatomegaly. Laboratory investigation revealed: Aspartate aminotransferase (AST) = 2400 U/L, Alanine aminotransferase (ALT) = 2500 U/L, Alkaline phosphatase = 210 U/L and total bilirubin = 8.6 mg/dL. Which of the following is the LEAST likely cause for the patient's condition?
- A. Acetaminophen poisoning
 - B. Acute Viral hepatitis
 - ☒ C. Alcoholic liver disease
 - D. Ischemic liver injury
30. A 56 year-old man presented with progressive abdominal distension and leg swelling of two months duration. He is a known patient with cirrhosis for the past one year on follow up but not on medication. On examination, he was not in distress and has moderate ascites and bilateral leg swelling. Which of the following is the best initial management?
- A. Furosemide
 - ☒ B. Furosemide and Spironolactone *(K⁺ sparing)* 5:1
 - C. Hydrochlorothiazide
 - D. Paracetamol and Furosemide
31. Which of the following is the most common complication of Peptic ulcer disease?
- ☒ A. Hemorrhage
 - B. Gastric outlet obstruction
 - C. Malignancy
 - D. Perforation
32. A 50 year-old woman presented with fever, cough and chest pain for two days. Chest X-ray showed pleural effusion. Pleural fluid analysis showed: WBC = 5,000/mm³, with 90% neutrophils, PH = 7.2, blood culture was positive for streptococcus pneumonia. What is the next best step in management of this patient besides antibiotics?
- A. Dexamethasone
 - B. Chest CT scan
 - ☒ C. Chest tube insertion
 - D. Lung function test

33. A 46 year-old man is brought to the Emergency Department because he was vomiting fresh blood. He reports that he has no abdominal pain or nausea. On physical examination; BP=70/40 mmHg, PR=118/min, RR= 24/min. He also has gynecomastia, dilated abdominal veins and ascites. Which of the following is the most likely cause of the bleeding?

- A. Esophageal cancer
- ☒ B. Esophageal varices

- C. Mallory Weiss
- D. Reflux esophagitis

34. A 47 year-old man comes to a Hospital complaining of epigastric pain that is exacerbated by eating. Antigenic test for *Helicobacter pylori* is positive. Endoscopic examination reveals a duodenal ulcer. Which of the following is the most appropriate therapy?

- ☒ A. Amoxicillin, clarithromycin and omeprazole
- B. Amoxicillin, metronidazole and anti-acid
- C. Omeprazole and anti-acid
- D. Ranitidine, amoxicillin and metronidazole

35. A 29 year-old man with HIV infection presents with chronic diarrhea associated with anorexia, abdominal cramps and significant weight loss. His recent CD4 count =20 cells/mm³. A modified acid-fast stain of a stool specimen shows 4-6 μ m oocysts. Which of the following is the most likely cause?

- ☒ A. *Cryptosporidium parvum*
- B. Microsporidia

- C. *Mycobacterium avium* complex
- D. *Pneumocystis jirovecii*

36. Which of the following is the best early indicator for diabetic nephropathy?

- A. Rising blood urea nitrogen
- ☒ B. Albuminuria
- C. Hypertension
- D. Rising creatinine level

37. A 35 year-old man presents with generalized body swelling of two months. Physical examination reveals BP=120/70mmHg, ascites and +3 pretibial edema. Protein = 4.5g/24hour urine collection. Which of the following is the most likely diagnosis?

- A. Chronic liver disease
- B. Heart failure
- ☒ C. Nephrotic syndrome
- D. Glomerulonephritis

38. A 55 year-old man comes to the Emergency OPD because of excruciating pain in his right big toe. He was diagnosed to have gouty arthritis. Which of the following is the most appropriate pharmacotherapy?

- A. Allopurinol
- B. Ceftriaxone

- ☒ C. Indomethacin
- D. Morphine

730
830
2930
1030
1130

39. A 28 year-old lactating woman presented to the emergency OPD with a complaint of right breast pain of five days duration. On examination body T= 38°C with right breast tenderness but no fluctuation. What is the most likely diagnosis?
- ☒ A. Breast Abscess
 - B. Breast Cancer
 - C. Fibroadenoma
 - D. Fibrocystic Change
40. A 32 year-old man presented to the surgical OPD with complaint of neck mass of one year duration. On examination he has a hoarse voice and a solitary 2cm x 2cm thyroid mass on the right side which has firm consistency and fixed to underlying structures. There was no lymphadenopathy. What is the most likely diagnosis?
- A. Thyroid adenoma
 - ☒ B. Thyroid cancer
 - C. Thyroiditis
 - D. Toxic Nodule
41. A 35 year-old man with a chronic peptic ulcer disease presented with features of acute abdomen and chest x-ray revealed pneumoperitoneum. His symptoms lasted for six hours and his vital signs were stable after infusion of one liter of Normal Saline solution. What should be the next step in the management of this case?
- A. Barium meal study
 - B. Endoscopy
 - C. Triple therapy
 - ☒ D. Laparotomy
42. A 50 year-old woman brought an ultrasound imaging result which showed Gall Bladder Stone. She has no abdominal pain and no known medical illness. Which of the following is the recommended management?
- A. Cholecystectomy
 - B. Extracorporeal Shockwave Lithotripsy (ESWL)
 - ☒ C. Reassurance
 - D. Ursodeoxycholic acid
43. A 68 year-old man is admitted to a Hospital with complaint of intermittent lower abdominal pain, bowel habit change and rectal bleeding of one month duration. He has tenesmus and incomplete sense of defecation and had lost weight. What is the most likely diagnosis?
- A. Amebic dysentery
 - ☒ B. Colorectal cancer
 - C. Sigmoid Volvulus
 - D. Ulcerative colitis
44. A 55 year-old man presented to the surgical OPD with a complaint of rectal bleeding of one month duration. He has no protruding mass per rectum and digital rectal examination was non-revealing. What is the investigation of choice?
- A. Abdominal CT-Scan
 - B. Barium Enema
 - ☒ C. Colonoscopy
 - D. Upper GI Endoscopy

45. A 30 year-old man was seen at the Emergency surgical OPD with perianal pain of two days duration. He has fever and chills. He has left perianal induration with tenderness but no fluctuation. What is the most likely diagnosis?

- A. Anal fissure
- ☒ B. Perianal Abscess
- C. Perianal fistula
- D. Thrombosed Hemorrhoids

46. A 20 year-old man was seen at the emergency department following road traffic accident of one hour duration. He has sustained trauma to the abdomen. On examination; Vital signs: BP=80/50 mmHg and PR=120/min. He has tire marks on the abdomen and it was tender. The patient is started on IV fluids. What is the next step in the management?

- A. Abdominal CT-scan
- ☒ B. Exploratory laparotomy
- C. Observation
- D. Plain abdominal x-ray

47. A 50 year- old man came complaining of fatigue and dyspepsia of three months duration. Physical examination revealed a pale conjunctiva with a palpable mass at the epigastrium that moves with respiration. What is the gold standard diagnostic modality?

- A. Abdominal Ultra sound
- B. CT scan of the abdomen
- ☒ C. Upper GI Endoscopy
- D. Plain abdominal film

48. A man was stabbed to the abdomen and presented to the Health Center with eviscerated small bowel which was getting dark. Which of the following is the correct initial measure before referral?

- ☒ A. Cover the intestine with saline gauze
- B. Refer to a nearby hospital without intervention
- C. Replace the intestine back into the peritoneal Cavity
- ☒ D. Widen the stab wound under local anesthesia

49. A 23 year-old farmer presented to the OPD with abdominal pain and frequent vomiting of one day duration. On examination the abdomen is slightly distended and has visible peristalsis. Plain abdominal x-ray showed step ladder pattern air-fluid level. Which of the following is the most likely diagnosis?

- A. Caecal Volvulus
- B. Compound Volvulus
- C. Sigmoid Volvulus
- ☒ D. Small bowel Volvulus

50. A 65 year-old man was seen at the Emergency OPD with abdominal pain, distension and failure to pass feces and flatus of two days duration. He had two similar attacks in the last three years and was successfully treated with rectal tube deflation. On Examination BP=90/60 mmHg and PR=110/min. The abdomen was tender and bowel sounds were absent. Which of the following measures is to be avoided?

- A. Analgesics
- B. Antibiotics
- C. IV fluid
- ☒ D. Rectal tube deflation

51. A 45 year-old housewife presented with progressive submandibular swelling of two days duration. She visited a dentist and undergone tooth extraction a day prior to the start of her symptoms. Examination showed tense, shiny bilateral submandibular swelling with no area of fluctuation. She also has drooling of the saliva and protruded tongue. What is the diagnosis of this patient?
- ☒ A. Ludwig's angina
 - ☐ B. Periodontal abscess
 - ☐ C. Submandibular cellulites
 - ☐ D. Submental abscess
52. A 20 year-old man presented to the OPD forty minutes after he sustained stab injury to his left posterior chest. He has shortness of breathing. He is acutely sick looking; BP= 110/60 mmHg, PR= 90/min, RR=26/min. He has decreased air entry on the lower left chest with visible air rush through the stab wound. What is the best first step in the management of this patient?
- ☐ A. Apply airtight dressing
 - ☒ B. Apply triangular dressing
 - ☐ C. Suture the wound
 - ☐ D. Take chest X-ray
53. A 35 year-old male patient came to emergency OPD 12 hours after he started to have sudden onset of epigastric pain. He has long standing history of dyspepsia for which he has been taking anti-acids repeatedly. Examination revealed tachycardia, rigid and tender abdomen. What is the appropriate imaging study to reach a definitive diagnosis?
- ☐ A. Barium follow through
 - ☐ B. Barium meal
 - ☒ C. Chest X-ray
 - ☐ D. Plain abdominal X-ray
54. A 27 year-old patient was involved in a road traffic accident. He was brought to the Hospital by the driver. He is agitated; PR=140/min and feeble, BP=80/40 mmHg. He has pale conjunctiva and his chest is clear with good air entry. He has an echymosis over his left flank area. Abdominal Ultrasonography showed a significant collection of fluid in his abdomen. What is the most likely diagnosis?
- ☐ A. Cardiogenic shock
 - ☒ B. Hemorrhagic shock
 - ☐ C. Neurogenic shock
 - ☐ D. Septic shock
55. A 64 year-old male patient presented to the Emergency OPD after he failed to pass urine for a day. He had difficulty of urination for the last one year. The previous night he took seven bottles of beer with friends. On examination he is acutely sick looking and in pain. Bladder is palpable 10 cm above the pubic bone and tender. What is the most likely cause for acute urinary retention?
- ☒ A. Benign Prostatic Hyperplasia
 - ☐ B. Bladder cancer
 - ☐ C. Bladder stone
 - ☐ D. Urethral Stricture

56. An 18 years old male presented with right scrotal swelling of 1 day duration. He also has associated pain and it started while he was asleep, it woke him up. He also has fever and pain during urination. On examination right testis is tender, hot to touch, swollen, and the testicular cord is not thickened. What is the most likely diagnosis?
- A. Testicular torsion
 - ☒ B. Epididymo-orchitis
 - C. Infected hydrocele
 - D. Testicular tumor
57. A 36 year-old male patient came with difficulty of urination for the last two years. He had previous history of whitish urethral discharge for which he was treated with tablets and injections. What is the most likely diagnosis?
- A. Benign prostatic hyperplasia
 - B. Prostatic Cancer
 - C. Sexually transmitted disease
 - ☒ D. Urethral Stricture
58. A 43 year-old male patient came to the Emergency OPD after being involved in a road traffic accident before a day. He has failed to urinate since the incident and is unable to move his right lower limb. On examination his vital signs are normal, there is right pubic tenderness and blood on the penile meatus. What should be the next step in management?
- A. Decompress with percutaneous needle puncture
 - B. Do image-guided catheterization
 - C. Insert a foley catheter to decompress the bladder
 - ☒ D. Perform emergency suprapubic cystostomy
59. A 24 year-old man came to the Emergency OPD with severe pain originating in the flank, radiating inferiorly and anteriorly. He also has nausea and vomiting. His vital signs are within normal limits. Abdominal examination showed marked costovertebral angle tenderness. Urinalysis showed microscopic hematuria. What is the most likely diagnosis?
- A. Cystitis
 - B. Pyelonephritis
 - C. Pyonephrosis
 - ☒ D. Ureteric colic
60. A 60 year-old female patient presented with right sided flank pain and hematuria of one month duration. She also has significant weight loss, night sweats and malaise. On physical examination, she is chronically sick looking with normal vital signs, a bimanually palpable mass on the right flank. What is the most likely diagnosis?
- A. Chronic pyelonephritis
 - B. Neuroblastoma
 - ☒ C. Renal cell cancer
 - D. Wilm's tumor

77. A 38 year-old primigravida comes for a routine visit at 39 weeks of gestation. Her blood pressure is persistently 140/90 mm Hg, and her urine protein is +2. Physical examination is otherwise unremarkable, and she is completely asymptomatic. Her cervix is 2 cm dilated and 90% effaced, vertex presentation and at station 0.

Which of the following is the most appropriate management option for this case?

- ☒ A. Admission for observation
- B. Immediate cesarean section
- C. Outpatient follow-up till 40 weeks
- D. Start induction of labor

78. A 34 year-old primigravida comes with cessation of fetal movement of four days duration at gestational age of 35 weeks from LNMP. She is otherwise asymptomatic and has no medical problem. Examination shows stable vital signs, symphysio-fundal height of 34cms, no uterine contraction and absent FHB.

Which of the following is the initial best step in the management of this patient?

- A. Consider Termination
- B. Do coagulation profile
- C. Prophylactic antibiotics
- ☒ D. Ultrasound for confirmation

79. A 63 year-old woman who came for regular gynecologic check-up was incidentally diagnosed to have cystocele. Which of the following is the most appropriate management for this patient?

- A. Anterior colporaphy
- B. Kegel's exercise
- C. Pessaries
- D. Vaginal hysterectomy

80. A 21 year-old GII PI mother at 24 weeks gestation complains of sever vulvar itching and cotton cheese like discharge. A wet mount slide show no clue cells but hyphae. Which of the following is the most likely diagnosis?

- A. Bacterial vaginosis
- ☒ B. Candida albicans
- ☒ C. Chlamydia trachomatis
- D. Trichomonas vaginalis

81. A 29 year-old woman presents with vaginal pruritus associated with white curd like discharge. On examination there is erythematous macular lesion but no vesicles. Investigation shows no clue cell, negative whiff test. Which of the following is the treatment of choice for this case?

- A. Azithromycin
- ☒ B. Fluconazole
- C. Metronidazole
- D. Oral acyclovir

82. A 26 year-old GIIPI woman at 26 weeks of gestation from her LNMP, comes for ANC follow up. She hasn't appreciated fetal movement yet. She had one episode of bleeding a month back and noticed failure of increase in uterine size. Her uterine size measures 20 weeks and you are unable to hear fetal heart tone.

Which of the following is the most descriptive diagnosis?

- A. Incomplete abortion
- B. Inevitable abortion
- ☒ C. Missed abortion
- D. Threatened abortion

90. A PIII mother who gave birth to an alive healthy term neonate vaginally before an hour was found to be restless and pale. On physical examination BP= 60/40 mmHg, PR=120/min, RR=24/min, T=36°C; uterus 22 week and soft and active vaginal bleeding. Which of the following is the most appropriate first step in the management of this patient?
- A. Administer crystalloids
 - B. Administer intra-nasal oxygen
 - C. Blood transfusion
 - ☒ D. Call for help
91. In the active management of third stage of labor, placenta failed to be delivered by controlled cord traction. Which of the following is the next best step in management?
- A. Fundal pressure
 - ☒ B. Manual removal of placenta
 - C. Pitocin IV drip
 - D. Wait for spontaneous expulsion
92. A 15 year-old girl came to Gynecology OPD with a complaint of abdominal pain of six months duration. The pain is crampy and recurrent. She had never seen menses before. On examination there is a central 14 week-sized abdomino-pelvic mass. Which of the following is the most likely diagnosis?
- A. Dermoid cyst
 - B. Full bladder
 - ☒ C. Imperforate hymen
 - D. Myoma
93. A 32 year-old nulligravida woman came to gynecology OPD with a complaint of heavy menstrual flow of one year duration. She also gave history of constipation and pelvic heaviness and failure to conceive of four years duration. On examination there is central, 12 weeks size firm abdomino-pelvic mass. Which one is the most likely diagnosis?
- A. Endometrial hyperplasia
 - B. Ovarian tumor
 - ☒ C. Uterine myoma
 - D. Uterine sarcoma
94. A 35 year-old GVPIV mother at gestational age of 8 weeks presented to gynecology OPD with a complaint of vaginal bleeding of four days duration. She had excessive vomiting her uterus is big-for-date. Which of the following is the most likely diagnosis?
- A. Cervical polyp
 - B. Ectopic pregnancy
 - C. Incomplete abortion
 - ☒ D. Molar pregnancy
95. Which of the following contraceptive methods is the most effective?
- A. Combined oral contraceptive pills
 - B. Female condom
 - ☒ C. Intrauterine contraceptive device
 - D. Medroxy progesterone acetate

96. A 17 year-old student came to a Clinic seeking pregnancy termination service at gestational age of 10 weeks. She gave a history of sexual assault. Which of the following is the best method of termination of pregnancy for this case?
- ☒ A. Dilatation and curettage
 - ☐ B. Intramuscular methotrexate
 - ☐ C. Manual vacuum aspiration
 - ☐ D. Mifepristone and misoprostol
97. A 24 year-old primigravida who came for a routine ANC visit tested positive for HIV. When should a health officer initiate ARV drugs for this woman?
- ☒ A. As soon as possible
 - ☐ B. At 20 weeks of gestation
 - ☐ C. At 28 weeks of gestation
 - ☐ D. At 36 weeks of gestation
98. A primigravida came to clinic with a complaint of lower abdominal pain of one day duration at GA of 22 weeks. She has previous history of excessive menstrual flow and uterine tumor which was not treated. On examination Vital signs-within normal limits. Abdomen localized tenderness, uterus 24 weeks size, No contraction FHR- 150bpm. Which of the following is the most likely diagnosis in this case?
- ☐ A. Abdominal pregnancy
 - ☐ B. Acute appendicitis
 - ☒ C. Degenerative myoma
 - ☐ D. Ovarian cyst torsion
99. A Para III mother who gave birth to an alive male neonate at home before one hour brought to a Hospital because of restlessness and protrusion of mass per vaginum. On examination she is acutely sick looking, BP is not recordable, PR=132/minute and feeble RR=20/min, T=36°C. Abdomen-uterus not palpable, globular fleshy mass with minimal bleeding protruding through the vaginum. Which of the following is the most appropriate initial management of this case?
- ☐ A. Intra nasal oxygen 6 L/minute
 - ☐ B. Reposition the uterus
 - ☒ C. Resuscitation with IV fluids
 - ☐ D. Start oxytocin drip
100. A 20 year-old primigravida at 38 weeks was admitted in second stage of labor and stayed for 90 minutes. Fetal heart beat is ranging between 80 – 90/min, vertex presentation with the occiput direct anterior, station +3, grade-I molding, no caput and grade-II meconium stained fluid. Which of the following is the next best step in management?
- ☒ A. Expedite delivery right away by vacuum
 - ☐ B. Refer her for cesarean section
 - ☐ C. Re-position her into the left lateral position
 - ☐ D. Secure IV line and await spontaneous vaginal delivery
101. A 6-year old diabetic patient is admitted to ER with the diagnosis of sever DKA and has received normal saline and insulin for the last 3 hours. Which of the following electrolyte abnormalities is expected?
- ☒ A. Hypocalcemia
 - ☐ B. Hypokalemia
 - ☐ C. Hyponatremia
 - ☐ D. Hypophosphatemia



61. A 56 year-old man presented with a scrotal mass of three months duration. Physical examination revealed a 3cm x 4 cm firm mass on the lower pole of the right testis. Scrotal ultrasound showed a well-defined hypoechoic lesion within the testis. Serum β -hCG was elevated. What is the most likely diagnosis?
- A. Orchitis
 - B. Hydrocele
 - ☒ C. Testicular Tumor
 - D. Testicular torsion
62. A 35 year-old male patient presented to the Emergency Room after sustaining a bullet injury to his left forearm three hours back. Up on evaluation vital signs were normal and a deformed and rotated forearm, intact distal neurovascular examination. X-ray showed comminuted displaced radio-ulnar fracture. Which of the following is a temporary means of immobilization?
- A. Apply POP cast
 - ☒ B. External fixation
 - C. Lateral arm traction
 - D. Plating and screw
63. A 65 year-old male patient sustained a fall down accident three hours prior to presenting to the Emergency Room. He fell down with the arms extended. On examination vital signs were normal with a right wrist deformity. X-ray showed displaced Colle's fracture. Which one is the best management option?
- A. Application of a POP Cast
 - ☒ B. Closed reduction and casting
 - C. Open reduction and plating
 - D. Skin traction for few days then casting
64. A 20 year-old male patient had an external fixator applied for a compound comminuted right tibio-fibular fracture three weeks prior to presentation. He now complains of pain at the pin site. On examination there is seropurulent discharge from the two distal pins. Which of the following is the best diagnosis?
- A. Cellulitis
 - B. Osteomyelitis
 - C. Osteonecrosis
 - ☒ D. Pin tract infection
65. A 20 year-old male patient came to you with right knee pain of five days duration. On examination BP=120/70 mmHg, PR= 84/min. The knee is swollen and tender with signs of knee effusion. X-ray showed a wide joint space. Joint fluid analysis revealed turbid fluid with $100,000 \text{ WBC/mm}^3$. Which of the following is the most likely diagnosis?
- A. Acute osteomyelitis
 - B. Osteoarthritis
 - ☒ C. Pyogenic arthritis
 - D. Tuberculous arthritis

102. Which of the following electrolytes is found in high concentrations in ReSOMal when compared with standard WHO oral rehydration solution?
- A. Chloride
 - B. Phosphorous
 - ☒ C. Potassium
 - D. Sodium
103. An 11-month old male infant is brought to the ER with watery diarrhea of 2 days duration. Physical examination reveals irritable infant, weight=10 kg, sunken eye balls and eagerness to drink. What is the correct type and volume of fluid for this infant?
- A. Ringer's lactate, 300ml
 - B. 0.45% Normal saline, 700ml
 - ☒ C. Oral rehydration solution, 750ml
 - D. Ringer's lactate 500ml then ORS 500ml
104. An 8-year old boy is diagnosed with acute rheumatic fever three weeks after an episode of sore throat. Which drug should have been given for the sore throat to prevent the rheumatic fever?
- A. Clindamycin within 11 days
 - B. Cloxacillin within 7 days
 - C. Erythromycin within 16 days
 - ☒ D. Penicillin within 9 days
105. An 8-month old infant is brought to ER with fever, cough, and fast breathing of three days duration. Further inquiry reveals history of repeated chest infection, interruption during breast feeding and poor weight gain. Precordial examination reveals active precordium and pansystolic murmur at left lower sternal border. Which of the following heart lesions most likely explains these findings?
- A. Atrial septal defect
 - B. Tetralogy of Fallot
 - C. Tricuspid atresia
 - ☒ D. Ventricular septal defect
106. A 9-week old girl who was apparently healthy two days ago is brought to the ER. She is pale and has dyspnea with RR=80/min, T=37.7°C, PR=195/min, heart sounds are distant, a gallop is heard, liver is 6 cm below right costal margin. Chest X-ray shows cardiomegaly and echocardiogram reveals poor ventricular function and dilated left heart chambers. Which of the following is the most likely diagnosis based on this clinical scenario?
- ☒ A. Dilated cardiomyopathy
 - B. Endocarditis
 - C. Myocarditis
 - D. Pericarditis
107. An 8-year old boy known cardiac patient on follow up has a 3-week history of low-grade fever of unknown source, fatigue, weight loss, myalgia, and headaches. On repeated examinations during this time, he is found to have developed a new heart murmur, petechiae, and mild splenomegaly. Which of the following is the most likely diagnosis?
- A. Acute rheumatic fever
 - B. Rheumatic recurrence
 - ☒ C. Infective endocarditis
 - D. Dilated cardiomyopathy

83. A 42 year-old healthy nulliparous woman has a firm, irregular shaped, 12 week-sized midline abdominal mass.

Which of the following is the best initial step in evaluating this patient?

- A. Abdominal x-ray
- B. CT scan
- C. MRI
- ☒ D. Ultrasound

84. Which of the following is NOT a component of active third stage management?

- A. Controlled cord traction
- ☒ B. Nipple stimulation
- C. Uterine massage
- D. Oxytocin provision

85. A GIII PII mother came to a clinic at gestational age of 16 weeks for ANC follow up.

Which of the following is LEAST likely to be requested as part of a routine work up?

- A. Blood group and Rh
- ☒ B. Ultrasound
- C. Urinalysis
- D. VDRL

86. A primigravida was admitted to a Health Center due to pushing down pain of 6 hours. Physical examination; vital signs were normal, uterine size 36 weeks, longitudinal, cephalic, FHR=146/min, two contractions in 10min each lasting for 30-45 seconds; cervix is 3 cm dilated and 60 % effaced. After 4 hours she was reevaluated and had same findings except the cervical effacement changed to 75 %.

Which of the following is the best management option?

- A. Augmentation
- B. Rehydration
- C. Referral to hospital
- ☒ D. Close follow up

87. Obstetric forceps was applied to effect delivery of an alive female neonate at station +2 in left occipito-anterior position with a 45° rotation.

Which of the following best describes this type of forceps delivery?

- A. High
- B. Low
- C. Midpelvic
- ☒ D. Outlet

88. A pregnant lady had a uterine size which is five centimeters less than the stated LMP.

Which of the following is LEAST likely as a possible cause for this discrepancy?

- A. Fetal infection
- B. Congenital malformation
- C. Wrong date
- ☒ D. Concealed abruption

89. Which of the following is an essential prerequisite for application of forceps?

- A. Cervix more than 8 cm
- B. Gestational age beyond 37 weeks
- ☒ C. Ruptured membranes
- D. Vertex presentation

108. A 6-year old boy is diagnosed with acute rheumatic fever with the following findings: migratory polyarthrititis, fever, elevated ESR, and evidence of preceding streptococcal infection. Which of the following is the appropriate regarding the management of the above patient?
- A. Amoxicillin for 7 days
 - B. Aspirin
 - ☒ C. Penicillin prophylaxis life long
 - D. Prednisolone
109. After a diagnosis of pertussis in a toddler, the parents are worried about the transmission to other house hold members. Which house hold members should receive erythromycin prophylaxis?
- A. All house hold members
 - B. Immuno-compromised
 - C. Unvaccinated against pertussis
 - ☒ D. Younger than 7 year
110. Which of the following clinical conditions that appear after pertussis vaccination is an absolute contraindication to subsequent administration of the vaccine?
- A. Fever and convulsion within three days
 - B. Immediate anaphylactic reaction
 - C. Inconsolable cry longer than two hours
 - D. Shock-like state after two days
111. Which of the following vaccine should NOT be given for an infant with severe immune suppression?
- A. Hepatitis B
 - B. Measles
 - C. Meningococcal
 - D. Pneumococcal vaccine
112. You are evaluating an HIV exposed neonate immediately after delivery. The mother had proper ANC follow up and was managed according PMTCT option B+. Which of the following is the appropriate recommendation for this newborn?
- A. Avoid breast feeding
 - B. Cotrimoxazole prophylaxis within the first week of life
 - C. DNA- PCR test within 48 hours
 - ☒ D. Nevirapine for 6 weeks
113. A 3 month old infant has rhinorrhea, cough and noisy breath. Physical examination reveals RR=72/min, PR=142/min and T= 37.8°C. He is irritable and has diffuse wheezes and chest retraction. What is the most likely diagnosis?
- ☒ A. Bronchiolitis
 - B. Gastro-esophageal reflux disease
 - C. Hyper active airway disease
 - D. Severe pneumonia

121. A 10 years old boy bled from multiple sites after he sustained an accident while riding a bicycle. On examination he has tachycardia and the blood pressure is not recordable. What will be the correct amount of crystalloid you will administer as the initial bolus?
- A. 10 ml/kg
☒ B. 20 ml/kg
 C. 30 ml/kg
 D. 40 ml/kg
122. A 5 year-old boy presented to the OPD with abdominal swelling of one month duration. On examination BP is normal, PR=100/min, RR= 42/min. He has stony dullness on the posterior right two third of the lung field, shifting dullness and fluid thrill on abdominal examination. He has pitting pedal edema. Urinalysis showed protein 4+. Which one of the following serum levels should you measure?
- A. Alanine transaminase
 B. Bilirubin
☒ C. Cholesterol
 D. Creatinine
123. A 36 month-old male infant presented with high grade fever and vomiting of two days duration. He has right costovertebral angle tenderness. White cell count showed leukocytosis with left shift and blood film is negative for hemoparasite. Urine is positive for nitrite and leukocyte esterase. What is the most likely diagnosis in this child?
- ☒ A. Pyelonephritis
 B. Hepatitis
 C. Appendicitis
 D. Lobar pneumonia
124. A 5-year old boy sustained a penetrating injury to his right leg. Two weeks later he developed swelling of the injured leg and high grade fever. X-ray of the affected limb showed features of acute osteomyelitis. What is the most likely etiology?
- A. ~~Pseudomonas~~ aeruginosa
 B. Klebsiella pneumonia
☒ C. Streptococcus pyogenes
 D. E. coli
125. A 2 year-old child has delay in all developmental mile stones for his age. On examination he has macrocephaly, severe stunting, protruded tongue, umbilical hernia and coarse facial features. The best initial test is to determine serum level of :
- A. Cortisol
 B. Growth Hormone
☒ C. Thyroid Stimulating Hormone
 D. Vitamin D
126. Which one of the following drugs is an appropriate antidote for Malathion poisoning?
- ☒ A. Atropine
 B. Flumazenil
 C. N acetyl cysteine
 D. Naloxone

114. Which of the following best describes clinical features of Guillain-Barré syndrome?
- ☒ A. Ascending symmetric paralysis without sensory change
 - ☐ B. Asymmetric paralysis and sensory change
 - ☐ C. Symmetric paralysis and cranial nerve palsy
 - ☐ D. Symmetric paralysis and sensory changes
115. A 5 year-old boy with HIV infection is receiving Zidovudine, Lamivudine and Nevirapine. He presents with progressive difficulty of breathing for three weeks. He has no history of fever. On examination PR=108/min, RR=72/min, T= 37°C and Oxygen saturation is 70 %. Chest radiograph reveals bilateral diffuse alveolar disease. What is the most likely diagnosis?
- ☐ A. Heart failure
 - ☒ B. Pneumocystis pneumonia
 - ☐ C. Pulmonary tuberculosis
 - ☐ D. Severe bacterial pneumonia
116. The recommended time to start folate supplementation for a mother who had a baby with myelomeningocele to prevent its recurrence in future pregnancies is:
- ☐ A. At the first missed period
 - ☐ B. End of first trimester
 - ☒ C. One month before conception
 - ☐ D. Third trimester
117. Which of the following CSF profile is expected in a child with meningoencephalitis?
- ☐ A. Leukocytosis with neutrophil predominance
 - ☐ B. Normal protein level
 - ☒ C. Normal or slightly reduced glucose level
 - ☐ D. Organism on Gram stain
118. A 12 hour old neonate presented with a boggy diffuse scalp swelling. Forceps was applied for prolonged second stage of labor. On examination he is in respiratory distress, grossly pale, and the capillary refill was 5 seconds. What is the most likely explanation for the above scenario?
- ☐ A. Caput succedaneum
 - ☐ B. Cephalhematoma
 - ☐ C. Intracranial hemorrhage
 - ☒ D. Subgaleal hemorrhage
119. A 6 week-old infant presented with high grade fever three hours after he took the first of his scheduled vaccines. He was well before the vaccine. On examination he is febrile and irritable otherwise there are no findings in the other system. What is the most appropriate management of this infant?
- ☐ A. Admit for observation
 - ☐ B. Initiate oral antibiotics
 - ☐ C. Investigate for infection
 - ☒ D. Paracetamol suppository
120. Which of the following CSF profiles suggests pyogenic meningitis in neonates?
- ☒ A. High cell count
 - ☐ B. High protein
 - ☐ C. Low glucose
 - ☐ D. Positive gram stain

32/min, temperature 39°C and has petechial rash on trunk. Which of the following is the best initial test for diagnosis?

- A. Blood film
- B. Complete blood count
- ☒ C. CSF analysis
- E. Widal test

133. Infectious diseases may have more than one mode of transmission. The number of modes of transmission has an effect on the control, elimination or eradication of that specific disease. From this perspective, which of the following diseases is more difficult to control?

- A. Pulmonary tuberculosis
- B. Relapsing fever
- C. Syphilis
- D. Trachoma

134. A health officer is delegated to lead a team travelling to West African countries to control the epidemic of Ebola Virus Disease (EVD). The role of the team includes giving supportive care to the cases of EVD and teaching them to avoid contact with the rest of the community members. To which prevention level (s) can this be categorized?

- A. Primary prevention both to the patients and the community
- B. Primary prevention to the patients but secondary prevention to the community
- C. Primary prevention to the community but secondary prevention to the patients
- D. Primary prevention to the community but tertiary prevention to the cases

135. An estimated 90 people out of 100 cases of Ebola die. What does this proportion show?

- A. Cause specific mortality rate
- B. Case fatality rate
- C. Proportionate mortality ratio
- D. Crude death rate

136. Suppose psychosis is a rare health problem in a specific country. Researchers want to study whether khat chewing causes psychosis in this country. Which one of the following study designs is most appropriate to assess this association?

- A. Cross sectional study
- ☒ B. Case control study
- C. Cohort study
- D. Experimental study

137. Considering the nature of the diseases, for which of the following health problems is screening LEAST recommended?

- A. Screening for HIV infection
- B. Screening pregnant women for syphilis
- ☒ C. Screening for relapsing fever
- D. Screening for cervical cancer

127. A 3 year-old boy presented with a complaint of intense pruritis worse at night. On examination there are vesicular and papular eruptions at the interdigital spaces. Which one is the best treatment for this child?

A. Acyclovir

B. Erythromycin

C. Mebendazole

☒ D. Permethrin

128. A 2-hour term neonate is delivered by elective C/S for placenta previa. On physical examination PR=170/min, RR=72/min and oxygen saturation of 80%.

What is the most likely cause of respiratory distress in this neonate?

A. Congenital pneumonia

B. Perinatal Asphyxia

C. Respiratory distress syndrome

☒ D. Transient tachypnea of the newborn

129. A 2-year old child presented with difficulty of breathing of 2 hours. Mother reported that he barking cough and hoarseness of voice in the last two days. On physical exam he is agitated, has audible stridor at rest and lower chest indrawing. What is the most likely diagnosis in this child?

☒ A. Croup syndrome

B. Foreign body aspiration

C. Pertussis

D. Severe pneumonia

130. A 7-year old female patient who is on follow-up for bronchial asthma presented with shortness of breath of three hours. On physical examination she is conscious, able to talk in complete sentence, RR= 46/min and PR=110/min. Chest examination reveals diffuse expiratory wheeze. What is the initial best step in the management of this patient?

A. Aminophylline

B. Magnesium sulphate

C. Oral prednisolone

☒ D. Salbutamol puff

131. A 3 year-old boy is brought to the ER with high grade fever and shortness of breath of two days. On physical examination RR= 62/min, T=39°C, lower chest indrawing with bilateral basal coarse crepitation. The chest X-ray showed left basal lung pneumatocele. Which of the following is the most likely etiology?

A. Haemophilus Influenzae

☒ B. Staphylococcus aureus

C. Streptococcus pneumonia

D. Streptococcus pyogenes

132. A 13-year old street boy presented to the ER with high grade fever of 5 days. He has epistaxis, fatigue and headache. On physical examination heart rate 120/min, respiratory rate

138. Suppose 'Test X' is used as a screening test for 'Disease Y'. The sensitivity and specificity of 'Test X' are 90% and 80% respectively. Which of the following is correct interpretation of these findings?
- A. 'Test X' can correctly identify 90% of individuals with 'Disease Y'
 - B. 'Test X' can correctly identify 90% of individuals without 'Disease Y'
 - C. 'Test X' can correctly identify 80% of individuals with 'Disease Y'
 - D. 20% of the individuals with 'Disease Y' will be missed by 'Test X'
139. Patients may be grouped based on different characteristics for reporting. Which of the following grouping represent ordinal data?
- A. Blood group
 - B. Ethnic group
 - C. Disease severity
 - D. Sex
140. A researcher is interested to determine whether there is malaria outbreak in a given zone of a region. Which method is the best to collect data for the intended purpose at community level?
- A. Conduct Sample Survey
 - B. Conduct Census
 - C. Review record of patients in all providers
 - D. Focus group discussions
141. Which of the following diagrammatic presentation turns cumulative frequency distribution in to graphs?
- A. Ogive curve
 - B. Frequency polygon
 - C. Stem and leaf plot
 - D. Histogram
142. The dietary assessment on patients from a refugee camp shows many of the patients subsist on a diet of corn meal and boiled greens. They have little appetite for food and alternate between diarrhea and constipation. They also have scaly dermatitis. What is the most probable dietary deficiency in these patients?
- A. Niacin
 - B. Pyridoxine
 - C. Vitamin E
 - D. Vitamin K
143. The serum level of 1, 25 di-hydroxy vitamin D in adolescent girls is believed to be normally distributed with mean 65 pg/ml and standard deviation 12.5 pg/ml. What percent of adolescent girls will have a level higher than 65 pg/ml?
- A. 10%
 - B. 25%
 - C. 50%
 - D. 100%



144. A medical director at a regional hospital wanted to assess prevalence of Malaria in the town in order to make sufficient preparation for a possible outbreak. The town is composed of urban and rural kebeles. Which sampling methods give better precision for estimating parameter of interest?

- A. Simple random sampling
- B. Cluster sampling
- C. Stratified sampling
- D. Quota sampling

145. A woman usually prepares food for her family from cereals and legumes. She usually uses wheat for making a number of food items. She hardly gets vegetables and fruits foods in the diet of the family. What dietary deficiency is this family likely to face?

- A. Cholecalciferol
- B. Folic acid
- C. Niacin
- D. Pyridoxine

146. A large number of under-five children were diagnosed with serious neurological problems and anemia in a certain Woreda. The families of these children were strictly vegetarian. What is the most likely nutritional deficiency of this community?

- A. Vitamin B12
- ☒ B. Folic Acid
- C. Niacin
- D. Pyridoxine

147. A four year-old child is brought to the Health Center with acute respiratory tract infection for the first visit. What dose of vitamin A will be given for this child?

- A. 50,000
- B. 100,000 IU
- C. 200,000 IU
- D. 400,000IU

148. A patient admitted to a Hospital ate a 57 gram serving of whole-wheat spaghetti containing 8 grams protein, 40 grams carbohydrate and 2 grams fat. What is the estimate of energy intake of this patient in Kcal?

- A. 200
- B. 210

C. 280

D. 400

149. A health officer aims to initiate a behavioral change among individuals by considering the socio-cultural factors after identifying the personal characteristics of the people. Which one is the best theory or model to apply?

- A. Health belief model
- B. Social cognitive theory
- C. Theory of planned behavior
- D. Trans-theoretical model

150. While giving health education to a group of mothers to introduce a new family planning method, which of the following learning materials are preferred?
- ☒ A. Model
 - B. Photograph
 - C. Poster
 - D. Real object
151. Which of the following sites is ideal place to display a message using information education communication materials to create awareness of a new family planning method in a rural village?
- A. Homes
 - ☒ B. Market places
 - C. Prisons
 - D. Religious institutions
152. Some community-based preventive interventions initiated by the government have failed to meet the desired objectives even though they have good plans. Which of the following is the main reason for the failure?
- ☒ A. Absence of appropriate technology
 - B. Higher cost of interventions
 - C. Lack of good policy in the country
 - D. Lack of planning with the community
153. A communal well water source development project was planned to be developed in a rural village. For the outcome of the project to be effective, what needs to be identified in the process?
- ☒ A. Acceptance of the water source by the community
 - B. Cost of materials for development purpose
 - C. Electrolyte composition of the water source
 - D. Level of trace minerals in the water source
154. A report from a health center showed most residents of that district were affected by water-related diseases. An assessment in the district revealed that most of the residents drink water from few sources of untreated wells by treating it with Water Agar. But the culture of keeping personal hygiene was poor. Which water related diseases would affect the residents of the district most?
- A. Water arthropod disease
 - ☒ B. Water borne infection
 - C. Water impounding disease
 - D. Water washed disease
155. Hazardous waste disposal was found to be a critical problem in a Health Center. A disposal options for hazardous wastes was required for a public health action. Which waste disposal option is the most appropriate for the health center?
- A. Burying
 - B. Composting
 - C. Incinerating
 - D. Offsite Disposal

162. Of the strategies to reduce maternal mortality, which of the following interventions is most effective in reducing the likelihood of developing pregnancy-related complications?
- ☒ A. Antenatal care
 - B. Emergency obstetric care
 - C. Family planning
 - D. Nutrition intervention
163. A manager of a Health Center wants to provide BCG vaccine for infants on Fridays. Which of the following is the best management function that the manager used?
- A. Organizing
 - B. Planning
 - ☒ C. Controlling
 - D. Implementing
164. A Hospital manager allocated a budget for three different activities. Which of the following is the next step in planning?
- A. Designing strategy
 - B. Prioritizing problem
 - C. Setting objective
 - D. Situational analysis
165. An EPI coordinator of a district and his subordinates were implementing the program for according to national standard of EPI coverage. Which one is the best method to assess their performance at the end of six months?
- A. Comparing performance against the standard
 - B. Implementing controlling process
 - C. Measuring actual performance
 - D. Standardization of performance
166. A zonal Health Department head assigned a budget for different tasks for a fiscal year. Which decisional role did he play?
- A. Negotiation role
 - B. Dissemination role
 - C. Resource allocation role
 - D. Monitoring role
167. The total population of certain locality was around 25, 000. Which of the following health facilities is suitable to provide health service to that community?
- A. District hospital
 - B. Health center
 - C. Health post
 - D. Health station
168. A college dean who has experts in different professions as employees left them to do their activities. What leadership style did he apply?
- A. Autocratic
 - B. Bureaucratic
 - C. Democratic
 - D. Laissez faire
169. There was interpersonal conflict in the same department of a certain health facility. What was the best management option that they do?
- A. Leave the issue for those involved
 - B. Manage the conflict after a month
 - C. Switch one of them to another department
 - D. Resolve the conflict as soon as possible

- 120 Sheets
9 1/2" x 7"
242mm x 178mm
SP03397
156. The malaria prevalence in a town is increasing from time to time. A team has been established to develop malaria epidemic control and prevention program. What priority intervention needs to be included in the program to effectively limit the progress of the epidemic?
- A. Application of outdoor residual spray
 - ☒ B. Destruction of mosquito breeding sites
 - C. Enforce usage of insect repellants
 - D. Timely treatment of cases
157. In a rural district, an ordinary millhouse is known for recruiting uneducated, economically poor individuals as permanent employees. The employees do not use any kind of Personal Protective Equipment. Which of the following is the most effective intervention to prevent noise-induced hearing loss?
- A. Early diagnosis and treatment
 - B. Enforce earmuff usage
 - C. Noise-proofing the mill
 - D. Reduce working hours
158. Reproductive health experts use maternal mortality as indicator of maternal health status although it has limitations. What is the main limitation of using this indicator?
- A. It is difficult to measure it at community level
 - B. It is not standard measure of deaths
 - C. It over-represents maternal health problems
 - D. It under-represents maternal health problems
159. The 2011 Ethiopian demographic and health survey reported that institutional delivery service utilization was 10% which was unacceptably low. What was the most important reason reported by women for not delivering in a health facility?
- A. High cost of delivery services
 - B. Lack of access to health services
 - C. Lack of awareness
 - D. Lack of transportation
160. If an organization wants to design a strategy targeting the most important direct cause of maternal mortality globally. Which cause should get the first priority?
- A. Abortion
 - ☒ B. Hemorrhage
 - C. Hypertension
 - D. Sepsis
161. A health officer wants to know risk of maternal death due to level of fertility and risk of maternal death per pregnancy or per birth. Which measure of maternal mortality should he/she use?
- A. Life time risk of maternal mortality
 - B. Maternal mortality rate
 - C. Maternal mortality ratio
 - D. Proportion maternal

- 120 Sheets
9 1/2" x 7"
242mm x 178mm
SP03597
170. A 24 year-old woman comes to your office for follow up on her depression. She has decided to try to get pregnant and wants to stop taking her fluoxetine. She had a history of suicidal attempt and severe peri-partum depression in the past. Which of the following is appropriate management?
- A. Continue to take fluoxetine
 - B. Shift to another antidepressant
 - C. Shift to antipsychotic drug
 - D. Start mood stabilizer
171. A 26 year-old woman brought by her sister to the Hospital with seven months history of unusual behavior. The patient gradually develops poor self-care, isolation and sleep problem. She hears the voices of several people talking to her. His sister says she talks to herself and laughs alone. On mental state examination she has blunt affect, auditory hallucination & has no insight to her illness. Which of the following is the most likely diagnosis?
- A. Delusional disorder
 - B. Schizoaffective disorder
 - ☒ C. Schizophrenia
 - D. Schizophreniform disorder
172. A 28 year-old man diagnosed as a case of Major depressive disorder with psychotic features. He is treated with haloperidol and amitriptyline. Two months later, his mood symptoms have resolved and he is no longer psychotic. Which of the following best describes the next step?
- A. Both drugs should be tapered and discontinued
 - B. Haloperidol should be tapered and discontinued
 - C. Amitriptyline should be tapered and discontinued
 - ☒ D. Both drugs should be continued for 6 months
173. A 53 year-old man comes to the Hospital complains of poor appetite, loss of pleasure in daily activities, sleeplessness, & decreased libido for 3 weeks. He is contemplating retiring, expressing concern that his poor performance in his job. On mental state examination he shows empty feeling & makes two errors in subtraction serial 3's from 20. Which of the following is the most likely diagnosis?
- A. Bipolar disorder
 - B. Cyclothymia
 - C. Dysthymic disorder
 - D. Major depressive disorder
174. A 35 year-old man diagnosed to have Major depressive disorder was given 75 mg amitriptyline at bed time. On the third day he starts hearing voices which tell him that "he is a bad person". On mental state examination he feels sad and has auditory hallucinations. Which of the following is the most appropriate treatment?
- A. Diazepam
 - B. Fluoxetine
 - ☒ C. Haloperidol
 - D. Sodium valporate

- 120 Sheets
9 1/2" x 7"
242mm x 178mm
SP03397
175. A 20 year-old male brought by his cousin to the Hospital. His cousin says that after smoking marijuana he became suspicious and started to hear voices. He talks alone and is easily irritable. These behavioral changes persisted for the last two days. On mental state examination he has thought & perceptual disturbance. He has no past psychiatric illness. Which of the following is the most likely diagnosis?
- A. Cannabis induced delirium
 - B. Cannabis induced psychosis
 - C. Cannabis intoxication
 - D. Cannabis withdrawal
176. A 55 year-old alcoholic is brought to the Emergency Room with a problem of irrelevant talking. Three days ago he quit alcohol after he was counseled. On mental state examination he is disoriented to time, place & person. He has visual hallucination and agitation. There is no history of head injury or medical illness. Which of the following is the most likely diagnosis?
- A. Alcohol induced psychosis
 - B. Alcohol intoxication
 - C. Delirium tremens
 - D. Wernicke's encephalopathy
177. A 30 year-old woman came to Medical OPD accompanied by her husband. She delivered a healthy baby three weeks ago at a Health center. Her husband reports that she has irritability, tearfulness, & sleeplessness for the last two weeks. The past week she has started to feel worthlessness and have loss of concentration. Which of the following is the most likely diagnosis?
- A. Dysthymia
 - B. Postpartum blues
 - C. Postpartum depression
 - D. Postpartum psychosis
178. A 55 year-old man comes to Emergency Room with acute retention of urine for 12 hours. He was suffering from depression and taking drugs for the last ten days. On examination there is distended bladder. No other physical findings. Which drug is the most likely culprit?
- A. Amitriptyline
 - B. Fluoxetine
 - C. Paroxetine
 - D. Sertraline
179. A patient comes to your clinic with a complaint of hearing difficulty of one month duration. There is only a set of tuning forks for hearing assessment at the Clinic. Which tuning fork should be used for the assessment?
- A. 256 Hz
 - B. 512 Hz
 - C. 1024 Hz
 - D. 2048 Hz
180. A patient comes demanding for ear wash. Which of the following would be a contraindication for ear wash?
- A. Beads
 - B. Chronic otitis media
 - C. Cockroach
 - D. Impacted wax

- 120 Sheets
9 1/2" x 17"
242mm x 432mm
SP03397
181. A 3 year-old female child presented to the Emergency OPD with foreign body in the right nasal cavity of three days duration. On physical examination, there was an alkaline small button battery inside with discharge and necrotic nasal mucosal surface. What is the best management plan?
- A. Give acidic nasal drops for two days and remove it
 - B. Give nasal decongestants for two days and remove it
 - C. Immediately removal
 - D. Irrigate with saline and give antibiotics
182. A 37 year- old female presented with a cockroach in the right ear canal. She is so much in discomfort because of the movement of the insect. On physical examination, the right ear has big dry central perforation of the tympanic membrane. What is the management?
- A. Add 3% hydrogen per oxide to kill the insect
 - B. Add ethyl alcohol to kill the insect
 - C. Irrigation with warm water
 - D. Take the insect out with a small artery while alive
183. A 22 year- old male patient has left sided ear pain with no discharge. He has no history of trauma and physical examination revealed tenderness over the tragus. What is the most likely diagnosis?
- A. Acute otitis externa
 - B. Acute otitis media
 - C. Cerumen impaction
 - D. Chronic otitis media
184. A 25 year-old man came to a Primary Hospital's OPD after a fertilizer was splashed into his eyes while working in his farm. He has redness on both eyes with photophobia. What is the best initial management?
- A. Take comprehensive history and ocular physical examination
 - B. Refer urgently to the nearest ophthalmologic center
 - C. Wash the eyes with normal saline
 - D. Give anti-inflammatory and antibiotic eye drops
185. Which of the following is a cause for constricted pupil?
- A. Atropine
 - B. Acute angle closure glaucoma
 - C. Oculomotor palsy
 - D. Organophosphate poisoning
186. During a school survey on refraction, a 6 year-old girl was found to have a refractive error of (+3) on the left eye while the right eye was (+6). If the refractive error is not corrected by spectacles, which of the following is LEAST expected to happen?
- A. Amblyopia
 - B. Floaters
 - C. Poor school performance
 - D. Strabismus
187. What is the leading cause of visual impairment globally?
- A. Cataract
 - B. Diabetic retinopathy
 - C. Glaucoma
 - D. Uncorrected refractive error

188. What is the most common cause of unilateral cataract in young adults?
- ☒ A. Metabolic/
 - B. Uveitis
 - C. Trauma
 - D. UV radiation
189. A 51 year-old man came to your OPD complaining he can't read contact numbers from his mobile phone since 6 months. He used to be able to read by holding the phone further away. He and no other ocular complaints. He had 6/6 distance vision on both eyes. What is the most likely diagnosis?
- A. Age-related macular degeneration
 - B. Astigmatism
 - C. Cataract
 - ☒ D. Presbyopia
190. A mother brought her 1 year old boy to the health center claiming the child has poor vision. Which of the following is the least likely to suggest congenital glaucoma?
- A. Blepharospasm
 - B. Epiphora
 - C. Increased corneal diameter /
 - D. Redness
191. A 55 year-old man came to medical OPD with a chief complaint of pain during mastication for the last one year. Intra-oral examination shows horizontal teeth mobility around 1mm, gingival recession with exposed root surfaces, intra-oral periapical x-ray shows alveolar bone resorption. Which of the following is the most likely diagnosis?
- A. Periodontitis
 - B. Gingivitis
 - C. Pulpitis
 - D. Stomatitis
192. A 30 year-old woman who had tooth extraction four days ago came with severe pain in the socket. On examination the socket is dry with no clot formation. The alveolar margin was tender. What is the best next step in management?
- A. Apply local paste
 - B. Do Debridement
 - C. Pack with gauze
 - D. Prescribe antibiotics
193. A 25 year-old women with gestational age of 11 weeks came to Medical OPD with recurrent toothache. On examination maxillary first molar hopelessly decayed and slightly tender to percussion. Which one of the following is the most appropriate management?
- A. Extraction
 - B. Pain control
 - C. PO tetracycline
 - D. Root canal therapy

194. Which one of the following is typical feature of bilateral anterior TMJ dislocation?
- Deviation of chin
 - Lateral cross bite
 - Posterior cross bite
 - Protrusion of chin
195. A 45 year-old man presented to the Emergency Department with a chief complaint of jaw swelling and difficulty of breathing. Physical examination showed non-pitting, firm, board like swelling over the primary mandibular facial spaces bilaterally with raised floor of mouth, dysphagia, tenderness on palpating the area and trismus of mouth. Which one of the following is most likely diagnosis?
- Ameloblastoma
 - Facial cellulitis
 - Ludwing's angina
 - Submandibular abscess
196. A 25 year-old woman presented with gum bleeding and bad breath odor for one month. Examination shows interdental papillae full of debris and multiple ulcers, easily bleeding gum, linear erythema and free gingival margin with full of ulcers. What is the most likely diagnosis?
- Acute necrotizing periodontitis
 - Acute necrotizing ulcerative gingivitis
 - Chronic gingivitis
 - Puberty induced gingivitis
197. A 25 year old women came with chief complaint of swelling related with lower right jaw of about one week duration. History and clinical examination confirms submandibular space abscess. Which of the following tooth is the most likely cause?
- Lower canine
 - Lower premolars
 - Lower 1st molar
 - Lower 2nd molar
198. A 35 year-old woman came to a hospital with a chief complaint of inability to close her eyes. Physical examination shows dropping of the corner of the mouth, leakage of saliva, inability to close the right eye. Deviation of mouth to the right side. Which one of the following is the most likely diagnosis?
- Trigeminal neuralgia
 - Facial nerve palsy
 - Glossopharyngeal neuralgia
 - Trigeminal nerve paralysis
199. A 30 year-old patient came with chief complaint of mobility of teeth. Which one of the following would suggest the diagnosis of periodontitis?
- False pocket
 - Gum growth
 - Gum recession
 - Short clinical crown
200. A 25 year-old man presented with fighting accident of 24 hours. Physical examination shows laceration of upper lip, fractured upper incisor. X-ray result shows apical half fracture of the offending tooth. Which one of the following is the best management option?
- Extraction
 - Inter-dental wiring
 - Root canal therapy
 - Splinting

1. A 26 year-old female presented with two months of excessive menses and gum bleeding. Physical examination revealed pale conjunctiva, petechiae and ecchymosis. Which of the following is the most likely diagnosis?

- A. Hemophilia ✗
- ☒ B. Idiopathic thrombocytopenic purpura ✓
- C. Scurvy ✗
- D. Vitamin K deficiency ✗

2. A 21 year-old woman presented with fatigue and light headedness. Physical examination showed marked conjunctival and palmar pallor. Which of the following physical findings is most likely to be seen in this patient?

- A. Apical diastolic murmur
- ☒ B. Ejection systolic murmur
- C. Narrow pulse pressure
- D. Rumbling mid diastolic murmur

Anemia - Ejection systolic murmur (for murmur)
Tachycardia

3. A health officer was evaluating an anemic patient. Laboratory test revealed: Hgb = 7 g/dl, MCV = 77 fl, WBC = 5400/ μ l, Platelet = 650,000/ μ l. Which of the following is the most likely diagnosis?

- A. Acute myelogenous leukemia
- B. Aplastic anemia
- ☒ C. Iron-deficiency anemia
- D. Megaloblastic anemia

Anemia of I. Defe. → microcytic
Hypochromic
Also: Eosinophilic vit. B12 def.

4. A 45 year-old woman presented with fatigue and orthostatic dizziness of two months duration. She has conjunctival pallor. Complete blood count revealed: Hgb = 8 g/dl, WBC = 4200/ μ l, and platelet = 170,000/ μ l. Which of the following co-morbidities is LEAST likely?

- A. Chronic kidney disease ✓
- ☒ B. Chronic obstructive lung disease
- C. Colorectal cancer ✓
- D. Disseminated tuberculosis ✓

Anemia
150 - 400,000

5. A 57 year-old man is admitted to the Medical Ward with hemorrhagic stroke. After a week the patient suddenly developed pleuritic chest pain and shortness of breath. Physical examination revealed signs of respiratory distress and a swelling involving his right calf and thigh. What is the most likely cause for his deterioration?

- A. Acute myocardial infarction
- B. Aspiration pneumonia
- C. Fat embolism
- ☒ D. Pulmonary embolism

DVT

1. A 26 year-old female presented with two months of excessive menses and gum bleeding. Physical examination revealed pale conjunctiva, petechiae and ecchymosis. Which of the following is the most likely diagnosis?

- A. Hemophilia ✗
- ☒ B. Idiopathic thrombocytopenic purpura ✓
- C. Scurvy ✗
- D. Vitamin K deficiency ✗

2. A 21 year-old woman presented with fatigue and light headedness. Physical examination showed marked conjunctival and palmar pallor. Which of the following physical findings is most likely to be seen in this patient?

- A. Apical diastolic murmur
- ☒ B. Ejection systolic murmur
- C. Narrow pulse pressure
- D. Rumbling mid diastolic murmur

Anemia - Ejection systolic murmur (for murmur)
Tachycardia

3. A health officer was evaluating an anemic patient. Laboratory test revealed: Hgb = 7 g/dl, MCV = 77 fl, WBC = 5400/ μ l, Platelet = 650,000/ μ l. Which of the following is the most likely diagnosis?

- A. Acute myelogenous leukemia
- B. Aplastic anemia
- ☒ C. Iron-deficiency anemia
- D. Megaloblastic anemia

Anemia of I.D. Def. → microcytic
Hypochromic
Also elevated vit. B12

4. A 45 year-old woman presented with fatigue and orthostatic dizziness of two months duration. She has conjunctival pallor. Complete blood count revealed: Hgb = 8 g/dl, WBC = 4200/ μ l, and platelet = 170,000/ μ l. Which of the following co-morbidities is LEAST likely?

- A. Chronic kidney disease ✓
- ☒ B. Chronic obstructive lung disease
- C. Colorectal cancer ✓
- D. Disseminated tuberculosis ✓

Anemia
150 - 400,000

5. A 57 year-old man is admitted to the Medical Ward with hemorrhagic stroke. After a week the patient suddenly developed pleuritic chest pain and shortness of breath. Physical examination revealed signs of respiratory distress and a swelling involving his right calf and thigh. What is the most likely cause for his deterioration?

- A. Acute myocardial infarction
- B. Aspiration pneumonia
- C. Fat embolism
- ☒ D. Pulmonary embolism

DVT

6. A 26 year-old man with rheumatic heart disease came for follow-up. He has active precordium and rumbling mid-diastolic murmur. What valvular lesion is most likely in this patient? *MS*

- A. Aortic regurgitation
- B. Mitral regurgitation
- ☒ C. Mitral stenosis
- D. Tricuspid regurgitation

7. A 65 year-old male with type 2 diabetes mellitus presents to the Emergency Department with a squeezing, retrosternal chest pain of one hour duration. He has vague discomfort in his right arm and shoulder. The physical examination is notable for diaphoresis and mild tachycardia. Which of the following is the most likely diagnosis?

- ☒ A. Acute coronary syndrome
- B. Acute pericarditis
- C. Aortic dissection
- D. Pulmonary embolism

8. Which of the following is a major criteria for the diagnosis of rheumatic fever?

- A. Elevated ESR
- B. Fever
- C. Prolonged PR segment
- ☒ D. Polyarthralgia

9. A 29 year-old woman came to the OPD with shortness of breath and body swelling of three weeks. On physical examination she has raised JVP, accentuated P2 and grade IV diastolic murmur at the apex. Which of the following drug delays disease progression?

- A. Frusemide
- B. Penicillin
- ☒ C. Enalapril
- D. Digoxin

10. A 45 year-old male with heart failure is comfortable at rest. He develops dyspnea and fatigue to walk to a nearby shop and is forced to rest for five minutes before symptoms improved. What is the stage and functional status of this patient?

- A. Stage A, Class II
- B. Stage B, Class III
- ☒ C. Stage C, Class IV
- D. Stage C, Class III

11. A 50 year-old man was found to have a BP=170/100. He has a history of asthma. Which one is the most appropriate drug of choice?

- A. Atenolol
- B. Enalapril
- ☒ C. Hydralazine
- D. Furosemide

12. A 40 year-old man presented with fever, shaking chills of three days duration. He also has non-productive cough and right-sided pleuritic chest pain. He has no significant past medical history. Physical examination revealed crepitation on the posterior right lower chest. Which one is the best initial investigation to confirm the diagnosis?
- ☒ A. Chest X-ray
 - B. Culture of sputum
 - C. Gram stain of sputum
 - D. White blood cell count
13. A 25 year-old woman who was diagnosed with pulmonary tuberculosis and started anti-tuberculosis treatment two weeks ago is found to be HIV positive on screening. On physical examination, she has pale conjunctiva and laboratory investigation shows CD4 count of 100cells/ μ L. Which one is the most appropriate first line antiretroviral regimen for this patient?
- A. ABC/3TC/NVP
 - ☒ B. TDF/3TC/EFV *1st (regimen)*
 - C. AZT/3TC/EFV *2nd (regimen)*
 - D. D4T/3TC/NVP
14. A 24 year-old man has generalized lymphadenopathy, aphthous ulcers, and gray-white plaques around his scrotum. Dark field microscopic examination demonstrates spirochetes. Which of the following is the treatment of choice?
- A. Ceftriaxone
 - B. Erythromycin
 - ☒ C. Penicillin
 - D. Tetracycline
15. A 30 year-old prisoner presents with intermittent fever and headache of two days duration. He claims that three of his roommates had similar symptoms and were treated in the prison. Blood film examination showed a spirochete. Which one of the following is the most appropriate antibiotic choice for this patient?
- A. Ceftriaxone
 - B. Ciprofloxacin
 - ☒ C. Procaine penicillin
 - D. Vancomycin
16. A 20 year-old male patient presents with abdominal swelling of three weeks duration. She lives in a rural village where the water source used is river water. Ultrasound examination reveals peri-portal fibrosis. Which one of the following is most likely cause for the patient's condition?
- A. River blindness
 - B. Ascariasis
 - C. Leishmaniasis
 - ☒ D. Schistosomiasis
17. A 22 year-old HIV positive patient who presented with shortness of breath was suspected to have Pneumocystis jirovecii pneumonia but he has no signs of respiratory distress and his CD4 count is 150/ μ L. Which of the following drugs is recommended for this patient?
- ☒ A. Co-trimoxazole
 - B. Fluconazole
 - C. Prednisolone
 - D. Pyrimethamine

23. A 30 year-old man presented with epistaxis, easy fatigability and intermittent fever for one month. Physical examination revealed hepatosplenomegaly. Which of the following is the best investigation to confirm the diagnosis?

- A. Blood culture
- ☒ B. Bone marrow aspiration
- C. Stool examination
- D. Ultrasound of abdomen

24. A 60 year-old man presented with sudden onset of right sided weakness on both upper and lower extremities before a day. The weakness is maximal at onset and during sleep. He is hypertensive but does not take medications. What is the most likely diagnosis?

- A. Hemorrhagic stroke
- ☒ B. Ischemic stroke
- C. Subarachnoid hemorrhage
- D. Transient ischemic attack

25. Which of the following is the most common cause of hemorrhagic stroke in Ethiopia?

- A. Aneurysm
- B. Anticoagulation
- C. Brain tumor
- ☒ D. Hypertension

26. A 24 year-old HIV positive woman presents with fever and skin rash over the left chest two days ago. The rash is vesicular and follows dermatomes (T4-5). She is confused and meningeal signs are positive. Which of the following is the most likely diagnosis?

- A. Fungal meningitis
- B. Pyogenic meningitis
- C. Tuberculous meningitis
- ☒ D. Viral Meningoencephalitis - *was Herpes Zoster*

27. A 30 year-old male presented with headache, fever and progressive weakness on the left side of his body two weeks ago. He has right ear discharge since two months. On physical examination BP=140/80 mmHg, PR=102/min, T=39°C. He is mentally alert and power on the left upper and lower extremities is 3/5. What is the most likely diagnosis?

- ☒ A. Brain abscess
- B. Brain tumor
- C. Ischemic stroke
- D. Subdural hematoma

18. A 60 year-old male had cough for five years with clear sputum production and shortness of breath. He has smoked a pack of cigarettes per day for 20 years. Laboratory examination reveals haemoglobin of 19 g/dl. X-ray of the chest shows hyperinflation without infiltrates. Which of the following is the most likely diagnosis?

- A. Bronchial asthma
- ☒ B. Chronic obstructive pulmonary disease
- C. Interstitial lung disease
- D. Pulmonary Tuberculosis

19. A 40 year-old alcoholic develops productive cough and fever. Chest x-ray shows an air-fluid level in the superior segment of the right lower lobe. Which of the following is the most appropriate pharmacotherapy?

- A. Ceftazidime and ciprofloxacin
- B. Ceftriaxone and gentamycin
- ☒ C. Metronidazole and crystalline penicillin (COPD + typhoid)
- D. Vancomycin and chloramphenicol

20. A 25 year-old man presents with abdominal pain, headache and fever of five days duration. The Health officer suspected the diagnosis of typhoid fever. Which of the following investigations confirms the diagnosis?

- A. Stool examination
- B. Blood culture
- C. Weil-Felix titer
- ☒ D. Widal test (typhoid)

21. A 27 year-old man complains of fever, rigors, productive cough with rusty-coloured sputum and pleuritic chest pain. Chest examination revealed increased tactile fremitus and dullness to percussion in the right lower lung field. What is the most likely diagnosis?

- ☒ A. Lobar pneumonia (S - 100)
- B. Bronchopneumonia
- C. Lung abscess
- D. Pleural effusion

22. A 35 year-old woman presented with trismus for a day. She also has shoulder and back pain. She reported that she sustained a puncture over the left foot a week ago. She is fully conscious. Which of the following is the most appropriate pharmacotherapy?

- A. Carbamazepine
- ☒ B. Diazepam
- C. Phenobarbital
- D. Phenytoin

28. A 60 year-old alcoholic patient presented with fever, headache and neck pain of two days duration. On examination body $T=39^{\circ}\text{C}$ and has neck stiffness. Bacterial meningitis was suspected but lumbar puncture was deferred because he has papilledema. Which of the following is the best empiric therapy for this patient in addition to Ampicillin?
- A. Vancomycin and Chloramphenicol
 - ☒ B. Ceftriaxone and Vancomycin
 - C. Ceftriaxone and Chloramphenicol
 - D. Cloxacillin and Ciprofloxacin
29. A 36-year-old male presented with fatigue and tea-colored urine for five days. Physical examination revealed jaundice and tender Hepatomegaly. Laboratory investigation revealed: Aspartate aminotransferase (AST) = 2400 U/L, Alanine aminotransferase (ALT) = 2500 U/L, Alkaline phosphatase = 210 U/L and total bilirubin = 8.6 mg/dL. Which of the following is the LEAST likely cause for the patient's condition?
- A. Acetaminophen poisoning
 - B. Acute Viral hepatitis
 - ☒ C. Alcoholic liver disease
 - D. Ischemic liver injury
30. A 56 year-old man presented with progressive abdominal distension and leg swelling of two months duration. He is a known patient with cirrhosis for the past one year on follow up but not on medication. On examination, he was not in distress and has moderate ascites and bilateral leg swelling. Which of the following is the best initial management?
- A. Furosemide
 - ☒ B. Furosemide and Spironolactone *(K⁺ saving)* 5:1
 - C. Hydrochlorothiazide
 - D. Paracetamol and Furosemide
31. Which of the following is the most common complication of Peptic ulcer disease?
- ☒ A. Hemorrhage
 - B. Gastric outlet obstruction
 - C. Malignancy
 - D. Perforation
32. A 50 year-old woman presented with fever, cough and chest pain for two days. Chest X-ray showed pleural effusion. Pleural fluid analysis showed: WBC = 5,000/mm³, with 90% neutrophils, PH = 7.2, blood culture was positive for streptococcus pneumonia. What is the next best step in management of this patient besides antibiotics?
- A. Dexamethasone
 - B. Chest CT scan
 - ☒ C. Chest tube insertion
 - D. Lung function test

33. A 46 year-old man is brought to the Emergency Department because he was vomiting fresh blood. He reports that he has no abdominal pain or nausea. On physical examination; BP=70/40 mmHg, PR=118/min, RR= 24/min. He also has gynecomastia, dilated abdominal veins and ascites. Which of the following is the most likely cause of the bleeding?

- A. Esophageal cancer
- ☒ B. Esophageal varices

- C. Mallory Weiss
- D. Reflux esophagitis

34. A 47 year-old man comes to a Hospital complaining of epigastric pain that is exacerbated by eating. Antigenic test for *Helicobacter pylori* is positive. Endoscopic examination reveals a duodenal ulcer. Which of the following is the most appropriate therapy?

- ☒ A. Amoxicillin, clarithromycin and omeprazole
- B. Amoxicillin, metronidazole and anti-acid
- C. Omeprazole and anti-acid
- D. Ranitidine, amoxicillin and metronidazole

35. A 29 year-old man with HIV infection presents with chronic diarrhea associated with anorexia, abdominal cramps and significant weight loss. His recent CD4 count =20 cells/mm³. A modified acid-fast stain of a stool specimen shows 4-6 μ m oocysts. Which of the following is the most likely cause?

- ☒ A. *Cryptosporidium parvum*
- B. Microsporidia

- C. *Mycobacterium avium* complex
- D. *Pneumocystis jirovecii*

36. Which of the following is the best early indicator for diabetic nephropathy?

- A. Rising blood urea nitrogen
- ☒ B. Albuminuria
- C. Hypertension
- D. Rising creatinine level

37. A 35 year-old man presents with generalized body swelling of two months. Physical examination reveals BP=120/70mmHg, ascites and +3 pretibial edema. Protein = 4.5g/24hour urine collection. Which of the following is the most likely diagnosis?

- A. Chronic liver disease
- B. Heart failure
- ☒ C. Nephrotic syndrome
- D. Glomerulonephritis

38. A 55 year-old man comes to the Emergency OPD because of excruciating pain in his right big toe. He was diagnosed to have gouty arthritis. Which of the following is the most appropriate pharmacotherapy?

- A. Allopurinol
- B. Ceftriaxone

- ☒ C. Indomethacin
- D. Morphine

730
830
2930
1030
1130

39. A 28 year-old lactating woman presented to the emergency OPD with a complaint of right breast pain of five days duration. On examination body T= 38°C with right breast tenderness but no fluctuation. What is the most likely diagnosis?
- ☒ A. Breast Abscess
 - B. Breast Cancer
 - C. Fibroadenoma
 - D. Fibrocystic Change
40. A 32 year-old man presented to the surgical OPD with complaint of neck mass of one year duration. On examination he has a hoarse voice and a solitary 2cm x 2cm thyroid mass on the right side which has firm consistency and fixed to underlying structures. There was no lymphadenopathy. What is the most likely diagnosis?
- A. Thyroid adenoma
 - ☒ B. Thyroid cancer
 - C. Thyroiditis
 - D. Toxic Nodule
41. A 35 year-old man with a chronic peptic ulcer disease presented with features of acute abdomen and chest x-ray revealed pneumoperitoneum. His symptoms lasted for six hours and his vital signs were stable after infusion of one liter of Normal Saline solution. What should be the next step in the management of this case?
- A. Barium meal study
 - B. Endoscopy
 - C. Triple therapy
 - ☒ D. Laparotomy
42. A 50 year-old woman brought an ultrasound imaging result which showed Gall Bladder Stone. She has no abdominal pain and no known medical illness. Which of the following is the recommended management?
- A. Cholecystectomy
 - B. Extracorporeal Shockwave Lithotripsy (ESWL)
 - ☒ C. Reassurance
 - D. Ursodeoxycholic acid
43. A 68 year-old man is admitted to a Hospital with complaint of intermittent lower abdominal pain, bowel habit change and rectal bleeding of one month duration. He has tenesmus and incomplete sense of defecation and had lost weight. What is the most likely diagnosis?
- A. Amebic dysentery
 - ☒ B. Colorectal cancer
 - C. Sigmoid Volvulus
 - D. Ulcerative colitis
44. A 55 year-old man presented to the surgical OPD with a complaint of rectal bleeding of one month duration. He has no protruding mass per rectum and digital rectal examination was non-revealing. What is the investigation of choice?
- A. Abdominal CT-Scan
 - B. Barium Enema
 - ☒ C. Colonoscopy
 - D. Upper GI Endoscopy

45. A 30 year-old man was seen at the Emergency surgical OPD with perianal pain of two days duration. He has fever and chills. He has left perianal induration with tenderness but no fluctuation. What is the most likely diagnosis?

- A. Anal fissure
- ☒ B. Perianal Abscess
- C. Perianal fistula
- D. Thrombosed Hemorrhoids

46. A 20 year-old man was seen at the emergency department following road traffic accident of one hour duration. He has sustained trauma to the abdomen. On examination; Vital signs: BP=80/50 mmHg and PR=120/min. He has tire marks on the abdomen and it was tender. The patient is started on IV fluids. What is the next step in the management?

- A. Abdominal CT-scan
- ☒ B. Exploratory laparotomy
- C. Observation
- D. Plain abdominal x-ray

47. A 50 year- old man came complaining of fatigue and dyspepsia of three months duration. Physical examination revealed a pale conjunctiva with a palpable mass at the epigastrium that moves with respiration. What is the gold standard diagnostic modality?

- A. Abdominal Ultra sound
- B. CT scan of the abdomen
- ☒ C. Upper GI Endoscopy
- D. Plain abdominal film

48. A man was stabbed to the abdomen and presented to the Health Center with eviscerated small bowel which was getting dark. Which of the following is the correct initial measure before referral?

- ☒ A. Cover the intestine with saline gauze
- B. Refer to a nearby hospital without intervention
- C. Replace the intestine back into the peritoneal Cavity
- ☒ D. Widen the stab wound under local anesthesia

49. A 23 year-old farmer presented to the OPD with abdominal pain and frequent vomiting of one day duration. On examination the abdomen is slightly distended and has visible peristalsis. Plain abdominal x-ray showed step ladder pattern air-fluid level. Which of the following is the most likely diagnosis?

- A. Caecal Volvulus
- B. Compound Volvulus
- C. Sigmoid Volvulus
- ☒ D. Small bowel Volvulus

50. A 65 year-old man was seen at the Emergency OPD with abdominal pain, distension and failure to pass feces and flatus of two days duration. He had two similar attacks in the last three years and was successfully treated with rectal tube deflation. On Examination BP=90/60 mmHg and PR=110/min. The abdomen was tender and bowel sounds were absent. Which of the following measures is to be avoided?

- A. Analgesics
- B. Antibiotics
- C. IV fluid
- ☒ D. Rectal tube deflation

51. A 45 year-old housewife presented with progressive submandibular swelling of two days duration. She visited a dentist and undergone tooth extraction a day prior to the start of her symptoms. Examination showed tense, shiny bilateral submandibular swelling with no area of fluctuation. She also has drooling of the saliva and protruded tongue. What is the diagnosis of this patient?
- ☒ A. Ludwig's angina
☐ B. Periodontal abscess
☐ C. Submandibular cellulites
☐ D. Submental abscess
52. A 20 year-old man presented to the OPD forty minutes after he sustained stab injury to his left posterior chest. He has shortness of breathing. He is acutely sick looking; BP= 110/60 mmHg, PR= 90/min, RR=26/min. He has decreased air entry on the lower left chest with visible air rush through the stab wound. What is the best first step in the management of this patient?
- ☐ A. Apply airtight dressing
☒ B. Apply triangular dressing
☐ C. Suture the wound
☐ D. Take chest X-ray
53. A 35 year-old male patient came to emergency OPD 12 hours after he started to have sudden onset of epigastric pain. He has long standing history of dyspepsia for which he has been taking anti-acids repeatedly. Examination revealed tachycardia, rigid and tender abdomen. What is the appropriate imaging study to reach a definitive diagnosis?
- ☐ A. Barium follow through
☐ B. Barium meal
☒ C. Chest X-ray
☒ D. Plain abdominal X-ray
54. A 27 year-old patient was involved in a road traffic accident. He was brought to the Hospital by the driver. He is agitated; PR=140/min and feeble, BP=80/40 mmHg. He has pale conjunctiva and his chest is clear with good air entry. He has an echymosis over his left flank area. Abdominal Ultrasonography showed a significant collection of fluid in his abdomen. What is the most likely diagnosis?
- ☐ A. Cardiogenic shock
☒ B. Hemorrhagic shock
☐ C. Neurogenic shock
☐ D. Septic shock
55. A 64 year-old male patient presented to the Emergency OPD after he failed to pass urine for a day. He had difficulty of urination for the last one year. The previous night he took seven bottles of beer with friends. On examination he is acutely sick looking and in pain. Bladder is palpable 10 cm above the pubic bone and tender. What is the most likely cause for acute urinary retention?
- ☒ A. Benign Prostatic Hyperplasia
☐ B. Bladder cancer
☐ C. Bladder stone
☐ D. Urethral Stricture

56. An 18 years old male presented with right scrotal swelling of 1 day duration. He also has associated pain and it started while he was asleep, it woke him up. He also has fever and pain during urination. On examination right testis is tender, hot to touch, swollen, and the testicular cord is not thickened. What is the most likely diagnosis?
- A. Testicular torsion
 - ☒ B. Epididymo-orchitis
 - C. Infected hydrocele
 - D. Testicular tumor
57. A 36 year-old male patient came with difficulty of urination for the last two years. He had previous history of whitish urethral discharge for which he was treated with tablets and injections. What is the most likely diagnosis?
- A. Benign prostatic hyperplasia
 - B. Prostatic Cancer
 - C. Sexually transmitted disease
 - ☒ D. Urethral Stricture
58. A 43 year-old male patient came to the Emergency OPD after being involved in a road traffic accident before a day. He has failed to urinate since the incident and is unable to move his right lower limb. On examination his vital signs are normal, there is right pubic tenderness and blood on the penile meatus. What should be the next step in management?
- A. Decompress with percutaneous needle puncture
 - B. Do image-guided catheterization
 - C. Insert a foley catheter to decompress the bladder
 - ☒ D. Perform emergency suprapubic cystostomy
59. A 24 year-old man came to the Emergency OPD with severe pain originating in the flank, radiating inferiorly and anteriorly. He also has nausea and vomiting. His vital signs are within normal limits. Abdominal examination showed marked costovertebral angle tenderness. Urinalysis showed microscopic hematuria. What is the most likely diagnosis?
- A. Cystitis
 - B. Pyelonephritis
 - C. Pyonephrosis
 - ☒ D. Ureteric colic
60. A 60 year-old female patient presented with right sided flank pain and hematuria of one month duration. She also has significant weight loss, night sweats and malaise. On physical examination, she is chronically sick looking with normal vital signs, a bimanually palpable mass on the right flank. What is the most likely diagnosis?
- A. Chronic pyelonephritis
 - B. Neuroblastoma
 - ☒ C. Renal cell cancer
 - D. Wilm's tumor

77. A 38 year-old primigravida comes for a routine visit at 39 weeks of gestation. Her blood pressure is persistently 140/90 mm Hg, and her urine protein is +2. Physical examination is otherwise unremarkable, and she is completely asymptomatic. Her cervix is 2 cm dilated and 90% effaced, vertex presentation and at station 0.

Which of the following is the most appropriate management option for this case?

- ☒ A. Admission for observation
- B. Immediate cesarean section
- C. Outpatient follow-up till 40 weeks
- D. Start induction of labor

78. A 34 year-old primigravida comes with cessation of fetal movement of four days duration at gestational age of 35 weeks from LNMP. She is otherwise asymptomatic and has no medical problem. Examination shows stable vital signs, symphysio-fundal height of 34cms, no uterine contraction and absent FHB.

Which of the following is the initial best step in the management of this patient?

- A. Consider Termination
- B. Do coagulation profile
- C. Prophylactic antibiotics
- ☒ D. Ultrasound for confirmation

79. A 63 year-old woman who came for regular gynecologic check-up was incidentally diagnosed to have cystocele. Which of the following is the most appropriate management for this patient?

- A. Anterior colporaphy
- B. Kegel's exercise
- C. Pessaries
- D. Vaginal hysterectomy

80. A 21 year-old GII PI mother at 24 weeks gestation complains of sever vulvar itching and cotton cheese like discharge. A wet mount slide show no clue cells but hyphae. Which of the following is the most likely diagnosis?

- A. Bacterial vaginosis
- ☒ B. Candida albicans
- ☒ C. Chlamydia trachomatis
- D. Trichomonas vaginalis

81. A 29 year-old woman presents with vaginal pruritus associated with white curd like discharge. On examination there is erythematous macular lesion but no vesicles. Investigation shows no clue cell, negative whiff test. Which of the following is the treatment of choice for this case?

- A. Azithromycin
- ☒ B. Fluconazole
- C. Metronidazole
- D. Oral acyclovir

82. A 26 year-old GIIPI woman at 26 weeks of gestation from her LNMP, comes for ANC follow up. She hasn't appreciated fetal movement yet. She had one episode of bleeding a month back and noticed failure of increase in uterine size. Her uterine size measures 20 weeks and you are unable to hear fetal heart tone.

Which of the following is the most descriptive diagnosis?

- A. Incomplete abortion
- B. Inevitable abortion
- ☒ C. Missed abortion
- D. Threatened abortion

90. A PIII mother who gave birth to an alive healthy term neonate vaginally before an hour was found to be restless and pale. On physical examination BP= 60/40 mmHg, PR=120/min, RR=24/min, T=36°C; uterus 22 week and soft and active vaginal bleeding. Which of the following is the most appropriate first step in the management of this patient?
- A. Administer crystalloids
 - B. Administer intra-nasal oxygen
 - C. Blood transfusion
 - ☒ D. Call for help
91. In the active management of third stage of labor, placenta failed to be delivered by controlled cord traction. Which of the following is the next best step in management?
- A. Fundal pressure
 - ☒ B. Manual removal of placenta
 - C. Pitocin IV drip
 - D. Wait for spontaneous expulsion
92. A 15 year-old girl came to Gynecology OPD with a complaint of abdominal pain of six months duration. The pain is crampy and recurrent. She had never seen menses before. On examination there is a central 14 week-sized abdomino-pelvic mass. Which of the following is the most likely diagnosis?
- A. Dermoid cyst
 - B. Full bladder
 - ☒ C. Imperforate hymen
 - D. Myoma
93. A 32 year-old nulligravida woman came to gynecology OPD with a complaint of heavy menstrual flow of one year duration. She also gave history of constipation and pelvic heaviness and failure to conceive of four years duration. On examination there is central, 12 weeks size firm abdomino-pelvic mass. Which one is the most likely diagnosis?
- A. Endometrial hyperplasia
 - B. Ovarian tumor
 - ☒ C. Uterine myoma
 - D. Uterine sarcoma
94. A 35 year-old GVIIV mother at gestational age of 8 weeks presented to gynecology OPD with a complaint of vaginal bleeding of four days duration. She had excessive vomiting her uterus is big-for-date. Which of the following is the most likely diagnosis?
- A. Cervical polyp
 - B. Ectopic pregnancy
 - C. Incomplete abortion
 - ☒ D. Molar pregnancy
95. Which of the following contraceptive methods is the most effective?
- A. Combined oral contraceptive pills
 - B. Female condom
 - ☒ C. Intrauterine contraceptive device
 - D. Medroxy progesterone acetate

96. A 17 year-old student came to a Clinic seeking pregnancy termination service at gestational age of 10 weeks. She gave a history of sexual assault. Which of the following is the best method of termination of pregnancy for this case?
- ☒ A. Dilatation and curettage
 - ☐ B. Intramuscular methotrexate
 - ☐ C. Manual vacuum aspiration
 - ☐ D. Mifepristone and misoprostol
97. A 24 year-old primigravida who came for a routine ANC visit tested positive for HIV. When should a health officer initiate ARV drugs for this woman?
- ☒ A. As soon as possible
 - ☐ B. At 20 weeks of gestation
 - ☐ C. At 28 weeks of gestation
 - ☐ D. At 36 weeks of gestation
98. A primigravida came to clinic with a complaint of lower abdominal pain of one day duration at GA of 22 weeks. She has previous history of excessive menstrual flow and uterine tumor which was not treated. On examination Vital signs-within normal limits. Abdomen localized tenderness, uterus 24 weeks size, No contraction FHR- 150bpm. Which of the following is the most likely diagnosis in this case?
- ☐ A. Abdominal pregnancy
 - ☐ B. Acute appendicitis
 - ☒ C. Degenerative myoma
 - ☐ D. Ovarian cyst torsion
99. A Para III mother who gave birth to an alive male neonate at home before one hour brought to a Hospital because of restlessness and protrusion of mass per vaginum. On examination she is acutely sick looking, BP is not recordable, PR=132/minute and feeble RR=20/min, T=36°C. Abdomen-uterus not palpable, globular fleshy mass with minimal bleeding protruding through the vaginum. Which of the following is the most appropriate initial management of this case?
- ☐ A. Intra nasal oxygen 6 L/minute
 - ☐ B. Reposition the uterus
 - ☒ C. Resuscitation with IV fluids
 - ☐ D. Start oxytocin drip
100. A 20 year-old primigravida at 38 weeks was admitted in second stage of labor and stayed for 90 minutes. Fetal heart beat is ranging between 80 – 90/min, vertex presentation with the occiput direct anterior, station +3, grade-I molding, no caput and grade-II meconium stained fluid. Which of the following is the next best step in management?
- ☒ A. Expedite delivery right away by vacuum
 - ☐ B. Refer her for cesarean section
 - ☐ C. Re-position her into the left lateral position
 - ☐ D. Secure IV line and await spontaneous vaginal delivery
101. A 6-year old diabetic patient is admitted to ER with the diagnosis of sever DKA and has received normal saline and insulin for the last 3 hours. Which of the following electrolyte abnormalities is expected?
- ☒ A. Hypocalcemia
 - ☐ B. Hypokalemia
 - ☐ C. Hyponatremia
 - ☐ D. Hypophosphatemia



61. A 56 year-old man presented with a scrotal mass of three months duration. Physical examination revealed a 3cm x 4 cm firm mass on the lower pole of the right testis. Scrotal ultrasound showed a well-defined hypoechoic lesion within the testis. Serum β -hCG was elevated. What is the most likely diagnosis?
- A. Orchitis
 - B. Hydrocele
 - ☒ C. Testicular Tumor
 - D. Testicular torsion
62. A 35 year-old male patient presented to the Emergency Room after sustaining a bullet injury to his left forearm three hours back. Up on evaluation vital signs were normal and a deformed and rotated forearm, intact distal neurovascular examination. X-ray showed comminuted displaced radio-ulnar fracture. Which of the following is a temporary means of immobilization?
- A. Apply POP cast
 - ☒ B. External fixation
 - C. Lateral arm traction
 - D. Plating and screw
63. A 65 year-old male patient sustained a fall down accident three hours prior to presenting to the Emergency Room. He fell down with the arms extended. On examination vital signs were normal with a right wrist deformity. X-ray showed displaced Colle's fracture. Which one is the best management option?
- A. Application of a POP Cast
 - ☒ B. Closed reduction and casting
 - C. Open reduction and plating
 - D. Skin traction for few days then casting
64. A 20 year-old male patient had an external fixator applied for a compound comminuted right tibio-fibular fracture three weeks prior to presentation. He now complains of pain at the pin site. On examination there is seropurulent discharge from the two distal pins. Which of the following is the best diagnosis?
- A. Cellulitis
 - B. Osteomyelitis
 - C. Osteonecrosis
 - ☒ D. Pin tract infection
65. A 20 year-old male patient came to you with right knee pain of five days duration. On examination BP=120/70 mmHg, PR= 84/min. The knee is swollen and tender with signs of knee effusion. X-ray showed a wide joint space. Joint fluid analysis revealed turbid fluid with $100,000 \text{ WBC/mm}^3$. Which of the following is the most likely diagnosis?
- A. Acute osteomyelitis
 - B. Osteoarthritis
 - ☒ C. Pyogenic arthritis
 - D. Tuberculous arthritis

102. Which of the following electrolytes is found in high concentrations in ReSOMal when compared with standard WHO oral rehydration solution?
- A. Chloride
 - B. Phosphorous
 - ☒ C. Potassium
 - D. Sodium
103. An 11-month old male infant is brought to the ER with watery diarrhea of 2 days duration. Physical examination reveals irritable infant, weight=10 kg, sunken eye balls and eagerness to drink. What is the correct type and volume of fluid for this infant?
- A. Ringer's lactate, 300ml
 - B. 0.45% Normal saline, 700ml
 - ☒ C. Oral rehydration solution, 750ml
 - D. Ringer's lactate 500ml then ORS 500ml
104. An 8-year old boy is diagnosed with acute rheumatic fever three weeks after an episode of sore throat. Which drug should have been given for the sore throat to prevent the rheumatic fever?
- A. Clindamycin within 11 days
 - B. Cloxacillin within 7 days
 - C. Erythromycin within 16 days
 - ☒ D. Penicillin within 9 days
105. An 8-month old infant is brought to ER with fever, cough, and fast breathing of three days duration. Further inquiry reveals history of repeated chest infection, interruption during breast feeding and poor weight gain. Precordial examination reveals active precordium and pansystolic murmur at left lower sternal border. Which of the following heart lesions most likely explains these findings?
- A. Atrial septal defect
 - B. Tetralogy of Fallot
 - C. Tricuspid atresia
 - ☒ D. Ventricular septal defect
106. A 9-week old girl who was apparently healthy two days ago is brought to the ER. She is pale and has dyspnea with RR=80/min, T=37.7°C, PR=195/min, heart sounds are distant, a gallop is heard, liver is 6 cm below right costal margin. Chest X-ray shows cardiomegaly and echocardiogram reveals poor ventricular function and dilated left heart chambers. Which of the following is the most likely diagnosis based on this clinical scenario?
- ☒ A. Dilated cardiomyopathy
 - B. Endocarditis
 - C. Myocarditis
 - D. Pericarditis
107. An 8-year old boy known cardiac patient on follow up has a 3-week history of low-grade fever of unknown source, fatigue, weight loss, myalgia, and headaches. On repeated examinations during this time, he is found to have developed a new heart murmur, petechiae, and mild splenomegaly. Which of the following is the most likely diagnosis?
- A. Acute rheumatic fever
 - B. Rheumatic recurrence
 - ☒ C. Infective endocarditis
 - D. Dilated cardiomyopathy

83. A 42 year-old healthy nulliparous woman has a firm, irregular shaped, 12 week-sized midline abdominal mass.

Which of the following is the best initial step in evaluating this patient?

- A. Abdominal x-ray
- B. CT scan
- C. MRI
- ☒ D. Ultrasound

84. Which of the following is NOT a component of active third stage management?

- A. Controlled cord traction
- ☒ B. Nipple stimulation
- C. Uterine massage
- D. Oxytocin provision

85. A GIII PII mother came to a clinic at gestational age of 16 weeks for ANC follow up.

Which of the following is LEAST likely to be requested as part of a routine work up?

- A. Blood group and Rh
- ☒ B. Ultrasound
- C. Urinalysis
- D. VDRL

86. A primigravida was admitted to a Health Center due to pushing down pain of 6 hours. Physical examination; vital signs were normal, uterine size 36 weeks, longitudinal, cephalic, FHR=146/min, two contractions in 10min each lasting for 30-45 seconds; cervix is 3 cm dilated and 60 % effaced. After 4 hours she was reevaluated and had same findings except the cervical effacement changed to 75 %.

Which of the following is the best management option?

- A. Augmentation
- B. Rehydration
- C. Referral to hospital
- ☒ D. Close follow up

87. Obstetric forceps was applied to effect delivery of an alive female neonate at station +2 in left occipito-anterior position with a 45° rotation.

Which of the following best describes this type of forceps delivery?

- A. High
- B. Low
- C. Midpelvic
- ☒ D. Outlet

88. A pregnant lady had a uterine size which is five centimeters less than the stated LMP.

Which of the following is LEAST likely as a possible cause for this discrepancy?

- A. Fetal infection
- B. Congenital malformation
- C. Wrong date
- ☒ D. Concealed abruption

89. Which of the following is an essential prerequisite for application of forceps?

- A. Cervix more than 8 cm
- B. Gestational age beyond 37 weeks
- ☒ C. Ruptured membranes
- D. Vertex presentation

108. A 6-year old boy is diagnosed with acute rheumatic fever with the following findings: migratory polyarthrititis, fever, elevated ESR, and evidence of preceding streptococcal infection. Which of the following is the appropriate regarding the management of the above patient?
- A. Amoxicillin for 7 days
 - B. Aspirin
 - ☒ C. Penicillin prophylaxis life long
 - D. Prednisolone
109. After a diagnosis of pertussis in a toddler, the parents are worried about the transmission to other house hold members. Which house hold members should receive erythromycin prophylaxis?
- A. All house hold members
 - B. Immuno-compromised
 - C. Unvaccinated against pertussis
 - ☒ D. Younger than 7 year
110. Which of the following clinical conditions that appear after pertussis vaccination is an absolute contraindication to subsequent administration of the vaccine?
- A. Fever and convulsion within three days
 - B. Immediate anaphylactic reaction
 - C. Inconsolable cry longer than two hours
 - D. Shock-like state after two days
111. Which of the following vaccine should NOT be given for an infant with severe immune suppression?
- A. Hepatitis B
 - B. Measles
 - C. Meningococcal
 - D. Pneumococcal vaccine
112. You are evaluating an HIV exposed neonate immediately after delivery. The mother had proper ANC follow up and was managed according PMTCT option B+. Which of the following is the appropriate recommendation for this newborn?
- A. Avoid breast feeding
 - B. Cotrimoxazole prophylaxis within the first week of life
 - C. DNA- PCR test within 48 hours
 - ☒ D. Nevirapine for 6 weeks
113. A 3 month old infant has rhinorrhea, cough and noisy breath. Physical examination reveals RR=72/min, PR=142/min and T= 37.8°C. He is irritable and has diffuse wheezes and chest retraction. What is the most likely diagnosis?
- ☒ A. Bronchiolitis
 - B. Gastro-esophageal reflux disease
 - C. Hyper active airway disease
 - D. Severe pneumonia

121. A 10 years old boy bled from multiple sites after he sustained an accident while riding a bicycle. On examination he has tachycardia and the blood pressure is not recordable. What will be the correct amount of crystalloid you will administer as the initial bolus?
- A. 10 ml/kg
☒ B. 20 ml/kg
 C. 30 ml/kg
 D. 40 ml/kg
122. A 5 year-old boy presented to the OPD with abdominal swelling of one month duration. On examination BP is normal, PR=100/min, RR= 42/min. He has stony dullness on the posterior right two third of the lung field, shifting dullness and fluid thrill on abdominal examination. He has pitting pedal edema. Urinalysis showed protein 4+. Which one of the following serum levels should you measure?
- A. Alanine transaminase
 B. Bilirubin
☒ C. Cholesterol
 D. Creatinine
123. A 36 month-old male infant presented with high grade fever and vomiting of two days duration. He has right costovertebral angle tenderness. White cell count showed leukocytosis with left shift and blood film is negative for hemoparasite. Urine is positive for nitrite and leukocyte esterase. What is the most likely diagnosis in this child?
- ☒ A. Pyelonephritis
 B. Hepatitis
 C. Appendicitis
 D. Lobar pneumonia
124. A 5-year old boy sustained a penetrating injury to his right leg. Two weeks later he developed swelling of the injured leg and high grade fever. X-ray of the affected limb showed features of acute osteomyelitis. What is the most likely etiology?
- A. ~~Pseudomonas~~ aeruginosa
 B. Klebsiella pneumonia
☒ C. Streptococcus pyogenes
 D. E. coli
125. A 2 year-old child has delay in all developmental mile stones for his age. On examination he has macrocephaly, severe stunting, protruded tongue, umbilical hernia and coarse facial features. The best initial test is to determine serum level of :
- A. Cortisol
 B. Growth Hormone
☒ C. Thyroid Stimulating Hormone
 D. Vitamin D
126. Which one of the following drugs is an appropriate antidote for Malathion poisoning?
- ☒ A. Atropine
 B. Flumazenil
 C. N acetyl cysteine
 D. Naloxone

114. Which of the following best describes clinical features of Guillain-Barré syndrome?
- ☒ A. Ascending symmetric paralysis without sensory change
 - ☐ B. Asymmetric paralysis and sensory change
 - ☐ C. Symmetric paralysis and cranial nerve palsy
 - ☐ D. Symmetric paralysis and sensory changes
115. A 5 year-old boy with HIV infection is receiving Zidovudine, Lamivudine and Nevirapine. He presents with progressive difficulty of breathing for three weeks. He has no history of fever. On examination PR=108/min, RR=72/min, T= 37°C and Oxygen saturation is 70 %. Chest radiograph reveals bilateral diffuse alveolar disease. What is the most likely diagnosis?
- ☐ A. Heart failure
 - ☒ B. Pneumocystis pneumonia
 - ☐ C. Pulmonary tuberculosis
 - ☐ D. Severe bacterial pneumonia
116. The recommended time to start folate supplementation for a mother who had a baby with myelomeningocele to prevent its recurrence in future pregnancies is:
- ☐ A. At the first missed period
 - ☐ B. End of first trimester
 - ☒ C. One month before conception
 - ☐ D. Third trimester
117. Which of the following CSF profile is expected in a child with meningoencephalitis?
- ☐ A. Leukocytosis with neutrophil predominance
 - ☐ B. Normal protein level
 - ☒ C. Normal or slightly reduced glucose level
 - ☐ D. Organism on Gram stain
118. A 12 hour old neonate presented with a boggy diffuse scalp swelling. Forceps was applied for prolonged second stage of labor. On examination he is in respiratory distress, grossly pale, and the capillary refill was 5 seconds. What is the most likely explanation for the above scenario?
- ☐ A. Caput succedaneum
 - ☐ B. Cephalhematoma
 - ☐ C. Intracranial hemorrhage
 - ☒ D. Subgaleal hemorrhage
119. A 6 week-old infant presented with high grade fever three hours after he took the first of his scheduled vaccines. He was well before the vaccine. On examination he is febrile and irritable otherwise there are no findings in the other system. What is the most appropriate management of this infant?
- ☐ A. Admit for observation
 - ☐ B. Initiate oral antibiotics
 - ☐ C. Investigate for infection
 - ☒ D. Paracetamol suppository
120. Which of the following CSF profiles suggests pyogenic meningitis in neonates?
- ☒ A. High cell count
 - ☐ B. High protein
 - ☐ C. Low glucose
 - ☐ D. Positive gram stain

32/min, temperature 39°C and has petechial rash on trunk. Which of the following is the best initial test for diagnosis?

- A. Blood film
- B. Complete blood count
- ☒ C. CSF analysis
- E. Widal test

133. Infectious diseases may have more than one mode of transmission. The number of modes of transmission has an effect on the control, elimination or eradication of that specific disease. From this perspective, which of the following diseases is more difficult to control?

- A. Pulmonary tuberculosis
- B. Relapsing fever
- C. Syphilis
- D. Trachoma

134. A health officer is delegated to lead a team travelling to West African countries to control the epidemic of Ebola Virus Disease (EVD). The role of the team includes giving supportive care to the cases of EVD and teaching them to avoid contact with the rest of the community members. To which prevention level (s) can this be categorized?

- A. Primary prevention both to the patients and the community
- B. Primary prevention to the patients but secondary prevention to the community
- C. Primary prevention to the community but secondary prevention to the patients
- D. Primary prevention to the community but tertiary prevention to the cases

135. An estimated 90 people out of 100 cases of Ebola die. What does this proportion show?

- A. Cause specific mortality rate
- B. Case fatality rate
- C. Proportionate mortality ratio
- D. Crude death rate

136. Suppose psychosis is a rare health problem in a specific country. Researchers want to study whether khat chewing causes psychosis in this country. Which one of the following study designs is most appropriate to assess this association?

- A. Cross sectional study
- ☒ B. Case control study
- C. Cohort study
- D. Experimental study

137. Considering the nature of the diseases, for which of the following health problems is screening LEAST recommended?

- A. Screening for HIV infection
- B. Screening pregnant women for syphilis
- ☒ C. Screening for relapsing fever
- D. Screening for cervical cancer

127. A 3 year-old boy presented with a complaint of intense pruritis worse at night. On examination there are vesicular and papular eruptions at the interdigital spaces. Which one is the best treatment for this child?

A. Acyclovir

B. Erythromycin

C. Mebendazole

☒ D. Permethrin

128. A 2-hour term neonate is delivered by elective C/S for placenta previa. On physical examination PR=170 /min, RR=72/min and oxygen saturation of 80%.

What is the most likely cause of respiratory distress in this neonate?

A. Congenital pneumonia

B. Perinatal Asphyxia

C. Respiratory distress syndrome

☒ D. Transient tachypnea of the newborn

129. A 2-year old child presented with difficulty of breathing of 2 hours. Mother reported that he barking cough and hoarseness of voice in the last two days. On physical exam he is agitated, has audible stridor at rest and lower chest indrawing. What is the most likely diagnosis in this child?

☒ A. Croup syndrome

B. Foreign body aspiration

C. Pertussis

D. Severe pneumonia

130. A 7-year old female patient who is on follow-up for bronchial asthma presented with shortness of breath of three hours. On physical examination she is conscious, able to talk in complete sentence, RR= 46/min and PR=110/min. Chest examination reveals diffuse expiratory wheeze. What is the initial best step in the management of this patient?

A. Aminophylline

B. Magnesium sulphate

C. Oral prednisolone

☒ D. Salbutamol puff

131. A 3 year-old boy is brought to the ER with high grade fever and shortness of breath of two days. On physical examination RR= 62/min, T=39°C, lower chest indrawing with bilateral basal coarse crepitation. The chest X-ray showed left basal lung pneumatocele. Which of the following is the most likely etiology?

A. Haemophilus Influenzae

☒ B. Staphylococcus aureus

C. Streptococcus pneumonia

D. Streptococcus pyogenes

132. A 13-year old street boy presented to the ER with high grade fever of 5 days. He has epistaxis, fatigue and headache. On physical examination heart rate 120/min, respiratory rate

138. Suppose 'Test X' is used as a screening test for 'Disease Y'. The sensitivity and specificity of 'Test X' are 90% and 80% respectively. Which of the following is correct interpretation of these findings?
- A. 'Test X' can correctly identify 90% of individuals with 'Disease Y'
 - B. 'Test X' can correctly identify 90% of individuals without 'Disease Y'
 - C. 'Test X' can correctly identify 80% of individuals with 'Disease Y'
 - D. 20% of the individuals with 'Disease Y' will be missed by 'Test X'
139. Patients may be grouped based on different characteristics for reporting. Which of the following grouping represent ordinal data?
- A. Blood group
 - B. Ethnic group
 - C. Disease severity
 - D. Sex
140. A researcher is interested to determine whether there is malaria outbreak in a given zone of a region. Which method is the best to collect data for the intended purpose at community level?
- A. Conduct Sample Survey
 - B. Conduct Census
 - C. Review record of patients in all providers
 - D. Focus group discussions
141. Which of the following diagrammatic presentation turns cumulative frequency distribution in to graphs?
- A. Ogive curve
 - B. Frequency polygon
 - C. Stem and leaf plot
 - D. Histogram
142. The dietary assessment on patients from a refugee camp shows many of the patients subsist on a diet of corn meal and boiled greens. They have little appetite for food and alternate between diarrhea and constipation. They also have scaly dermatitis. What is the most probable dietary deficiency in these patients?
- A. Niacin
 - B. Pyridoxine
 - C. Vitamin E
 - D. Vitamin K
143. The serum level of 1, 25 di-hydroxy vitamin D in adolescent girls is believed to be normally distributed with mean 65 pg/ml and standard deviation 12.5 pg/ml. What percent of adolescent girls will have a level higher than 65 pg/ml?
- A. 10%
 - B. 25%
 - C. 50%
 - D. 100%



144. A medical director at a regional hospital wanted to assess prevalence of Malaria in the town in order to make sufficient preparation for a possible outbreak. The town is composed of urban and rural kebeles. Which sampling methods give better precision for estimating parameter of interest?

- A. Simple random sampling
- B. Cluster sampling
- C. Stratified sampling
- D. Quota sampling

145. A woman usually prepares food for her family from cereals and legumes. She usually uses wheat for making a number of food items. She hardly gets vegetables and fruits foods in the diet of the family. What dietary deficiency is this family likely to face?

- A. Cholecalciferol
- B. Folic acid
- C. Niacin
- D. Pyridoxine

146. A large number of under-five children were diagnosed with serious neurological problems and anemia in a certain Woreda. The families of these children were strictly vegetarian. What is the most likely nutritional deficiency of this community?

- A. Vitamin B12
- ☒ B. Folic Acid
- C. Niacin
- D. Pyridoxine

147. A four year-old child is brought to the Health Center with acute respiratory tract infection for the first visit. What dose of vitamin A will be given for this child?

- A. 50,000
- B. 100,000 IU
- C. 200,000 IU
- D. 400,000IU

148. A patient admitted to a Hospital ate a 57 gram serving of whole-wheat spaghetti containing 8 grams protein, 40 grams carbohydrate and 2 grams fat. What is the estimate of energy intake of this patient in Kcal?

- A. 200
- B. 210

- C. 280
- D. 400

149. A health officer aims to initiate a behavioral change among individuals by considering the socio-cultural factors after identifying the personal characteristics of the people. Which one is the best theory or model to apply?

- A. Health belief model
- B. Social cognitive theory
- C. Theory of planned behavior
- D. Trans-theoretical model

150. While giving health education to a group of mothers to introduce a new family planning method, which of the following learning materials are preferred?
- ☒ A. Model
 - B. Photograph
 - C. Poster
 - D. Real object
151. Which of the following sites is ideal place to display a message using information education communication materials to create awareness of a new family planning method in a rural village?
- A. Homes
 - ☒ B. Market places
 - C. Prisons
 - D. Religious institutions
152. Some community-based preventive interventions initiated by the government have failed to meet the desired objectives even though they have good plans. Which of the following is the main reason for the failure?
- ☒ A. Absence of appropriate technology
 - B. Higher cost of interventions
 - C. Lack of good policy in the country
 - D. Lack of planning with the community
153. A communal well water source development project was planned to be developed in a rural village. For the outcome of the project to be effective, what needs to be identified in the process?
- ☒ A. Acceptance of the water source by the community
 - B. Cost of materials for development purpose
 - C. Electrolyte composition of the water source
 - D. Level of trace minerals in the water source
154. A report from a health center showed most residents of that district were affected by water-related diseases. An assessment in the district revealed that most of the residents drink water from few sources of untreated wells by treating it with Water Agar. But the culture of keeping personal hygiene was poor. Which water related diseases would affect the residents of the district most?
- A. Water arthropod disease
 - ☒ B. Water borne infection
 - C. Water impounding disease
 - D. Water washed disease
155. Hazardous waste disposal was found to be a critical problem in a Health Center. A disposal options for hazardous wastes was required for a public health action. Which waste disposal option is the most appropriate for the health center?
- A. Burying
 - B. Composting
 - C. Incinerating
 - D. Offsite Disposal

162. Of the strategies to reduce maternal mortality, which of the following interventions is most effective in reducing the likelihood of developing pregnancy-related complications?
- ☒ A. Antenatal care
 - B. Emergency obstetric care
 - C. Family planning
 - D. Nutrition intervention
163. A manager of a Health Center wants to provide BCG vaccine for infants on Fridays. Which of the following is the best management function that the manager used?
- A. Organizing
 - B. Planning
 - ☒ C. Controlling
 - D. Implementing
164. A Hospital manager allocated a budget for three different activities. Which of the following is the next step in planning?
- A. Designing strategy
 - B. Prioritizing problem
 - C. Setting objective
 - D. Situational analysis
165. An EPI coordinator of a district and his subordinates were implementing the program for according to national standard of EPI coverage. Which one is the best method to assess their performance at the end of six months?
- A. Comparing performance against the standard
 - B. Implementing controlling process
 - C. Measuring actual performance
 - D. Standardization of performance
166. A zonal Health Department head assigned a budget for different tasks for a fiscal year. Which decisional role did he play?
- A. Negotiation role
 - B. Dissemination role
 - C. Resource allocation role
 - D. Monitoring role
167. The total population of certain locality was around 25,000. Which of the following health facilities is suitable to provide health service to that community?
- A. District hospital
 - B. Health center
 - C. Health post
 - D. Health station
168. A college dean who has experts in different professions as employees left them to do their activities. What leadership style did he apply?
- A. Autocratic
 - B. Bureaucratic
 - C. Democratic
 - D. Laissez faire
169. There was interpersonal conflict in the same department of a certain health facility. What was the best management option that they do?
- A. Leave the issue for those involved
 - B. Manage the conflict after a month
 - C. Switch one of them to another department
 - D. Resolve the conflict as soon as possible

- 120 Sheets
9 1/2" x 7"
242mm x 178mm
SP03397
156. The malaria prevalence in a town is increasing from time to time. A team has been established to develop malaria epidemic control and prevention program. What priority intervention needs to be included in the program to effectively limit the progress of the epidemic?
- A. Application of outdoor residual spray
 - ☒ B. Destruction of mosquito breeding sites
 - C. Enforce usage of insect repellants
 - D. Timely treatment of cases
157. In a rural district, an ordinary millhouse is known for recruiting uneducated, economically poor individuals as permanent employees. The employees do not use any kind of Personal Protective Equipment. Which of the following is the most effective intervention to prevent noise-induced hearing loss?
- A. Early diagnosis and treatment
 - B. Enforce earmuff usage
 - C. Noise-proofing the mill
 - D. Reduce working hours
158. Reproductive health experts use maternal mortality as indicator of maternal health status although it has limitations. What is the main limitation of using this indicator?
- A. It is difficult to measure it at community level
 - B. It is not standard measure of deaths
 - C. It over-represents maternal health problems
 - D. It under-represents maternal health problems
159. The 2011 Ethiopian demographic and health survey reported that institutional delivery service utilization was 10% which was unacceptably low. What was the most important reason reported by women for not delivering in a health facility?
- A. High cost of delivery services
 - B. Lack of access to health services
 - C. Lack of awareness
 - D. Lack of transportation
160. If an organization wants to design a strategy targeting the most important direct cause of maternal mortality globally. Which cause should get the first priority?
- A. Abortion
 - ☒ B. Hemorrhage
 - C. Hypertension
 - D. Sepsis
161. A health officer wants to know risk of maternal death due to level of fertility and risk of maternal death per pregnancy or per birth. Which measure of maternal mortality should he/she use?
- A. Life time risk of maternal mortality
 - B. Maternal mortality rate
 - C. Maternal mortality ratio
 - D. Proportion maternal

- 120 Sheets
9 1/2" x 7"
242mm x 178mm
SP03597
170. A 24 year-old woman comes to your office for follow up on her depression. She has decided to try to get pregnant and wants to stop taking her fluoxetine. She had a history of suicidal attempt and severe peri-partum depression in the past. Which of the following is appropriate management?
- A. Continue to take fluoxetine
 - B. Shift to another antidepressant
 - C. Shift to antipsychotic drug
 - D. Start mood stabilizer
171. A 26 year-old woman brought by her sister to the Hospital with seven months history of unusual behavior. The patient gradually develops poor self-care, isolation and sleep problem. She hears the voices of several people talking to her. His sister says she talks to herself and laughs alone. On mental state examination she has blunt affect, auditory hallucination & has no insight to her illness. Which of the following is the most likely diagnosis?
- A. Delusional disorder
 - B. Schizoaffective disorder
 - ☒ C. Schizophrenia
 - D. Schizophreniform disorder
172. A 28 year-old man diagnosed as a case of Major depressive disorder with psychotic features. He is treated with haloperidol and amitriptyline. Two months later, his mood symptoms have resolved and he is no longer psychotic. Which of the following best describes the next step?
- A. Both drugs should be tapered and discontinued
 - B. Haloperidol should be tapered and discontinued
 - C. Amitriptyline should be tapered and discontinued
 - ☒ D. Both drugs should be continued for 6 months
173. A 53 year-old man comes to the Hospital complains of poor appetite, loss of pleasure in daily activities, sleeplessness, & decreased libido for 3 weeks. He is contemplating retiring, expressing concern that his poor performance in his job. On mental state examination he shows empty feeling & makes two errors in subtraction serial 3's from 20. Which of the following is the most likely diagnosis?
- A. Bipolar disorder
 - B. Cyclothymia
 - C. Dysthymic disorder
 - D. Major depressive disorder
174. A 35 year-old man diagnosed to have Major depressive disorder was given 75 mg amitriptyline at bed time. On the third day he starts hearing voices which tell him that "he is a bad person". On mental state examination he feels sad and has auditory hallucinations. Which of the following is the most appropriate treatment?
- A. Diazepam
 - B. Fluoxetine
 - ☒ C. Haloperidol
 - D. Sodium valporate

- 120 Sheets
9 1/2" x 7"
242mm x 178mm
SP03397
175. A 20 year-old male brought by his cousin to the Hospital. His cousin says that after smoking marijuana he became suspicious and started to hear voices. He talks alone and is easily irritable. These behavioral changes persisted for the last two days. On mental state examination he has thought & perceptual disturbance. He has no past psychiatric illness. Which of the following is the most likely diagnosis?
- A. Cannabis induced delirium
 - B. Cannabis induced psychosis
 - C. Cannabis intoxication
 - D. Cannabis withdrawal
176. A 55 year-old alcoholic is brought to the Emergency Room with a problem of irrelevant talking. Three days ago he quit alcohol after he was counseled. On mental state examination he is disoriented to time, place & person. He has visual hallucination and agitation. There is no history of head injury or medical illness. Which of the following is the most likely diagnosis?
- A. Alcohol induced psychosis
 - B. Alcohol intoxication
 - C. Delirium tremens
 - D. Wernicke's encephalopathy
177. A 30 year-old woman came to Medical OPD accompanied by her husband. She delivered a healthy baby three weeks ago at a Health center. Her husband reports that she has irritability, tearfulness, & sleeplessness for the last two weeks. The past week she has started to feel worthlessness and have loss of concentration. Which of the following is the most likely diagnosis?
- A. Dysthymia
 - B. Postpartum blues
 - C. Postpartum depression
 - D. Postpartum psychosis
178. A 55 year-old man comes to Emergency Room with acute retention of urine for 12 hours. He was suffering from depression and taking drugs for the last ten days. On examination there is distended bladder. No other physical findings. Which drug is the most likely culprit?
- A. Amitriptyline
 - B. Fluoxetine
 - C. Paroxetine
 - D. Sertraline
179. A patient comes to your clinic with a complaint of hearing difficulty of one month duration. There is only a set of tuning forks for hearing assessment at the Clinic. Which tuning fork should be used for the assessment?
- A. 256 Hz
 - B. 512 Hz
 - C. 1024 Hz
 - D. 2048 Hz
180. A patient comes demanding for ear wash. Which of the following would be a contraindication for ear wash?
- A. Beads
 - B. Chronic otitis media
 - C. Cockroach
 - D. Impacted wax

- 120 Sheets
9 1/2" x 17"
242mm x 432mm
SP03397
181. A 3 year-old female child presented to the Emergency OPD with foreign body in the right nasal cavity of three days duration. On physical examination, there was an alkaline small button battery inside with discharge and necrotic nasal mucosal surface. What is the best management plan?
- A. Give acidic nasal drops for two days and remove it
 - B. Give nasal decongestants for two days and remove it
 - C. Immediately removal
 - D. Irrigate with saline and give antibiotics
182. A 37 year- old female presented with a cockroach in the right ear canal. She is so much in discomfort because of the movement of the insect. On physical examination, the right ear has big dry central perforation of the tympanic membrane. What is the management?
- A. Add 3% hydrogen per oxide to kill the insect
 - B. Add ethyl alcohol to kill the insect
 - C. Irrigation with warm water
 - D. Take the insect out with a small artery while alive
183. A 22 year- old male patient has left sided ear pain with no discharge. He has no history of trauma and physical examination revealed tenderness over the tragus. What is the most likely diagnosis?
- A. Acute otitis externa
 - B. Acute otitis media
 - C. Cerumen impaction
 - D. Chronic otitis media
184. A 25 year-old man came to a Primary Hospital's OPD after a fertilizer was splashed into his eyes while working in his farm. He has redness on both eyes with photophobia. What is the best initial management?
- A. Take comprehensive history and ocular physical examination
 - B. Refer urgently to the nearest ophthalmologic center
 - C. Wash the eyes with normal saline
 - D. Give anti-inflammatory and antibiotic eye drops
185. Which of the following is a cause for constricted pupil?
- A. Atropine
 - B. Acute angle closure glaucoma
 - C. Oculomotor palsy
 - D. Organophosphate poisoning
186. During a school survey on refraction, a 6 year-old girl was found to have a refractive error of (+3) on the left eye while the right eye was (+6). If the refractive error is not corrected by spectacles, which of the following is LEAST expected to happen?
- A. Amblyopia
 - B. Floaters
 - C. Poor school performance
 - D. Strabismus
187. What is the leading cause of visual impairment globally?
- A. Cataract
 - B. Diabetic retinopathy
 - C. Glaucoma
 - D. Uncorrected refractive error

188. What is the most common cause of unilateral cataract in young adults?
- ☒ A. Metabolic/
 - B. Uveitis
 - C. Trauma
 - D. UV radiation
189. A 51 year-old man came to your OPD complaining he can't read contact numbers from his mobile phone since 6 months. He used to be able to read by holding the phone further away. He and no other ocular complaints. He had 6/6 distance vision on both eyes. What is the most likely diagnosis?
- A. Age-related macular degeneration
 - B. Astigmatism
 - C. Cataract
 - ☒ D. Presbyopia
190. A mother brought her 1 year old boy to the health center claiming the child has poor vision. Which of the following is the least likely to suggest congenital glaucoma?
- A. Blepharospasm
 - B. Epiphora
 - C. Increased corneal diameter /
 - D. Redness
191. A 55 year-old man came to medical OPD with a chief complaint of pain during mastication for the last one year. Intra-oral examination shows horizontal teeth mobility around 1mm, gingival recession with exposed root surfaces, intra-oral periapical x-ray shows alveolar bone resorption. Which of the following is the most likely diagnosis?
- A. Periodontitis
 - B. Gingivitis
 - C. Pulpitis
 - D. Stomatitis
192. A 30 year-old woman who had tooth extraction four days ago came with severe pain in the socket. On examination the socket is dry with no clot formation. The alveolar margin was tender. What is the best next step in management?
- A. Apply local paste
 - B. Do Debridement
 - C. Pack with gauze
 - D. Prescribe antibiotics
193. A 25 year-old women with gestational age of 11 weeks came to Medical OPD with recurrent toothache. On examination maxillary first molar hopelessly decayed and slightly tender to percussion. Which one of the following is the most appropriate management?
- A. Extraction
 - B. Pain control
 - C. PO tetracycline
 - D. Root canal therapy

194. Which one of the following is typical feature of bilateral anterior TMJ dislocation?
- Deviation of chin
 - Lateral cross bite
 - Posterior cross bite
 - Protrusion of chin
195. A 45 year-old man presented to the Emergency Department with a chief complaint of jaw swelling and difficulty of breathing. Physical examination showed non-pitting, firm, board like swelling over the primary mandibular facial spaces bilaterally with raised floor of mouth, dysphagia, tenderness on palpating the area and trismus of mouth. Which one of the following is most likely diagnosis?
- Ameloblastoma
 - Facial cellulitis
 - Ludwing's angina
 - Submandibular abscess
196. A 25 year-old woman presented with gum bleeding and bad breath odor for one month. Examination shows interdental papillae full of debris and multiple ulcers, easily bleeding gum, linear erythema and free gingival margin with full of ulcers. What is the most likely diagnosis?
- Acute necrotizing periodontitis
 - Acute necrotizing ulcerative gingivitis
 - Chronic gingivitis
 - Puberty induced gingivitis
197. A 25 year old women came with chief complaint of swelling related with lower right jaw of about one week duration. History and clinical examination confirms submandibular space abscess. Which of the following tooth is the most likely cause?
- Lower canine
 - Lower premolars
 - Lower 1st molar
 - Lower 2nd molar
198. A 35 year-old woman came to a hospital with a chief complaint of inability to close her eyes. Physical examination shows dropping of the corner of the mouth, leakage of saliva, inability to close the right eye. Deviation of mouth to the right side. Which one of the following is the most likely diagnosis?
- Trigeminal neuralgia
 - Facial nerve palsy
 - Glossopharyngeal neuralgia
 - Trigeminal nerve paralysis
199. A 30 year-old patient came with chief complaint of mobility of teeth. Which one of the following would suggest the diagnosis of periodontitis?
- False pocket
 - Gum growth
 - Gum recession
 - Short clinical crown
200. A 25 year-old man presented with fighting accident of 24 hours. Physical examination shows laceration of upper lip, fractured upper incisor. X-ray result shows apical half fracture of the offending tooth. Which one of the following is the best management option?
- Extraction
 - Inter-dental wiring
 - Root canal therapy
 - Splinting

- 12F40
- Regurgitant murmur
6. A 25 year-old man with rheumatic heart disease came for follow-up. He has active precordium and rumbling mid-diastolic murmur. What valvular lesion is most likely in this patient? old - sten
- ~~C. MS~~
- A. Aortic regurgitation
 - B. Mitral regurgitation
 - ☒ C. Mitral stenosis
 - D. Tricuspid regurgitation
7. A 65 year-old male with type 2 diabetes mellitus presents to the Emergency Department with a squeezing, retrosternal chest pain of one hour duration. He has vague discomfort in his right arm and shoulder. The physical examination is notable for diaphoresis and mild tachycardia. Which of the following is the most likely diagnosis?
- ☒ A. Acute coronary syndrome
 - B. Acute pericarditis
 - C. Aortic dissection
 - D. Pulmonary embolism
8. Which of the following is a major criteria for the diagnosis of rheumatic fever?
- A. Elevated ESR
 - B. Fever
 - C. Prolonged PR segment
 - ☒ D. Polyarthralgia
9. A 29 year-old woman came to the OPD with shortness of breath and body swelling of three weeks. On physical examination she has raised JVP, accentuated P2 and grade IV diastolic murmur at the apex. Which of the following drug delays disease progression?
- A. Frusemide
 - B. Penicillin
 - ☒ C. Enalapril
 - D. Digoxin
10. A 45 year-old male with heart failure is comfortable at rest. He develops dyspnea and fatigue to walk to a nearby shop and is forced to rest for five minutes before symptoms improved. What is the stage and functional status of this patient?
- A. Stage A, Class II
 - B. Stage B, Class III
 - C. Stage C, Class IV
 - ☒ D. Stage C, Class III
11. A 50 year-old man was found to have a BP=170/100. He has a history of asthma. Which one is the most appropriate drug of choice?
- A. Atenolol
 - B. Enalapril
 - ☒ C. Hydralazine
 - D. Furosemide

1. A 26 year-old female presented with two months of excessive menses and gum bleeding. Physical examination revealed pale conjunctiva, petechiae and ecchymosis. Which of the following is the most likely diagnosis?
 A. Hemophilia
 B. Idiopathic thrombocytopenic purpura.
 C. Scurvy (pt)
 D. Vitamin K deficiency
2. A 21 year-old woman presented with fatigue and light headedness. Physical examination showed marked conjunctival and palmar pallor. Which of the following physical findings is most likely to be seen in this patient?
 A. Apical diastolic murmur
 B. Ejection systolic murmur
 C. Narrow pulse pressure
 D. Rumbling mid diastolic murmur
 Anemia - (E) - S/S murmur. (for murmur)
 Tachycardia
3. A health officer was evaluating an anemic patient. Laboratory test revealed: Hgb = 7 g/dl, MCV = 77 fl, WBC = 5400/ μ l, Platelet = 650,000/ μ l. Which of the following is the most likely diagnosis?
 A. Acute myelogenous leukemia
 B. Aplastic anemia
 C. Iron-deficiency anemia
 D. Megaloblastic anemia
 80 - 88
 Anemia of I. Defe.
 -> microcytic
 -> Hypochromic
 Also cobalamin vit. B12 def.
4. A 45 year-old woman presented with fatigue and orthostatic dizziness of two months duration. She has conjunctival pallor. Complete blood count revealed: Hgb = 8 g/dl, WBC = 4200/ μ L and platelet = 170,000/ μ l. Which of the following co-morbidities is LEAST likely?
 A. Chronic kidney disease (✓) (Anemia)
 B. Chronic obstructive lung disease
 C. Colorectal cancer (✓) (bleeding)
 D. Disseminated tuberculosis (✓) (liver, spleen, BM)
5. A 57 year-old man is admitted to the Medical Ward with hemorrhagic stroke. After a week the patient suddenly developed plueritic chest pain and shortness of breath. Physical examination revealed signs of respiratory distress and a swelling involving his right calf and thigh. What is the most likely cause for his deterioration?
 A. Acute myocardial infarction
 B. Aspiration pneumonia
 C. Fat embolism
 D. Pulmonary embolism (DVT)

28. A 60 year-old alcoholic patient presented with fever, headache and neck pain of two days duration. On examination body $T=39^{\circ}\text{C}$ and has neck stiffness. Bacterial meningitis was suspected but lumbar puncture was deferred because he has papilledema. Which of the following is the best empiric therapy for this patient in addition to Ampicillin?
- A. Vancomycin and Chloramphenicol
 - ☒ B. Ceftriaxone and Vancomycin
 - C. Ceftriaxone and Chloramphenicol
 - D. Cloxacillin and Ciprofloxacin
29. A 36-year-old male presented with fatigue and tea-colored urine for five days. Physical examination revealed jaundice and tender Hepatomegaly. Laboratory investigation revealed: Aspartate aminotransferase (AST) = 2400 U/L, Alanine aminotransferase (ALT) = 2500 U/L, Alkaline phosphatase = 210 U/L and total bilirubin = 8.6 mg/dL. Which of the following is the LEAST likely cause for the patient's condition?
- A. Acetaminophen poisoning
 - B. Acute Viral hepatitis
 - ☒ C. Alcoholic liver disease
 - D. Ischemic liver injury
30. A 56 year-old man presented with progressive abdominal distension and leg swelling of two months duration. He is a known patient with cirrhosis for the past one year on follow up but not on medication. On examination, he was not in distress and has moderate ascites and bilateral leg swelling. Which of the following is the best initial management?
- A. Furosemide
 - ☒ B. Furosemide and Spironolactone
 - C. Hydrochlorothiazide
 - D. Paracentesis and Furosemide
31. Which of the following is the most common complication of Peptic ulcer disease?
- ☒ A. Hemorrhage
 - B. Gastric outlet obstruction
 - C. Malignancy
 - D. Perforation
32. A 50 year-old woman presented with fever, cough and chest pain for two days. Chest X-ray showed pleural effusion. Pleural fluid analysis showed: $\text{WBC} = 5,000/\text{mm}^3$, with 90% neutrophils, $\text{PH} = 7.2$, blood culture was positive for streptococcus pneumonia. What is the next best step in management of this patient besides antibiotics?
- A. Dexamethasone
 - B. Chest CT scan
 - ☒ C. Chest tube insertion
 - D. Lung function test

32/min, temperature 39°C and has petechial rash on trunk. Which of the following is the best initial test for diagnosis?

- A. Blood film
- B. Complete blood count
- ☒ C. CSF analysis
- E. Widal test

133. Infectious diseases may have more than one mode of transmission. The number of modes of transmission has an effect on the control, elimination or eradication of that specific disease. From this perspective, which of the following diseases is more difficult to control?

- A. Pulmonary tuberculosis
- B. Relapsing fever
- C. Syphilis
- D. Trachoma

134. A health officer is delegated to lead a team travelling to West African countries to control the epidemic of Ebola Virus Disease (EVD). The role of the team includes giving supportive care to the cases of EVD and teaching them to avoid contact with the rest of the community members. To which prevention level (s) can this be categorized?

- A. Primary prevention both to the patients and the community
- B. Primary prevention to the patients but secondary prevention to the community
- C. Primary prevention to the community but secondary prevention to the patients
- D. Primary prevention to the community but tertiary prevention to the cases

135. An estimated 90 people out of 100 cases of Ebola die. What does this proportion show?

- A. Cause specific mortality rate
- B. Case fatality rate
- C. Proportionate mortality ratio
- D. Crude death rate

136. Suppose psychosis is a rare health problem in a specific country. Researchers want to study whether khat chewing causes psychosis in this country. Which one of the following study designs is most appropriate to assess this association?

- A. Cross sectional study
- ☒ B. Case control study
- C. Cohort study
- D. Experimental study

137. Considering the nature of the diseases, for which of the following health problems is screening LEAST recommended?

- A. Screening for HIV infection
- B. Screening pregnant women for syphilis
- ☒ C. Screening for relapsing fever
- D. Screening for cervical cancer

138. Suppose 'Test X' is used as a screening test for 'Disease Y'. The sensitivity and specificity of 'Test X' are 90% and 80% respectively. Which of the following is correct interpretation of these findings?
- A. 'Test X' can correctly identify 90% of individuals with 'Disease Y'
 - B. 'Test X' can correctly identify 90% of individuals without 'Disease Y'
 - C. 'Test X' can correctly identify 80% of individuals with 'Disease Y'
 - D. 20% of the individuals with 'Disease Y' will be missed by 'Test X'
139. Patients may be grouped based on different characteristics for reporting. Which of the following grouping represent ordinal data?
- A. Blood group
 - B. Ethnic group
 - C. Disease severity
 - D. Sex
140. A researcher is interested to determine whether there is malaria outbreak in a given zone of a region. Which method is the best to collect data for the intended purpose at community level?
- A. Conduct Sample Survey
 - B. Conduct Census
 - C. Review record of patients in all providers
 - D. Focus group discussions
141. Which of the following diagrammatic presentation turns cumulative frequency distribution in to graphs?
- ☒ A. Ogive curve
 - ☐ B. Frequency polygon
 - C. Stem and leaf plot
 - D. Histogram
142. The dietary assessment on patients from a refugee camp shows many of the patients subsist on a diet of corn meal and boiled greens. They have little appetite for food and alternate between diarrhea and constipation. They also have scaly dermatitis. What is the most probable dietary deficiency in these patients?
- ☒ A. Niacin
 - B. Pyridoxine
 - C. Vitamin E
 - D. Vitamin K
143. The serum level of 1, 25 di-hydroxy vitamin D in adolescent girls is believed to be normally distributed with mean 65 pg/ml and standard deviation 12.5 pg/ml. What percent of adolescent girls will have a level higher than 65 pg/ml?
- A. 10%
 - B. 25%
 - C. 50%
 - D. 100%



- Which of the following best describes clinical features of Guillain-Barré syndrome?
114. ☒ A. Ascending symmetric paralysis without sensory change
 - ☐ B. Asymmetric paralysis and sensory change
 - ☐ C. Symmetric paralysis and cranial nerve palsy
 - ☐ D. Symmetric paralysis and sensory changes
115. A 5 year-old boy with HIV infection is receiving Zidovudine, Lamivudine and Nevirapine. He presents with progressive difficulty of breathing for three weeks. He has no history of fever. On examination PR=108/min, RR=72/min, T= 37°C and Oxygen saturation is 70 %. Chest radiograph reveals bilateral diffuse alveolar disease. What is the most likely diagnosis?
- ☐ A. Heart failure
 - ☒ B. Pneumocystis pneumonia
 - ☐ C. Pulmonary tuberculosis
 - ☐ D. Severe bacterial pneumonia
116. The recommended time to start folate supplementation for a mother who had a baby with myelomeningocele to prevent its recurrence in future pregnancies is:
- ☐ A. At the first missed period
 - ☐ B. End of first trimester
 - ☒ C. One month before conception
 - ☐ D. Third trimester
117. Which of the following CSF profile is expected in a child with meningoencephalitis?
- ☐ A. Leukocytosis with neutrophil predominance
 - ☐ B. Normal protein level
 - ☒ C. Normal or slightly reduced glucose level
 - ☐ D. Organism on Gram stain
118. A 12 hour old neonate presented with a boggy diffuse scalp swelling. Forceps was applied for prolonged second stage of labor. On examination he is in respiratory distress, grossly pale, and the capillary refill was 5 seconds. What is the most likely explanation for the above scenario?
- ☐ A. Caput succedaneum
 - ☐ B. Cephalhematoma
 - ☐ C. Intracranial hemorrhage
 - ☒ D. Subgaleal hemorrhage
119. A 6 week-old infant presented with high grade fever three hours after he took the first of his scheduled vaccines. He was well before the vaccine. On examination he is febrile and irritable otherwise there are no findings in the other system. What is the most appropriate management of this infant?
- ☐ A. Admit for observation
 - ☐ B. Initiate oral antibiotics
 - ☐ C. Investigate for infection
 - ☒ D. Paracetamol suppository
120. Which of the following CSF profiles suggests pyogenic meningitis in neonates?
- ☒ A. High cell count
 - ☐ B. High protein
 - ☐ C. Low glucose
 - ☐ D. Positive gram stain

45. A 30 year-old man was seen at the Emergency surgical OPD with perianal pain of two days duration. He has fever and chills. He has left perianal induration with tenderness but no fluctuation. What is the most likely diagnosis?

- A. Anal fissure
- ☒ B. Perianal Abscess

- C. Perianal fistula
- D. Thrombosed Hemorrhoids

46. A 20 year-old man was seen at the emergency department following road traffic accident of one hour duration. He has sustained trauma to the abdomen. On examination; Vital signs: BP=80/50 mmHg and PR=120/min. He has tire marks on the abdomen and it was tender. The patient is started on IV fluids. What is the next step in the management?

- A. Abdominal CT-scan
- ☒ B. Exploratory laparotomy

- C. Observation
- D. Plain abdominal x-ray

47. A 50 year- old man came complaining of fatigue and dyspepsia of three months duration. Physical examination revealed a pale conjunctiva with a palpable mass at the epigastrium that moves with respiration. What is the gold standard diagnostic modality?

- A. Abdominal Ultra sound
- B. CT scan of the abdomen

- ☒ C. Upper GI Endoscopy
- D. Plain abdominal film

48. A man was stabbed to the abdomen and presented to the Health Center with eviscerated small bowel which was getting dark. Which of the following is the correct initial measure before referral?

- ☒ A. Cover the intestine with saline gauze
- B. Refer to a nearby hospital without intervention
- C. Replace the intestine back into the peritoneal Cavity
- ☒ D. Widen the stab wound under local anesthesia

49. A 23 year-old farmer presented to the OPD with abdominal pain and frequent vomiting of one day duration. On examination the abdomen is slightly distended and has visible peristalsis. Plain abdominal x-ray showed step ladder pattern air-fluid level. Which of the following is the most likely diagnosis?

- A. Caecal Volvulus
- B. Compound Volvulus

- C. Sigmoid Volvulus
- ☒ D. Small bowel Volvulus

50. A 65 year-old man was seen at the Emergency OPD with abdominal pain, distension and failure to pass feces and flatus of two days duration. He had two similar attacks in the last three years and was successfully treated with rectal tube deflation. On Examination BP=90/60 mmHg and PR=110/min. The abdomen was tender and bowel sounds were absent. Which of the following measures is to be avoided?

- A. Analgesics
- B. Antibiotics

- C. IV fluid
- ☒ D. Rectal tube deflation

A 10 years old boy bled from multiple sites after he sustained an accident while riding a bicycle. On examination he has tachycardia and the blood pressure is not recordable. What will be the correct amount of crystalloid you will administer as the initial bolus?

A. 10 ml/kg

☒ B. 20 ml/kg

C. 30 ml/kg

D. 40 ml/kg

122. A 5 year-old boy presented to the OPD with abdominal swelling of one month duration. On examination BP is normal, PR=100/min, RR= 42/min. He has stony dullness on the posterior right two third of the lung field, shifting dullness and fluid thrill on abdominal examination. He has pitting pedal edema. Urinalysis showed protein 4+. Which one of the following serum levels should you measure?

A. Alanine transaminase

B. Bilirubin

☒ C. Cholesterol

D. Creatinine

123. A 36 month-old male infant presented with high grade fever and vomiting of two days duration. He has right costovertebral angle tenderness. White cell count showed leukocytosis with left shift and blood film is negative for hemoparasite. Urine is positive for nitrite and leukocyte esterase. What is the most likely diagnosis in this child?

☒ A. Pyelonephritis

B. Hepatitis

C. Appendicitis

D. Lobar pneumonia

124. A 5-year old boy sustained a penetrating injury to his right leg. Two weeks later he developed swelling of the injured leg and high grade fever. X-ray of the affected limb showed features of acute osteomyelitis. What is the most likely etiology?

A. *Pseudomonas aeruginosa*

B. *Klebsiella pneumonia*

☒ C. *Streptococcus pyogenes*

D. *E. coli*

125. A 2 year-old child has delay in all developmental mile stones for his age. On examination he has macrocephaly, severe stunting, protruded tongue, umbilical hernia and coarse facial features. The best initial test is to determine serum level of :

A. Cortisol

B. Growth Hormone

☒ C. Thyroid Stimulating Hormone

D. Vitamin D

126. Which one of the following drugs is an appropriate antidote for Malathion poisoning?

☒ A. Atropine

B. Flumazenil

C. N acetyl cysteine

D. Naloxone

51. A 45 year-old housewife presented with progressive submandibular swelling of two days duration. She visited a dentist and undergone tooth extraction a day prior to the start of her symptoms. Examination showed tense, shiny bilateral submandibular swelling with no area of fluctuation. She also has drooling of the saliva and protruded tongue. What is the diagnosis of this patient?
- ☒ A. Ludwig's angina
☐ B. Periodontal abscess
☐ C. Submandibular cellulites
☐ D. Submental abscess
52. A 20 year-old man presented to the OPD forty minutes after he sustained stab injury to his left posterior chest. He has shortness of breathing. He is acutely sick looking; BP= 110/60 mmHg, PR= 90/min, RR=26/min. He has decreased air entry on the lower left chest with visible air rush through the stab wound. What is the best first step in the management of this patient?
- ☐ A. Apply airtight dressing
☒ B. Apply triangular dressing
☐ C. Suture the wound
☐ D. Take chest X-ray
53. A 35 year-old male patient came to emergency OPD 12 hours after he started to have sudden onset of epigastric pain. He has long standing history of dyspepsia for which he has been taking anti-acids repeatedly. Examination revealed tachycardia, rigid and tender abdomen. What is the appropriate imaging study to reach a definitive diagnosis?
- ☐ A. Barium follow through
☐ B. Barium meal
☒ C. Chest X-ray
☒ D. Plain abdominal X-ray
54. A 27 year-old patient was involved in a road traffic accident. He was brought to the Hospital by the driver. He is agitated; PR=140/min and feeble, BP=80/40 mmHg. He has pale conjunctiva and his chest is clear with good air entry. He has an echymosis over his left flank area. Abdominal Ultrasonography showed a significant collection of fluid in his abdomen. What is the most likely diagnosis?
- ☐ A. Cardiogenic shock
☒ B. Hemorrhagic shock
☐ C. Neurogenic shock
☐ D. Septic shock
55. A 64 year-old male patient presented to the Emergency OPD after he failed to pass urine for a day. He had difficulty of urination for the last one year. The previous night he took seven bottles of beer with friends. On examination he is acutely sick looking and in pain. Bladder is palpable 10 cm above the pubic bone and tender. What is the most likely cause for acute urinary retention?
- ☒ A. Benign Prostatic Hyperplasia
☐ B. Bladder cancer
☐ C. Bladder stone
☐ D. Urethral Stricture

33. A 46 year-old man is brought to the Emergency Department because he was vomiting fresh blood. He reports that he has no abdominal pain or nausea. On physical examination; BP=70/40 mmHg, PR=118/min, RR= 24/min. He also has gynecomastia, dilated abdominal veins and ascites. Which of the following is the most likely cause of the bleeding?

A. Esophageal cancer

☒ B. Esophageal varices

C. Mallory Weiss

D. Reflux esophagitis

34. A 47 year-old man comes to a Hospital complaining of epigastric pain that is exacerbated by eating. Antigenic test for *Helicobacter pylori* is positive. Endoscopic examination reveals a duodenal ulcer. Which of the following is the most appropriate therapy?

☒ A. Amoxicillin, clarithromycin and omeprazole

B. Amoxicillin, metronidazole and anti-acid

C. Omeprazole and anti-acid

D. Ranitidine, amoxicillin and metronidazole

35. A 29 year-old man with HIV infection presents with chronic diarrhea associated with anorexia, abdominal cramps and significant weight loss. His recent CD4 count =20 cells/mm³. A modified acid-fast stain of a stool specimen shows 4-6 μ m oocysts. Which of the following is the most likely cause?

☒ A. *Cryptosporidium parvum*

B. *Microsporidia*

C. *Mycobacterium avium* complex

D. *Pneumocystis jiroveci*

36. Which of the following is the best early indicator for diabetic nephropathy?

A. Rising blood urea nitrogen

☒ B. Albuminuria

C. Hypertension

D. Rising creatinine level

37. A 35 year-old man presents with generalized body swelling of two months. Physical examination reveals BP=120/70mmHg, ascites and +3 pretibial edema. Protein = 4.5g/24hour urine collection. Which of the following is the most likely diagnosis?

A. Chronic liver disease

B. Heart failure

☒ C. Nephrotic syndrome

D. Glomerulonephritis

38. A 55 year-old man comes to the Emergency OPD because of excruciating pain in his right big toe. He was diagnosed to have gouty arthritis. Which of the following is the most appropriate pharmacotherapy?

A. Allopurinol

B. Ceftriaxone

☒ C. Indomethacin

D. Morphine

39. A 28 year-old lactating woman presented to the emergency OPD with a complaint of right breast pain of five days duration. On examination body T= 38°C with right breast tenderness but no fluctuation. What is the most likely diagnosis?

- ☒ A. Breast Abscess
- B. Breast Cancer

- C. Fibroadenoma
- D. Fibrocystic Change

40. A 32 year-old man presented to the surgical OPD with complaint of neck mass of one year duration. On examination he has a hoarse voice and a solitary 2cm x 2cm thyroid mass on the right side which has firm consistency and fixed to underlying structures. There was no lymphadenopathy. What is the most likely diagnosis?

- A. Thyroid adenoma
- ☒ B. Thyroid cancer

- C. Thyroiditis
- D. Toxic Nodule

41. A 35 year-old man with a chronic peptic ulcer disease presented with features of acute abdomen and chest x-ray revealed pneumoperitoneum. His symptoms lasted for six hours and his vital signs were stable after infusion of one liter of Normal Saline solution. What should be the next step in the management of this case?

- A. Barium meal study
- B. Endoscopy

- C. Triple therapy
- ☒ D. Laparotomy

42. A 50 year-old woman brought an ultrasound imaging result which showed Gall Bladder Stone. She has no abdominal pain and no known medical illness. Which of the following is the recommended management?

- A. Cholecystectomy
- B. Extracorporeal Shockwave Lithotripsy (ESWL)
- ☒ C. Reassurance
- D. Ursodeoxycholic acid

43. A 68 year-old man is admitted to a Hospital with complaint of intermittent lower abdominal pain, bowel habit change and rectal bleeding of one month duration. He has tenesmus and incomplete sense of defecation and had lost weight. What is the most likely diagnosis?

- A. Amebic dysentery
- ☒ B. Colorectal cancer
- C. Sigmoid Volvulus
- D. Ulcerative colitis

44. A 55 year-old man presented to the surgical OPD with a complaint of rectal bleeding of one month duration. He has no protruding mass per rectum and digital rectal examination was non-revealing. What is the investigation of choice?

- A. Abdominal CT-Scan
- B. Barium Enema
- ☒ C. Colonoscopy
- D. Upper GI Endoscopy

77. A 38 year-old primigravida comes for a routine visit at 39 weeks of gestation. Her blood pressure is persistently 140/90 mm Hg, and her urine protein is +2. Physical examination is otherwise unremarkable, and she is completely asymptomatic. Her cervix is 2 cm dilated and 90% effaced, vertex presentation and at station 0. Which of the following is the most appropriate management option for this case?
- ☒ A. Admission for observation
 - B. Immediate cesarean section
 - C. Outpatient follow-up till 40 weeks
 - D. Start induction of labor
78. A 34 year-old primigravida comes with cessation of fetal movement of four days duration at gestational age of 35 weeks from LNMP. She is otherwise asymptomatic and has no medical problem. Examination shows stable vital signs, symphysio-fundal height of 34cms, no uterine contraction and absent FHB. Which of the following is the initial best step in the management of this patient?
- A. Consider Termination
 - B. Do coagulation profile
 - C. Prophylactic antibiotics
 - ☒ D. Ultrasound for confirmation
79. A 63 year-old woman who came for regular gynecologic check-up was incidentally diagnosed to have cystocele. Which of the following is the most appropriate management for this patient?
- A. Anterior colporaphy
 - B. Kegel's exercise
 - C. Pessaries
 - D. Vaginal hysterectomy
80. A 21 year-old GII PI mother at 24 weeks gestation complains of sever vulvar itching and cotton cheese like discharge. A wet mount slide show no clue cells but hyphae. Which of the following is the most likely diagnosis?
- A. Bacterial vaginosis
 - ☒ B. Candida albicans
 - ~~C. Chlamydia trachomatis~~
 - D. Trichomonas vaginalis
81. A 29 year-old woman presents with vaginal pruritus associated with white curd like discharge. On examination there is erythematous macular lesion but no vesicles. Investigation shows no clue cell, negative whiff test. Which of the following is the treatment of choice for this case?
- A. Azithromycin
 - ☒ B. Fluconazole
 - C. Metronidazole
 - D. Oral acyclovir
82. A 26 year-old GIPI woman at 26 weeks of gestation from her LNMP, comes for ANC follow up. She hasn't appreciated fetal movement yet. She had one episode of bleeding a month back and noticed failure of increase in uterine size. Her uterine size measures 20 weeks and you are unable to hear fetal heart tone. Which of the following is the most descriptive diagnosis?
- A. Incomplete abortion
 - B. Inevitable abortion
 - ☒ C. Missed abortion
 - D. Threatened abortion

56. An 18 years old male presented with right scrotal swelling of 1 day duration. He also has associated pain and it started while he was asleep, it woke him up. He also has fever and pain during urination. On examination right testis is tender, hot to touch, swollen, and the testicular cord is not thickened. What is the most likely diagnosis?
- A. Testicular torsion
☒ B. Epididymo-orchitis
 C. Infected hydrocele
 D. Testicular tumor
57. A 36 year-old male patient came with difficulty of urination for the last two years. He had previous history of whitish urethral discharge for which he was treated with tablets and injections. What is the most likely diagnosis?
- A. Benign prostatic hyperplasia
 B. Prostatic Cancer
 C. Sexually transmitted disease
☒ D. Urethral Stricture
58. A 43 year-old male patient came to the Emergency OPD after being involved in a road traffic accident before a day. He has failed to urinate since the incident and is unable to move his right lower limb. On examination his vital signs are normal, there is right pubic tenderness and blood on the penile meatus. What should be the next step in management?
- A. Decompress with percutaneous needle puncture
 B. Do image-guided catheterization
 C. Insert a foley catheter to decompress the bladder
☒ D. Perform emergency suprapubic cystostomy
59. A 24 year-old man came to the Emergency OPD with severe pain originating in the flank, radiating inferiorly and anteriorly. He also has nausea and vomiting. His vital signs are within normal limits. Abdominal examination showed marked costovertebral angle tenderness. Urinalysis showed microscopic hematuria. What is the most likely diagnosis?
- A. Cystitis
 B. Pyelonephritis
 C. Pyonephrosis
☒ D. Ureteric colic
60. A 60 year-old female patient presented with right sided flank pain and hematuria of one month duration. She also has significant weight loss, night sweats and malaise. On physical examination, she is chronically sick looking with normal vital signs, a bimanually palpable mass on the right flank. What is the most likely diagnosis?
- A. Chronic pyelonephritis
 B. Neuroblastoma
☒ C. Renal cell cancer
 D. Wilm's tumor

61. A 56 year-old man presented with a scrotal mass of three months duration. Physical examination revealed a 3cm x 4 cm firm mass on the lower pole of the right testis. Scrotal ultrasound showed a well-defined hypoechoic lesion within the testis. Serum β -hCG was elevated. What is the most likely diagnosis?

- A. Orchitis
- B. Hydrocele
- ☒ C. Testicular Tumor
- D. Testicular torsion

62. A 35 year-old male patient presented to the Emergency Room after sustaining a bullet injury to his left forearm three hours back. Up on evaluation vital signs were normal and a deformed and rotated forearm, intact distal neurovascular examination. X-ray showed comminuted displaced radio-ulnar fracture. Which of the following is a temporary means of immobilization?

- A. Apply POP cast
- ☒ B. External fixation
- C. Lateral arm traction
- D. Plating and screw

63. A 65 year-old male patient sustained a fall down accident three hours prior to presenting to the Emergency Room. He fell down with the arms extended. On examination vital signs were normal with a right wrist deformity. X-ray showed displaced Colle's fracture. Which one is the best management option?

- A. Application of a POP Cast
- ☒ B. Closed reduction and casting
- C. Open reduction and plating
- D. Skin traction for few days then casting

64. A 20 year-old male patient had an external fixator applied for a compound comminuted right tibio-fibular fracture three weeks prior to presentation. He now complains of pain at the pin site. On examination there is seropurulent discharge from the two distal pins. Which of the following is the best diagnosis?

- A. Cellulitis
- B. Osteomyelitis
- C. Osteonecrosis
- ☒ D. Pin tract infection

65. A 20 year-old male patient came to you with right knee pain of five days duration. On examination BP=120/70 mmHg, PR= 84/min. The knee is swollen and tender with signs of knee effusion. X-ray showed a wide joint space. Joint fluid analysis revealed turbid fluid with $100,000 \text{ WBC/mm}^3$. Which of the following is the most likely diagnosis?

- A. Acute osteomyelitis
- B. Osteoarthritis
- ☒ C. Pyogenic arthritis
- D. Tuberculous arthritis

83. A 42 year-old healthy nulliparous woman has a firm, irregular shaped, 12 week-sized midline abdominal mass.

Which of the following is the best initial step in evaluating this patient?

- A. Abdominal x-ray
- B. CT scan
- C. MRI
- ☒ D. Ultrasound

84. Which of the following is NOT a component of active third stage management?

- A. Controlled cord traction
- ☒ B. Nipple stimulation
- C. Uterine massage
- D. Oxytocin provision

85. A GIII PII mother came to a clinic at gestational age of 16 weeks for ANC follow up.

Which of the following is LEAST likely to be requested as part of a routine work up?

- A. Blood group and Rh
- ☒ B. Ultrasound
- C. Urinalysis
- D. VDRL

86. A primigravida was admitted to a Health Center due to pushing down pain of 6 hours. Physical examination; vital signs were normal, uterine size 36 weeks, longitudinal, cephalic, FHR=146/min, two contractions in 10min each lasting for 30-45 seconds; cervix is 3 cm dilated and 60 % effaced. After 4 hours she was reevaluated and had same findings except the cervical effacement changed to 75 %.

Which of the following is the best management option?

- A. Augmentation
- B. Rehydration
- C. Referral to hospital
- ☒ D. Close follow up

87. Obstetric forceps was applied to effect delivery of an alive female neonate at station +2 in left occipito-anterior position with a 45° rotation.

Which of the following best describes this type of forceps delivery?

- A. High
- B. Low
- C. Midpelvic
- ☒ D. Outlet

88. A pregnant lady had a uterine size which is five centimeters less than the stated LMP.

Which of the following is LEAST likely as a possible cause for this discrepancy?

- A. Fetal infection
- B. Congenital malformation
- C. Wrong date
- ☒ D. Concealed abruption

89. Which of the following is an essential prerequisite for application of forceps?

- A. Cervix more than 8 cm
- B. Gestational age beyond 37 weeks
- ☒ C. Ruptured membranes
- D. Vertex presentation

188. What is the most common cause of unilateral cataract in young adults?
- ☒ A. Metabolic/
 - B. Uveitis
 - C. Trauma
 - D. UV radiation
189. A 51 year-old man came to your OPD complaining he can't read contact numbers from his mobile phone since 6 months. He used to be able to read by holding the phone further away. He and no other ocular complaints. He had 6/6 distance vision on both eyes. What is the most likely diagnosis?
- A. Age-related macular degeneration
 - B. Astigmatism
 - C. Cataract
 - ☒ D. Presbyopia
190. A mother brought her 1 year old boy to the health center claiming the child has poor vision. Which of the following is the least likely to suggest congenital glaucoma?
- A. Blepharospasm
 - B. Epiphora
 - C. Increased corneal diameter /
 - D. Redness
191. A 55 year-old man came to medical OPD with a chief complaint of pain during mastication for the last one year. Intra-oral examination shows horizontal teeth mobility around 1mm, gingival recession with exposed root surfaces, intra-oral periapical x-ray shows alveolar bone resorption. Which of the following is the most likely diagnosis?
- A. Periodontitis
 - B. Gingivitis
 - C. Pulpitis
 - D. Stomatitis
192. A 30 year-old woman who had tooth extraction four days ago came with severe pain in the socket. On examination the socket is dry with no clot formation. The alveolar margin was tender. What is the best next step in management?
- A. Apply local paste
 - B. Do Debridement
 - C. Pack with gauze
 - D. Prescribe antibiotics
193. A 25 year-old women with gestational age of 11 weeks came to Medical OPD with recurrent toothache. On examination maxillary first molar hopelessly decayed and slightly tender to percussion. Which one of the following is the most appropriate management?
- A. Extraction
 - B. Pain control
 - C. PO tetracycline
 - D. Root canal therapy

194. Which one of the following is typical feature of bilateral anterior TMJ dislocation?
- Deviation of chin
 - Lateral cross bite
 - Posterior cross bite
 - Protrusion of chin
195. A 45 year-old man presented to the Emergency Department with a chief complaint of jaw swelling and difficulty of breathing. Physical examination showed non-pitting, firm, board like swelling over the primary mandibular facial spaces bilaterally with raised floor of mouth, dysphagia, tenderness on palpating the area and trismus of mouth. Which one of the following is most likely diagnosis?
- Ameloblastoma
 - Facial cellulitis
 - Ludwing's angina
 - Submandibular abscess
196. A 25 year-old woman presented with gum bleeding and bad breath odor for one month. Examination shows interdental papillae full of debris and multiple ulcers, easily bleeding gum, linear erythema and free gingival margin with full of ulcers. What is the most likely diagnosis?
- Acute necrotizing periodontitis
 - Acute necrotizing ulcerative gingivitis
 - Chronic gingivitis
 - Puberty induced gingivitis
197. A 25 year old women came with chief complaint of swelling related with lower right jaw of about one week duration. History and clinical examination confirms submandibular space abscess. Which of the following tooth is the most likely cause?
- Lower canine
 - Lower premolars
 - Lower 1st molar
 - Lower 2nd molar
198. A 35 year-old woman came to a hospital with a chief complaint of inability to close her eyes. Physical examination shows dropping of the corner of the mouth, leakage of saliva, inability to close the right eye. Deviation of mouth to the right side. Which one of the following is the most likely diagnosis?
- Trigeminal neuralgia
 - Facial nerve palsy
 - Glossopharyngeal neuralgia
 - Trigeminal nerve paralysis
199. A 30 year-old patient came with chief complaint of mobility of teeth.
Which one of the following would suggest the diagnosis of periodontitis?
- False pocket
 - Gum growth
 - Gum recession
 - Short clinical crown
200. A 25 year-old man presented with fighting accident of 24 hours. Physical examination shows laceration of upper lip, fractured upper incisor. X-ray result shows apical half fracture of the offending tooth. Which one of the following is the best management option?
- Extraction
 - Inter-dental wiring
 - Root canal therapy
 - Splinting

194. Which one of the following is typical feature of bilateral anterior TMJ dislocation?
D A. Deviation of chin
B. Lateral cross bite
C. Posterior cross bite
D. Protrusion of chin
195. A 45 year-old man presented to the Emergency Department with a chief complaint of jaw swelling and difficulty of breathing. Physical examination showed non-pitting, firm, board like swelling over the primary mandibular facial spaces bilaterally with raised floor of mouth, dysphagia, tenderness on palpating the area and trismus of mouth. Which one of the following is most likely diagnosis?
C A. Ameloblastoma
B. Facial cellulitis
C. Ludwig's angina
D. Submandibular abscess
196. A 25 year-old woman presented with gum bleeding and bad breath odor for one month. Examination shows interdental papillae full of debris and multiple ulcers, easily bleeding gum, linear erythema and free gingival margin with full of ulcers. What is the most likely diagnosis?
C A. Acute necrotizing periodontitis
B. Acute necrotizing ulcerative gingivitis
C. Chronic gingivitis
D. Puberty induced gingivitis
197. A 25 year old women came with chief complaint of swelling related with lower right jaw of about one week duration. History and clinical examination confirms submandibular space abscess. Which of the following tooth is the most likely cause?
D A. Lower canine
B. Lower premolars
C. Lower 1st molar
D. Lower 2nd molar
198. A 35 year-old woman came to a hospital with a chief complaint of inability to close her eyes. Physical examination shows dropping of the corner of the mouth, leakage of saliva, inability to close the right eye. Deviation of mouth to the right side. Which one of the following is the most likely diagnosis?
B A. Trigeminal neuralgia
B. Facial nerve palsy
C. Glossopharyngeal neuralgia
D. Trigeminal nerve paralysis
199. A 30 year-old patient came with chief complaint of mobility of teeth.
Which one of the following would suggest the diagnosis of periodontitis?
C A. False pocket
B. Gum growth
C. Gum recession
D. Short clinical crown
200. A 25 year-old man presented with fighting accident of 24 hours. Physical examination shows laceration of upper lip, fractured upper incisor. X-ray result shows apical half fracture of the offending tooth. Which one of the following is the best management option?
A A. Extraction
B. Inter-dental wiring
C. Root canal therapy
D. Splinting

170. A 24 year-old woman comes to your office for follow up on her depression. She has decided to try to get pregnant and wants to stop taking her fluoxetine. She had a history of suicidal attempt and severe peri-partum depression in the past. Which of the following is appropriate management?
- A. Continue to take fluoxetine
 - B. Shift to another antidepressant
 - C. Shift to antipsychotic drug
 - D. Start mood stabilizer
171. A 26 year-old woman brought by her sister to the Hospital with seven months history of unusual behavior. The patient gradually develops poor self-care, isolation and sleep problem. She hears the voices of several people talking to her. His sister says she talks to herself and laughs alone. On mental state examination she has blunt affect, auditory hallucination & has no insight to her illness. Which of the following is the most likely diagnosis?
- A. Delusional disorder
 - B. Schizoaffective disorder
 - ☒ C. Schizophrenia
 - D. Schizophreniform disorder
172. A 28 year-old man diagnosed as a case of Major depressive disorder with psychotic features. He is treated with haloperidol and amitriptyline. Two months later, his mood symptoms have resolved and he is no longer psychotic. Which of the following best describes the next step?
- A. Both drugs should be tapered and discontinued
 - B. Haloperidol should be tapered and discontinued
 - C. Amitriptyline should be tapered and discontinued
 - ☒ D. Both drugs should be continued for 6 months
173. A 53 year-old man comes to the Hospital complains of poor appetite, loss of pleasure in daily activities, sleeplessness, & decreased libido for 3 weeks. He is contemplating retiring, expressing concern that his poor performance in his job. On mental state examination he shows empty feeling & makes two errors in subtraction serial 3's from 20. Which of the following is the most likely diagnosis?
- A. Bipolar disorder
 - B. Cyclothymia
 - C. Dysthymic disorder
 - D. Major depressive disorder
174. A 35 year-old man diagnosed to have Major depressive disorder was given 75 mg amitriptyline at bed time. On the third day he starts hearing voices which tell him that "he is a bad person". On mental state examination he feels sad and has auditory hallucinations. Which of the following is the most appropriate treatment?
- A. Diazepam
 - B. Fluoxetine
 - ☒ C. Haloperidol
 - D. Sodium valporate

162. Of the strategies to reduce maternal mortality, which of the following interventions is most effective in reducing the likelihood of developing pregnancy-related complications?
- ☒ A. Antenatal care
 - B. Emergency obstetric care
 - C. Family planning
 - D. Nutrition intervention
163. A manager of a Health Center wants to provide BCG vaccine for infants on Fridays. Which of the following is the best management function that the manager used?
- A. Organizing
 - B. Planning
 - ☒ C. Controlling
 - D. Implementing
164. A Hospital manager allocated a budget for three different activities. Which of the following is the next step in planning?
- A. Designing strategy
 - B. Prioritizing problem
 - C. Setting objective
 - D. Situational analysis
165. An EPI coordinator of a district and his subordinates were implementing the program for according to national standard of EPI coverage. Which one is the best method to assess their performance at the end of six months?
- A. Comparing performance against the standard
 - B. Implementing controlling process
 - C. Measuring actual performance
 - D. Standardization of performance
166. A zonal Health Department head assigned a budget for different tasks for a fiscal year. Which decisional role did he play?
- A. Negotiation role
 - B. Dissemination role
 - C. Resource allocation role
 - D. Monitoring role
167. The total population of certain locality was around 25, 000. Which of the following health facilities is suitable to provide health service to that community?
- A. District hospital
 - B. Health center
 - C. Health post
 - D. Health station
168. A college dean who has experts in different professions as employees left them to do their activities. What leadership style did he apply?
- A. Autocratic
 - B. Bureaucratic
 - C. Democratic
 - D. Laissez faire
169. There was interpersonal conflict in the same department of a certain health facility. What was the best management option that they do?
- A. Leave the issue for those involved
 - B. Manage the conflict after a month
 - C. Switch one of them to another department
 - D. Resolve the conflict as soon as possible

181. A 3 year-old female child presented to the Emergency OPD with foreign body in the right nasal cavity of three days duration. On physical examination, there was an alkaline small button battery inside with discharge and necrotic nasal mucosal surface. What is the best management plan?
- Give acidic nasal drops for two days and remove it
 - Give nasal decongestants for two days and remove it
 - Immediately removal
 - Irrigate with saline and give antibiotics
182. A 37 year- old female presented with a cockroach in the right ear canal. She is so much in discomfort because of the movement of the insect. On physical examination, the right ear has big dry central perforation of the tympanic membrane. What is the management?
- Add 3% hydrogen per oxide to kill the insect
 - Add ethyl alcohol to kill the insect
 - Irrigation with warm water
 - Take the insect out with a small artery while alive
183. A 22 year- old male patient has left sided ear pain with no discharge. He has no history of trauma and physical examination revealed tenderness over the tragus. What is the most likely diagnosis?
- Acute otitis externa
 - Acute otitis media
 - Cerumen impaction
 - Chronic otitis media
184. A 25 year-old man came to a Primary Hospital's OPD after a fertilizer was splashed into his eyes while working in his farm. He has redness on both eyes with photophobia. What is the best initial management?
- Take comprehensive history and ocular physical examination
 - Refer urgently to the nearest ophthalmologic center
 - Wash the eyes with normal saline
 - Give anti-inflammatory and antibiotic eye drops
185. Which of the following is a cause for constricted pupil?
- Atropine
 - Acute angle closure glaucoma
 - Oculomotor palsy
 - Organophosphate poisoning
186. During a school survey on refraction, a 6 year-old girl was found to have a refractive error of (+3) on the left eye while the right eye was (+6). If the refractive error is not corrected by spectacles, which of the following is LEAST expected to happen?
- Amblyopia
 - Floater
 - Poor school performance
 - Strabismus
187. What is the leading cause of visual impairment globally?
- Cataract
 - Diabetic retinopathy
 - Glaucoma
 - Uncorrected refractive error

175. A 20 year-old male brought by his cousin to the Hospital. His cousin says that after smoking marijuana he became suspicious and started to hear voices. He talks alone and is easily irritable. These behavioral changes persisted for the last two days. On mental state examination he has thought & perceptual disturbance. He has no past psychiatric illness. Which of the following is the most likely diagnosis?
- A. Cannabis induced delirium
 - B. Cannabis induced psychosis
 - C. Cannabis intoxication
 - D. Cannabis withdrawal
176. A 55 year-old alcoholic is brought to the Emergency Room with a problem of irrelevant talking. Three days ago he quit alcohol after he was counseled. On mental state examination he is disoriented to time, place & person. He has visual hallucination and agitation. There is no history of head injury or medical illness. Which of the following is the most likely diagnosis?
- A. Alcohol induced psychosis
 - B. Alcohol intoxication
 - C. Delirium tremens
 - D. Wernicke's encephalopathy
177. A 30 year-old woman came to Medical OPD accompanied by her husband. She delivered a healthy baby three weeks ago at a Health center. Her husband reports that she has irritability, tearfulness, & sleeplessness for the last two weeks. The past week she has started to feel worthlessness and have loss of concentration. Which of the following is the most likely diagnosis?
- A. Dysthymia
 - B. Postpartum blues
 - C. Postpartum depression
 - D. Postpartum psychosis
178. A 55 year-old man comes to Emergency Room with acute retention of urine for 12 hours. He was suffering from depression and taking drugs for the last ten days. On examination there is distended bladder. No other physical findings. Which drug is the most likely culprit?
- A. Amitriptyline
 - B. Fluoxetine
 - C. Paroxetine
 - D. Sertraline
179. A patient comes to your clinic with a complaint of hearing difficulty of one month duration. There is only a set of tuning forks for hearing assessment at the Clinic. Which tuning fork should be used for the assessment?
- A. 256 Hz
 - B. 512 Hz
 - C. 1024 Hz
 - D. 2048 Hz
180. A patient comes demanding for ear wash. Which of the following would be a contraindication for ear wash?
- A. Beads
 - B. Chronic otitis media
 - C. Cockroach
 - D. Impacted wax

144. A medical director at a regional hospital wanted to assess prevalence of Malaria in the town in order to make sufficient preparation for a possible outbreak. The town is composed of urban and rural kebeles. Which sampling methods give better precision for estimating parameter of interest?
- Simple random sampling
 - Cluster sampling
 - Stratified sampling
 - Quota sampling
145. A woman usually prepares food for her family from cereals and legumes. She usually uses wheat for making a number of food items. She hardly gets vegetables and fruits foods in the diet of the family. What dietary deficiency is this family likely to face?
- Cholecalciferol
 - Folic acid
 - Niacin
 - Pyridoxine
146. A large number of under-five children were diagnosed with serious neurological problems and anemia in a certain Woreda. The families of these children were strictly vegetarian. What is the most likely nutritional deficiency of this community?
- Vitamin B12
 - ☒ Folic Acid
 - Niacin
 - Pyridoxine
147. A four year-old child is brought to the Health Center with acute respiratory tract infection for the first visit. What dose of vitamin A will be given for this child?
- 50,000
 - 100,000 IU
 - 200,000 IU
 - 400,000IU
148. A patient admitted to a Hospital ate a 57 gram serving of whole-wheat spaghetti containing 8 grams protein, 40 grams carbohydrate and 2 grams fat. What is the estimate of energy intake of this patient in Kcal?
- | | |
|--------|--------|
| A. 200 | C. 280 |
| B. 210 | D. 400 |
149. A health officer aims to initiate a behavioral change among individuals by considering the socio-cultural factors after identifying the personal characteristics of the people. Which one is the best theory or model to apply?
- Health belief model
 - Social cognitive theory
 - Theory of planned behavior
 - Trans-theoretical model

156. The malaria prevalence in a town is increasing from time to time. A team has been established to develop malaria epidemic control and prevention program. What priority intervention needs to be included in the program to effectively limit the progress of the epidemic?
- A. Application of outdoor residual spray
 - ☒ B. Destruction of mosquito breeding sites
 - C. Enforce usage of insect repellants
 - D. Timely treatment of cases
157. In a rural district, an ordinary millhouse is known for recruiting uneducated, economically poor individuals as permanent employees. The employees do not use any kind of Personal Protective Equipment. Which of the following is the most effective intervention to prevent noise-induced hearing loss?
- A. Early diagnosis and treatment
 - B. Enforce earmuff usage
 - C. Noise-proofing the mill
 - D. Reduce working hours
158. Reproductive health experts use maternal mortality as indicator of maternal health status although it has limitations. What is the main limitation of using this indicator?
- A. It is difficult to measure it at community level
 - B. It is not standard measure of deaths
 - C. It over-represents maternal health problems
 - D. It under-represents maternal health problems
159. The 2011 Ethiopian demographic and health survey reported that institutional delivery service utilization was 10% which was unacceptably low. What was the most important reason reported by women for not delivering in a health facility?
- A. High cost of delivery services
 - B. Lack of access to health services
 - C. Lack of awareness
 - D. Lack of transportation
160. If an organization wants to design a strategy targeting the most important direct cause of maternal mortality globally. Which cause should get the first priority?
- A. Abortion
 - ☒ B. Hemorrhage
 - C. Hypertension
 - D. Sepsis
161. A health officer wants to know risk of maternal death due to level of fertility and risk of maternal death per pregnancy or per birth. Which measure of maternal mortality should he/she use?
- A. Life time risk of maternal mortality
 - B. Maternal mortality rate
 - C. Maternal mortality ratio
 - D. Proportion maternal

138. Suppose 'Test X' is used as a screening test for 'Disease Y'. The sensitivity and specificity of 'Test X' are 90% and 80% respectively. Which of the following is correct interpretation of these findings?
- A. 'Test X' can correctly identify 90% of individuals with 'Disease Y'
 - B. 'Test X' can correctly identify 90% of individuals without 'Disease Y'
 - C. 'Test X' can correctly identify 80% of individuals with 'Disease Y'
 - D. 20% of the individuals with 'Disease Y' will be missed by 'Test X'
139. Patients may be grouped based on different characteristics for reporting. Which of the following grouping represent ordinal data?
- A. Blood group
 - B. Ethnic group
 - C. Disease severity
 - D. Sex
140. A researcher is interested to determine whether there is malaria outbreak in a given zone of a region. Which method is the best to collect data for the intended purpose at community level?
- A. Conduct Sample Survey
 - B. Conduct Census
 - C. Review record of patients in all providers
 - D. Focus group discussions
141. Which of the following diagrammatic presentation turns cumulative frequency distribution in to graphs?
- A. Ogive curve
 - B. Frequency polygon
 - C. Stem and leaf plot
 - D. Histogram
142. The dietary assessment on patients from a refugee camp shows many of the patients subsist on a diet of corn meal and boiled greens. They have little appetite for food and alternate between diarrhea and constipation. They also have scaly dermatitis. What is the most probable dietary deficiency in these patients?
- A. Niacin
 - B. Pyridoxine
 - C. Vitamin E
 - D. Vitamin K
143. The serum level of 1, 25 di-hydroxy vitamin D in adolescent girls is believed to be normally distributed with mean 65 pg/ml and standard deviation 12.5 pg/ml. What percent of adolescent girls will have a level higher than 65 pg/ml?
- A. 10%
 - B. 25%
 - C. 50%
 - D. 100%

150. While giving health education to a group of mothers to introduce a new family planning method, which of the following learning materials are preferred?
- ☒ A. Model
 - B. Photograph
 - C. Poster
 - D. Real object
151. Which of the following sites is ideal place to display a message using information education communication materials to create awareness of a new family planning method in a rural village?
- A. Homes
 - ☒ B. Market places
 - C. Brisons
 - D. Religious institutions
152. Some community-based preventive interventions initiated by the government have failed to meet the desired objectives even though they have good plans. Which of the following is the main reason for the failure?
- ☒ A. Absence of appropriate technology
 - B. Higher cost of interventions
 - C. Lack of good policy in the country
 - D. Lack of planning with the community
153. A communal well water source development project was planned to be developed in a rural village. For the outcome of the project to be effective, what needs to be identified in the process?
- ☒ A. Acceptance of the water source by the community
 - B. Cost of materials for development purpose
 - C. Electrolyte composition of the water source
 - D. Level of trace minerals in the water source
154. A report from a health center showed most residents of that district were affected by water-related diseases. An assessment in the district revealed that most of the residents drink water from few sources of untreated wells by treating it with Water Agar. But the culture of keeping personal hygiene was poor. Which water related diseases would affect the residents of the district most?
- A. Water arthropod disease
 - ☒ B. Water borne infection
 - C. Water impounding disease
 - D. Water washed disease
155. Hazardous waste disposal was found to be a critical problem in a Health Center. A disposal options for hazardous wastes was required for a public health action. Which waste disposal option is the most appropriate for the health center?
- A. Burying
 - B. Composting
 - C. Incinerating
 - D. Offsite Disposal

1. A 26 year-old female presented with two months of excessive menses and gum bleeding. Physical examination revealed pale conjunctiva, petechiae and ecchymosis. Which of the following is the most likely diagnosis?
- A. Hemophilia
 B. Idiopathic thrombocytopenic purpura.
 C. Scurvy (pt)
 D. Vitamin K deficiency
2. A 21 year-old woman presented with fatigue and light headedness. Physical examination showed marked conjunctival and palmar pallor. Which of the following physical findings is most likely to be seen in this patient?
- A. Apical diastolic murmur
 B. Ejection systolic murmur
 C. Narrow pulse pressure
 D. Rumbling mid diastolic murmur
3. A health officer was evaluating an anemic patient. Laboratory test revealed: Hgb = 7 g/dl, MCV = 77 fl, WBC = 5400/ μ l, Platelet = 650,000/ μ l. Which of the following is the most likely diagnosis? 80 - 88
- A. Acute myelogenous leukemia
 B. Aplastic anemia
 C. Iron-deficiency anemia
 D. Megaloblastic anemia
4. A 45 year-old woman presented with fatigue and orthostatic dizziness of two months duration. She has conjunctival pallor. Complete blood count revealed: Hgb = 8 g/dl, WBC = 4200/ μ L and platelet = 170,000/ μ l. Which of the following co-morbidities is LEAST likely?
- A. Chronic kidney disease
 B. Chronic obstructive lung disease
 C. Colorectal cancer
 D. Disseminated tuberculosis
5. A 57 year-old man is admitted to the Medical Ward with hemorrhagic stroke. After a week the patient suddenly developed plueritic chest pain and shortness of breath. Physical examination revealed signs of respiratory distress and a swelling involving his right calf and thigh. What is the most likely cause for his deterioration?
- A. Acute myocardial infarction
 B. Aspiration pneumonia
 C. Fat embolism
 D. Pulmonary embolism

Anemia - (E) - sys. murmur.
 (for murmur)
 Tachycardia

Anemia of I. Defe.
 -> microcytic
 -> Hypochromic
 Also cobalamin vit. B12 def.

Anemia

Emphysema

breast

liver, spleen, BM

DVT

National licensing exam 2011 first round medicine MCQ

1. First line treatment for PID(inpatient) **a. ceftra + metro b. cefotetan +doxy c.ampicillin + gent**
2. Management for CO poisoning
3. PID (case asking for diagnosis)
4. WHO growth curve
5. What is the age(if the child can wear with help, go upstairs ,know his age and gender, feed himself) **a.2yrs b.3yr c.4yr**
6. Case asking for acute bacterial rhinosinitis
7. Case asking for PSGN complication
8. Best diagnostic modality for loculated pleural effusion **a.CT b. PA CXR c.lateral CXR d.chest U/S**
9. Dm screening method in pregnant mother **a. 50g glucose tolerance test b. 100g po tolerance test**
10. Best time to start fetal surveillance in DM mother **a. 26 b.32 c.33 d.36**
11. Type of epidemics **a. mixed b. propagated c.continues common source d.common source**
12. Which study design is best for rare occurrence **a.cohort b.case control**
13. Case control
14. Ulnar shaft fracture + radial head dislocation **a.galazie b.montagia**
15. Strangulated SBO(hx suggest SBO + mild tenderness around the umbilicus)
16. First management in pt SBO.
17. Gunshot case ...protruded Bowel .what is management a. conservative b. explore c. local exploration under local anesthesia
18. Pelvic trauma ...after 4bags of crystalloid still BP <90/60 what is next a.send for CT b. transfuse whole blood c. more crystalloid
19. 20 yrs old young presented with sudden onset of scrotal pain of 3hrs, cremasteric reflex absent , pain exacerbated with elevation what is next a. explore b. send for Doppler c. antibiotics and analgesic
20. 24 yrs old present with rummy nose, low grade fever , cough ..P/E erythematous nasal fold . what is management **a. reassurance**
21. Complication of parapneumonic effusion.. it was tapped and frank pus
a. Thoracentesis b. thoracostomy
22. Hx suggestive of Acute pyelonephritis: U/A leucocyte +ve , nitrite +ve ..antibiotics of choice
a.cephalexin b.cipro c.cotrimoxazole d. amoxicillin
23. Acute otitis media,Antibiotics **a.augmentin**
24. Case asking Tb lymphadenitis
25. **65yrs old male known asthmatic pt presented with recurrent attack of Right middle lobe pneumonia ... he is alcoholic, he smoke 30 pack yr,**
What is the top cause of his recurrent pneumonia
a. Asthma b. airway obstruction c. immunocompromise
26. 24 yrs old para I on her 4th post op(CS) day presented with sign and symptoms peripartum infection .. labor lasts 28hrs and rupture of membrane is 20hr
What is the single most important cause of her endometritis

- a. CS delivery b. prolonged rupture of membrane c. multiple digital exam c. prolonged labor
27. Cause otitis media in Swimmers
a. **Pseudomonas** b. **S.aureas** c. **streptococcus** d. **E.choli**
28. Nasal bleeding from anterior plexus casebleeding doesn't stop after cautery ..what is next step
A. Anterior nasal Packing b. **Embolization** c. **surgery**
29. Stage IIA cervical ca mgt **a. radical hysterectomy** c. **radiation therapy** c.
30. 34yrs old women unable to conceive for the last 4yrs 1.which investigation you sent first U/S is normal
a. Follicle stimulating hormone b. **progesterone** c. **salpingohysteroscopy** d. **CT**
31. 50yr old with breast lump ...swelling in the right upper outer quadrant, involve the skin ..what is the minimum stage
a. IIA b. IIB
32. pt with head trauma scalp laceration sutured with lidocaine . What is the earliest sign lidocaine toxicity
A. SEIZURE B. DISORIENTATION C. SLURRED SPEECH D. tongue
33. bilateral submandibular swelling.....leading angina
34. 50yr old male presented with painless preauricular swelling w/c is mobile, firm, non tender
A. PLEOMORPHIC PAROTID ADENOMA B. PAROTID CA C. WARFENG tumor of parotid D. TUMOR OF THE PAROTID
35. Case Diabetic retinopathy: w/c needs urgent referral to ophthalmologist
A. Neovascularization B. Retinal aa narrowing C. Hemorrhage D. Occular spotting
36. A 25yr old pt presented with loss of vision 5 days duration BP=170/110mmhg w/c of the ff is save his vision if managed first
A. controlling systemic BP B. decreasing ICP C. Refer to ophthalmologist
37. Case of s.malaria with COMA + p.falciparum
A. COMA CARE +ARTISUNATE+ANTIBIOTICS
B. COMA CARE+ ARTISUNATE
C. COMA CARE+ QUININE
D. COMA CARE+ ARTEMETER
38. A 12wk pregnant mother wants go to malaria endemic area. w/c prophylaxis you prefer
A. Doxycycline B. Quinine C. chloroquine D. Mefloquine

39. A 9 day neonate surgery was done for meningocele, after 4 days he presented with bulged fontanel, increased H.C, sunset appearance
- A. Hydrocephalus . B. Meningitis
40. 20yrs male pt fall from 10m high, presented with head CT. Showed brain contusion. After 4 days he developed persistent headache, analgesia, vomiting. P/E no focal neurological deficit, he has bradycardia and irregular respiration
- A. ICP B. brain herniation C. Subdural hematoma
41. A 30yr old known RVI pt presented with fever, headache 15 days duration. CT normal, CD4 100, CSF – opening pressure 200, glucose 20, protein 100, WBC 100. The most likely DX
- A. Toxoplasmosis B. Cryptococcus C. Bacterial meningitis
42. Newly RVI DX pregnant 3wks back mother gave birth to neonates, she is on HAART 3wks back
43. a 7 YRS OLD MALE PT partially treated meningitis presented with same complaint. CT shows 2x2 enhancing lesion on frontal lobe. What is mgt
- A. Ceftriaxone + vanco + metro
- B. Ceftriaxone + vanco
44. A 19yr old female presented with the complaint of amenorrhea of 2 month duration after she got pregnant from her brother's close friend without her willing. What is legal base to terminate?
- A. Rape B. Incest C. age
45. A 60 yr old known hypertensive on HCT and strict diet modification presented to OPD. On PE BP is 159/100mmHg. What is next plan?
- A. Increase dose of HCT B. salt restriction same dose of HCT C. HCT + Amlodipine D. Enalapril
46. Safety triangle case
47. Case of left orbit fracture
48. A 45 yr old female presented complaining with failure to close her mouth that occurs after yawning. What is the Dx
- A. TMJ dislocation B. Mandibular fracture
49. A 13 day old male neonate presented with right leg and thigh swelling. PE swollen right thigh and leg with limitation of knee joint. DX

A. septic arthritis B myositis C. Pyomyositis

50. A 20 yr old male pt presented with complain of preumbilical pain of 4days wich shifted to RLQ later. He treated with antibiotics .PE 4x5cm firm slightly well demarcated mass

A appendiceal mass B Appendiceal abscess

51.perforated PUD case

52. testiculat torsion Case

52 which is not routine in basic neonatal care

A. suction B, breart feeding, C TTC

53.Neonatal sepsis Case

54. Case of apnea of prematurity

54. Neonatal hypoglycemia with Seizure mgt. Ten percent glucose 4mg/kg then 10% glucose maintenance

55. A 30yr Fe pt presented with diarrhea vomiting abdominal pain of 20days. Lab Na is 120meq/mol

What is the cause of hyponatremia

A. vomiting B. Diarrhea

56. Lab K 3meq/l what is abnormality

A. hypokalemia B hyponatremimia

57 case of acute pyscotic disorder

58. case of post truomatic depression

59. A25 yr old para II mother who is on levothyroxine presented with hypothyroid symptoms

Lab TSH is 6(high) what is mgt

A. increase levothroxine B. PTU

60.Which antipsychotic drug needs regular TFT checkup

A. Lithium B.Sodium valporiate C. Carbamezapine

61. Side effects of anti depressent

62. Pt presented with tonic clonic sz of right side of the body late involved whole body, loss of consciousness. What is Dx

A primary seizure B. secndory generalized C. status epilepticus

63. Gardiasis Rx A. Tinidazole B. paromycine

64. pt presented bloody diarrhea + vomiting , S/E: puss cells

what is Rx

a. ORS + cipro b. albendazole c. ORS alone

65. case of brief psychotic disorder

66. **Delirium**, case

67. Case of vaginal candidiasis .. What is treatment?

68. 34wks pregnant mother come for ANC, no bleeding and other complain. U/S shows placenta previa
Totalis.. plan

a. do CS at 37weeks b.do Cs at 38 weeks c. wait for labor

69. 20yrs old mother comes with pushing down pain,p/E membrane intact, cervix 6cm , contraction
3'10'35'' what is Dx a. latent b.active first stage c.2nd stage

70. 24yrs old female presented with pushing down pain

P/E on digital examination you palpate mouth and the chin is toward the rectum

What is the presentation

a. mentum anterior face presentation

b. mentum, posterior face presentation

c. mentum anterior brow presentation

d. mentum posteriot brow presentation

71. 20yr old male presented with perianal pain and fever P/E ... there tender, indurated mass on the
posterior part of rectum ..what is management

a. I and D b. antibiotic analgesic with close follow up c. Send home with Po antibiotics

72. 3month baby presented with recurrent vomiting, significant weight loss .P/E visible peristalsis with
palpable mass on epigastric area ... which investigation do you send first

a. abdominal U/s b. plan abdominal X-ray c. barium enema D.CT

73. 40yrs old presented with abdominal pain and melena .Which of the following the best investigation to
confirm **a.sigmodoscopy b. colonoscopy c.CT d. Endoscopy**

74. Case DVT in post operative

75. HIV with DVT management

76. Case of CHF....which of the following drug decrease progression of disease in CHF

a. Beta blocker .b. digitalis c. loop diuretics c. calcium channel blockers

77. 7yr old male pt presented with swelling on un erupted upper canines...it was tapped and mucus
content

Management a. marsupialization and follow b. disimpaction c. enucleation

78. 21 yrs old male pt presented with swelling and pain on the 3rd molar. On exam there is an erupted wisdom tooth, tender, swollen. What is management

a. antibiotics, analgesic + follow up

79. Drug reaction case (see type of drug reaction like fixed drug reaction, TEN, Erythema multiforme)

80. case of cervical insufficiency

81. the neonate born at 32 weeks and wt 900g ... diagnosis

a. preterm + extremely very low birth weight

National Licensing Exam for Med - second round November, 2012 E.C

SECOND ROUND COC EXAM FOR MEDICAL STUDENTS

NOVEMBER, 2012

“MED STAR 6”



National Licensing Exam for Med - second round November, 2012 E.C

1. A 9 month old infant presented with execive crying, bloody diarrhea, abdominal distention and swelling, at health center given cotri and ors and no improvement... Dx

- A. Idiopathic intussusception
- B. Intussusception with mekel diverticulitis
- C. Intussusception with polyp
- D. Intussusception with lymphoma

Part 2

2. Appropriate time for Hepatitis Immunoglobulin for newborn

- A. 72 hrs
- B. 12Hrs
- C. 1 hrs
- D. 24hrs

3. which of the following confirm Dx of CML

- A. Philadelphia chromosome
- B. aures rod
- C. ***

4. about Tx of neonatal meningitis..

5. 11yr old male child who presented with 8month duration of progressive hoarseness of voice, difficulty of breathing and hx of decrease air entry, on indirect laryngoscopy there is strawberry soft tissue lesion on vocal cord

- A. laryngeal papilloma
- B. laryngeal carcinoma
- C. laryngeal Web
- D. laryngocele

6. 40 years old Patient with **POP**, a physician wants to see the maximum protrusion of the mass, what position is preferred

- A. Lying on her back
- B. Trendelburg
- C. Left lateral
- D. standing

National Licensing Exam for Med - second round November, 2012 E.C

7. female pt referred from health center after presented with cc of dyspnea. All P/E, cxr and Echo are normal, what investigation rule out heart failure

- A, d dimer
- B, c reactive protein
- C, Cardiac troponin
- D, natriuretic peptide

8. pregnant mother presented with urine frequency, dysuria on urine analysis pyuria and on culture no organism

What is the causative agent?

- A. E. coli
- B. Proteus
- C. Klebsiella pneum
- D. Chl.trochamitis

9. 10 years old child come to ER complaining SOB, no history of fever. WBC 7000/ml with 40 so2 85 percent neutrophils HGB 25 mg/dl PLT 600,000, what is the best management

- A. Give INo2
- B. Phlebotomy with N/S
- C. Partial volume Transfusion
- D. Antibiotics

10. A 45 yrs old male pt was presented with a complint of dyspnea on minimal exertion and cough productive of tenacious sputum. He has sneezing and hx of allergy. He has smoked for 30 pack yrs. The GP had difficulty to differentiate b/n Asthma and COPD, what favors dx of COPD.

- A. Seasonal variation of z symptoms.
- B. marked response to bronchodilator.
- C. lung function after bronchodilator FEV/FVC<70.
- D. marked response to prednisolone for 2 weeks duration

11. couples came to gyn OPD for a compliant of failure to conceive of 2yr duration. z female has prior hx. of STI txt 1yr back. z husband semen analysis is normal. wat is best next lx modality for z girl...

- A. estrogen.
- B. hysterosalpingography

National Licensing Exam for Med - second round November, 2012 E.C

C. progesterone.

12. 70 yrs man with tinnitus, vertigo and easy fatigability was admitted and transfused with 3 unit blood for severe anemia overnight and then he develop distress and irritability, on CXR he has peripheral haziness &...

Whats the likely Diagnosis...

A. ARDS

B. acute lung injury

C. volume overload

D.....

13. A 55year old known Parkinson's disease on medication presented with lower abdominal pain and failure to pass feaces and flatus of one day duration...most likely DX

A SBO

B, oglives syndrome

B LBO

Part – I

14. A 67 year old male patient on his seventh post op day after right hemicolectomy plus ileotransverse anastomosis done who has been on ceftriaxone 1gm IV BID and metronidazole 500 mg IV TID developed fever, diarrhea and abdominal distension. What is the initial step in managing this patient?

A) Stop ceftriaxone and metronidazole

B) Obtain abdominal ultrasound

C) Do stool culture

D) Provide loperamide

15. Patient diagnosed with TB meningitis treated with antitb and steroid for three weeks. what most likely complication is anticipated

A, hypernatremia

B, hydrocephalus

C, hypokalemia

D, brain abscess

16. 75 yrs old male pt presented with bloody vomiting of 24 hrs duration his vs at presentation bp 60/30 pr 110 rr 25 t 37 he was resussusutated with NS, catheterized, NG tube inserted, transfused. his V/S

National Licensing Exam for Med - second round November, 2012 E.C

become BP 90/60 PR 100 RR 20 has adequate UOP. endoscopy done shows clean base ulcer with visible blood vessel at posterior part of first part of duodenum. what is the most appropriate next step management

- A. high dose PPI infusion
- B. emergency laparotomy
- C. close observation at hospital
- D. do emergency upper GI endoscopy

17. 32 yrs old female pt comes to derma opd complaining rash which occurs after she takes self prescribed PO medication for her respiratory symptom. the rash is erythematous maculopapular. which one is most likely drug she took.

- A. ampicillin
- B. diclofenac
- C. metronidazole
- D. bisacodyl

18. 45 yrs old, an epileptic pt presented with productive cough, low grade fever, with poor epilepsy control.... chest normal, x-ray multiple cavities and air fluid level..... which one is choice of AB..

- A, Clindamycin
- B, ceftria
- C, metro
- D, penicillin G

19. 33-A 4 day neonate present with failure to pass meconium and abdominal distension, he urinates a clear urine which is multiple times, he has features of down syndrome, no anal opening, Dx?

- A, imperforate Anus with no fistulas
- B, Imperforate Anus with rectovestibular fistula
- C, Imperforate Anus with rectovaginal fistula
- D, Imperforate Anus with perianal fistula

20. G4p3 mother with pushing down pain of 6hr duration, cervix-6cm, station -1 after 4hr evaluations cervix still 6cm, station 0. dx?

- A, protracted cervical dilatation
- B, Arrest of Cervical dilatation
- C, protracted descent

National Licensing Exam for Med - second round November, 2012 E.C

D, Arrest of descent

Part two

21. 4 Yrs old male child presented with new onset of shortness of breath. He has no hx of body swelling, fever, chills. On examination he has clubbing, cyanosis with saturation of 80% at atmospheric pressure. He has follow up for congenital heart disease. Lab result is wbc-7000 with N-70 %, Hg-25mg/dl, with MCV - 71fl,

Which of the following is the best management

A, starting 5mg/kg of iron

B, phlebotomy

C, put on INO2

D,

22. A four yr old male child was brought by his mother with a complaint of decrease hearing

Of voice of four days duration. the mother also told that her child has snoring and opening of mouth during night time for the past two years. on otoscopic examination there is dull, red bulged tympanic membrane bilaterally. What is the diagnosis for this patient ????

(A) Chronic otitis media

(B) Acute otitis media

(C) Cholesteatoma

(D) Glue ear

23. A 60 yr old known hypertensive and type 2 diabetic who was on antihypertensive and oral hypoglycemia agent presented with progressive decrement of his visual function.

Of the following one is LEAST likely cause of his visual impairment???

(A) Cataract

(B) Glaucoma

(C) Macular degeneration

(D) Hypertensive retinopathy

24. A three yr old male child was presented with in 15 minute after a single episode of tonic clonic seizure, on examination T-38.4'c and runny nose. CSF analysis was done with protein of 35mg/dl with no cell seen. What is the most likely diagnosis?

(A) Febrile seizure

(B) Encephalitis

National Licensing Exam for Med - second round November, 2012 E.C

(C) Meningitis

(D) Malaria

25. 3 yr old male child with weight for height less than -3 SD. most likely diagnosis

A) SAM

B) MAM

C)

26. A 23yr old female patient presented with inability to open mouth.14 yr back he had history of facial trauma where he was not treated promptly. he had also characteristic bird's eye appearance

What is the most likely in this patient??

(A) TMJ subluxation

(B) TMJ ankyloses

(C) Myofacial pain syndrome

(D) TMJ dislocation

27. A post partuem women was reffered from district hospital after 15hr labor. then after 1hr of delivery she develope PR-120, BP-70/50, GCS-15/15. What is most likely

(A) Eclampsia

(B) Syncopal attack

(C) PPH

(D)

28. A 10yr old female child presented with itchy crusted scaly skin lesion over the head. she had also history of lymphadenopathy over the neck. The most likely diagnosis

(A) Black dot tinea captis

(B) Gray patch tinea captis

(C) Superinfected tinea captis

(D) Superinfected S.dermatitis

29. HIV positive patient while he was in the ward he develops shortness of breath and cough. he also develops violaceous papule, macule, plaque, vesicle. What is most likely diagnosis??

(A) Histoplasmosis

(B) Kaposi sarcoma

(C) M.tuberculosis

National Licensing Exam for Med - second round November, 2012 E.C

(D) Avium mycobacteria infection

30. A three years old lady presented with white spots on the eye. he had also leukoria, strabismus.....

(A) PVPV

(B) Cataract

(C) Coats disease

(D) Retinoblastoma

31. A girl took self-prescribed drug. For her URTI... and presented with maculo papular rash on her body...most probable drug is...?

- A. aspirin
- B. diclofenac
- C. Ampicillin
- D. Metronidazole

32. a 36 years old female pt with congenital nevus on her eye lid recently noticed increment in size and irregular border.

What malignancy may arise from this lesion

A, marjolin's ulcer

B, squamous cc

C, malignant melanoma

D, me.....

33. a 28 years old primigravida mother peresented to center of labor and delivery with pushing down pain of 12 hours, on P/E vital signs are in normal range, 38 weeks garavid uterus, cervix is 6 centimeter, grade 1 Meconium and being followed by parthograph after 4 hour on PV cervix is still 6 cm, grade 3 molding, and Excessive caput

What is the DX and next mng.t

A, protracted cx dilatation and

B, CPD and c/s

C, arrest of CX dilatation and augmentation

D, arrest of CX dilatation n c/s

34. A 30 yr old p5 mother was reffered from health center who had a diagnosis of polyhydraminos antenatally after One hr of delivery at which time she develops vaginal bleeding.

The most likely cause of her presentation??

National Licensing Exam for Med - second round November, 2012 E.C

A. Cervical tear

B. Uterine atony

C. polyp

35. 34 year old G-III P-II mother from rural area present at GA of 44 weeks from LNMP. GA from obs U/S 36 weeks with RBPP. What is most likely cause for the discrepancy in GA?

A) Male sex

B) wrong dating

C) multiparity

D) Previous history

36. Child on 3rd day of antibiotics for pneumonia. Still febrile, dullness on percussion and decreased air entry, anticipated complication.

A, lung abscess

B, empyema

C, pneumothorax

37. mother on coc..☹. Havin discomfort..mild symptoms..mgt?

A, switch TO...

B, Stop...

C, tx mild sx.....

38. 65 years old male pt presented with a c/c of loin pain and hematuria of 2 weeks duration. V/S BP 160/110, PR 86, T 37.9 U/A;- blood, S/E;- hypocalcemia. Best Dxs is?

A, Pyelonephritis

B, Renal cell Carcinoma

C, Renal failure

thanks to all the AMU Medical Graduates of

2012 E.C who participated in collecting this

Questions. Specially Dr. H. A. and

MEMBERS OF DORM NO-09, BLOCK-303

H@nke 2012

GOD BLESS ETHIOPIA!



National Licensing Exam – Medicine, 2019 G.C



“XANTHRONS” – St. Paul’s Class of 2019

1. A pt in minor OR given LA-lidocaine. In which system is the earliest complication (Toxicity) seen?
A. GIT B. CVS C. CNS
2. A mother of 2 yrs old male child notices the absence of Rt testis in the scrotum and she brought him to the hospital. Upon examination he has an inguinal Rt testis which descend to the scrotum up on manipulation. Dx?
A. Undescended testis B. Retractable testis
3. A case of perforated PUD (Hx & PE). What is the best initial step in approaching this pt?
A. Erect chest x-ray B. Resuscitate with Ns C. Take the pt to OR
4. A case of breast abscess in a lactating mother. Dx?
5. A pt presented to u with a complaint of sensory loss in the superiomedial aspect of the thigh after he had Bassini inguinal hernia repair. Which nerve was injured during the surgery?
A. Genital branch of genitofemoral nerve B. Femoral bran of genitofemoral nerve



National Licensing Exam – Medicine, 2019 G.C

6. A case of internal hemorrhoid. Dx ?
7. A pt came to u after he sustained RTA of 2 hrs back and sustained injury to his head. The pt is conscious with stable V/S and scalp laceration. Immediately after admission, he deteriorates with GCS - 10/15 and hemiparesis. Dx ? **A. AEDH** **B. ASDH** **C. SAH**
8. Which one of the f.f is true about x-ray features of SBO?
- A. Valvulae conventies** **B. Haustral markings** **C. Coffe bean appearance...**
9. A pt started to have carpopedal spasm after he had undergone thyriodectomy. Which of the f.f signs will elicit this? **A. Trousser's sign** **B.**
10. A case of Neonatal sepsis. How would u like to manage this (ampi and Genta - route, dose, frequency and duration)
11. A 7 days old preterm neonate admitted to NICU and after 3rd day of admission V/S - tachycardic, tachypenic and started to develop abdominal distension, hypertympanic to percussion. CBC; WBC - 1,200, Plt-100,000. What is the most likely dx?
- A. HAI** **B. NEC** **C. Sepsis**
12. A 9 yrs old male child presented with easy fatigability of one month duration. Otherwise he has no melena, or any bleeding. His dad claims that he has an optimal nutrition. Hgb- 7 g/dl. What Ix would you like to send? **A. Occult stool exam**
13. A 2 yrs old child with coarse face, unable to speak and umbilical hernia. What initial Ix would u like to send for this pt? **A. Cortisol** **B. TSH & T 4**
14. A child with signs of raised ICP. What is the dose of mannitol that you will order (it doesn't specify; loading vs maintenance)? **A. 0.25-0.5 g/kg** **B. 0.5-1** **C. 1-2**
15. A 12 months old infant with anthropometric measurement of WFL, LFA and HC for age in Z-score results were mentioned. What is the interpretation respectively?
- A. Moderate wasting, Severe stunting, Microcephaly**
16. A neonate born from a mother with BG- A neg. The newborn presented with jaundice with D-bilirubin of 18 mg/dl with BG-B positive. What is the cause of hyperbilirubinemia?
- A. ABO incompatibility** **B. Rh incompatibility**



National Licensing Exam – Medicine, 2019 G.C

17. The earliest time to diagnose RVI in RVI exposed infant with Ix modality.
- A. Antibody test 6th week
B. DNA test at 6th week
C. Antibody test at 9 month
D. Antibody test at 1 8 month
18. A case of RVI exposed neonate. How would you like to manage this neonate?
19. A 5 yrs old child dx with DM with DKA and currently he is out of DKA & u r planning to discharge him. By how much dose of insulin r u going to discharge him?
- A. 0.25-0.5 u/kg/day B. 0.5-0.75 C. 1-1.2
20. A 5 yrs old child presented with sudden onset cough and SOB while he was playing at school. What is ur dx? A. Bronchiolitis B. Asthma
21. Which one of the f.f are components of Basic newborn care?
22. Case - a newborn who is unable to cry immediately after delivery. True about neonatal resuscitation steps?
23. A case about preterm feeding
24. A pt who was taking phenobarbital for seizure d/o currently came to you for follow-up. She has no seizure for the last 8 years. What is the next best step in the mgt of this pt?
- A. Discontinue phenobarb B. Taper phenobarb C. Continue same mgt
25. A pt came to u with abnormal body movement (tonic-clonic type) on one side of the body which subsequently involves the whole body. In which class of seizure d/o do u categorize this pt? A. GTC B. complex partial seizure C.
26. A case of CNS toxoplasmosis. Dose & duration of Rx- cotrimoxazole?
27. Type of anemia on a pt with an Ix; Hgb-9, mcv-73, mch- 24 and RDW- 15. Dose & duration of Rx with iron?
28. A pt with CKD have Hgb-6 and symptoms of anemia. How do you manage it?
- A. Erythropoietin B. Fe C. Blood transfusion
29. A woman from rural area presented with massive bleeding per mouth of 1 day duration. No other pertinent hx. Hgb-8 g/dl. What is the most likely dx?
- A. Variceal bleed B. Leech infestation C. PUD



National Licensing Exam – Medicine, 2019 G.C

- 30.** A pt who was Rtcd for T. Corporis long time ago currently dx with RVI and initiated with HAART for the last 3 weeks but the skin lesion worsen. What is the next step in the mgt of this pt?
- A.** Stop ART
B. Start antifungal
C. Start new regimen of ART
- 31.** Case-IE : 25 yrs old pt with fever, hematuria, splenomegaly. What is the dx?
- 32.** A pt with uncrossed hemiplegia comes to u with sudden onset focal neurologic deficit. What Ix will u send to confirm your dx?
- A.** Brain CT scan **B.** Brain CT with contrast **D.** Brain MRI
- 33.** A case of sagittal sinus thrombosis. Best Ix modality?
- A.** MRI angiography **B.** MRI venography **C.** Brain CT with contrast
- 34.** A pt with rheumatoid arthritis on NSAIDs for 2 years currently presented with decreased UOP and raised cr. What is the possible mechanism?
- A.** Tubular necrosis **B.** Impaired renal autoregulation
- 35.** A case about portal hypertension causes.
- 36.** A case of cirrhotic ascites. How would you like to manage?
- 37.** A case of RVI. When do we say treatment failure and how to manage it?
- 38.** A 60 yrs old patient presented with sudden onset of heaviness in the chest and diaphoresis, no hx of HTN, DM; ECG- showed ST- segment depression on lead..., RBS-174. What is the initial best management option? **A.** Give insulin **B.** Aspirin...
- 39.** A case about nerve injury (nerve supplying Achilles tendon). We were given clinical features and asked the nerve?
- 40.** A spinal nerve root responsible for knee reflex? **A.** L-4 **B.** ...
- 41.** A sero neg pt with sx & symptom of meningitis (no evidence of raised ICP was mentioned). Then the intern did LP and subsequently the pt deteriorate. What is the possible cause of sudden deterioration?
- A.** Brain herniation.... **B.** The Intern was not focusing b/c he was thinking about strike
- 42.** Which of the following results of pleural fluid analysis necessitate insertion of chest tube in a pt with pleural effusion? **A.** PH>7.4 **B.** ...



National Licensing Exam – Medicine, 2019 G.C

43. A pt who was on anticoagulant-warfarin, currently came to OPD for follow-up with coagulation profile; INR-5, no bleeding compliant. What is the best initial step of mgt?
A. Hold warfarin & send INR **B.** Give vit K **C.** Continue same mgt
44. A case of Horner's syndrome with ptosis, dry face and miosis. Dx?
45. A PG lady with GA-1 2 weeks who is a known type-I DM pt came to u for ANC follow-up. In which category of focused ANC should she be classed?
A. Low risk **B.** Basic **C.** Specialized **D.** High risk
46. A 22 yrs old lady who had 2 ex-sexual partners currently presented with Vx discharge. Currently she has a smoker boyfriend and she is taking OCP. What is the most important risk factor for STI in this pt?
A. Having smoker boyfriend **B.** Multiple sexual partner **C.** OCP
47. A PG lady with GA 30 weeks presented with RUQ pain and urine protien +2. What is the dx? **A.** Preclampsia **B.** severe GHTN
48. MG with GDM at 39 weeks of gestation. Her FBS-169 mg/dl. What is the best option of mgt?
A. Immediate Induction **B.** Push the px till 40 weeks **C.** Do an immediate Cs
49. A PG lady with GA-30 weeks came to u with anterior neck swelling and toxic symptoms. Ix TSH-0.03, low; T4-11, high. How do u manage this pt?
A. Lugol's Iodine **B.** Diazepam **C.** PTU **D.** Thyroxin
50. A case of AUB. Dx and mgt?
51. Diagnosis and management of normal and abnormal labor(4/5 qns)
52. A case of bipolar-I d/o. Dx? **A.** BP-I **B.** Cyclothymia
53. A case of MDD d/o Dx? **A.** Persistent DD **B.** Dysthymia
54. A case of anxiety d/o for a pt who presented with excessive fearfulness in gathering activities. How would u like to manage it? **A.** BDZ **B.**
55. A case of ADHD (Attention deficit Hyperactive disorder). Dx?
56. Case upon psychiatric interview the physician said ' I understand that u have suffered a lot'. What does the physician felt to his pt? **A.** Empathy **B.** sympathy...



National Licensing Exam – Medicine, 2019 G.C

57. In a country who has 372 maternal deaths in a year. In the same year, there were 1 00,000 live births and 2500 stillbirths. If there were no pregnancy, how many of maternal deaths can be prevented? **A.** 372/1 00,000 **B.**
58. Doubling time of population growth in a country with CBR - 50/1000 and CDR - 12/1 000, no migration or immigration.
59. A case of Ludwig's angina. Dx?
60. A case of pericoronitis. Dx?
61. A case of parotid tumor (malignant one). What to do at first encounter (as GP)?
A. Referral to specialist **B.** Do excisional biopsy **C.** Do incisional biopsy
62. A case of Basal cell carcinoma- wound with delayed healing and with irregular margin and raised. Dx? **A.** BCC **B.** SCC
63. About localized cutaneous leishmaniasis on Rx with an intralesional antimonial but doesn't respond to it. What is the next option of mgt?
64. A case of Leprosy. Rx?
65. A case about Irritant vs allergic contact dermatitis. Dx?
66. A case of Psoriasis. Dx?
67. A case of T.versicolor, pityriasis. Dx?
68. Fall down accident which result in injury to his eyes....initial best mgt (there were 4 or 5 similar trauma related qns...focusing on best initial mgt options)
69. A case assessing about types of conjunctivitis.
70. A case of chronic otitis media with an ear discharge. Dx?

DORM MEMBERS OF 134

THANK YOU !!!