

45. A 27-year-old schizophrenic patient, who has attempted to kill himself a day before, was brought to a hospital due to his aggressiveness. He has delusional belief and low self-esteem. The episode lasted nine months, but he did not get any treatment.

What is the most necessary criterion consistent with the diagnosis?

- (A) Aggressiveness
- (B) Suicidal attempt
- (C) Low self-esteem
- (D) Delusional belief

46. A male psychotic patient was admitted to a hospital and stabilized with oral antipsychotics. However, he refused to take medication again after he discharged. He had a previous history of drug non-compliance.

What is the most appropriate management in this condition?

- (A) Psycho education
- (B) Depot antipsychotics
- (C) Motivational interviewing
- (D) Advise patient to be re-admitted

47. A 19-year-old woman was brought to a clinic by her husband who reported that she has been showing persistently extreme happiness, grandiosity, distractibility and talkativeness for the last five weeks. She has also hearing of voices in the absence of external stimuli for the last one week.

What is the most likely diagnosis of this woman?

- (A) Brief psychotic disorder
- (B) Bipolar I disorder manic episode
- (C) Bipolar I disorder depressive episode
- (D) Bipolar I disorder manic episode with psychotic feature

48. A 42-year-old woman came to a hospital with a complaint of being worried about many issues for the last six months. As a result, she left her job for the last two weeks. The nurse suspects that she is suffering from generalized anxiety disorder.

What additional criterion is most important to support this diagnosis?

- (A) Difficulty to control worry
- (B) Poor concentration
- (C) Sleep disturbance
- (D) Loss of interest

49. A 35-year-old man came to a psychiatry clinic with complaints of irritable mood and reduced need for sleep for one week. During the interview, he moves frequently from one topic to another, singing and dancing.

What is the most likely thought disturbance experienced by the patient?

- (A) Disorganized thought
- (B) Loose association
- (C) Flight of ideas
- (D) Tangentiality

18. A seven-year-old child was brought to a clinic after thermal burn on the chest, abdomen and left thigh. On arrival, the health care team started fluid administration, cleaning and dressing the wound. However, the child suddenly developed shortness of breathing, edema on the lower leg and rapid and weak pulse.

What could be the immediate nursing management for this child?

- (A) Stop IV fluid
- (B) Increase IV fluid flow
- (C) Put on 100% oxygen
- (D) Relieve the pain using ice pack

19. A two-day-old neonate was brought to a hospital with a complaint of unable to breastfeed of one-day duration. On physical examination, the neonate has brown eyes, yellowish palms and bluish mouth and lips. All neonatal reflexes were active. The nurse has suspected hyperbilirubinemia.

Which finding is the most confirmatory manifestation for this diagnosis?

- (A) Yellow palms
- (B) Bluish mouth
- (C) Brown eyes
- (D) Bluish lips

20. A five-year-old child came to a hospital with a complaint of diarrhea of three days duration. The diarrhea is not bloody, but it is watery, rice water color and has no mucus.

What is the most likely cause of diarrhea in this case?

- (A) Salmonella species
- (B) Vibrio cholera
- (C) Rota virus
- (D) Giardiasis

21. A known HIV positive mother brought her six-month-old infant to a hospital for growth monitoring and checkup. The baby was on exclusive breast-feeding. The nurse assessed the infant and there was poor weight gain. The child's antibody test was positive at six months of age and the DNA PCR test result was pending.

What is the primary indicator to say HIV exposed child in this case?

- (A) Poor weight gain
- (B) Positive antibody test
- (C) Exclusive breast feeding
- (D) Born from positive mother

22. A two-month-old infant was brought to a hospital with a complaint of cough of two days duration. The infant is alert and can feed breast milk. The infant respiration is 50/min, pulse rate is 110/min and temperature is 38 °C. The nurse wants to classify this illness based on IMNCI protocol.

What is the most likely diagnosis of the child?

- (A) Very severe pneumonia
- (B) Severe pneumonia
- (C) Pneumonia
- (D) Cough/Cold

32. A newborn baby was admitted to a hospital due to severe respiratory distress and severe jaundice. Prenatal history shows the presence of hydropsfetalis. On physical examination, there is pallor, hepatosplenomegaly, edema, petechiae and ascites. The baby is also positive for coombs test and hemoglobin value of 9 g/dl.

What is the most likely clinical sign indicating hemolytic disease in this case?

- (A) Hepatosplenomegaly
- (B) Edema and ascites
- (C) Severe jaundice
- (D) Hydropsfetalis

33. A nurse had been assessing a mother on breast-feeding clues for adequate breast-feeding. On observation, the infant sucked five times before pausing and gurgle sound was heard during swallowing.

What is the most important additional finding that can show the adequacy of breast-feeding?

- (A) Chin of the infant should touch the breast
- (B) The infant's mouth should be widely open
- (C) More areola seen on the upper part of breast
- (D) Dimpling of cheeks may be seen while suckling

34. A 10-month-old child with clinical features of severe wasting, edema of lower extremities and palmar pallor came to a hospital. The nurse admitted and put the child on phase one management of severe acute malnutrition.

What is the priority nursing intervention for this child?

- (A) Iron folate
- (B) Deworming
- (C) IV fluid infusions
- (D) Folic acid supplement

35. A child with complicated severe acute malnutrition was on treatment in the inpatient treatment program of severe acute malnutrition. On admission, the child had severe wasting, pitting edema and failed the appetite test. However, the child did not respond to treatments and classified under failure to treatments.

What is the most likely indicator of failure to treatment the nurse assessed?

- (A) Failure to lose edema on day 10
- (B) Failure to regain appetite on day 10
- (C) If edema persists for more than 10 day
- (D) Failure to gain at least 5 g/kg bodyweight in day four

68. A psychiatrist wished to examine the influence of strong family support on reducing the suicidal death with acute schizophrenia after hospital discharge. He selected 145 series of schizophrenic patients who had family support and 435 series of schizophrenic patients who had no family support. After one year, the researcher reported 12 suicidal deaths among schizophrenic patients who had family support and 18 suicidal deaths among schizophrenic patients who had no family support.

What is the most likely study design used by the researcher?

- (A) Cohort study design  
(B) Ecological study design  
(C) Case control study design  
(D) Cross-sectional study design

69. A hospital head nurse wishes to know satisfaction level of patients and the proportion of discharged patients who received during hospitalization. The nurse wants to calculate the sample size by using 95% confidence interval and 5% margin of error.

How large should be the sample size?

- (A) 324  
(B) 344  
(C) 384  
(D) 394

70. A health researcher wants to study the impact of folic acid supplementation on pregnancy outcome and its side effect among pregnant mothers. Before the actual data collection, the researcher planned to collect data on 5% of the sample size in another area to improve the sequence and layout of the questionnaire and to know the total time required for the interview.

What is the most likely data quality control method used by of the researcher?

- (A) Training  
(B) Pretesting  
(C) Pilot study  
(D) Supervision

71. A researcher interested to conduct a study on knowledge and practice of farmers in a district on COVID-19 prevention mechanism. The researcher developed a data collection tool appropriate for the farmers who cannot read and write.

What is the most likely data collection method used by the researcher?

- (A) Record review  
(B) Direct observation  
(C) Face to face interview  
(D) Self-administered questionnaire

72. A nurse, who is working in a health center at adult OPD, wants to know the age distribution of patients at his service. The age of adult patients who visited the OPD in a day is 35, 60, 52, 33, 25, 22, 70, 66, 45 years.

What is the mean age of these patients?

- (A) 36.3 years  
(B) 39.3 years  
(C) 40.3 years  
(D) 45.3 years

54. A nurse working in a school health office is planning to design substance use prevention plan as part of routine school health services. During preliminary assessment in the school, many young students use *khat* because of abundant availability of it in the surrounding areas.

What is the most recommended effective measure of prevention of *khat* use in this school?

- (A) Push policy makers to impose sanction by researching the health impact
- (B) Work with community police to clear school surrounding from *Khat* shops
- (C) Discuss with the school administration to impose serious punishment on users
- (D) Teach the students and the community around the school regarding its health impact

55. A home nurse evaluates a 70-year-old male during her routine weekly home visit. His families are concerned that sometimes he forgets appointments and have difficulty to find things that he places. Except recent onset of forgetfulness, he is doing well.

What is the most likely diagnosis of this patient?

- (A) Aging
- (B) Dementia
- (C) Amnesia
- (D) Depression

56. A seven-year-old boy came to a health center with his mother with a complaint of language difficulty. He lacks interest in interacting with other children and often prefers to stay with nonliving materials. On examination, he flaps his finger and is having difficulty to show his emotion to parents.

What is the most appropriate intervention for this child?

- (A) Consult with specialist
- (B) Teach simple self-care skills
- (C) Send him to special need school
- (D) Anticipate and fulfill the child's needs

57. A head nurse of a hospital collected and analyzed data about the services, staff shifting and inventory of medical materials. This procedure is done routinely in different departments to prevent unwise usage of organizational resources.

What is the most likely management concept applied by the head nurse?

- (A) Efficiency
- (B) Effectiveness
- (C) Work relations
- (D) Economy of scarce resources

58. The ministry of health of a country in partnership with stakeholders had prepared annual research conference. The conference is aimed at promoting a joint concern and responsibility of public offices to identify problems and undertake tasks that have importance for the health and wellbeing of the community.

Which principle of primary health care was the focus of the ministry in the case scenario?

- (A) Decentralization
- (B) Community participation
- (C) Focus on preventive services
- (D) Intersectoral collaboration

40. A nurse is providing vaccine for children. She checks the health status of children prior to administering the vaccines. She has remembered that presence of some clinical conditions prohibit administering vaccines.

What is the most likely contraindication to be ruled out by the nurse prior to vaccination?

- (A) Malnutrition
- (B) Symptomatic HIV infection
- (C) Chronic non-communicable diseases
- (D) Family history of allergy for vaccines

41. A nurse working in an immunization unit is responsible to maintain the potency of vaccines. To do so, monitoring vaccine refrigerator temperature is one important mechanism.

What has to been done to maintain the potency of vaccines?

- (A) Measuring vaccine refrigerator temperature daily for seven days
- (B) Measuring vaccine refrigerator temperature twice a day for seven days
- (C) Measuring vaccine refrigerator temperature once per day in working days
- (D) Measuring vaccine refrigerator temperature twice per day in working days

42. A mother brought her seven-week-old child to an immunization clinic. The nurse assessed the overall clinical condition of the child and decided to give vaccines accordingly. The child did not take any vaccine because the mother gave birth at home.

What are the most appropriate vaccines that the nurse can give for this child?

- (A) BCG, rota 1, PCV1, and Penta 1, measles
- (B) OPV 1, BCG, rota 1, PCV1, and Penta 1
- (C) OPV 1, IPV, BCG, rota 1, PCV1, and Penta 1
- (D) OPV 0, OPV 1, BCG, rota 1, PCV1, and Penta 1

43. A nine-year-old child, who was recently diagnosed with hearing impairment due to medication side effect, was brought to HEENT clinic. Despite proper medication treatment of the problem, the child's hearing is limited that he is unable to identify the meaning of words.

What is the most likely nursing diagnosis for this case?

- (A) Impaired psychological status
- (B) Impaired activities of daily living
- (C) Impaired communication skill
- (D) Impaired daily activities

44. A six-year-old child was brought to a hospital for physical rehabilitation due to apparent physical disability. On assessment, the nurse identifies that the child has bowed legs, delayed denotation and a history of gross motor developmental delay.

What is the most likely cause of the disability in this case?

- (A) Iron deficiency
- (B) Calcium deficiency
- (C) Vitamin D deficiency
- (D) Phosphorus deficiency

99. A nurse working in an ICU always enters earlier to assess the order and revises cares for all patients. One day, the nurse gets wrong order and doubtful medication for her critically ill patient. Then, the nurse immediately consults the doctor who ordered the medication before she carries out the medication.

What is the most likely reason that the nurse wants to consult the doctor?

- (A) Fears not to take accountability for any bad consequence
- (B) Lacks confidence to work independently as professional nurse
- (C) Since he is newly assigned, he is not clear with ICU procedures
- (D) To prevent mal practices that risk the patients and to safeguard them

100. A male nurse has been working in medical ward where there were patients who need close follow up. The nurse always tries to listen and understand the patient's need and perspective. He also feels patients' emotion in his daily patient care without being tired. The patients feel happier every time they see this nurse than other nurses in the wards.

What is the most likely reason that this nurse is preferred than other nurses in the ward?

- (A) He gives compassionated care
- (B) He uses good communication skill
- (C) He has good knowledge on patient care
- (D) He has good skillful nurse on patient care

# THE END

4. A woman using a combined oral contraceptive come to a family planning clinic with complaints of nausea, mood changes and less sex-drive after taking this contraceptive. The woman said that she is gaining weight and she wants to change to other contraceptive method.

What is the most likely sign that indicates the side effect of the contraceptive?

- (A) Nausea
  - (B) Weight gain
  - (C) Mood change
  - (D) Decreased sex-drive
5. A 35-year-old woman was admitted to a gynecology ward with vesicovaginal fistula. The nurse identified that she has been isolated from the society and humiliated because of her urine incontinence. Laboratory investigation results show HCT level of 30%, urine chemistry and microscopy are in normal range.

What is the priority nursing diagnosis of this woman?

- (A) Acute confusion
  - (B) Sexual dysfunction
  - (C) Post-trauma syndrome
  - (D) Disturbed personal identity
6. A 45-year-old mother came to a hospital with the complaints of bearing down sensation, dragging discomfort and low back pain. She was admitted to a gynecology ward for a diagnosis of second-degree uterine prolapse. On pelvic examination, soft and reducible mass bulging into the vagina and distending through vaginal introitus. She is now scheduled for surgical procedure.

What is the most appropriate nursing care for this mother at this time?

- (A) Teach about patient self-care
  - (B) Advise the mother about exercise
  - (C) Physical and psychological preparation
  - (D) Prescribe and administer sedative drugs
7. A 43-year-old woman came to a hospital with a complaint of vaginal bleeding for two months duration. On assessment, she has bleeding after sexual intercourse and profuse vaginal discharge. On pathological examination, there was stromal invasion of the cervix 3 mm in depth and horizontal extension of 5mm.

What is the most likely stage of the cervical cancer?

- (A) Stage-IB<sub>2</sub>
- (B) Stage-IB<sub>1</sub>
- (C) Stage-IA<sub>2</sub>
- (D) Stage-IA<sub>1</sub>

60. A head of health center discussed the bottlenecks of health care insurance service utilization with community leaders who came from the catchment *kobele*. During discussion, the community leaders urged the health center head to address frequent hospital supply stock-out issues.

What is the most likely problem prioritization criterion addressed by the head?

- (A) Feasibility
- (B) Magnitude
- (C) Sustainability
- (D) Community concern

61. A staff recruitment committee in a hospital wants to develop human resource planning. During the course of planning, the committee anticipates changes in the present work force in terms of entry of new professionals and departures of the employees for different reasons.

What is the most appropriate step of human resource planning done by the committee?

- (A) Profiling
- (B) Estimating
- (C) Forecasting
- (D) Inventorying

62. A matron nurse who works as a member of a hospital management wants to increase the nursing service performance for the coming year. Likewise, he conducted a meeting with nurses from different wards to deal with achievements and failed activity based on the previously designed objective. Finally, the matron designed a new plan based on the feedback of the report.

What is the most likely principle of management used by the matron?

- (A) Division of labor
- (B) Management by objective
- (C) Learning from experience
- (D) Management by exception

63. A matron nurse of a hospital assigns some of the activities to one of the ward head nurses to control, coordinate the work, and give feedbacks. However, the matron repeatedly dictates the ward head nurse not to solve problems own her own.

What is the **MISSED** characteristic of effective delegation by the matron?

- (A) Not taking back delegated tasks
- (B) Giving the autonomy and freedom
- (C) Training and developing subordinates
- (D) Following up by visiting subordinates

64. In a health center, conflict arisen between OPD case team staffs on the issue of managing patients with COVID-19 infection. However a, head of the health center wants to resolve the conflict by considering assertiveness and cooperation between parties.

What is the most likely conflict resolution strategy used by the head?

- (A) Avoidance
- (B) Competition
- (C) Collaboration
- (D) Accommodation

BOOKLET CODE: 012

27. A neonate, who was diagnosed with neonatal sepsis, was admitted to a hospital. The nurse in charge examined the baby and the findings showed central cyanosis, chest in drawing, tachycardia and tachypnea.

What is the most likely nursing diagnosis of this case?

- (A) Ineffective breathing pattern
- (B) Ineffective airway clearance

- (C) High risk for hypoventilation
- (D) Ineffective tissue perfusion

28. A 35-week-old neonate was admitted to a hospital for a diagnosis of neonatal jaundice. Baseline temperature is  $36.2^{\circ}\text{C}$  and pulse rate is 120/min. Currently, the neonate's random blood sugar is 35 mg/dl. Then, the nurse diagnosed hypothermia and treated the neonate accordingly.

What is the most likely risk factor for hypothermia of this neonate?

- (A) Being preterm
- (B) Having hypoglycemia

- (C) Adverse effect of phototherapy
- (D) Undressing it for phototherapy

29. A 34-week-old neonate, with a diagnosis of neonatal jaundice and sepsis, was admitted to a NICU. The neonate has difficulty of feeding. His temperature is  $36.2^{\circ}\text{C}$ , pulse rate is 120/min and random blood sugar is 34 mg/dl.

What is the most priority nursing intervention for this neonate?

- (A) Treat hypothermia
- (B) Treat hypoglycemia

- (C) Treat neonatal sepsis
- (D) Treat neonatal jaundice

30. A two-day-old preterm male neonate with neonatal conjunctivitis was admitted to a NICU. Physical examination shows that he is irritable with chest in drawing. The neonate has an expiratory grunt and his lips started to change to blue. His pulse rate is 170/min, respiratory rate is 80/min and temperature is  $36.9^{\circ}\text{C}$ .

What is the most appropriate initial step in the management of this patient?

- (A) Advice on oral feeding
- (B) Administer maintenance fluid

- (C) Administer supplementary oxygen
- (D) Maintain the temperature range

31. A newborn baby, with a clinical manifestation of respiratory distress, was admitted to a hospital. Prenatal history shows that there was fetal stress as manifested by Meconium stained amniotic fluid. Central cyanosis and chest in drawing were the pertinent findings during physical examination. Vital sign shows that the child's pulse rate was 170/min and the respiratory rate was 80/min.

What is the most likely complication this infant may develop?

- (A) Meningitis
- (B) Jaundice

- (C) Shock
- (D) Anemia

assessment, she has been  
diagnosed her as having breast cancer and admitted to

What is the priority nursing intervention for this mother?

- (A) Giving analgesics
- (B) Giving cytotoxic agent

- (C) Advice to take caffeine
- (D) Aspiration of cysts

9. A toddler was brought to an emergency department after an initial first aid management of drowning incident. On arrival, the child was shivering, grunting and had severe intercostal muscle retraction with cyanosis on lips. The nurse has confirmed that the child is in respiratory distress.

What is the most likely nursing diagnosis of this child?

- (A) Risk for aspiration
- (B) Risk for hypothermia

- (C) Ineffective airway clearance
- (D) Ineffective breathing pattern

10. A child was admitted to a hospital after Salicylate poisoning with convulsions, dehydration and hyperventilation. The child was managed with lavage and charcoal, 3 ml/kg/hr isotonic fluid, 2 mEq/kg/h of sodium bicarbonate and 30 mEq/L  $K^+$  over the first six hours and the ward nurse assessed the status of the child.

What laboratory value most likely indicates the progress of the child?

- (A) Low serum  $K^+$  values
- (B) Urine pH of greater than 7.0
- (C) Urine flow of 1 mL/kg/h established
- (D) Laboratory value indicating metabolic acidosis

11. A 12-year-old child came to a clinic with complaints of swelling of knees, ankles, and elbows joints. On examination, the child had hot, red, tender and migratory swelling. The nurse suspected acute rheumatic fever.

What additional major manifestation is needed to confirm the diagnosis?

88. A nurse, who works as public health emergency surveillance expert, solicits the reports of disease from different health facility and centrally organizes the information and writes a report.

What type of surveillance system is manifested in the above case scenario?

- (A) Passive surveillance system
- (B) Active surveillance system
- (C) Sentinel surveillance system
- (D) Cross-over surveillance system

89. A nurse was giving wound care on BID base for a 24-year-old male patient. She explained the rationale of the care to the patient. However, the patient refused the care plan and requested the nurse to give him only medication and wanted to go home. Though the nurse tried to explain more, the patient insisted to refuse. The nurse took informed consent, informed to concerned body and cancelled the planned wound care.

Which ethical principle did the nurse apply in this case scenario?

- (A) Non-maleficence
- (B) Beneficence
- (C) Autonomy
- (D) Justice

90. A nurse in a labor ward attended nine deliveries during his night shift and became exhausted. In the morning, a new client came with a complaint of pushing down pain of three-hour duration. The nurse told the woman to wait until the next shift starts because he was tired. He then left the ward because his shift was over while the woman was requesting for help due to sever back pain.

What is the type of compassionate quality that the nurse **MISSED** to show?

- (A) Caring
- (B) Sympathy
- (C) Being sensitive
- (D) Distress tolerance

91. A patient with falling accident visited a private clinic and a nurse diagnosed the case, as it was incomplete fracture and informed the patient that it could be managed in the clinic. The patient accepted the nurse's decision. The nurse immobilized the fractured hand, gave anti pain, counseled home management principles and told the patient to come back after three days. After one day, the patient came to the clinic with complications and he accused the nurse. The patient was sent to x-ray and the result confirmed that the patient had a complete closed radial bone fracture.

What is the most likely cause of professional malpractice in this case?

- (A) Practicing beyond scope
- (B) Having much moral value
- (C) Lack of diagnostic material
- (D) Negligence during practice

Ethiopian Health Professionals' Licensure Examination

Nursing Exam  
Nehase 2014/August 2022  
PART TWO

BOOKLET CODE: 012  
Number of Items: 100

SUBJECT CODE: 02  
Time Allowed: 2:30 hours

**DIRECTIONS:** Each of the following questions is followed by four possible alternative answers. Read each question carefully and **blacken** the letter of your best choice on the separate answer sheet provided.

1. A woman came to a family planning unit with complaints of vomiting and severe migraine for five days. She was using combined oral contraceptive for the past two months. Her blood pressure is 110/70 mmHg, pulse rate is 82/min, and temperature is 36.4°C.

What is the likely nursing diagnosis for this woman?

- (A) Risk for shock  
(B) Risk for infection  
(C) Risk for acute confusion  
(D) Risk for electrolyte imbalance

2. A 24-year-old woman came to a hospital to utilize a contraceptive method. The nurse provided counseling regarding the different contraceptive options available and their side effects. The nurse identified that the woman had understood their discussion. The woman showed a desire to use COC.

What most likely response of the woman that shows her understanding?

- (A) She decides to change her idea ✓  
(B) She takes another appointment  
(C) She lists contraceptive methods  
(D) She asks a question about pregnancy

3. A 30-year-old woman came to a family planning room to use contraceptive. The health care provider in the unit counseled her about methods, side effects, effectiveness and time to return. Finally, she chose to take depo-Provera.

What is the most appropriate site to administer this contraceptive?

- (A) IM on the thigh  
(B) IM on the abdomen  
(C) IM on the right side deltoid muscle  
(D) IM on the left side deltoid muscle ✓

92. A nurse saw a researcher who was collecting data from an admitted patient during the COVID-19 pandemic time. The researcher had ethical approval letter to conduct the research. On observation, the researcher was sitting close to the patient without wearing facemask and he was taking blood sample without minimum recommended precautions.

How should the nurse manage this case?

- (A) Advocates for patient and restrain data collection
- (B) Let the researcher to continue data collection
- (C) Assist the researcher in collecting data
- (D) Report the case to the management

93. The morning shift nurse left the ward by giving and recoding the 8 o'clock medication earlier. The afternoon shift nurses checked the patient card and found that the medications were administered earlier than the schedule.

What is the most appropriate reason to accuse the morning shift nurse?

- (A) Negligence
- (B) Breach of duty
- (C) Omission of care
- (D) Commission of care

94. A newly graduated female nurse planned to start professional services and applied to a private hospital. The human resource of the hospital asked her to bring the evidence that indicates legal permit that the government agency grants her to practice as a BSC nurse.

What is the most likely requirement the nurse must fulfill?

- (A) Accreditation evidence
- (B) Registration evidence
- (C) Certification evidence
- (D) License evidence

95. A 15-year-old adolescent patient was admitted to a surgical ward with a second-degree burn. While the nurse was giving wound care, the patient cried and tried to guard due to pain and the nurse insulted him. The nurse also said "Be careful, if you cry and try to guard me again, I will hit you, I will inject a syringe to your mouth, I will discharge you and you will die". After three hours, other patients reported the case to the ward head nurse. The ward head nurse had other similar record on the nurse for which he advised but no change.

What is the most appropriate next step that the head nurse should do?

- (A) Reported to concerned body as the nurse committed malpractice
- (B) Reported to concerned body as the nurse committed negligence
- (C) Reported to concerned body as the nurse committed battery
- (D) Report to concerned body as the nurse committed assault

23. A two-year-old child was brought to a hospital with a complaint of cough of two days duration. The child has poor feeding, inspiratory stridor, and convulsion during examination and nasal flaring. The child's respiratory rate is 50/min, pulse rate is 110/min and temperature is  $37.5^{\circ}\text{C}$ .

What is the most appropriate nursing management for this child?

- (A) Administer 0.5 mg/kg Diazepam injection solution per rectum
- (B) Administer first dose of IV/IM Ampicillin and Gentamycin
- (C) Administer oral Amoxicillin for five days
- (D) Administer maintenance fluid IV

24. A two-year-old child was brought to a hospital with a complaint of diarrhea of three days duration. The diarrhea is watery, but not bloody. The child has sunken eyes and was not able to drink, skin pinch goes back within four seconds, and weight for length z-score is  $< -3$  standard deviation.

What is the most appropriate drug management for this child?

- (A) Administer 75 ml/kg ORS for four hours
- (B) Administer 5 ml/kg 5% glucose solution for two hours
- (C) Administer 5 ml/kg ReSoMal solution per 30 minutes two hours
- (D) Administer 100 ml/kg IV Ringer's lactate solution for three hours

25. A known HIV positive woman gave a birth for her first baby. She was not on follow up and the infant was exposed to HIV which was identified through infant HIV antibody testing. The infant is on breastfeeding and the nurse has anticipated the case and intended to give proper care to prevent the infant from actual infection.

What is the most appropriate action that the nurse should take in this case?

- (A) Provide Abacavir prophylaxis for the baby
- (B) Provide Nevirapine prophylaxis for six week
- (C) Recommend mother to stop breast feeding immediately
- (D) Provide Stavudine prophylaxis for the baby for six month

26. A nurse in the PMTCT unit has prepared a nursing care plan to provide a holistic care to HIV exposed infants. A mother planned to stop breast-feeding today following the advice from the nurse. On the follow up visit schedule, the nurse planned to do a rapid antibody test.

When is the best time to do the antibody test in this case scenario?

- (A) At birth
- (B) At 6 weeks
- (C) At 10 weeks
- (D) At 12 weeks

36. A three-year-old child has been admitted due to severe acute malnutrition. The child has cough of six days duration. Physical examination during admission shows that the child has bilateral pitting edema (++), bulged abdomen, and skin lesion (++). The child has been on treatment for severe acute malnutrition. At six day of admission, the child has, good appetite edema decreased (+) with no IV or NG tube.

What will be the progress of a child based on the current situation?

- (A) Should stay in phase one
  - (B) Transfer him to transition phase
  - (C) Based on the progress should be phase two
  - (D) Transfer him to outpatient therapeutic program
37. A child has been brought to a hospital with a complaint of diarrhea for five days duration. The nurse wants to assess the growth and development status of the child. The anthropometric result shows weight for height  $< -1$  standard deviation, mid upper circumference (MUAC) is 12 cm, height for age is  $< -2$  standard deviation and weight for age is  $< -1$  standard deviation.

What is the most appropriate interpretation of the child's anthropometric measurement?

- (A) Under weight
  - (B) Moderate wasting
  - (C) Acute malnutrition
  - (D) Chronic malnutrition
38. An 18-month-old child, who was brought to a hospital for growth checkup, was assessed for developmental status. The nurse has identified that the child can walk alone, crawls up stairs, makes tower of 4 cubes, speaks an average 10 words, and can feed himself. From assessments, the nurse has noticed that the child has red flags of under development.

What is the most likely delayed development domain noticed by the nurse?

- (A) Social
- (B) Motor
- (C) Language
- (D) Emotional

39. A nurse who is working in the EPI unit, wanted to keep the potency of vaccines through appropriate storage in fridge. Then, the nurse plan to put vaccines orderly based on their level of sensitivity putting heat sensitive vaccines near to the ice in the fridge and freezing sensitive vaccines away from it.

Which vaccines are placed away from the heat sources?

- (A) OPV
- (B) Measles
- (C) Penta
- (D) Rotavirus

83. A hospital head nurse, who is working in a district, observes that there are a number of pregnant street children around the district. Therefore, she decides to open a mobile health center around the area where they live to give family planning services.

What strategy is taken by the nurse to solve the problem of the street children?

- (A) Making health care facilities available
- (B) Provision of Street based health education
- (C) Looking for possibilities to reunite with their families
- (D) Providing medication so that they will live healthy life

84. A nurse wanted to intervene on a family who lost their son due to road traffic accident while walking to school. During assessment, the nurse asked how the family handled the conflicts, perception and response to stressors.

What is the most likely positive outcome of behavior change in this scenario?

- (A) Family function
- (B) Family coping pattern
- (C) Family structure and role
- (D) Family communication pattern

85. A health center head assigned a nurse to a school to conduct scabies screening among the school students. During the screening, the nurse identified a number of scabies cases.

What is the most likely rationale of conducting screening among school students?

- (A) To detect and treat the disease early
- (B) To increase basic knowledge and skill of children
- (C) To develop right attitude towards health and illness
- (D) To increase children's use of educational facilities

86. In a district, a conflict happened between individuals on an issue. Unfortunately, the conflict aggravated and led to loss of lives and damaged large amount of properties.

What is the most likely type of disaster in this scenario?

- (A) Mixed disaster
- (B) Natural disaster
- (C) Epidemic disaster
- (D) Manmade disaster

87. A nurse works as surveillance team member of a national COVID-19 prevention and control. The team detected COVID-19 case on each day and reported immediately to the ministry of health.

What is the most likely good feature of the surveillance done by the team?

- (A) The team had comprehensive action
- (B) The team had timely reporting of cases
- (C) The team used combined passive and active mechanisms
- (D) The team had strong laboratory services for accurate diagnosis

50. A 60-year-old woman with depression has started initial treatment of Amitriptyline 75 mg/day. After two weeks of treatment, she reports dry mouth, blurring of vision and constipation. She has no history of substance use.

What is the most likely cause for these symptoms?

- (A) Toxic effect of her medication
- (B) Cardiovascular effect of her medication
- (C) Anticholinergic effect of her medication
- (D) Leftover symptoms of under treated depression

51. A 35-year-old married woman came to a hospital with complaints of numbness and throat tightness, which began within two months after her husband died unexpectedly. Every time she sees his picture, she cries a lot but she is able to maintain her functioning. Symptoms do not persist beyond those two months.

What is the most likely diagnosis of this woman?

- (A) Bereavement
- (B) Anniversary reactions
- (C) Traumatic bereavement
- (D) Complicated bereavement

52. A 25-year-old male patient came to a hospital with complaints of sad mood, difficulty of falling asleep, and problem of concentrating on his daily activities for the last 15 days. He reported that he had been, tested HIV positive one month ago. The patient also told the nurse about having an idea to end his life. The nurse planned to improve the patient's mood.

What is the most appropriate indicator that shows a reduced self-harm behavior?

- (A) Develop an honest, nonjudgmental relationship with the patient
- (B) Becoming conscious with the painful memories and seems coping
- (C) He is seeking emergency assistance when he is in immediate danger
- (D) Family members become aware of the disorder and treatment measures

A 46-year-old man, with a history of excessive alcohol drinking for 10 years, was hospitalized because of seizure and increased hand tremor. Symptoms follow three days cessation of alcohol as part of his repeated unsuccessful effort to stop. He has no identified medical illness.

What is the most likely admission diagnosis for this patient?

- (A) Alcohol abuse
- (B) Alcohol dependency
- (C) Alcohol withdrawal
- (D) Alcohol intoxication

64. A nurse, who works in a nursing association, thinks beyond the horizon, influences constituents beyond jurisdiction, puts heavy emphasis on development of vision, copes with changes and inspires young nurses.

What is the most likely role-played by the nurse?

- (A) Management
- (B) Leadership

- (C) Governance
- (D) Administration

65. A comprehensive specialized hospital formed an independent working committee to check service quality. The committee made a series of discussions to encourage staffs, to have shared goals and objectives during their visits. However, the top performing professionals were not credited for their contributions.

What is the most likely principle of supervision NOT considered by the hospital?

- (A) Sharing the overall goals
- (B) Creating good communication
- (C) Respecting your staff and the give praise
- (D) Rewarding those with better performances

66. A health science college dean and department heads of a teaching hospital made discussions to determine the goals of the organization. They started by clarifying the risk and the extent of which to confidence and reliability have been developed for the organizational aim.

What step of evolution was done in this situation?

- (A) Assessing the evaluation situation
- (B) Developing indicator for evaluation
- (C) Gathering and analysis of data on evaluation
- (D) Using evaluation results for learning and action

67. A research methodology instructor prepared an assignment for graduating class students to write proposal on identified problem. Utilization of ANC is one of the lowest performance of the health services in a district.

What is the most likely objective of the proposal?

- (A) Study the reasons for low utilization of ANC at the district
- (B) Identify the reasons for low utilization of ANC at the district
- (C) Appreciate the reasons for low utilization of ANC at the district
- (D) Understand the reasons for low utilization of ANC at the district

73. A nurse is part of a surveillance team at an area where acute watery diarrhea is epidemic. He wants to observe the number and progress of the cases during the epidemic period by using diagrammatic data presentation methods.

What is the most likely diagrammatic data presentation method to be used by the nurse?

- (A) Bar chart
- (B) Pie chart
- (C) Line graph
- (D) Scatter diagram

74. Graduating class students were conducting community health assessment in a district. The students collected information on age-sex distribution of the community. Based on the information collected, they drew population pyramid, wherein the base and the top of the pyramid illustrate wider, and the center is narrow.

What is the most likely interpretation of the population pyramid?

- (A) There is fluctuations of fertility
- (B) There is age-selective migration
- (C) There is a large proportion of productive age
- (D) There is a good economic state of the community

75. A nurse was working in a district with catchment population of 30,000. That year, there were 600 women in child bearing age and from them, 240 gave live birth and 12 infants died within 28 days of birth and 24 died before they celebrate their 1<sup>st</sup> birthday.

What is the infant mortality rate in this scenario?

- (A) 20 infants died per 1000 live births
- (B) 24 infants died per 1000 live births
- (C) 50 infants died per 1000 live births
- (D) 100 infants died per 1000 live births

76. A nurse participated in a curriculum development of nursing specialty with other nursing professionals who came from different institutions. The curriculum development team decided that final exam, mid exam, quiz, individual and group assignments with presentation are the mechanism by which the instructor of a specific course should use to decide the students grade level.

What is the most likely component of a curriculum listed by the nurse?

- (A) Strategy to ignite learning
- (B) Delivery method of the course
- (C) Evaluation method of the course
- (D) Recourse needed for the course

77. An infection prevention team wants to use a waste management technique in order to reduce the quantity as well as to destroy the microbial content of infectious waste from the hospital.

What is the most likely waste disposal technique used by the team?

- (A) Incineration
- (B) Land filling
- (C) Autoclaving
- (D) Compaction

96. A hospital nursing director always tells the nurses that nurses are always responsible to give all necessary care for dying patients and the patients should get peaceful death irrespective of their background and the possible outcomes.

What is the most likely rationale of the director?

- (A) It is the nurses' professional responsibility to give care for all patients equally
  - (B) Nursing director is compassionate person and planned to help all patients till death
  - (C) It is the nurses' professional responsibility to give dignified care for all human beings
  - (D) Nursing director intended to decrease the possible family's complaint after patient death
97. A hospital nursing director has his own daily and weekly evaluation of nursing services with the hospital nursing supervisors. During their supervision and evaluation, they identified many gaps. The nursing performance was found to be below the expected. Few diploma nurses were performing advanced nursing procedures, beyond their scope. Generally, the nursing director and supervisors concluded that the nursing services in the hospital are below the expected implying the need for a training.

What is the most appropriate training area needed for this hospital nursing staffs?

- (A) Standard of practice
  - (B) Attitude and perception
  - (C) Nursing code of ethics
  - (D) Nursing knowledge and skill
98. A 23-year-old pregnant client came to a health center for ANC follow up. She complained perianal pain. There were two nurses and two examination beds in one ANC room. The two nurses were working in parallel for different clients at the same time. The pregnant woman asked the female nurse to examine her. Unfortunately, the second client came and the male nurse started examination side by side. Then, the pregnant woman with perianal pain told the nurse as she came only for routine ANC follow-up and returned home without treatment for perianal pain.

What is the most likely reason for the pregnant woman to return home without treatment?

- (A) Her main need was ANC follow up
- (B) The environment was inconvenient
- (C) She wants to be examined by male nurse
- (D) She was not satisfied with the nurse communication approach

78. A nurse who is works in scabies prevalent community wants to design a different means to prevent the disease in the community in collaboration with stakeholders.

What is the most appropriate prevention strategy to be used by the nurse?

- (A) Avoidance drinking of contaminated water
- (B) Discharging the accumulated water from the area
- (C) Provision of adequate water for personal hygiene
- (D) Prevention swimming and crossing in the infected water

79. A 29-year-old woman came to a health center to deliver a baby. Then, she delivered a stillbirth female child. On physical examination, the nurse showed the mother neuroskeletal abnormality signs from the stillbirth baby.

What is the most likely nutritional problem that the mother faced?

- (A) Iodine deficiency
- (B) Calcium deficiency
- (C) Vitamin A deficiency
- (D) Iron foliate deficiency

80. A prison administrator in a *woreda* wrote a letter to a health office requesting for health professional to inspect the prison facility. A nurse was assigned to inspect, and she found out that overcrowded living condition, shortage of water and bedding and prevalent cases of relapsing fever were the most common problems at the prison.

What is most likely mechanism of solving the health problems of the prison?

- (A) Treating the sick
- (B) Latrine construction
- (C) Proper waste disposal
- (D) Delousing of the prisoners

81. A nurse works as a nutrition expertise in a health center. In the catchment area of the health center, many mothers were not feeding their children properly because they are busy in their day-to-day activities. Therefore, the nurse wants to assess the nutritional status of the children by measuring the physical dimensions and gross composition of the children's body.

What is the most likely nutritional assessment method that the nurse uses?

- (A) Dietary method
- (B) Clinical method
- (C) Biochemical method
- (D) Anthropometric method

82. A nurse came across different cases where the grooms severely beat their very young brides and drag them out of houses. Then, the nurse discussed the issue with local leaders and noted that since most girls get married early and if a girl is not virgin, she will be considered as unfit and she will disgrace her family.

What is the most likely cause of child abuse in this scenario?

- (A) Social norm
- (B) Family conflict
- (C) Economic disparity
- (D) Poor function of social institution

12. A newborn baby was admitted to a hospital due to severe respiratory distress and anemia. Prenatal history shows the presence of hydrops fetalis. On physical examination, there is pallor, hepatosplenomegaly, edema, petechiae and ascites. The baby is also positive for coombs test and hemoglobin value of 9 g/dl.

What is the most likely clinical sign indicating hemolytic disease in this case?

- (A) Hepatosplenomegaly
- (B) Edema and ascites
- (C) Severe jaundice
- (D) Hydrops fetalis

13. A nurse had been assessing a mother on breast-feeding clues for adequate breast-feeding. On observation, the infant sucked five times before pausing and gurgle sound was heard during swallowing.

What is the most important additional finding that can show the adequacy of breast-feeding?

- (A) Chin of the infant should touch the breast
- (B) The infant's mouth should be widely open
- (C) More areola seen on the upper part of breast
- (D) Dimpling of cheeks may be seen while suckling

14. A 10-month-old child with clinical features of severe wasting, edema of lower extremities and palmar pallor came to a hospital. The nurse admitted and put the child on phase one management of severe acute malnutrition.

What is the priority nursing intervention for this child?

- (A) Iron folate
- (B) Deworming
- (C) IV fluid infusions
- (D) Folic acid supplement

15. A child with complicated severe acute malnutrition was on treatment in the inpatient treatment program of severe acute malnutrition. On admission, the child had severe wasting, pitting edema and failed the appetite test. However, the child did not respond to treatments and classified under failure to treatments.

What is the most likely indicator of failure to treatment the nurse assessed?

- (A) Failure to lose edema on day 10
- (B) Failure to regain appetite on day 10
- (C) If edema persists for more than 10 day
- (D) Failure to gain at least 5 g/kg bodyweight in day four