

30 Scenario-Based MCQs on First Aid for NORCET Exam

1. A 45-year-old male collapses at a community center during a meeting, unresponsive and not breathing normally. You check for responsiveness and call for help. What is the immediate next step in first aid?

- A. Start chest compressions at 100-120 per minute
- B. Give 2 rescue breaths before compressions
- C. Check for a pulse for 10 seconds
- D. Place the person in recovery position

Correct Answer: A

Rationale: In adult cardiac arrest (unresponsive, not breathing normally), the priority is to start high-quality CPR with chest compressions immediately (100-120/min, 5-6 cm depth) to restore circulation. Pulse checks delay CPR and are not recommended by BLS guidelines for lay rescuers; rescue breaths follow after 30 compressions in the CAB sequence.

2. A child at school chokes on a grape, cannot speak or cough, and turns blue. You perform abdominal thrusts, but the obstruction persists. What should you do next?

- A. Perform back blows alternating with abdominal thrusts
- B. Give the child water to drink
- C. Lay the child flat and perform CPR
- D. Wait for the child to cough it out

Correct Answer: A

Rationale: For choking in children (conscious, severe), continue with 5 back blows followed by 5 abdominal thrusts until the object is expelled or the child becomes unresponsive. Water can worsen obstruction; CPR starts only if unresponsive. This follows pediatric first aid protocols to dislodge foreign bodies safely.

3. A factory worker suffers a deep laceration on his arm from machinery, with bright red blood spurting rhythmically. You apply direct pressure, but bleeding continues. What is the next action?

- A. Elevate the arm above heart level
- B. Apply a tourniquet 5-7 cm above the wound
- C. Pack the wound with gauze and wrap tightly
- D. Apply ice to constrict vessels

Correct Answer: B

Rationale: Rhythmic spurting indicates arterial bleeding, a life-threatening emergency. After failed direct pressure, a tourniquet is indicated for limb injuries to control hemorrhage quickly. Elevation aids venous bleeding; packing is for non-arterial wounds. Tourniquets must be noted with time applied per ATLS guidelines.

4. An elderly woman slips in the kitchen and fractures her hip, complaining of severe pain and inability to bear weight. She is conscious but pale. What is the first aid priority?

- A. Splint the leg to the other leg
- B. Apply a cold pack to the hip
- C. Help her walk to a chair
- D. Monitor for shock signs and keep her warm

Correct Answer: D

Rationale: Suspected hip fracture risks shock due to pain and immobility. Prioritize ABCs, keep the patient still, warm, and reassure while monitoring vital signs. Splinting is secondary; assisting to walk can worsen displacement. Shock management (warmth, elevation if no fracture suspicion) prevents hypovolemic complications.

5. A hiker is stung by a bee and develops rapid swelling, hives, and difficulty breathing within minutes. You recognize anaphylaxis. What is the immediate first aid intervention if epinephrine is unavailable?

- A. Apply a cold compress to the sting site
- B. Position the person supine with legs elevated
- C. Encourage deep breathing exercises
- D. Administer oral antihistamine if available

Correct Answer: B

Rationale: Anaphylaxis causes airway compromise and hypotension; supine position with legs elevated improves venous return and oxygenation. Epinephrine is first-line, but if unavailable, positioning buys time. Cold compresses are for local reaction only; antihistamines are adjunctive and slower-acting per anaphylaxis algorithms.

6. During a family picnic, a child pulls boiling water, sustaining a deep partial-thickness burn on the hand (blisters forming). What is the correct initial first aid?

- A. Apply butter to soothe the burn

DOWNLOAD NURSES HUB APP FOR MORE MCQS

- B. Run cool (not cold) water over the burn for 20 minutes
- C. Pop the blisters to drain fluid
- D. Cover with a dry, sterile dressing immediately

Correct Answer: B


Rationale: Cool running water (15-20°C) for 20 minutes reduces tissue damage by dissipating heat and minimizing burn progression in partial-thickness burns. Home remedies like butter trap heat; blisters protect against infection and should not be popped. Dressing follows cooling per burn first aid protocols.

7. A construction worker falls from a ladder, hitting his head, and is found unconscious with irregular breathing. No visible bleeding. What do you do first?

- A. Tilt the head back to open airway
- B. Check for neck stability before moving
- C. Perform a rapid neurological assessment
- D. Apply a cervical collar if available

Correct Answer: B

Rationale: Suspected head/neck trauma requires manual in-line stabilization to prevent spinal cord injury before airway management. Head-tilt can exacerbate cervical injury; assessments are secondary. Collars are ideal but not always first if untrained, per trauma first aid (C-spine precautions).



8. A diabetic patient at a clinic becomes hypoglycemic, sweating profusely, confused, and shaky. Blood sugar is 50 mg/dL. She is conscious. What is the best first aid?

- A. Give insulin to correct the low
- B. Administer 15g fast-acting carbohydrate (e.g., glucose tabs)
- C. Encourage a large meal immediately
- D. Lay her down without oral intake

Correct Answer: B

Rationale: Conscious hypoglycemia requires the "15-15 rule": 15g quick carbs (juice, glucose), recheck in 15 min. Insulin worsens it; large meals delay absorption. Recumbent position risks aspiration if vomiting occurs. This follows ADA guidelines for emergency glucose management.

9. A swimmer is pulled from the pool, gasping but not breathing adequately, with blue lips. Bystanders report sudden collapse. What is the priority?

DOWNLOAD NURSES HUB APP FOR MORE MCQS

- A. Give 5 rescue breaths
- B. Start CPR with compressions
- C. Warm the body with blankets
- D. Check for water in the lungs

Correct Answer: B

Rationale: Drowning victims often have hypoxic cardiac arrest; treat as standard cardiac arrest with CPR (compressions first). Rescue breaths are integrated but not initial. Rewarming is for hypothermia, secondary here. Lung checks delay intervention per drowning resuscitation guidelines.

10. An office worker cuts her finger deeply while opening mail, with steady bleeding. No spurting. You apply pressure for 5 minutes, but it soaks through. Next step?

- A. Remove the dressing to reapply pressure
- B. Add more layers of dressing and continue pressure
- C. Elevate and apply ice
- D. Use a tourniquet

Correct Answer: B

Rationale: For venous bleeding, reinforce pressure without removing soaked dressings to avoid disturbing clots. Elevation/ice aids but pressure is primary. Tourniquets are for arterial/life-threatening only. This prevents exsanguination per wound care basics.

11. A motorcyclist crashes, with a deformed forearm and exposed bone. He screams in pain. What first aid action takes precedence?

- A. Realign the bone manually
- B. Immobilize in current position with sling/splint

- C. Give oral pain medication
- D. Flush the wound with water

Correct Answer: B

Rationale: Open fractures risk neurovascular damage; immobilize without realignment to avoid further injury/infection. Pain meds/flushing are secondary; splinting (rigid support) stabilizes for transport per orthopedic first aid.

Let's Join Nurses Hub Online Test Series
AIIMS NORCET TEST SERIES
250+ Pre + Mains TEST
25000+ MCQs Test Series Join करने के लिए Nurses Hub App Download करें

- ◆ Subject Wise Test Series
- ◆ OT ICU Emergency
- ◆ NCLEX Based Test
- ◆ AIIMS OLD Paper
- ◆ Image Based Question
- ◆ Grand Test
- ◆ Non Nursing Subject
- ◆ All India Live Rank
- ◆ Pdf Notes (Validity > 6 Months)

The advertisement also shows a hand holding a smartphone displaying the app interface.

12. During a heatwave, an elderly man at a market collapses, hot/dry skin, rapid pulse, confused (heat stroke). What immediate first aid?

- A. Give ice-cold water to drink
- B. Move to shade, remove excess clothing, fan vigorously
- C. Apply alcohol rubs to skin
- D. Encourage walking to cool down

Correct Answer: B

Rationale: Heat stroke requires rapid cooling: shade, evaporative cooling (fanning wet skin), and light clothing removal. Cold drinks risk aspiration; alcohol evaporates too quickly. Activity worsens core temperature per hyperthermia management.

13. A toddler ingests household cleaner, vomiting and drooling excessively. Parents call you. First response?

- A. Induce vomiting with syrup of ipecac
- B. Give milk or water to dilute (if conscious)
- C. Rinse mouth and call poison control
- D. Observe for 30 minutes

Correct Answer: C

Rationale: Chemical ingestion: rinse mouth gently, avoid inducing vomiting (risks aspiration/esophageal burns), and contact poison center immediately. Dilution is case-specific; observation delays care per toxicology first aid.

14. A pregnant woman in her third trimester has sudden severe abdominal pain and vaginal bleeding. She is stable but anxious. Priority first aid?

- A. Perform pelvic exam
- B. Place in left lateral position and monitor fetal heart
- C. Give oral fluids
- D. Apply heat to abdomen

Correct Answer: B

Rationale: Suspected placental abruption/ectopic: left lateral reduces aortocaval compression, improving uteroplacental flow. No exams/fluids/heat without assessment. Urgent transport follows per obstetric emergencies.

15. An athlete twists his ankle during a game, with swelling and inability to bear weight. What is the initial first aid (RICE protocol)?

- A. Rest, Ice, Compression, Elevation

DOWNLOAD NURSES HUB APP FOR MORE MCQS

- B. Run, Immobilize, Cold pack, Exercise
- C. Rub, Ice, Compress, Extend
- D. Rest, Heat, Immobilize, Elevate

Correct Answer: A

Rationale: RICE is standard for acute soft tissue injuries: Rest prevents aggravation, Ice reduces swelling, Compression limits edema, Elevation promotes drainage. Heat increases inflammation initially.

16. A patient with chest pain, radiating to jaw, sweating (suspected MI). Conscious, alert. You are a nurse bystander. First action?

- A. Encourage coughing forcefully
- B. Position semi-Fowler's, give aspirin if no allergy
- C. Apply oxygen via mask
- D. Massage the chest

Correct Answer: B

Rationale: Acute coronary syndrome: aspirin (162-325mg chewed) inhibits platelets; semi-Fowler's eases breathing. Oxygen only if hypoxic; coughing/massage ineffective/dangerous per ACS first aid.

17. A burn victim from a house fire has singed nasal hairs and hoarseness. No visible skin burns. Concern?

- A. Inhalation injury; high-flow oxygen, monitor airway
- B. Apply topical antibiotic
- C. Reassure and discharge
- D. Give cough syrup

Correct Answer: A

Rationale: Signs of smoke inhalation (singed hairs, voice change) indicate airway edema risk; prioritize oxygen and intubation readiness. Skin treatment secondary; discharge risky per thermal injury protocols.

18. A seizure patient in a store convulses for 2 minutes, then postictal confusion. Bystander care?

- A. Insert tongue blade

Let's Join Nurses Hub Online Test Series

AIIMS NORCET TEST SERIES

250+ TEST

25000+ MCQs

Test Series Join करने के लिए Nurses Hub App Download करें

- ◆ Subject Wise Test Series
- ◆ OT ICU Emergency
- ◆ NCLEX Based Test
- ◆ AIIMS OLD Paper
- ◆ Image Based Question
- ◆ Grand Test
- ◆ Non Nursing Subject
- ◆ All India Live Rank
- ◆ Pdf Notes (Validity :- 6 Months)



- B. Time the seizure, protect from injury, recover position after
- C. Restrain arms/legs
- D. Offer water during event

Correct Answer: B

Rationale: Epilepsy first aid: clear hazards, time duration (>5 min = emergency), lateral position post-seizure prevents aspiration. Tongue blades/restraints risk injury; fluids during contraindicated.

19. A child with fever (40°C) is lethargic, stiff neck, rash (meningitis suspicion). Home first aid?

- A. Give paracetamol and observe
- B. Urgent transport; keep comfortable, no fluids by mouth if vomiting
- C. Apply cold sponging
- D. Isolate from family

Correct Answer: B

Rationale: Bacterial meningitis is life-threatening; immediate medical evacuation key. Antipyretics secondary; cold worsens shivering. Isolation not first aid priority per pediatric emergencies.

20. A cyclist hit by car has compound leg fracture, pale, rapid breathing (shock). First aid focus?

- A. Splint and elevate leg
- B. Control bleeding, keep warm, reassure, elevate legs if no fracture
- C. Give oral rehydration salts
- D. Walk to safety

Correct Answer: B

Rationale: Hypovolemic shock from fracture: stop bleeding, prevent heat loss, elevate trunk/legs (if spinal cleared). Splinting secondary; no walking/oral fluids if unstable per shock management.

21. Frostbite on a mountaineer's fingers after exposure: white, numb. Immediate action?

- A. Rub vigorously to warm
- B. Immerse in warm (not hot) water (40-42°C), protect from refreezing
- C. Apply dry heat
- D. Thaw with ice

Correct Answer: B

DOWNLOAD NURSES HUB APP FOR MORE MCQS

Rationale: Rewarm frostbite gradually in warm water to restore circulation without burns. Rubbing damages tissue; dry heat uneven. Prevent refreezing critical per cold injury guidelines.

22. A woman hyperventilates during anxiety attack, tingling hands, rapid breathing. What to guide her?

- A. Breathe into paper bag
- B. Slow breathing with pursed lips, reassure it's temporary
- C. Lie flat and elevate legs
- D. Give sedatives

Correct Answer: B

Rationale: Respiratory alkalosis from hyperventilation: rebreathing CO₂ via pursed-lip breathing corrects pH. Bags outdated/risky; no meds/positioning needed for anxiety-induced.

23. Electrocution victim from power line, unconscious, burns on hands/feet. First step?

- A. Start CPR immediately
- B. Ensure power off, check safety, then assess ABCs
- C. Pour water on burns
- D. Move without checking pulse

Correct Answer: B

Rationale: Electrical injuries risk arrhythmia; scene safety first (power isolation) prevents rescuer harm. CPR if asystole; water conducts electricity. ABCs follow per electrical trauma protocol.

24. Snakebite on leg during hike: swelling, pain, no neuro symptoms yet. First aid?

- A. Apply tourniquet above bite
- B. Immobilize limb, keep below heart, mark swelling, urgent evacuation
- C. Incise and suck venom
- D. Apply ice pack

Correct Answer: B

Rationale: Venomous snakebite: immobilize to slow spread, monitor progression, no tourniquet/incision (worsens tissue damage). Ice contraindicated per WHO snakebite management.

25. Allergic reaction to shellfish: facial swelling, wheezing. EpiPen given, improving. Next?

- A. Discharge after 10 minutes
- B. Monitor for 4-6 hours, transport to ER
- C. Give more epinephrine
- D. Apply cold to face

Correct Answer: B

Rationale: Biphasic anaphylaxis possible; observation ensures stability. Repeat epi only if worsening; cold symptomatic only. ER for steroids/antihistamines.

26. Dehydrated child after diarrhea: sunken eyes, dry mouth, lethargic. Oral rehydration?

- A. Give plain water freely
- B. ORS solution (50-100 ml/kg over 4 hours)
- C. IV fluids immediately
- D. Stop all fluids

Correct Answer: B

Rationale: Mild-moderate dehydration: WHO ORS replaces electrolytes/glucose efficiently. Water dilutes sodium; IV for severe only. Prevents shock per pediatric dehydration protocol.

27. Eye chemical splash (acid): burning pain, tearing. Immediate first aid?

- A. Rub eye to flush
- B. Irrigate with saline/water for 20 minutes, evert lid
- C. Neutralize with base
- D. Patch both eyes

Correct Answer: B

Rationale: Copious irrigation dilutes/removes chemical; 20 min minimum. Rubbing spreads it; neutralization risks heat reaction. Patch one eye only per ocular emergency care.

28. Asthma attack in park: wheezing, using accessory muscles, peak flow low. Inhaler empty. What now?

- A. Encourage coughing
- B. Sit forward, loose clothing, call emergency
- C. Lie supine
- D. Give coffee as bronchodilator

DOWNLOAD NURSES HUB APP FOR MORE MCQS

Correct Answer: B

Rationale: Severe asthma: upright position aids breathing, reduce anxiety; EMS for nebulizer/oxygen. Supine worsens; coffee ineffective/delaying.

29. Animal bite (dog) on arm: puncture wounds, bleeding controlled. Risk?

- A. Clean with soap/water, irrigate, tetanus check, rabies prophylaxis
- B. Apply honey dressing
- C. Ignore if pet known
- D. Stitch immediately

Correct Answer: A

Rationale: Bites infect easily (Pasteurella); thorough irrigation reduces risk. Stitching traps bacteria; rabies post-exposure key per wound infection prevention.

30. Heat exhaustion at marathon: cramps, nausea, cool/clammy skin. Runner alert. First aid?

- A. Force vomiting
- B. Rest in shade, small sips of cool fluids, cool with wet cloths
- C. Continue race slowly
- D. Give salt tablets dry

Correct Answer: B

Rationale: Heat exhaustion: rehydrate orally (electrolyte drinks), active cooling prevents progression to stroke. Vomiting risks dehydration; salt alone inadequate per exertional heat illness guidelines.



NH NURSES HUB

Let's Join Nurses Hub Online Test Series

AIIMS NORCET TEST SERIES


250+ Pre + Mains TEST

25000+ MCQs

Test Series Join करने के लिए Nurses Hub App Download करें

- ◆ Subject Wise Test Series
- ◆ OT ICU Emergency
- ◆ NCLEX Based Test
- ◆ AIIMS OLD Paper
- ◆ Image Based Question
- ◆ Grand Test
- ◆ Non Nursing Subject
- ◆ All India Live Rank
- ◆ Pdf Notes

(Validity :- 6 Months)



NH
NURSES

Let's Join Nurses Hub Online Test Series


AIIMS NORCET TEST SERIES

250+ Pre + Mains TEST

25000+ MCQs

Test Series Join करने के लिए Nurses Hub App Download करे

- ◆ Subject Wise Test Series
- ◆ OT ICU Emergency
- ◆ NCLEX Based Test
- ◆ AIIMS OLD Paper
- ◆ Image Based Question
- ◆ Grand Test
- ◆ Non Nursing Subject
- ◆ All India Live Rank
- ◆ Pdf Notes (Validity :- 6 Months)



Download App

Popular courses [See All →](#)

Free All India Test Series

FREE TESTS

Free Test Series

✓ FREE COURSE

Get this course