

63. A 47-year-old female patient was admitted in medical ward with a diagnosis of COPD. Oxygen therapy was ordered as a treatment. The nurse in the ward made a nursing diagnosis of impaired gas exchange related to excessive mucus in the airway.

What is the most likely expected outcome criterion the nurse should set?

- (A) Blood pH above 7.45
- (B) Pulseoximetry reading above 90%
- (C) Partial pressure of arterial oxygen 90 mmHg
- (D) Partial pressure of arterial CO₂ below 35 mmHg

64. A head nurse working in an operating room wants to orient nursing students about the areas in the operating theatre. He gives a focus that "traffic flow is limited and the area is also used as access corridors to the operating rooms".

What is the most likely area explained by the head nurse?

- (A) Restricted
- (B) Scrubbing
- (C) Unrestricted
- (D) Semi restricted

65. A nurse, who has the responsibility to control the traffic of the operating room, wants a patient's family to stay in area near to operation room where surgery is ongoing.

To what area will the nurse allow the patient's family to stay with their usual cloth?

- (A) Semi restricted area
- (B) Unrestricted area
- (C) Restricted area
- (D) Operating area

66. A 43-year-old female was transferred from an OR to a recovery room after having a surgery on her abdomen with general anesthesia. After four hours of procedure, she had snoring, stridor, labored chest movement, intercostal muscle retractions while breathing. She also showed cyanosis, apnea and was not oriented to time, place, and person.

What is the most likely postoperative complication the patient developed?

- (A) Hypotension
- (B) Internal bleeding
- (C) Unconsciousness
- (D) Respiratory distress

67. The nurse is administering a postoperative medication to a male patient who had aortobifemoral bypass surgery. The patient has a previous history of deep vein thrombosis. On assessment, he developed moderate dyspnoea, chest pain, diaphoresis, restlessness, rapid respirations and pulse, cough and cyanosis.

What is the most likely postoperative complication the nurse could anticipate?

- (A) Pulmonary embolism
- (B) Deep vein thrombosis
- (C) Pneumonia
- (D) Shock

BOOKLET CODE: 011

NIS 29 Item SUBJECT CODE: 01

16. A nurse was providing discharge instructions to a 55-year-old man who was being discharged from a hospital after cardiac artery bypass surgery. The surgery was for the treatment of atherosclerosis. The patient was obese and diabetic for 15 years.

What should be the priority-teaching plan the nurse should consider for this patient?

- (A) Eat grains and vegetables more
- (B) Promote cessation of tobacco use

- (C) Perform physical activity daily
- (D) Minimize vegetable oils

17. A 65-year-old patient was admitted to a cardiac unit with a diagnosis of myocardial infarction secondary to the narrowing of the coronary artery disease. The patient was instructed to take nitroglycerine.

What is the most likely predicted effect of this drug on this patient?

- (A) Improve pulmonary circulation
- (B) Improve systemic circulations

- (C) Dilate coronary blood vessels
- (D) Dilate systemic blood vessels

18. An adult female patient was admitted to a medical ward with a diagnosis of angina pectoris. On history, chest pain occurs both at rest and at extraneous activities with no radiation pattern. During examination, there was an ST segment elevation over ECG that reversed in second measurement of four hours later and her body mass index was 29.

What is the type of angina manifested in this patient?

- (A) Stable angina
- (B) Unstable angina

- (C) Variant angina
- (D) Intractable angina

19. A 46-year-old male daily laborer patient came to a clinic with a severe acute chest pain. On physical examination, he held left side of a chest during breathing and he had BMI of 25 kg/m². His blood pressure was 110/85 mmHg and respiratory rate was 26/min. On laboratory investigation, he had total cholesterol of 250 mg/dl and cardiac specific Troponin was increased.

What is the most appropriate initial step in the management of this patient?

- (A) Administer morphine
- (B) Administer propranolol

- (C) Administer nitroglycerine
- (D) Administer large doses of aspirin

BOOKLET CODE: 011

81. A group of people working in animal bone processing factory were brought to a hospital with increasing fever, dyspnea, stridor, hypoxia, and hypotension after eating raw meat prepared for the cultural celebration.

What is the most likely diagnosis of these patients?

- (A) Inhalation anthrax
- (B) Cutaneous Anthrax

- (C) Oropharyngeal anthrax
- (D) Gastrointestinal Anthrax

82. A 30-year-old pregnant woman has come to an ANC clinic with complaints of headache, abdominal pain, vomiting and difficulty of breathing during sleeping for two weeks duration. Her blood pressure is 100/70mmHg, Pulse rate is 80/min and fetal heartbeat is 140/min. On physical examination, she has markedly enlarged abdomen and gestational age of 34 weeks. The nurse has suspected that the woman is at greater risk for amniotic fluid disorder.

Which is the most pertinent characteristic that indicates the disorder?

- (A) Vomiting
- (B) Abdominal pain

- (C) Enlarged abdomen
- (D) Difficulty of breathing

83. A 25-year-old pregnant woman has come to a nearby hospital for antenatal visit. Her blood pressure is 110/60 mmHg, respiratory rate is 24/min, GA is 28 weeks and fetal heart rate is 140/min. The fetal weight is 10 percentile below the given gestation. The nurse suspects IUGR for her pregnancy.

What is the most appropriate assessment method in this case?

- (A) Maternal weight gain
- (B) Maternal abdominal girth

- (C) Symphysis fundal height
- (D) Serial uterine fundal height

84. A 30th week pregnant woman came to a health center for her first ANC visit. Her blood pressure is 120/80mmHg with positive fetal heart beat by Doppler ultrasound. The woman has been provided with an appropriate routine antenatal care including micronutrient supplementation for three months without any other precaution.

What is the most appropriate micronutrient supplement that is given for this woman?

- (A) Zinc
- (B) Calcium
- (C) Vitamin A
- (D) Iron folate

85. A 24-year-old pregnant woman came to an ANC clinic with complaints of headache and dizziness for two weeks. Her pulse rate is 70/min, gestational age 20 weeks and fetal heart rate is 150/min. Laboratory investigation shows that Hgb 10g/dl and serum iron level less than 4 µg/dl (Normal range of serum iron = 65-75 µg/100mL).

What is the most appropriate management for this woman?

- (A) Diet therapy
- (B) Fluid therapy

- (C) Ferrous sulphate
- (D) Blood transfusion

4. A 35-year-old female patient came to a hospital with complaints of chest tightness and difficulty of breathing for two hours. On physical examination, she had wheezing and hyper resonant chest sound on percussion. Chest x-ray revealed narrowing of the bronchus.

What is the most likely diagnosis of this patient?

- (A) Bronchial asthma
- (B) Allergic rhinitis

- (C) Plugged mucus
- (D) Pneumonia

5. A 45-year-old male patient, working in a factory, was admitted to a medical ward with a diagnosis of chronic bronchitis. After a full nursing assessment, the nurse developed a nursing diagnosis of "impaired gas exchange due to chronic inhalation of irritants".

What is the most likely predicted goal for this nursing diagnosis?

- (A) Improvement in blood oxygenation
- (B) Independency in self-care activities

- (C) Improvement in activity tolerance
- (D) Achievement of airway clearance

6. A 45-year-old female patient was admitted to a surgical ward after having extensive surgery on her abdomen. She stayed unconscious for several days and was with multiple IV lines for medication and fluid administration. After performing a full assessment, the nurse has suspected that the patient may develop pulmonary embolism.

What is the most likely assessment finding that has made the nurse to suspect the diagnosis?

- (A) Increased blood pressure, chills, jugular vein distension and painful cough
- (B) Lethargy, nausea, vomiting and diminished heart sounds
- (C) Fever, right upper quadrant abdominal pain and dyspnea
- (D) Chest pain, shortness of breath and tachypnea

7. A nurse working at a health center has administered the first dose of procaine penicillin injection based on the prescription. Few minutes later, the patient starts to have flushing, warmth, itching and appears anxious.

What will be the most appropriate immediate management to be taken in this case?

- (A) Administer appropriate dose of epinephrine
- (B) Take history about hypersensitivity reaction
- (C) Notifying the senior staffs about the incident
- (D) Immediately referring to hospital for better care

68. A 47-year-old female patient came to an emergency care unit with a complaint of breathing difficulty. Her assessment revealed pain in inhalation and shifting of trachea in unaffected side. Chest x-ray result shows large fluid accumulation in the pleural space.

What is the most likely cause for the tracheal shift?

- (A) Pleurisy
- (B) Pneumonia
- (C) Pneumothorax
- (D) Pleural effusion

69. A 46-year-old male patient came to a hospital with complaints of weakness of all extremities. Assessment findings revealed extensive trauma on cervical bone (C2 and C3) and loss of consciousness.

What is the priority nursing diagnosis of this patient?

- (A) Impaired tissue perfusion related to disease process
- (B) Impaired physical mobility motor system paralysis
- (C) Ineffective breathing related to muscle paralysis
- (D) Neuropathic pain related to nerve irritation

70. A 39-year-old female patient was admitted to orthopedic unit with a diagnosis of pelvic bone fracture. Physical examination and x-ray findings showed excessive bleeding and fragmented fracture with scanty blood around the site of injury. The nurse gave immobilization care and prepared for additional intervention.

What is the most appropriate next nursing intervention for this patient after immobilization?

- (A) Protecting further injury
- (B) Insuring airway patency
- (C) Administering analgesia
- (D) Administering IV fluids

71. A 36-year-old man was brought to an emergency unit after he encountered motor vehicle accident. He has severe pain to his back, fast breathing and difficulty of speaking due to shortness of breath but no open wound. The injury was on his neck and chest area. The emergency care team suspected cervical spinal cord injury.

What is the immediate nursing intervention for this patient?

- (A) Be prepared to manage seizures
- (B) Make neurologic examinations
- (C) Immobilize the cervical spine
- (D) Keep the patient warm

20. A 24-year-old male patient came to a hospital with complaints of severe headache after having subtotal gastrectomy. On examination, the patient had numbness and tingling sensation over his fingers, toes, and he has pale conjunctiva. The patient's pulse rate was 112/min and blood pressure was 110/65 mmHg. On laboratory investigation, his hematocrit level was 35% and increased mean corpuscular volume (MCV) of red blood cells.

What is the most appropriate management for this patient?

- (A) Monthly IM injections of vitamin B₁₂
- (B) Improve the nutritional status of patient
- (C) Ferrous sulphate, 325 mg (65 mg elemental iron)
- (D) Give 1mg mg of folic acid daily with intramuscular route

21. A 34-year-old man came to a dermatology clinic with painful grouped vesicles. The patient also complained malaise and headache before the eruption of vesicles. The nurse diagnosed the patient with herpes zoster.

What additional sign and symptoms help to reach at this diagnosis?

- (A) The vesicles that affect the genitalia
- (B) The ruptured vesicles which left yellow crust
- (C) The recurrent and persistent wound over the thorax
- (D) Unilateral band like configuration over the cervical nerve

22. A 35-year-old male, who is a known diabetic patient, comes to a health center with a complaint of painful deep abscess on the back of his neck of six days duration. On history, he reported that the lesion started from painful small, red, and raised lesions. On physical examination, there was a hot, erythematous, firm nodule with multiple openings to the skin surface.

What is the most likely diagnosis of this patient?

- (A) Carbuncle
- (B) Folliculitis
- (C) Cellulitis
- (D) Furuncle

A known epileptic cook had a burn accident to her chest with boiled oil after an epileptic attack. She was shouting in pain. On examination, the epidermis and papillary dermis were destroyed; and pink, moist, and blistered open wound was observed.

What is the type of burn this patient encountered?

- (A) Superficial
- (B) Deep partial
- (C) Full thickness
- (D) Superficial partial

8. A 70-year-old woman was referred to a hospital with allergic reaction to her entire skin for unknown reason. On history, she reported that she did not know the things that made her itchy. The nurse prepared a nursing care plan to solve the occurrence of the problem for the woman.

What is the most appropriate outcome criteria expected in preparing nursing care plan?

- (A) Identifying reportable signs and symptoms of allergic reaction
- (B) Expressing increased sense of comfort and decreased pain
- (C) Identifying means of preventing contact with allergens
- (D) Maintaining intact skin integrity by the patient

9. A 19-year-old female patient came to a hospital with complaints of pruritus, burning sensation and erythema on her face and neck. The symptoms appeared after four days of cosmetics exposure. The patient revealed she has been utilizing cosmetics in the past four days.

What type of hypersensitivity has the patient developed?

- (A) Type-I hypersensitivity
- (B) Type-II hypersensitivity
- (C) Type-III hypersensitivity
- (D) Type-IV hypersensitivity

10. A 31-year-old patient came to a hospital with complaints of cough, night sweats and loss of appetite. After examination, the patient was diagnosed with TB and HIV co-infection.

What is the most appropriate management for this patient?

- (A) Start anti TB treatment and ART at the same time
- (B) Initiate ART after the completion of anti TB treatment
- (C) Start anti TB followed by ART within 8 weeks of treatment
- (D) Start ART and followed by anti TB within 12 weeks of treatment

11. A 45-year-old female with chronic otitis media came to a health center with a complaint of pain to the ear and around the ear. She has fever, sneezing and headache. On physical examination, there is tenderness on her nose.

What is the most likely diagnosis of this patient?

- (A) Acute otitis media
- (B) Mastoiditis
- (C) Rhinitis
- (D) Sinusitis

72. A 35-year-old female patient came to a hospital with a complaint of bloody diarrhea of three days duration. She has also abdominal cramps, nausea, vomiting, tenesmus and fatigue. She ate uncooked tomato five days ago. Her laboratory result shows presence of immunohistochemical test in stool.

What is the priority nursing diagnosis for this patient?

- (A) Sustained nausea
- (B) Diarrhea volume

- (C) Recurrent fatigue
- (D) Deficient fluid volume

73. A nurse, who is working at a health center, assessed a male patient and diagnosed him for acute watery diarrhea. Multiple cases came in morning and the nurse thought that cholera outbreak might happen in the catchment area.

What clinical finding is the most likely suggestive of the case?

- (A) Extreme weakness and malaise
- (B) Severely increased body temperature
- (C) Abrupt watery diarrhea accompanied with pain
- (D) Explosive watery diarrhea that looks like rice-water

74. A 54-year-old female patient came to a hospital with a complaint of lesion on her right leg of one-month duration. On assessment, she has numerous and extensive nodules that diffused bilaterally and symmetrically. The laboratory finding revealed that AFB in skin smear is positive for *Mycobacterium leprae*.

What is the best prevention method in this case?

- (A) Providing immunization
- (B) Administration of antibiotics
- (C) Identification and treatment for the patient
- (D) Isolation of patient after treatment is initiated

75. A 23-year-old patient was brought to a hospital with complaints of cough, and shortness of breath. On arrival, the nurse conducted holistic assessment and diagnosed the patient with measles.

What is the most likely manifestation the nurse used for the diagnosis of this case?

- (A) Rash preceding the fever
- (B) Rash characterized by yellow crusts
- (C) Rash starts immediately days after fever
- (D) Rash starts from the face and work down

Ethiopian Health Professionals' Licensure Examination

Nursing Exam Nehase 2014/August 2022 PART ONE

BOOKLET CODE: 011
Number of Items: 100

SUBJECT CODE: 01
Time Allowed: 2:30 hours

DIRECTIONS: Each of the following questions is followed by four possible alternative answers. Read each question carefully and **blacken** the letter of your best choice on the separate answer sheet provided.

1. A 45-year-old male patient has come to an outpatient department with a complaint of loss of voice. Though he had common cold, currently the symptom has improved. He has no history of smoking and exposure to known irritants.

What is the immediate nursing intervention to regain voice of the patient?

- (A) Give Anti-pain
- (B) Give Antibiotics
- (C) Take vocal rest
- (D) Increase fluid intake

2. A 25-year-old male patient came to a clinic with complains of sense of irritation and fullness on the throat of four months duration. His sore throat worsens during swallowing and coughing. Additional investigations revealed that he has no thyroid gland disorder.

What is the most appropriate intervention for this patient?

- (A) Applying cold compression to support the sore throat
- (B) Advising the patient to make oral gargling
- (C) Providing antihistamine medication
- (D) Providing pain killers medication

3. A 50-year-old male patient has been admitted to a medical ward with a diagnosis of pneumonia. The patient has fever, shortness of breathing, tachycardia, tachypnea and altered mental status. His vital signs are in a normal range.

What will be the priority nursing diagnosis?

- (A) Imbalance nutrition less than body requirement
- (B) Risk for deficient fluid volume
- (C) Ineffective airway clearance
- (D) Activity intolerance

76. A 35-year-old male patient has come to a health center with complaints of headache and chills of three days duration. He also complains rigors and fever that vary within a short period. The blood film result shows presence of parasite.

What is the most important clinical feature to diagnose this case?

- (A) Nature of chills
- (B) Nature of fever
- (C) Parasite in the blood
- (D) Presence of headache

77. A 47-year-old female patient came to a medical outpatient department with complaints of itching papules and local edema around the abdomen. After one week, the patient developed fever, abdominal pain and transient generalized urticaria. The nurse highly suspected for *schistosoma haematobium*.

What is the most appropriate laboratory investigation to confirm the diagnosis?

- (A) Widal test
- (B) Liver biopsy
- (C) Urine analysis
- (D) Stool examination

78. A 30-year-old female patient came to a medical OPD with a complaint of malodorous, yellowish vaginal discharge of one-month duration. On history, she has vulvar erythema and itching, dysuria and dyspareunia. On microscopic finding, there is motile trichomonad on the discharge.

What most likely action is to be taken in order to prevent the disease transmission?

- (A) Treatment of underlying conditions
- (B) Educate public to seek medical help
- (C) Clotrimazole vaginal suppository
- (D) Case detection and treatment

79. A 34-year-old man has come to a dermatology clinic with a shallow and non-indurate tender ulcer to his penis, fever, headache, malaise and muscular pain.

What is the most likely etiology of this problem?

- (A) *Chlamydia trachomatis*
- (B) *Haemophilus ducreyi*
- (C) *Mycoplasma genitalium*
- (D) *Neisseria gonorrhoea*

80. A 29-year-old female patient has come to a health facility with a complaint of edematous lesion on her tonsils of three days duration. She also complains fever, sore throat, dysphagia, and painful regional lymphadenopathy. Mostly, she eats meat product. She has no previous history of drug allergic reaction.

What is the most appropriate drug management for this patient?

- (A) Penicillin
- (B) Tetracycline
- (C) Ciprofloxacin
- (D) Erythromycin

58. A 42-year-old female patient was admitted to a surgical ward with a diagnosis of small bowel obstruction. The physician ordered insertion of nasogastric tube for decompression.

What is the most appropriate NG tube the nurse should prepare?

- (A) Levin tube with single lumen
- (B) Small-bore tubes with single lumen
- (C) Large-bore tubes with single lumen
- (D) Miller-Abbott tube with double lumen

59. A 55-year-old male postoperative prostatectomy patient has a three way-indwelling catheter for continuous bladder irrigation. The duty nurse prepared the necessary materials including fluid to give bladder irrigation care for the patient.

What is the most appropriate type of fluid for this procedure?

- (A) Dextrose 5%
- (B) Saline 0.9%
- (C) Saline 0.45%
- (D) Ringer Lactates

60. A clinical nursing staff soaked used surgical equipment for 10 minutes in 0.5% chlorine solution. Then, she washed with soapy and clean water before packing for the sterilization purpose. A nursing student asked the clinical staff "why equipment are soaked before cleaning?"

What is the most appropriate response the clinical staff should give?

- (A) Socking helps to have safe cleaning
- (B) Socking is helpful for rinsing equipment
- (C) Socking helps to release clots and tissue from equipment
- (D) Socking is optional we can also directly clean if we do not have enough time

61. A 32-year-old man is in a balanced suspension skeletal traction for fracture of the femur. The man reports burning sensation and pain at the pin insertion site. His caregiver requests the nurse to help the patient in order to alleviate his problem.

What is the most appropriate nursing intervention in this case?

- (A) Put a support for the traction weight
- (B) Give the prescribed analgesics
- (C) Put a support beneath his foot
- (D) Massage the area

62. A 33-year-old male patient was admitted to a medical inpatient unit with a diagnosis of pneumonia. His previous medical history shows that the patient has valvular heart problem and the nurse has decided to do postural drainage to clear excessive bronchial secretions.

What is the most appropriate precaution the nurse should consider during the procedure?

- (A) Patient positioning
- (B) Effective coughing technique
- (C) Hydration of the thick mucus
- (D) Drainage amount and consistency

98. A mother on her seventh day of postpartum came to a health center with a complaint of restlessness and confusion for two days. She stated that her baby was changed and she refused to care the baby. She had also visual and auditory hallucinations and impaired judgment and insight.

What is the most likely diagnosis of this mother?

(A) Postpartum blue

(B) Postpartum phobia

(C) Postpartum depression

(D) Postpartum psychosis

99. A 20-year-old mother gave a birth for 3.4 kg male neonate vaginally at hospital. The nurse provided active third stage management of labor and transferred the mother to the postnatal ward.

What is the most likely next nursing intervention for this mother?

(A) Discharge the mother

(B) Encourage fluid intake/urination

(C) Monitor vital signs every 15 minutes

(D) Encourage mobilization of the mother

100. A postpartum mother has developed bleeding after three hours of delivery. On assessment, she has profuse vaginal bleeding, hypotension, dizziness, and pallor/palms. The uterus is soft and boggy. The nurse has given her IV fluid and oxytocin as prescribed.

What is the most likely sign that helps to predict her improvement?

(A) Pink palms

(B) Contracted uterus

(C) Increase level of comfort

(D) Profuse vaginal discharge

THE END

86. A pregnant woman came to an ANC clinic with complaint of vaginal bleeding after four months of amenorrhea for three days. On physical examination, the woman had Serosanguineous vaginal discharge, big abdomen and uterus to date and FHR tone is absent. The ultrasonography finding revealed that there was a typical snowstorm appearance without gestational sac in uterus.

What is the most appropriate nursing intervention for this woman?

- (A) Perineal care
- (B) Decreasing anxiety
- (C) Dilation and curettage
- (D) Maintaining fluid volume

87. An HIV positive pregnant woman came to an ANC clinic. She has been taking HAART for the past six months. Her baseline findings show that she had oral thrush, unexplained fever, CD4 count of 240 and Hgb is 12g/dl. Currently, she is taking her medication consistently.

What is the most appropriate indicator that shows her improvement?

- (A) Decreased viral load
- (B) Decreased CD4 count
- (C) Increase WHO treatment stage
- (D) Similar weight with baseline result

88. A primigravida pregnant woman was admitted to a gynecology ward with hyperemesis gravidarum. During admission, her blood pressure was 80/50mmHg and Pulse rate was 100/min. She had dry mouth and poor concentration. Laboratory investigation results show urine ketone 3+, urine proteins 2+ and her HCT is 35%. She has been managed with two bags of ringer lactate with two vials of 40% dextrose.

What is the most appropriate sign that indicates her improvement?

- (A) Protein free urine
- (B) Ketone free urine
- (C) Increase blood glucose
- (D) Increased hematocrit level

89. A 35-year-old pregnant woman came to a hospital with complaints of headache and fainting for one hour. She has family history of diabetic mellitus. On examination, her gestational age is 26 weeks, random plasma glucose concentration is 210mg/dl. She has been resuscitated in the emergency OPD. The duty nurse plans to determine her glucose level in order to know her prognosis.

What is the most likely intervention outcome used to predict the woman's prognosis?

- (A) One hour oral glucose test 140mg/dl
- (B) Two hours oral glucose test 140mg/dl
- (C) Three hours oral glucose test 140mg/dl
- (D) Four hours oral glucose test 140mg/dl

12. A 40-year-old male patient with mastoiditis had a surgery. He was well informed about the procedure and he coped adequately with every situation after the surgery. The nurse provided postoperative nursing care and prepared the patient a expected outcome.

What is the most likely expected outcome prepared for the patient?

- (A) Demonstrates reduced anxiety about surgical procedure
- (B) Attainment of a positive self-concept
- (C) Remains free of discomfort or pain
- (D) Remains free of injury and trauma

13. A 22-year-old man came to an ENT clinic having ear problem that persisted for seven weeks. A nurse performed holistic assessment and diagnosed the patient with Chronic Otitis Media.

What is the typical manifestation that helps the nurse for the diagnosis?

- (A) Sensation of fullness in the ear
- (B) Foul-smelling Otorrhea
- (C) Sensorineural hearing loss
- (D) Tenderness over the ear

14. A 26-year-old known sinusitis patient came to an ENT clinic with complaints worsening of his symptoms. Based on the patients' complaint, a duty nurse performs a holistic assessment and develops nursing diagnoses.

What is the appropriate collaborative nursing diagnosis of this patient?

- (A) Fever
- (B) Headache
- (C) Brain abscess
- (D) Pain and tenderness

15. A 36-year-old woman gave birth to twin babies by elective caesarian section. After four days of the procedure, the patient developed a Deep Venous Thrombosis (DVT). On nursing assessment, the patient was not mobile due to fear of incision site pain. Her BMI was 30 kg/m^2 .

What was the main risk factor that possibly caused the development of DVT?

- (A) Endothelial blood vessel trauma due to surgery
- (B) Altered coagulation due to her twin pregnancy
- (C) Increased emboli due to body fat and obesity
- (D) Venous stasis due to bed immobilization

24. A 62-year-old man, with HIV/AIDS, has come to a hospital with lesions around his right shoulder that do not pass the midline of the body. On assessment, the lesions have blister and tingling sensation. He experiences severe pain over the entire dermatome supplied by the affected nerve root. The patient avoids movement and does not want clothing or bed linen to touch the affected shoulder.

What is the most appropriate management for the patient?

- (A) Administer Tramadol
- (B) Administer antibiotics
- (C) Administer Acyclovir
- (D) Administer Calamine lotion

25. A 59-year-old woman came to a medial OPD with complains of abdominal pain, feeling of epigastric fullness, nausea, and belching. She was taking NSAIDs for several years as part of management for osteoarthritis. The nurse at the OPD was assessing the patient to reach at a proper diagnosis.

What would be the most important additional question the nurse should ask?

- (A) 'Have you ever noticed change in odor of breath?'
- (B) 'Have you noticed any problems with your breathing?'
- (C) 'Have you ever vomited blood or had very dark stools?'
- (D) 'Did you experience nausea always whenever you take the drug?'

26. A 22-year-old male patient has come to a health center with severe generalized abdominal pain that worsens at the right lower quadrant after 24 hours duration. Physical examination reveals board like rigid abdomen and normal bowel sounds. His body temperature is 37.9°C . A nurse has suspected appendicitis.

What is the most likely complication that would happen in this patient?

- (A) Perforation
- (B) Abscess
- (C) Peritonitis
- (D) Infection

27. A 58-year-old female hypertensive patient has been admitted to a medical ward for a stroke. Currently she is in a coma. The nurse has planned to insert NG tube feeding since she was unable to eat or drink.

What is the indicator used to confirm if the NG tube is properly measured and placed?

- (A) Bubbling
- (B) Clear fluid aspiration
- (C) Aspiration with PH value of 4
- (D) Litmus paper color change from red to blue

38. A 48-year-old female patient was admitted to a medical ward to rule out meningitis. On assessment, she is feeling of pain and spasm on hip flexion, has irritability, but no pain on neck flexion two days after lumbar puncture. The nurse suspected brain irritation secondary to the diagnostic procedure.

What is the most likely additional sign that confirms the nurse's suspect?

- (A) Pain and stiffness
- (B) High grade fever
- (C) Mental status change
- (D) Nausea and vomiting

39. A 26-year-old male patient was presented to an emergency care unit after sustaining spinal cord injuries from T₁ to T₆. The nurse and the emergency physician discussed the priority management option to prevent worsening of a neurologic deficit.

What is the most likely priority nursing intervention in this case?

- (A) Reduction and stabilization
- (B) Oxygen administration
- (C) Application of traction
- (D) Immobilization

40. A 73-year-old male patient was admitted to a medical inpatient unit with complaints of severe headache, confusion, dysphasia and aphasia. Diagnostic test result showed obstructed cerebral vessel that supply the brain tissue.

What is the most appropriate nursing intervention for this patient?

- (A) Administer pain medication
- (B) Instruct client to speak slowly
- (C) Supportive care for rehabilitation
- (D) Elevate client's head for eating or drinking

41. A 25-year-old male patient was transferred from an outpatient department to a medical inpatient unit for admission after the diagnosis of Type-I diabetes mellitus. Results from chart review showed the presence of polydipsia, polyphagia and polyuria. The nurse wants to explore additional clinical features for better nursing interventions.

What is the most likely additional clinical manifestation the nurse wants to include?

- (A) Cognitive changes
- (B) Skin infections
- (C) Blurred vision
- (D) Weight loss

52. A 45-year-old female known diabetic patient came to a clinic with dehydration and hypoglycemia after doing a strenuous exercise for longer duration. A physician prescribed a rehydration with isotonic IV fluid containing glucose.

What is the most appropriate fluid for this patient?

(A) Dextrose 5% water

(B) Dextrose 5% in RL

(C) Dextrose 5% in 0.9% saline

(D) Dextrose 10% in 0.9% saline

53. A nurse working in an operating theatre immersed contaminated surgical equipment immediately after surgical procedure in 5% chlorine solution for 10 minutes.

What is the most likely explanation for the nurse's action?

(A) Sorting

(B) Cleaning

(C) Sterilization

(D) Decontamination

54. A 28-year-old male patient came to a medical emergency unit with a sudden loss of consciousness. On immediate assessment, the nurse noticed irregular, weak and rapid radial pulse. Then, the nurse assessed the apical pulse.

What is the most appropriate rationale for assessing apical pulse in this case?

(A) Best site for weak pulse

(B) Best for irregular pulse

(C) Best site for rapid pulse validation

(D) Best site to palpate and auscultator

55. A 43-year-old female patient came to a surgical referral clinic for routine appointment. After clinical examination, the physician decided to admit the patient for surgical interventions as per the Ethiopian National Patient Admission Protocol.

What is the initial nursing admission intervention in this case?

(A) Prepare bed for the patient

(B) Familiarize environment

(C) Document admission date

(D) Provide hygiene care

56. A 40-year-old female patient started to complain lower extremity pain and spasm three days after admission. Pain assessment result showed moderate level of pain. The nurse in charge has given heat application care as a nursing intervention.

What is the most likely outcome of the nursing intervention?

(A) Relaxed muscles

(B) Increased blood flow

(C) Decreased blood flow

(D) Decreased blood pressure

57. A physician ordered cleansing enema for 34-year-old patient with diagnosis of abdominal distension as a part of pre-operative preparation.

What is the calculated amount of fluid required for this procedure?

(A) 100-150 ml

(B) 350 to 500 ml

(C) 500-1000 ml

(D) 750-1000ml

42. A 43-year-old female patient was admitted to a surgical ward. The patient reported discomfort has flushing of face. Assessment findings showed protruded eyes, normal tracheal alignment and her temperature was 38.3°C , pulse rate was 104/min, respiratory rate was 22/min, and blood pressure was 130/80 mmHg.

What is the priority nursing diagnosis of this patient?

- (A) Hyperthermia related to increased metabolic rate
- (B) Imbalanced nutrition greater than body requirements
- (C) Impaired verbal communication related to hoarseness
- (D) Ineffective breathing pattern related to compression of the trachea

43. A 65-year-old female patient was enrolled to a surgical follow-up clinic with a diagnosis of hyperthyroidism. The patient has stress due to psychosocial problems in the family. The physician has ordered propylthiouracil (PTU) as a management.

What is the most likely expected complication if the patient is not compliant to treatment?

- (A) Thyrotoxicosis
- (B) Hypothyroidism
- (C) Hypercalcemia
- (D) Hypothermia

44. A 41-year-old female has come to a hospital with complaints of polyphagia, progressive weight loss, fatigue, amenorrhea, diarrhea and nervousness. She is tolerating heat poorly and has sweating even at rest. Her eyeballs are abnormally protruded and bulging. Her pulse rate is 140/min.

What is the most likely diagnosis of this patient?

- (A) Hyperthyroidism
- (B) Diabetes mellitus
- (C) Hypothyroidism
- (D) Endemic goiter

45. A 72-year-old man has been presented to a clinic with complaints of painless and blurry vision. The patient perceives that surroundings are dimmer and the problem is worsening progressively. The Snellen visual acuity test shows decreased vision and ophthalmoscopic examination shows opacity of the lenses, cloudy pupil and absence of red reflex.

What is the most likely problem of this patient?

- (A) Conjunctivitis
- (B) Glaucoma
- (C) Trachoma
- (D) Cataract

46. A 32-year-old female patient has been presented to an ophthalmologic unit with complaints of severe recurrent headache and blurred vision. On physical examination, the patient needs to hold things close to the eyes to see them well.

What is the most appropriate management for this patient?

- (A) Eye surgery
- (B) Antibiotics
- (C) Eye glasses
- (D) Eye irrigation

33. A nurse was caring for an admitted patient with renal failure. During assessment, the patient reported difficulty of breathing, metallic taste, nausea and vomiting. On physical examination, the patient was lethargic, had ankle edema, and crackle on auscultation was identified.

What serious complication was the patient experiencing?

- (A) Uremia
- (B) Hypovolemia
- (C) Hypervolemia
- (D) Hyponatremia

34. A 45-year-old female patient was admitted to a medical ward with a diagnosis of acute renal failure. She lost 6 kg of weight within a week. On assessment, the nurse revealed oliguria, decreased skin turgor, dry mucous membrane and hypothermia.

What is the most appropriate immediate management for this patient?

- (A) Parental fluid replacement
- (B) Fluid and sodium restriction
- (C) Measuring input and output
- (D) Oxygen administration

35. A nursing student was caring for a patient with an end stage renal disease. On history, the patient reported difficulty of breathing, nausea, vomiting and decrease urine output. On physical examination, edema below ankle, crackle on auscultation and weak rapid pulse were observed.

What is most likely body fluid and electrolyte status of this patient?

- (A) Hypovolemia
- (B) Hypervolemia
- (C) Hyperkalemia
- (D) Hyponatremia

36. A 35-year-old male patient has been admitted to a surgical ward after a suspect of having a renal stone. He has non-radiating pain, nausea, vomiting, abdominal distention and oliguria.

What is the most priority nursing diagnosis of the patient?

- (A) Acute pain
- (B) Risk for infection
- (C) Deficient knowledge
- (D) Altered fluid volume

37. A 22-year-old male patient suddenly felt down on the ground, his hands were clenched and all his body parts were rigid. His assessment findings showed that period of apnea in breathing pattern and loss of consciousness, and the nurse suspected seizure disorder.

What is the most likely type of seizure the patient developed?

- (A) Tonic-clonic seizures
- (B) Myoclonic seizures
- (C) Atonic seizures
- (D) Absence seizures

47. A 36-year-old female patient came to an emergency ophthalmologic unit with complaints of cough, extreme tearing and redness of both eyes. On physical examination, watery discharge and extreme photophobia were identified.

What is the most recommended nursing intervention for this patient?

- (A) Administer prescribed antiviral ointment
- (B) Teach the patient to wear dark glasses
- (C) Administer prescribed antibiotics
- (D) Apply eye irrigations

48. A nurse is caring for a new post-operative patient with above knee amputation. The patient feels discomfort, burning, tingling, throbbing, and itching on the missing limb. The physical examination reveals red, warm and slightly edematous amputated dressed wound.

What is the most likely complication this patient is experiencing?

- (A) Infection
- (B) Phantom limb pain
- (C) Compartment syndrome
- (D) Fat embolism syndrome

49. A 28-year-old male patient was admitted to an orthopedic inpatient unit with a diagnosis of left knee joint dislocation. The patient has a history of rheumatoid arthritis treatment. The nurse decided to manage his complain in order to relieve his pain.

What is the most likely positive care outcome for this patient?

- (A) Verbalize increase in comfort
- (B) Perform range of motion exercises
- (C) Report the current level of discomfort
- (D) Be free from joint swelling and inflammation

50. A 20-year-old female patient came to a hospital with a complaint of pain on the hamstring after participating on circus showing festival. Diagnostic test result showed muscle rupture and tear.

What is the most likely diagnosis of this client?

- (A) Sprain
- (B) Contusion
- (C) Strain
- (D) Dislocation

51. A 23-year-old male patient came to an emergency unit of the health center few hours after falling accident. A nurse applied splint as an immediate intervention and referred him to a hospital for better management.

What is the most likely indication for the above nursing intervention?

- (A) Bursitis
- (B) Fracture
- (C) Strains
- (D) Sprains