

Ministry of Health
Ethiopian Health Professionals' Licensure Examination

NURSING
Tikimt 2013/October 2020

Full Name _____ Registration No. _____

BOOKLET CODE: 012

SUBJECT CODE: 02

PART: TWO

TIME ALLOWED: 2:30 HOURS

GENERAL DIRECTIONS

IN THIS EXAMINATION, THERE ARE A TOTAL OF **100** MULTIPLE CHOICE QUESTIONS. SELECT THE BEST ANSWER AND **BLACKEN** ONLY THE LETTER OF YOUR CHOICE ON THE SEPARATE ANSWER SHEET PROVIDED. FOLLOW THE INSTRUCTIONS IN THE ANSWER SHEET AND THE EXAMINATION PAPER CAREFULLY. USE ONLY PENCIL TO MARK YOUR ANSWERS. YOUR ANSWER MARK SHOULD BE **HEAVY** AND **DARK** COVERING THE ANSWER SPACE COMPLETELY. IF YOU CHANGE YOUR ANSWER, PLEASE ERASE IT COMPLETELY AND NEATLY.

YOU ARE ALLOWED TO WORK ON THE EXAMINATION FOR **2:30** HOURS. WHEN TIME IS CALLED, YOU MUST IMMEDIATELY STOP WORKING, PUT YOUR PENCIL DOWN, AND WAIT FOR FURTHER INSTRUCTIONS.

ANY FORM OF CHEATING OR AN ATTEMPT TO CHEAT IN THE EXAMINATION WILL RESULT IN AN AUTOMATIC DISMISSAL FROM THE EXAMINATION HALL AND CANCELLATION OF YOUR SCORE(S). PLEASE MAKE SURE THAT YOU HAVE WRITTEN ALL THE REQUIRED INFORMATION IN THE ANSWER SHEET BEFORE YOU START TO WORK ON THE EXAMINATION. YOUR ANSWER SHEET WILL NOT BE MARKED UNLESS YOUR QUESTION PAPER IS RETURNED.

DO NOT TURN THIS PAGE OVER UNTIL YOU ARE TOLD TO DO SO.

Ethiopian Health Professionals' Licensure Examination

Nursing Exam Tikimt 2013/October 2020 PART TWO

BOOKLET CODE: 012

Number of Items: 100

SUBJECT CODE: 02

Time Allowed: 2:30 hours

DIRECTIONS: Each of the following questions is followed by four possible alternative answers. Read each question carefully and **blacken** the letter of your best choice on the separate answer sheet provided.

1. A lactating mother came to the family planning clinic seeking contraception for the first time. The nurse discussed about contraception options, the benefits, and side effects. But the mother was confused to decide the type of contraception she wants to use.

What is the priority counseling approach to be applied in this condition?

- (A) Keep two-way communication (C) Ensure privacy
(B) Remain non judgmental (D) Show empathy

2. A six month lactating mother came to health center for family planning service. After assessing her eligibility and providing counseling, the nurse inserts Jadelle in September 2012 E.C. Before the mother left the clinic, the nurse gave her a reminder card for return visit for removal purpose.

What would be the most appropriate appointment date for the next visit?

- (A) September 2014 (C) September 2016
(B) September 2015 (D) September 2017

3. A mother who uses an Implant contraceptive came with complaints of changed bleeding pattern, headaches, abdominal pain, breast tenderness and nausea. The mother reported that her menstruation period become sometimes absent and little in amount on other time.

What is the priority intervention for this woman?

- (A) Switch contraception X (C) Check for pregnancy
(B) Provide counseling (D) Remove the contraception X

4. A 25-year-old mother came with complaints of excessive and irregular monthly bleeding after six month of taking contraception. She also complains about increments of her weight. Her body mass index was 26.4 and blood pressure 140/80mmhg. The nurse counsels her to shift the contraceptive she uses.

What is the priority side effect of the contraception?

- (A) Weight gain (C) Irregular bleeding
(B) Excessive bleeding (D) Increased blood pressure

5. A 31-year-old woman came to gynecologic unit with complaints of lower abdominal pain, pain during coitus, abnormal vaginal discharge, and fever. On physical examination, the nurse found uterine and adnexal tenderness.

What is the most likely confirmatory manifestation for this patient?

- (A) Pain during coitus (C) Abnormal vaginal discharge
(B) Lower abdominal pain (D) Uterine and adnexal tenderness

6. A woman came to a gynecologic OPD with complaints of excessive and irregular uterine bleeding. The woman cannot remember when her menstrual period started. The amount and duration of bleeding also vary during each cycle.

What is the most likely explanation of this finding?

- (A) Menorrhagia (C) Polymenorrhea
(B) Metrorrhagia (D) Menometrorrhagia

7. An adolescent girl came with complaints of difficulty and pain during sexual intercourse. She has regular menstrual period with abdominal cramps. On pelvic examination: a nurse identified loses of hymen membrane and a membrane like structure with small perforation on middle third of vagina. The nurse suspected that her problem is vaginal congenital anomalies.

What is the most likely manifestation for diagnosis of this condition?

- (A) Abdominal cramp (C) Painful sexual intercourse
(B) Small perforation (D) Membrane like structure

8. A 20 weeks pregnant woman came with the complaint of vaginal bleeding for three hours. She had abdominal and back pain. On examination, her hemoglobin level was 9g/dl (NR: 10.5-13.7 g/dl), blood pressure 90/50mmHg and pulse rate 110/min. The woman was diagnosed for inevitable abortion.

What is the priority nursing intervention for this woman?

- (A) Provide analgesia
(B) Reassure the woman
(C) Prepare for blood transfusion
(D) Resuscitate with normal saline
9. A 35-year-old Multi-Para mother came with complaint of protrusion of mass on her vagina. She had also sense of pelvic pressure, backache and fatigue. After assessment, she was diagnosed for uterine prolapse. The Uterus is repositioned by gynecologists.

What is the priority intervention for this mother?

- (A) Resuscitate the woman
(B) Administer pain inhibitors
(C) Educate about Pelvic exercises
(D) Teach how to control urine incontinence
10. A nurse was providing care for a girl with abnormal excessive uterine bleeding, breast tenderness, anorexia and vomiting. After 24 hours of care, her blood pressure became to 110/70mmHg, pulse rate 76/min, respiratory rate 22/min, T^0 37.8°C and hemoglobin level 9g/dl.

Which finding shows effectiveness of the patient's management?

- (A) Blood Pressure
(B) Body Temperature
(C) Hemoglobin Level
(D) Respiration Rate
11. A nulli-parous woman came to nearby hospital with weight loss and difficulty of breathing over the past several months. Physical examination result showed lymph gland enlargement, tender nodular, firm liver, bilateral mass in hypogastrium with tender and irregular appearance. The cytological examination also detected ovarian malignancy.

What is the most likely stage of the cancer?

- (A) Stage IIA
(B) Stage IIB
(C) Stage III
(D) Stage IV

12. A 28-year-old woman came to a hospital with complaints of dysmenorrhea and dyspareunia that occurs pre-menstrual and subsides after menses begins. She also had constipation and rectal bleeding with defecation. Physical examination shows that she had tender adnexal masses and pain upon uterine motion. She is diagnosed for endometriosis.

What is the most likely manifestation to confirm the diagnosis?

- (A) Pain
- (B) Constipation
- (C) Dysmenorrhea
- (D) Adnexal masses

13. The nurse evaluates a two-year-old child who was admitted for the treatment of failure to thrive secondary to severe acute malnutrition. The child had good appetite and no edema for 10 consecutive days. The nurse decided to discharge the child.

What is the most likely additional finding that could support the nurse's decision?

- (A) Weight for age >85%
- (B) Weight for age >90%
- (C) Weight for height >80%
- (D) Weight for height > 85%

14. A 15-month-old child was brought to health center for growth and developmental check up. On examination, the infant can walk independently but cannot grasp of an object using thumbs and index finger. The child responds to a simple command like "wave bye bye".

What was the most likely developmental problem of this child?

- (A) Emotional developmental problem
- (B) Language developmental problem
- (C) Cognitive developmental problem
- (D) Fine motor developmental problem

15. A nurse in immunization room found that the refrigerator is out of order. The technician of the health center confirmed that maintenance of the refrigerator could take a maximum of four to five days. So, the nurse wants to store vaccines till the refrigerator maintained.

What is the most likely type of cold chain used to store the vaccines in this time?

- (A) Cold box with cool water pack
- (B) Cold box with frozen water pack
- (C) Vaccine carrier with cool water pack
- (D) Vaccine carrier with frozen water pack

16. A health center has a total of 20,000 populations in the catchment area, of which infants account 20%. In this health center, the planned annual coverage for PCV vaccine is 90% with vaccine wastage multiplication factor of 1.11.

What is the annual maximum stock level of PCV vaccine in the health center?

- (A) 11, 988 doses
(B) 12, 988 doses
(C) 14, 985 doses
(D) 15, 985 doses
17. A nurse wants to initiate feeding for 1.5kg neonate admitted in neonatal intensive care unit. The infant had a very weak sucking reflex and on risk for necrotized enterocolitis.

What is the most appropriate initial step in this patient's care?

- (A) Pouring the expressed breast milk via NG tube continuously as demanded
(B) Inject 4.5ml expressed breast milk using 10ml syringe at a time per feeding
(C) Give bolus 3ml expressed breast milk via NG tube over 20-25 min per feeding
(D) Give bolus the expressed breast milk using 10ml syringe over 5-10min per feeding
18. A nurse assessed the progress of a two-year-old child who was on phase one management of severe acute malnutrition. The child was improved clinically with an evidence of reduced edema to grade two and complications were completely treated.

What is the most likely additional finding used to proceed to the transition phase?

- (A) The edema has to be reduced to grade for a day
(B) The edema has to be lost completely for consecutive days
(C) Improved appetite evidenced by finishing F-75 food easily
(D) Improved appetite evidenced by finishing F-100 food easily
19. A 16-hour-old preterm neonate who was born at home was admitted to the NICU due to inability to feed and lethargy. On examination, the neonate had body temperature of 36.2°C , pulse rate 140/min and blood glucose level of 35 mg/dl.

What is the best nursing intervention to manage the neonate's problem?

- (A) Provide intravenous glucose
(B) Put skin-to-skin with mother
(C) Provide milk through NG tube
(D) Warm the child using hot water bottle

20. A nurse who is giving immediate newborn care dries and stimulates the newborn. Unfortunately, the newborn do not cry.

What is the most appropriate next step for this newborn?

- (A) Suck fluid from the nose
- (B) Bag and mask ventilation
- (C) Observe and evaluate breathing
- (D) Back rubbing and flicking soles

21. A six-day-old neonate was admitted to a hospital for management of breathing difficulty. The neonate had spontaneous breathing at birth but breathing difficulty started on second day of birth and worsened on the fourth day. The neonate was put on Continuous Positive Airway Pressure Ventilation.

What is the best and safest method of checking the progress of this neonate?

- (A) Pulse oximetry
- (B) Blood gas analysis
- (C) Pulmonary function test
- (D) Carbon dioxide monitoring

22. A 15-day-old neonate was brought to a hospital with a complaint of difficulty of feeding of one day duration. The neonate had history of convulsion. On examination, the neonate had also trismus and irritability.

What is the most likely diagnosis of this neonate?

- (A) Sepsis
- (B) Tetanus
- (C) Meningitis
- (D) Hypoglycemia

23. A nurse was giving resuscitation for a newborn by using bag and mask ventilation. After a minute of resuscitation, the neonate didn't start spontaneous breathing and had pulse rate of 90/min.

What condition makes the nurse to stop resuscitation?

- (A) No spontaneous breathing after 10 minute
- (B) Breathing less than 30 breath/min after 10 minute
- (C) Breathing less than 30 breath/min after 20 minute
- (D) Pulse rate of less than 60 beats/min after 20 minute

24. A 12-day-old neonate was brought to a hospital with a complaint of petechial rash all over the trunk for one day duration. On examination, the neonate had normal platelet count, increased prothrombin time and increased partial thromboplastin time.

What additional information is the most likely used to identify the neonate's problem?

- (A) Neonatal history of bleeding
 - (B) Maternal history of infection
 - (C) Neonatal history of vitamin K injection at birth
 - (D) Maternal history of warfarin use during pregnancy
25. A low birth weight neonate was brought to NICU with the clinical features of unable to breast feed, difficulty of breathing and yellowish discoloration of a body part. The neonatal nurse classifies the neonate as jaundice according to IMNCI.

What is the most likely finding used to confirm this classification?

- (A) Yellowish discoloration of palm and sole
 - (B) Yellow skin and eye, and baby is six-day-old
 - (C) Yellow skin and eye, and baby is 16-hour-old
 - (D) Yellow skin and eye, and baby is seven-day-old
26. A four-year-old child was presented to the health center with diarrhea of five days duration. On physical examination, the skin pinch goes back within two seconds, and the child was very eager to drink water and had WFH z-score >-1 .

What is the most appropriate management of this child according to IMNCI?

- (A) Give ORS over 4-hours period
 - (B) Give ORS solution in each loss
 - (C) Give ReSoMAL solution in each loss
 - (D) Give RL solution intravascular over three hours
27. An eight-month-old infant was brought to a pediatric OPD with the clinical features of cough and chest indrawing. The respiration rate of the child was 48/min.

What is the most likely classification of the infant according to IMNCI?

- (A) Pneumonia
- (B) Cough or cold
- (C) Severe pneumonia
- (D) Very severe disease

28. An infant who was born from a known HIV positive mother is brought to a pediatric OPD with the clinical features of cough, stridor and oral thrush. The HIV status of the infant was positive for antibody test and unknown for DNA/PCR test due to unavailability of the test.

What is the appropriate management for the child according to IMNCI?

- (A) Initiate PMTCT care
(B) Initiate ARV treatment
(C) Advise on HIV prevention
(D) Advise to stop breastfeeding
29. A 30-month-old child was brought to the health center with a complaint of cough of three weeks duration. The nurse identified that the father of the child is on initiation phase of TB treatment and the child also had low grade fever.

What is the most appropriate initial step to do for this child?

- (A) Refer urgently to the hospital
(B) Initiate INH as TB prophylaxis
(C) Put the child on anti-TB treatment
(D) Counsel on the diagnosis of TB exposure
30. A 10-month-old HIV exposed infant was brought to the under-five clinic for routine checkup. The nurse immediately took DBS sample and found DNA PCR test positive.

What is the most appropriate initial step the nurse should do?

- (A) Take DBS to repeat viral test
(B) Start ART as soon as possible
(C) Start cotrimoxazole immediately
(D) Start ARV prophylaxis immediately
31. A four-year-old child was brought to a health center with a complaint of cough of three days duration. On assessment, the nurse observed that the child had inspiratory stridor and barking cough.

What is the most likely diagnosis of this patient?

- (A) Croup
(B) Pneumonia
(C) Pulmonary TB
(D) Tonsillo-pharyngitis
32. A child was brought to a pediatric OPD with the clinical features of nasal congestion, sneezing, lacrimation and conjunctival suffusion. The nurse diagnosed the child has pertussis.

What is the most likely finding that could support the nurse's suspicion?

- (A) Dyspnea
(B) Rhinorrhea
(C) Low grade fever
(D) whooping cough

33. A six-week-old infant was brought to pediatric OPD with the clinical features of high grade fever, neck stiffness, bulging fontanel and high pitch cry.

What is the most likely cause of this infant's problem?

- (A) Escherichia coli (C) Group A streptococcus
(B) Group B streptococcus (D) Staphylococcus aureus

34. A child was admitted to pediatric ward for the treatment of femoral bone fracture. The child was on traction with a cast and complained severe unpleasant sensation at the fracture site.

What is priority nursing diagnosis for this case?

- (A) Acute pain (C) Activity intolerance
(B) Impaired mobility (D) Impaired activities of daily living

35. A three-year-old child who was brought to a health center for vaccination aspirated coin while playing.

What is the next most appropriate nursing intervention to be done for this child?

- (A) Give artificial respiration
(B) Perform Heimlich maneuver
(C) Dislodge the coin with little finger
(D) Refer the child to nearby hospital urgently

36. A 20-day-old male baby was brought to pediatric OPD with a complaint of being unable to feed and convulsion. On examination his temperature was 38°C , heart rate 98/min and CSF findings showed that his Leukocytes were $30/\text{mm}^3$, Protein 110 mg/dL and 60% of the blood glucose.

What is the most likely diagnosis of this baby?

- (A) Neonatal seizure (C) Neonatal asphyxia
(B) Neonatal sepsis (D) Neonatal meningitis

37. A six-year-old child came to pediatric emergency with a complaint of sudden loss of consciousness. The child was found to be in diabetic ketoacidosis. The nurse planned to administer fluid for the child. Child's weight was 16kg.

What is the most appropriate fluid to be given in the first hour of admission?

- (A) 160-320ml IV DNS (C) 160-320ml IV 0.45% NS
(B) 160-320ml IV 0.9% NS (D) 160-320ml IV 5% dextrose

38. A seven-year-old patient came to pediatric OPD with a complaint of rapid, shock-like and non-rhythmic convulsion of four episodes with two days duration.

What is the most likely diagnosis of this child?

- (A) Clonic seizure
(B) Petitmal seizure

- (C) Grandmal seizure
(D) Myoclonic seizure

39. A child weighting 5kg was admitted to pediatric ward with a complaint of unable to feed. On examination, the baby had moderate respiratory distress and his hematocrit level was 25% (HCT packed RBC to be given was 70%).

What is the amount of packed RBC to be transfused for this patient?

- (A) 160 ml
(B) 182 ml

- (C) 198 ml
(D) 204 ml

40. An adolescent girl came to pediatric clinic and diagnosed with uncomplicated nephrotic syndrome.

What is the most appropriate nursing intervention for this patient?

- (A) Administration of prednisone
(B) Administration of cyclosporine
(C) Administration of cephalosporins
(D) Administration of cyclophosphamide

41. A two-year-old female child was brought to a health center with difficulty of feeding and fever of three days duration. On examination, she has rash around the face and neck and the clinician suspected measles.

What could be the pathognomonic sign to substantiate this diagnosis?

- (A) Red eyes
(B) Rose spots

- (C) Runny nose
(D) Koplik spots

42. An infant with confirmed hemophilia B developed hematuria and admitted to the hospital. The nurse provided the appropriate amount of maintenance fluid with bed rest but hematuria continued for consecutive days.

What is the most appropriate next step in the management of this infant?

- (A) Giving 20IU/kg factor IX
(B) Giving 40IU/kg factor IX

- (C) Giving 40IU/kg factor VIII
(D) Giving 20IU/kg factor VIII

43. A three-year-old male child goes to supermarket with his mother and asked his mother to buy him extra chocolate. The mother told him that she didn't have sufficient money and promised him to buy next time. However, the baby didn't agree, dropped to the floor and started crying, screaming and kicking in rage at the middle of the supermarket.

What is the most likely behavioral characteristic of this child?

- (A) Dawdling (C) Regression
(B) Negativism (D) Temper Tantrums

44. A three-year-old child with the complaints of change in social interaction and repetitive behavior came to a hospital. On assessment, the child loves to play alone, used multiple nonverbal behaviors, plays with the same toys and did not speak words appropriately.

What is the most likely diagnosis of this child?

- (A) Attention deficit (C) Autism spectrum disorders
(B) Down syndrome (D) Childhood depression disorder

45. A 20-year-old woman came to a clinic and reported that she has not been getting asleep for one week. She was more talkative than usual. On mental status examination, she is uncooperative, easily annoyed, and seductive and has flight of idea.

What is the most important mental status finding to support her diagnosis?

- (A) Irritable (C) Flight of idea
(B) Seductive (D) Uncooperative

46. A 25-year-old woman came to a clinic and complained about persistent feeling of sadness, hopelessness, sleep disturbance and loss of interest. The mental status examination revealed that she had difficulty to sustain attention.

What are the core symptoms in the diagnosis of the above patient?

- (A) Hopelessness and/or distractibility
(B) Loss of interest and/or hopelessness
(C) Distractibility and/or sleep disturbance
(D) Depressed mood and/or loss of interest

47. A 17-year-old boy gets 5 mg of haloperidol to control his aggression. After an hour of his first injection, he starts to experience brief contractions of muscles that result in obviously abnormal movements, tongue protrusion and difficulty of swallowing.

What is the most likely diagnosis of the above patient?

- (A) Dystonia
(B) Akathesia
(C) Parkinsonism
(D) Epilepsy
48. A 19-year-old married woman, recently diagnosed with bipolar I disorder manic episode, came to a clinic crying loudly. Her nose bleed and there is visible laceration on her face. Her mother reports that her husband has beaten her due to her unacceptable behavior with other man.

What is the priority nursing action for the above case?

- (A) Tell her she has to avoid such behavior in the future
(B) Report to legal service unit for necessary investigation
(C) Stop bleeding and reassure her she will be safe at hospital
(D) Calm the patient and call her husband for further communication
49. A 65-year-old patient from middle income family, with a diagnosis of dementia, was brought to a clinic by his daughter. She reports that despite she has been giving prescribed medication timely, his behavior is becoming worse. Currently, he lives alone and smokes three pieces of cigarette/day.

What is the best predictor of risk for poor adherence in this patient?

- (A) Income
(B) No insight
(C) Substance use
(D) Forgetfulness
50. A 55-year-old woman came to a clinic for psychiatric evaluation. At the start of the interview, she focuses on issues that the nurse consider irrelevant.

What is the next best step that the nurse should follow?

- (A) Ask her if she understood what he is asking her
(B) Guide the patient to come to direction of his point
(C) Tell her he will give another time to talk the details
(D) Listen for a while even things he doesn't want to hear

51. A 35-year-old patient with a diagnosis of delusional disorder was brought to a hospital. He stopped his hazardous drinking habit before one year. Despite his false belief which interferes with his functioning, he performs well during the mental status examination. He has never been met criteria for schizophrenia.

What is the most appropriate explanation for his diagnosis?

- (A) He has abnormal false belief
- (B) He has functional impairment
- (C) He has history of excessive alcohol use
- (D) He has never met criteria for schizophrenia

52. A 17-year-old student has persistent fear of speaking in front of his class mates. He fears that he will act in a way that will be negatively evaluated by his colleagues. Because of his fear, he failed to score good mark in class activities for the last one year.

What is the most likely diagnosis?

- (A) Specific phobia
- (B) Social anxiety disorder
- (C) Separation anxiety disorder
- (D) Generalized anxiety disorder

53. A 32-year-old soldier saw terrorist attack on television and starts to experience feelings of panic. The panic disappeared within a few hours; however, for the next few nights, he had night mares with frequent thoughts about dying. Before 10 years, he survived ambush which killed three of his close friends in front of him.

What is the most likely diagnosis for the above case?

- (A) Complicated grief
- (B) Acute stress disorder
- (C) Adjustment disorder
- (D) Post traumatic stress disorder

54. The nurse working in university students clinic meet a 19-year-old fresh student fulfilling the diagnostic criteria for adjustment disorder with depressed mood. He occasionally had difficulty of sleeping but function optimally.

What is the most appropriate nursing intervention for this case?

- (A) Give medication to help him sleep well
- (B) Tell him to come back if symptoms worsen
- (C) Reassure and advise to contact counseling unit
- (D) Reassure him symptoms will subside with time

55. A 54-year-old man had been able to stop drinking many times. Currently, he is experiencing tremor, anxiety, and problems sleeping in his trial for one more times. He is typically self-medicating these symptoms with three 10-mg tablets of diazepam per day.

What is the most likely diagnosis of the above case?

- (A) Alcohol abuse
- (B) Alcohol dependency
- (C) Alcohol intoxication
- (D) Alcohol withdrawal

56. A group of patients with substance use disorder were admitted to substance rehabilitation center. Most of them deny their problem. The nurse is preparing a teaching plan for them to help their recovery.

What is the most appropriate information to be included in the nursing plan?

- (A) Communicate one can use substance for their stress relieving effect
- (B) Focus on change in the patient's motivation probably has the best impact
- (C) Persons with substance-related problems can recover without formal treatment
- (D) Communicate that substance use is a problem not related to individual weakness

57. A 67-year-old woman was brought to a hospital due to progressive deterioration of memory and language ability in the last two years. Her son reports that she couldn't stay at home, wandering on the street and often miss to return to home. During examination, she is alert, has memory problem and talks alone appearing as if discussing with someone.

What is the most likely outcome of treating this patient?

- (A) Peaceful death within a short period of time
- (B) Eventual full return to pre morbid level of functioning
- (C) Eventual partial return of pre morbid level of functioning
- (D) Behavioral control with delaying of cognitive deterioration

58. A 50-year-old dementia patient was admitted to a hospital. On assessment, he had forgetfulness, fluctuating mood and impaired decision making ability. His son is struggling to prevent him from removing the IV line.

What is the most expected immediate outcome to be achieved after hours of intervention?

- (A) Improved memory
- (B) Reduced disturbance
- (C) Reduced mood fluctuation
- (D) Improved ability to make own decision

59. A nurse who is working as a ward head in a hospital developed annual plan with hospital staff teams. However, they failed to achieve the plan because of inability of mobilizing the human and material resources from the hospital.

What is the most likely management function that causes low achievement of this ward?

- (A) Planning
- (B) Organizing
- (C) Directing
- (D) Controlling

60. A university board assigned a committee in order to review curriculum vitae of applicants for a managerial position as president of a university, the committee understands that an applicant is experienced in analyzing the external environment of institutions and engaging in problem solving activities that can bring major changes.

What is the most appropriate managerial skill the applicant had based on this scenario?

- (A) Technical Skill
- (B) Conceptual Skill
- (C) Human relation Skill
- (D) Communication skill

61. A Head of the health center and planning team decided to develop a plan. The planning team developed a plan which will be used in case when unexpected problems and events occurred even if the health center developed its annual plan.

What is the most appropriate type of planning developed by the team?

- (A) Strategic planning
- (B) Tactical planning
- (C) Operational planning
- (D) Contingency planning

62. A newly appointed head of the health center had discussion with catchment *kebele* leaders to identify real health problems that occurred in the community. The head identified trachoma as major health problem following poor latrine utilization though they have constructed latrine.

What would be the most appropriate good care delivery approach used by the head?

- (A) Disease control programs
- (B) Conventional care in clinics
- (C) People-centered primary care
- (D) Convectional care in referral hospitals

63. A newly appointed CEO of a hospital decides to decentralize activities to subordinates after performing situational analysis. The process was done using the service regulations and consistent discussion with community members in the catchment area.

What is the most likely health service building block focused by the CEO?

(A) Governance (C) Human resource
(B) Financing (D) Service delivery

64. Job Evaluation Grading (JEG) implementation committee in zonal health administration works in the new staff job appointment. During the course, a nurse who is working in disease prevention directorate raised complains about his current appointment and takes the case to the court. The court accepted employee's complaint and ordered the organization to cancel the decision.

What is the most likely conflict resolution used in this scenario?

(A) Win- Lose (C) Win -Win
(B) Lose-Lose (D) Lose- Win

65. Head of a health center lead a staff meeting on the issue of improving latrine utilization of the community in the catchment area. Participants of the meeting arrived at their own conclusion and stated that the head frequently relinquish responsibility and had minimal involvement in this activity.

What is the most likely leadership style the head used?

- (A) Authoritarian leadership
(B) Democratic leadership
(C) Laissez faire leadership
(D) Transformational leadership

66. An experienced public health expert comes to Ethiopia from abroad and wants information about the current health delivery system of the country. During their discussion, the *woreda* health manager explained about the current health care tier system used by the nation.

What is the most likely explanation expected from the manager?

- (A) Ethiopia has four tier system
(B) Primary hospital serves 15,000-25,000 people
(C) General hospitals are grouped under tertiary level
(D) Specialized Hospital serves 3.5- 5.0 million people

67. A hospital board discusses with the management team on how to mobilize a resource to provide quality health service. They decided to improve income by collecting service cost from health care, diagnostic services, consultancies and trainings.

What is the most likely part of health care financing reform used by the board?

- (A) Health Insurance
- (B) Revenue Retention
- (C) Fee Waiver System
- (D) Exemption System

68. The Ministry of health organized a review meeting with representatives of health professionals. On the meeting, nurses from different hospitals discussed on the difference between monitoring and evaluation but they could not reach an agreement.

What is the most likely characteristic that differentiate the two concepts?

- (A) Monitoring is continuous, where as evaluation is periodic
- (B) Monitoring is intermittent, where as evaluation is continuous
- (C) Monitoring can be done after a project ends, but not evaluation
- (D) Monitoring used to have a lesson, where as evaluation prevent distraction

69. A 28-year-old female patient was diagnosed with ovarian cancer and the oncologist decided the need for radiation therapy. The nurse in charge explained the risk of infertility secondary to radiation therapy. She refused radiation therapy and at the same time she had strong desire to have children. The procedure was done to save her life; she was cured of ovarian cancer and develop infertility

What ethical principle most likely forced the health care team's action?

- (A) Autonomy
- (B) Beneficence
- (C) Justice
- (D) Non-maleficence

70. A 28-year-old female patient was admitted to surgical ward with ovarian cyst. The preoperative assessment identified hepatitis B infection. To prevent professionals, a nurse in charge communicated with all health care providers who might take care of the patient. The patient volunteered to share information about diagnosis and blood screening finding.

What principle is most likely at risk of violation?

- (A) Advocacy
- (B) Confidentiality
- (C) Informed Consent
- (D) Privacy

71. A 48-year-old male patient who was diagnosed with hypovolumic shock refused blood transfusion irrespective of repeated counseling by the nurse in charge.

What is the most appropriate legal framework in dealing with this patient?

- (A) Continue patient resuscitation
- (B) Take verbal consent and let him go
- (C) Have written consent and let him go
- (D) Look for family permission and let him go

72. A 32-year-old female patient came to hospital with sudden onset of sharp knife like chest pain accompanied by her family. On assessment, she pronounced dyspnea and central cyanosis. The physician and the nursing team in charge performed emergency surgery. The patient and her family presented the case to court because she developed iatrogenic pleural effusion as complication.

What nursing action most likely prevents accusation?

- (A) Keep fidelity
- (B) Take consent
- (C) Keep Veracity
- (D) Assess patient capacity

73. An experienced nurse, who was not happy with the working environment, accepted a 56-year-old male patient that came with complaint of joint pain. The nurse looks at the eye of the patient, listens and feels the pain of the patient. The nurse touches the patient's shoulder while teaching and reassuring him about the case. The patient was happy with nurse's approach.

What is the most likely principle of compassion the nurse lacks?

- (A) Affection
- (B) Attention
- (C) Acceptance
- (D) Acknowledgement

74. A 32-year-old female patient came to health center for wound care. A nurse in charge provided the wound care. The patient developed infection and referred to a hospital. In the hospital, she got treated and cured. After six years, she is informed that the infection she developed in the health center was due to inappropriate dressing. The patient filed malpractice act against the nurse in health center and prosecuted the nurse for malpractice act.

What is the first thing the nurses' attorney should have as a defense?

- (A) Assumption of risk
- (B) Comparative negligence
- (C) Contributory negligence
- (D) Statute of limitations

75. A 30-year-old nurse was working in a hospital oncology unit. Every day, he cares for patients from different parts of Ethiopia by respectful and dignified manner. Every day he tried to maintain the provision of dignified care.

What principle of respectful care helps to achieve nurses' dream?

- (A) Recognize diversity and individual uniqueness
 - (B) Uphold responsibility to shape care and support service
 - (C) Communicate with individuals in a meaningful way to them
 - (D) Value workplace culture that promote respect for every body
76. A 28-year-old nurse has been doing every routine nursing activity using nursing process. The nurse also participates in monitoring activities at her institution that improve quality of nursing practice, in order to keep her knowledge base and skills, and current practice with the guidelines, statutes, rules and regulations.

What is the most likely nursing law that this nurse is practicing?

- (A) The scope of nursing practice
 - (B) The standards of practice
 - (C) The nursing policy statements
 - (D) The position statements
77. A nurse working in a medical ward administered drug intramuscularly that should have been given through the vein. Another nurse working in the same ward reported the medication administration error to the director.

What is the first thing the nurse's director should do?

- (A) Disclose verification
 - (B) Confidential verification
 - (C) Report to ethical committees
 - (D) Direct confrontation and supervision
78. A 58-year-old female patient was on screening for breast cancer. The patient is worried about her diagnosis and fate of her family. She doesn't want to hear bad news. The nurse in charge is worried about how to disclose the diagnosis of metastatic breast cancer to the patient.

What is the most likely condition that worried the nurse?

- (A) Contract between confidentiality and autonomy
- (B) Contrast between veracity and nonmaleficence
- (C) Contrast between beneficence and justice
- (D) Contrast between autonomy and veracity

79. A nurse works in a district where health service is 100% available and with more than 80% expected coverage of rotavirus vaccination among under-five children. However, at the end of the year rotavirus vaccination coverage in the district was found to be 16%.

What is the most likely research problem needed to study in the district?

- (A) Expected rotavirus vaccination coverage for under-five children
 - (B) Availability of rotavirus vaccination for under-five children
 - (C) Rotavirus vaccination coverage among under-five children
 - (D) Factors influencing low rotavirus vaccination coverage
80. A researcher wants to conduct a study on the prevalence and associated factors of trachoma among under-five children in a district. Then, the researcher wants to set appropriate objectives.

What is the most likely specific objective of this research?

- (A) To study the prevalence of trachoma among under-five children
 - (B) To appreciate the prevalence of trachoma among under-five children
 - (C) To understand associated factors of trachoma among under-five children
 - (D) To determine factors associated with trachoma among under-five children
81. A researcher wants to conduct a study among 867 samples of households to determine access to safe water supply in a district. The district has 10,000 households, of which 20% are from urban and 80% from rural households.

What is the most likely sampling technique the researcher will use?

- (A) Systematic sampling
 - (B) Stratified sampling
 - (C) Multi-stage sampling
 - (D) Simple random sampling
82. A nurse working in ART clinic of a hospital wants to summarize CD4 T cell counts for 13 HIV positive patients who were treated with highly active antiretroviral therapy (HAART) for the last 6 months. Their CD4 T cell counts were 230, 205, 313, 207, 227, 245, 173, 58, 103, 181, 105, 301, and 169.

Calculate the median CD4 T cell counts of these HIV positive patients?

- (A) 173
- (B) 181
- (C) 205
- (D) 207

83. A research finding on diarrhea status among children 6-23 months showed 47% mild diarrhea, 36 % moderate diarrhea and 17% severe diarrhea. Finally, the researcher wants to present the data in graphs.

What is the most likely type of graph to be used by this researcher?

- (A) Bar graph (C) Histogram
(B) Line graph (D) Ogive curve

84. A nurse working in a district with a total population of 45,000. In the area, there was an epidemic of acute watery diarrhea (AWD) that lasts for three consecutive weeks. The nurse observed 26 cases in the 1st week, 10 cases in 2nd and 5 cases in the 3rd week and 4 deaths over the three weeks.

Calculate attack rate of AWD over the three weeks?

- (A) 89 AWD cases per 1,00,0000 people at risk
(B) 911 AWD cases per 1,00,0000 people at risk
(C) 578 AWD cases per 1,00,0000 people at risk
(D) 333 AWD cases per 1,00,0000 people at risk

85. A nurse was assigned in a *woreda* health office where immigration and out migration of the people is high from different parts of the country. Monthly health report of the office indicated that tuberculosis incidence is stable. But, the prevalence of tuberculosis increased highly in the *woreda*.

What is the most likely reason for increment of tuberculosis patient's in the *woreda*?

- (A) Out-migration of cases
(B) Improved cure rate of disease
(C) Out-migration of healthy people
(D) High case-fatality rate from disease

86. The nurse wants to test whether the mean age of adults who visited the youth friendly service is not different from 25 years. The nurse took sample of 196 adult patients from population. The mean age of sampled adults was 20 years and the given population standard deviation was 6 years.

What is the most appropriate statistical test used for the given hypothesis?

- (A) Paired t-test (C) Chi-square test
(B) Student t-test (D) Standard normal Z- test

87. A nurse works in a district at which most of the communities cannot read and write. He delivered health education program on how prevent diarrheal disease on the community of the district. When the nurse delivers health education session, he used locally spoken words and local proverbs.

What is the most likely health education principle applied by the nurse?

- (A) Health education aims at change of behavior
- (B) Health Education is based on scientific findings
- (C) The health educators have to make themselves acceptable
- (D) The health educator has to adjust his talk and action to suit the group

88. The Health Professional's Competency Assessment and Licensure Directorate of Ministry of Health want to assess the nursing professionals' curriculum together with concerned bodies. In the process of evaluation, the team gave emphasis on description, specifically in relation to program objectives.

What is the most likely curriculum evaluation model used by the team?

- (A) Second-generation model
- (B) First-generation model
- (C) Third-generation model
- (D) Fourth-generation model

89. A 24-year-old pregnant woman with gestational age of 30 weeks came to ANC clinic with a chief complaint of small fetal size. On assessment, a nurse found a mother has a problem of dietary intake and weight. Then, the nurse told the mother about the possible outcome of the fetus.

What is the most likely outcome of the fetus in this scenario?

- (A) Birth of infant with low weight for gestational age
- (B) Birth of infant with large weight for gestational age
- (C) Birth of infant with normal weight for gestational age
- (D) Birth of infant who initiates breast feeding successfully

90. A four-year-old child came to a health facility with a chief complaint of abdominal pain and irritability followed by fatigue, anorexia, anemic manifestations, gait disturbance, and slurred speech. During history taking, the mother claims that the baby was frequently playing using wires and metallic objects.

What is the most likely cause of poisoning to this child?

- (A) Lead
- (B) Arsenic
- (C) Cyanide
- (D) Mercury

91. A 29-year-old woman had fever and visited the nearby health center. The health care provider on duty at that day was very busy and shouted at her, "Do you want us to waste our time for a mild fever? Come back when we are less busy." She did not like being shouted at. Since then, she never visited the health center even if she is sick.

What is the most likely behavior motivator for the woman?

- (A) Norms
- (B) Values
- (C) Belief
- (D) Attitudes

92. An epidemic investigation team in the regional health bureau found an epidemic disease in a community. The investigators found that the disease occurred due to consumption of contaminated community water supply and the disease has an epidemic curve with a wide peak.

What is the most likely type of epidemic in this community?

- (A) Mixed epidemic
- (B) Progressive epidemic
- (C) Point source epidemic
- (D) Continuous common source epidemic

93. Ministry of health of Ethiopia wanted to get epidemiologic data for decision making purpose. Based on the situation, the Ministry decided to use sentinel surveillance and wanted to use reports for surveillance purpose.

What is the most likely source of report to be used by the ministry?

- (A) Solicit reports of diseases gained from professionals
- (B) Report sent by professionals with a known set of rules
- (C) Pre-arranged sample of reporting sources about all cases
- (D) Report sent about a disease when it reaches to epidemic level.

94. Nurses working in a health center were assigned to give school health service. During designing training, the nurses disagreed on the importance of school health services to the students and for teaching learning process.

What is the most likely importance of school health to be mentioned by the nurses?

- (A) Helping to build and use of educational facilities
- (B) Making teachers and students anxious about disease
- (C) Providing a healthy and safe environment in all-round
- (D) Treating communicable and non-communicable disease

95. A nurse who was assigned as member of malaria epidemic management team wanted to use a permanent measure to remove the mosquito breeding site with the participation of the villagers in the *woreda*.

What is the most appropriate vector control method to be used by the team?

- (A) Indoor residual sprayer
- (B) Use of window screens
- (C) Drainage of stagnant water
- (D) Use of insecticide treated bed net

96. Members of rural community in a district asked a nurse about "how to keep the safety of our water for drinking ". A nurse wanted to instruct the community about the most cost effective and reliable method of disinfecting water at household level.

What is the most likely water treatment technique instructed by the nurse?

- (A) Cloth filtration
- (B) Boiling of water
- (C) Home Candle Filters
- (D) Chlorination of water

97. A nurse who is the member of infection prevention team in a health center collected a package of disposal needles, syringes and scalpels from care provision rooms. Then, the collected waste transported for the proper disposal in the facility.

What is the most appropriate disposal technique for this waste?

- (A) Boiling
- (B) Autoclaving
- (C) Incineration
- (D) Chemical disinfection

98. A three-month-old child came with his mother to under-five children OPD with the complaint of diarrhea for three days. During history taking, the mother claims that "this disease might associate with my practice of giving only breast milk and also he cried a lot now. I think the milk is not enough, it is better to start giving a complementary food". Then, the nurse wants to advice a woman about breast feeding.

What is most likely feeding advice for this mother?

- (A) Continue breast feeding until 2years
- (B) Use good positioning and attachment techniques
- (C) Continue exclusive breastfeeding until six months
- (D) Have to empty one breast before switching to the other

99. A 12-year-old male child who came to town by a child trafficker frequently asked for help. During the interview, the child claims that he does not have any relatives and frequently abused by the gangs in the street.

What is the most likely priority intervention for this child?

- (A) Look for possibilities to reunite with his families
 - (B) Reassure the child about availability of health facility
 - (C) Attach the child to credible children home care providers
 - (D) Provide street based health education on personal hygiene
100. A 24-year-old patient came to adult OPD with complaint of polyphagia, polydipsia and polyuria. On history, the patient reported that "previously a doctor told me that I have diabetes but I am not taking drug till now because I believe God can cure me soon". However, the clinical and lab finding indicate the worsening of the patient's condition and he needs to start insulin injection.

What is the most appropriate health education to influence his behavior?

- (A) Educate him about indication of medication use
- (B) Tell him about the consequences of resisting medication
- (C) Educate him about use of medication despite God's help
- (D) Encourage him to practice spiritual action without use of drugs

THE END