

[Back](#)**Question 5**Not yet
answeredMarked out of
1.00

Flag question

A five year female child brought to Health center from low malaria risk area with fever, convulsion and lethargic. Which of the following is the best category of this child?

- ☐ a. Fever: no malaria
- ☐ b. very severe febrile disease
- ☐ c. Malaria
- ☐ d. severe complicated malaria

[Previous page](#)[Next page](#)

Time left 2:00:39

[Hide](#)

Exam na

1	2
10	11
19	20
28	
37	
46	
55	
64	
73	
8	

[Back](#)**Question 15**Not yet
answeredMarked out of
1.00

Flag question

A patient with a comminuted fracture of the femur is to have an open reduction with internal fixation (ORIF) of the fracture. The nurse explains that ORIF is indicated when:

- ☐ a. Adequate alignment cannot be obtained by other nonsurgical methods.
- ☐ b. The patient cannot tolerate the discomfort of a closed reduction.
- ☐ c. Cast would be too large to provide normal mobility.
- ☐ d. The patient is able to tolerate long-term immobilization.

[Previous page](#)

Time left 1:51:33

[Hide](#)**Exam navigation**

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50
55	56	57	58	59
64	65	66	67	68
73	74	75	76	77
82	83	84	85	86
91	92	93	94	95
100				

[Next page](#)



MoEEP

Back

Time left 2:01:34

Hide

Question 4

Not yet
answeredMarked out of
1.00

Flag question

A 10-year-old male child falls headfirst from a tree and lands on the ground. He experienced immediate loss of consciousness, vomiting, and have unequal pupil size upon regaining consciousness. What is the most appropriate nursing intervention for this patient?

- ☐ a. Performing neurologic assessments.
- ☐ b. Administer antiemetic medications
- ☐ c. Keep the head elevated at a 30-degree angle
- ☐ d. Stabilize the child's neck and spine.

Previous page

Next page

Exam navigation

1	2	3	4
10	11	12	13
19	20	21	22
28	29	30	31
37	38	39	40
46	47	48	49
55	56	57	58
64	65	66	67
73	74	75	76
82	83	84	85
91	92	93	94

Back

Time left 2:06:27

Hide

Question 3

Not yet
answered

Marked out of
1.00

Flag question

Nursing management of the patient with hemiplegia during the acute phase of a stroke includes:

- ☐ a. Performing passive range of motion on all limbs every 4 hours
- ☐ b. Maintaining the patient in a recumbent, side-lying position
- ☐ c. Restricting active movement
- ☐ d. Positioning each joint higher than the proximal joint

Previous page

Next page

[Back](#)


Time left 2:07:40

[Hide](#)**Question 2**Not yet
answeredMarked out of
1.00 Flag question

During the postoperative period, the patient with an above-the-knee amputation should be instructed that the residual limb should not be routinely elevated because (2 extra

- ☐ a. The flexed position can promote hip flexion contracture
- ☐ b. This position promotes clot formation at the incision site and thigh
- ☐ c. This position reduces the development of phantom limb sensation
- ☐ d. Unnecessary movement of the extremity can cause wound dehiscence

[Previous page](#)[Next page](#)



Back

Time left 1:29:13

Hide

Question 36

Not yet
answered
Marked out of
1.00
Flag question

Researchers sometimes use a member of a hidden population to provide names of other members of that population to recruit participants in research. The resulting sample would involve

- ☐ a. Probability sample
- ☐ b. Self-selected sampling
- ☐ c. Chain-referral method
- ☐ d. Sampling frame

Previous page

Next page

Exam navigation

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74

[Back](#)

Time left 1:29:44

[Hide](#)

Question 35

Not yet
answeredMarked out of
1.00[Flag question](#)

Which of the following is an example of a strategic leadership role for a nurse leader?

- ☐ a. Developing and implementing organizational goals
- ☐ b. Conducting research studies
- ☐ c. Providing direct patient care
- ☐ d. Training new nursing staff

[Previous page](#)[Ne](#)



MoEEP

Back

Question 3

Not yet
answered

Marked out of
1.00

Flag question

Nursing management of the patient with hemiplegia during the acute phase of a stroke includes:

- ☐ a. Performing passive range of motion on all limbs every 4 hours
- ☐ b. Maintaining the patient in a recumbent, side-lying position
- ☐ c. Restricting active movement
- ☐ d. Positioning each joint higher than the proximal joint

Time left 2:02:30

Hide

Previous page

Next page

Exam

1

10

19

28

3

[Back](#)**Question 6**Not yet
answeredMarked out of
1.00

Flag question

Which of the following is an example of quantitative research?

- ☐ a. Conducting in-depth interviews
- ☐ b. Administering a structured questionnaire to a large sample
- ☐ c. Analyzing textual data from open-ended survey questions
- ☐ d. Observing and recording behaviors in a natural setting

Time left 1:59:53

[Hide](#)[Previous page](#)[Next page](#)

Exam

10

19

28

37

46

55

64

7

8

[Back](#)

Time left 1:58:53

[Hide](#)

Question 7

Not yet
answeredMarked out of
1.00[Flag question](#)

After surgery, Ms. Hanna is having difficulty voiding. Which nursing action would most likely lead to an increased difficulty with voiding?

- ☐ a. Using a warm bedpan when Ms. Hanna feels the urge to void
- ☐ b. Having Ms. Hanna ignore the urge to void until her bladder is full
- ☐ c. Stroking Ms. Hanna's leg or thigh
- ☐ d. Pouring warm water over Ms. Hanna's fingers

[Previous page](#)[Next page](#)

Exam na

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65



MoEEP

[Back](#)**Question 89**Not yet
answeredMarked out of
1.00

Flag question

Time left 0:36:03

[Hide](#)

Mr. Tadese is a head nurse at a health center. His major focus as head is building and maintaining relationships among staffs who are working under his unit. Which one of the following best described the role of Mr. Tadese?

- ☐ a. Task roles
- ☐ b. individual roles
- ☐ c. Social-emotional roles
- ☐ d. Procedural roles

Exam navigation

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50
55	56	57	58	59
64	65	66	67	68
73	74	75	76	77
82	83	84	85	86
91	92	93	94	95

[Previous page](#)[Next page](#)

Question 11

et
ered

d out of

g question

The nurse is teaching a patient with a moderate hearing impairment in preparation for hospital discharge. To facilitate communication, the nurse should:

- ☐ a. Over articulate his/her speech
- ☐ b. Raise the voice to a higher pitch
- ☐ c. Use simple sentences
- ☐ d. Write out all questions and responses



[Back](#)**Question 12**Not yet
answeredMarked out of
1.00

Flag question

Time left 1:54:42

[Hide](#)

Mr. Abera is receiving continuous tube feedings through a small-bore nasogastric tube. Which method is the most accurate in verifying correct tube placement?

- ☐ a. Measurement of the gastric aspirate pH
- ☐ b. Auscultatory method
- ☐ c. Radiographic examination
- ☐ d. Measurement of the amount of residual

[Previous page](#)[Next page](#)

Exam

1

10

19

20

3

4



Back

Question 8

yet
vered
ed out of
question

s page

Time left 1:58:05

Hide

When a nurse assesses a patient for pain and offers a plan to manage the pain, which principle is used to encourage the nurse to monitor the patient's response to the pain?

- ☐ a. Non-maleficence
- ☐ b. Fidelity
- ☐ c. Respect for autonomy
- ☐ d. Beneficence

Next page

Exam r

1	2
10	
19	
28	
37	
4	

[Back](#)**Question 1**

Answer saved

Marked out of
1.00

Flag question

Time left 2:09:07

[Hide](#)

Which one of the following is the most likely nursing intervention benefit for schizotypal personality disorder clients?

- ☐ a. Teaching social skills
- ☐ b. Providing emotional support
- ☐ c. Cognitive restructuring techniques
- ☒ d. Improving community functioning

[Clear my choice](#)[Next page](#)

Exam naviga

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75
82	83	84
91	92	93
100		

Finish att



MoEEP

Back

Time left 1:40:05

Question 26

Not yet
answered

Marked out of
1.00

Flag question

Nursing care for a patient with an indwelling catheter includes which of the following?

- ☐ a. Telling the patient that burning and irritation are normal, subsiding within a few days
- ☐ b. Encouraging a generous fluid intake if not contraindicated by the patient's condition.
- ☐ c. Irrigation of the catheter with 30 mL of normal saline solution every 4 hours
- ☐ d. Disconnecting and reconnecting the drainage system quickly to obtain a urine specimen

Previous page



MoEEP

[Back](#)

Time left 1:57:16

[Hide](#)**Question 9**Not yet
answeredMarked out of
1.00

Flag question

Performing the assessment while the baby is still wet and unclothed When providing postpartum care, the nurse should instruct the patient on proper breastfeeding technique, which of the following best describe this technique?

- ☐ a. Ensuring a deep latch with the baby's mouth covering the nipple and areola
- ☐ b. Applying heat to the breasts before feeding to promote milk production
- ☐ c. Taking the baby off the breast abruptly to avoid overfeeding
- ☐ d. Offering the baby only one breast during each feeding session

[Previous page](#)[Next page](#)**Exam navig**

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75
82	83	84
91	92	93

Back

Question 21

Not yet
answered

Marked out of
1.00

Flag question

Which of the following is a compensatory response to decreased cardiac output?

- ☐ a. Decreased vascular resistance
- ☐ b. Alteration in LOC
- ☐ c. Increased vascular resistance and fluid retention
- ☐ d. Decreased BP and diuresis

Previous page

Time left 0:35:01

Which action demonstrates proper technique for conducting a newborn assessment immediately after delivery?

- ☐ a. Focusing solely on physical assessment and omitting the neurologic examination
- ☐ b. Delaying the assessment until the baby is clothed and in a warmer
- ☐ c. Using a non-sterile stethoscope to auscultate the baby's heart sounds
- ☐ d. Performing the assessment while the baby is still wet and unclothed

Next

[Back](#)

Time left

Question 19Not yet
answeredMarked out of
1.00[Flag question](#)

In research, what does the acronym "IRB" stand for?

- ☐ a. Investigative Research Bureau
- ☐ b. Institute of Research and Biomedicine
- ☐ c. International Review Board
- ☐ d. Institutional Review Board

[Previous page](#)

[Back](#)

Question 22

Not yet
answeredMarked out of
10[Flag question](#)

Time left 1:46:14

[Hide](#)

A 25 year old male patient with Obsessive Compulsive Disorder is presented with ritualistic hand washing that last several hours each day. The skin on his hands is red and cracked, has minor bleeding. Which one of the following is the nursing goal for this client?

- ☐ a. Providing good skin care
- ☐ b. Eliminating the hand washing rituals
- ☐ c. Decreasing the time spent washing hands
- ☐ d. Providing milder soap for hand washing

[Previous page](#)[Next page](#)

Time left 1:47:51

Hide

A 28-year-old woman who is 6 months pregnant presents to the emergency department with a temperature of 38.2°C (100.8°F) and complains of shaking chills and pain on her right side, which she locates by pointing to the area above her right iliac crest. During the examination she is tender to percussion at the junction of the lower ribs and the thoracic vertebrae. Urinalysis reveals WBC casts. Which of the following is the most likely diagnosis?

- ☐ a. Pyelonephritis
- ☐ b. Localized cystitis
- ☐ c. Ectopic pregnancy
- ☐ d. Appendicitis

 Next page

Exam na

1

2

10

11

19

20

28

29

37

38

46

47

55

56

64

65

73

74

82

83

91

92



MoEEP

[Back](#)

Time left 1:39:41

[Hide](#)

Question 27

Not yet
answeredMarked out of
1.00

Flag question

A 10-year-old male was recently admitted to hospital with signs of increased intracranial pressure (ICP). He presented with severe headaches, vomiting, and altered levels of consciousness. A CT scan was sent and it revealed a space-occupying lesion. Finally, brain tumor was diagnosed. What is the most appropriate nursing interventions for this patient?

- ☐ a. Administering antipyretics
- ☐ b. Encouraging deep breathing exercises
- ☐ c. Administering antibiotics
- ☐ d. Elevating the head of the bed at 30 degree

[Previous page](#)[Next page](#)

Exam navig

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75
82	83	84
91	92	93

Question 14

out of

Question

page

Time left 1:52:11

Hide

Mr. Molla has a fecal impaction. The nurse correctly administers an oil-retention enema by doing which of the following?

- ☐ a. Instructing the patient to retain the enema for at least 30 minutes
- ☐ b. Administering the enema while the patient is sitting on the toilet
- ☐ c. Mixing milk and molasses in equal parts for an enema
- ☐ d. Administering a large volume of solution (500–1,000 mL)

Next page



MoEEP

Back

Question 86

Not yet
answered

Marked out of
1.00

Flag question

Time left 0:38:53

Hide

During assessment of a patient the nurse notes an area of red, sharply defined plaques covered with silvery scales that are mildly itchy on the patient's knees and elbows. The nurse recognizes this finding as

- ☒ a. Vertigo
- ☐ b. Psoriasis
- ☐ c. Actinic keratosis
- ☐ d. Seborrheic keratosis

Previous page

Next page

ack

Question 10

et
red
d out of
question


Time left 1:56:31

How can nurses incorporate creativity in promoting breastfeeding success?

- ☐ a. Providing formula supplementation without exploring other options
- ☐ b. Restricting breastfeeding practices to a strict schedule and regimen
- ☐ c. Discouraging mothers from seeking lactation support or consulting professionals
- ☐ d. Encouraging the use of different breastfeeding positions and techniques

s page

Ne



Back

Time left 1:38:22

Hide

Exam navigation

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75

Question 28

Not yet
answered

Marked out of
1.00

Flag question

Which of the following actions demonstrates the nurse's commitment to providing compassionate care during the postpartum period?

- ☐ a. Dismissing the mother's emotional distress as common postpartum blues
- ☐ b. Rushing through postpartum education and discharge planning
- ☐ c. Focusing solely on physical health assessments and ignoring emotional needs
- ☐ d. Showing genuine interest and actively listening to the concerns of the new mother

Next page

Previous page

[Back](#)**Question 13**Not yet
answeredMarked out of
1.00 Flag question

Time left 1:53:25

Patients with eye inflammation or an eye infection should be taught:

- ☐ a. Those acute conditions commonly lead to chronic problems
- ☐ b. That regular, careful hand washing may prevent the infection from spreading
- ☐ c. To wear dark glasses to prevent irritation from UV light
- ☐ d. To apply a cold washcloth with pressure to the inflamed area frequently

[Previous page](#)

Back

Question 16

Not yet
answered

Marked out of
1.00

Flag question

Which one of the following is the diagnostic method of endemic typhus?

- ☐ a. blood film
- ☐ b. sputum smear
- ☐ c. Weil Felix agglutination test (serology)
- ☐ d. Widal test

Previous page





MoEEP

Back

Time left 1:43:20

Hide

Question 25

Not yet
answered

Marked out of
1.00

Flag question

72-year-old woman is taking several medications on a regular basis. Which of the following categories of drugs would be most likely to increase her surgical risk?



- ☐ a. Antacids
- ☐ b. Anticoagulants
- ☐ c. Sedatives
- ☐ d. Laxatives

Previous page

Next page

Exam nav

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74
82	83



Back

Time left 0:39:39

Hide

Question 85

Not yet
answered

Marked out of
1.00

Flag question

Which of the following measures is used to assess the strength of association between exposure and disease in a case-control study?

- ☐ a. Attributable risk
- ☐ b. Odds ratio
- ☐ c. Relative risk
- ☐ d. Hazard ratio

Previous page

Next page

Exam na

1 2

10 1

19

28

37

46

5



Back

Question 92

Not yet
answered

Marked out of
1.00

Flag question

A researcher who is providing therapy for a person should not develop a close friendship with the client because such dual relationships can compromise the success of the therapy. This problem relates to which General Ethical Principle?

- ☐ a. Respect for people's rights and dignity
- ☒ b. Fidelity and responsibility
- ☐ c. Justice
- ☐ d. Beneficence and nonmaleficence

Clear my choice

Time left 0:33:41

Hide

Previous page

Next page

Exam navigation

1 2 3

10 11

19 20

28 29

37 38


46 47

55 56

64 65

73 74

82 83



Back

Question 94

Not yet answered

Marked out of 1.00

Flag question

A 35 year old mother brought a 4 month baby to under 5 OPD where you are working. When you evaluate the baby the baby is not well attached to breast & not suckling effectively, has more than 18 breastfeeds in 24 hours and didn't start other foods or drinks. Which of the following classification best describe this condition?

- ☐ a. Sever developmental delay
- ☐ b. No feeding problem and not underweight
- ☐ c. No developmental delay
- ☐ d. Feeding problem or underweight

Time left 0:32:15

Hide

Previous page

Next page

Exam navigation

1	2	3	4
10	11	12	13
19	20	21	22
28	29	30	31
37	38	39	40
46	47	48	49
55	56	57	58
64	65	66	67
73	74	75	76
82	83	84	85

Question 17

yet
wered

ked out of

ag question

Which of the following is an example of a nurse leader demonstrating ethical behavior?

- ☐ a. Prioritizing their own interests over the well-being of patients
- ☐ b. Engaging in dishonest practices for personal gain
- ☐ c. Advocating for patient rights and autonomy
- ☐ d. Ignoring ethical guidelines and standards



Back

Question 91

yet
ered
d out of
question

s page

Which of the following action demonstrates the nurse's creative problem-solving skills in a postpartum complication scenario?

- ☐ a. Dismissing alternative treatment options without consideration
- ☐ b. Ignoring the signs and symptoms of complications
- ☐ c. Maintaining a rigid treatment plan without adjustments
- ☐ d. Seeking input from the healthcare team to explore different solutions

Time left 0:34:20

Hide

Next page

Exam

1

10

19

28

37

46

55

64

73

Back

Time left 0:33

Question 93

Not yet
answered

Marked out of
1.00

Flag question

Which one of the following is priority nursing intervention for HIV exposed neonate?

- ☐ a. Give Co-trimoxazole from 5 weeks
- ☐ b. Give Co-trimoxazole from 4 weeks
- ☐ c. Assess feeding and counsel
- ☐ d. Link only neonate to ART clinic and start ART

Previous page

[Back](#)

Time left 0:30:21

[Hide](#)**Question 98**Not yet
answeredMarked out of
1.00[Flag question](#)

Which of the following best describes your attitude towards patients who are non-compliant with their treatment plan?

- ☐ a. I become annoyed and judgmental towards these patients.
- ☐ b. I try to understand the reasons behind the non-compliance and find alternative solutions.
- ☐ c. I feel frustrated and give up on these patients, believing they will never change.
- ☐ d. I dismiss these patients as uncooperative and give up on trying helping them.

[Previous page](#)[Next page](#)



MoEEP

Back

Time left 0:27:25

Hide

Question 99

Not yet
answered

Marked out of
1.00

Flag question

The nurse demonstrates to a patient scheduled for a total hip replacement that it is important after surgery to avoid:

- ☐ a. Bearing weight on the affected leg after 4 to 6 weeks
- ☐ b. Sitting with the legs crossed
- ☐ c. Sleeping on the abdomen
- ☐ d. Abduction exercises of the affected leg

Previous page

Next page



MoEEP

Back

Question 58

Not yet
answered

Marked out of
1.00

Flag question

Bladder training interventions in a male patient who has urinary incontinence after a stroke includes:

- ☐ a. Keeping a urinal in place at all times
- ☐ b. Catheterizing the patient every 4 hours
- ☐ c. Limiting fluid intake
- ☐ d. Assisting the patient to stand to void

Time left 1:10:59

Hide

Previous page

Next page

Exam navigation

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75



MoEEP

Back

Time left 1:10:08

Hide

Question 59

Not yet
answered

Marked out of
1.00

Flag question

Administering or providing opioids to a dying patient who is experiencing moderate to severe pain:

- ☐ a. Will likely hasten the person's death.
- ☐ b. Is an appropriate nursing action.
- ☐ c. May cause addiction.
- ☐ d. Will probably be ineffective.

Previous page

Next page


(Comprehensive)

Time left 1:30:37

Which of the following terms best describes the synthesis of major features of decreasing risk of transmission of blood borne pathogens and body substance isolation?

- ☐ a. Universal Precautions
- ☐ b. Medical Asepsis
- ☐ c. Standard Precautions
- ☐ d. Protective Isolation

Next page



Back

Time left 1:11:32

Hide

Question 57

Not yet
answered

Marked out of
1.00

Flag question

The factor related to cerebral blood flow that most often determines the extent of cerebral damage from a stroke is the:

- ☐ a. Oxygen content of the blood
- ☐ b. Level of carbon dioxide in the blood
- ☐ c. Amount of cardiac output
- ☐ d. Degree of collateral circulation

Previous page

Next page

Exam nav

1

2

10

11

19

20

28

29

37

38

46

47

55

56

64

65

73

74

82

83



MoEEP

[Back](#)

Time left 1:07:13

[Hide](#)**Question 66**Not yet
answeredMarked out of
1.00[Flag question](#)

A patient has recently been diagnosed with early stages of breast cancer. Which of the following is most appropriate intervention for the nurse to focus on?

- ☐ a. Discussing the patient's past experiences with her grandmother's cancer
- ☐ b. Preparing a will and advance directives
- ☐ c. Discussing replacement child care for patient's children
- ☐ d. Maintaining patient's hope

[Previous page](#)[Next page](#)

Exam na

1

2

10

11

19

20

28

29

37

38

46

47

55

56

64

65

73


74

82

83

91

92



MoEEP

Back

Time left 1:16:14

Hide

Question 51

Not yet
answered

Marked out of
1.00

Flag question

In planning nursing interventions to increase bladder control in the patient with urinary incontinence, the nurse includes:

- ☐ a. Teaching the patient biofeedback mechanisms to suppress the urge to void
- ☐ b. Counseling the patient concerning choice of incontinence containment device
- ☐ c. Teaching the patient to use Kegel exercises
- ☐ d. Clamping and releasing a catheter to increase bladder tone

Previous page

Next page

Exam na

1	2
10	11
19	20
28	29
37	38
46	47



MoEEP

Back

Question 43

Not yet
answered

Marked out of
1.00

Flag question

A patient with psoriasis tells the nurse that she has quit her job as a receptionist because she feels her appearance is disgusting to customers. The nursing diagnosis that best describes this patient response is

- ☐ a. Social isolation related to fear of rejection secondary to presence of lesions
- ☐ b. Ineffective coping related to lack of social support.
- ☐ c. Anxiety related to lack of knowledge of the disease process.
- ☐ d. Impaired skin integrity related to presence of lesions.

Time left 1:21:59

Hide

Previous page

Next page

Exam navigation

1	2	3	4	5	6	7
10	11	12	13	14	15	16
19	20	21	22	23	24	25
28	29	30	31	32	33	34
37	38	39	40	41	42	43
46	47	48	49	50	51	52
55	56	57	58	59	60	61
64	65	66	67	68	69	70
73	74	75	76	77	78	79
82	83	84	85	86	87	88
91	92	93	94	95	96	97



MoEEP

Back

Time left 1:35:08

Hide

Question 34

Not yet
answered

Marked out of
1.00

Flag question

Which of the following describes an odds ratio of 1?

- ☐ a. The exposure is not associated with the outcome
- ☐ b. There is no association between the exposure and the outcome
- ☐ c. The exposure is associated with a decreased risk of the outcome
- ☐ d. The exposure is associated with an increased risk of the outcome

Previous page

Next page

Exam n

1


2

10

19

21

3

[Back](#)

Time left 1:15:24

[Hide](#)**Question 52**Not yet
answeredMarked out of
1.00[Flag question](#)

When language differences exist between the nurse and client, the nurse should do which of the following first:

- ☐ a. Use a bilingual family member to facilitate assessment/care
- ☐ b. Act in the client's best interest
- ☐ c. Obtain a key informant to navigate patient's cultural cue
- ☐ d. Establish whether interpreter is needed

[Previous page](#)[Next page](#)

Exam navigation

1

2

10

11

19

20

28

29

37

46

55

64

73

Back

Question 33

Not yet
answered

Marked out of
1.00

Flag question

Time left 1:36:01

Hide

Which of the following is defined as death of a woman while pregnant, or within 42 days of termination of pregnancy, from any cause related to, or aggravated by, the pregnancy or its management, but not from accidental or incidental death?

- ☐ a. Direct maternal death
- ☐ b. Maternal death
- ☐ c. Indirect maternal death
- ☐ d. Maternal mortality rate

Previous page

Next page



MoEEP

[Back](#)

Time left 1:19:54

[Hide](#)**Question 47**Not yet
answeredMarked out of
1.00[Flag question](#)

A patient who has fallen and injured his wrist carefully cradles it with the other hand. The patient is demonstrating which of the following responses to pain?

- ☐ a. Physiologic
- ☐ b. Affective
- ☐ c. Behavioral
- ☐ d. Involuntary

[Previous page](#)[Next page](#)

Exam naviga

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75
82	83	84
91	92	93

[Back](#)

Time left 1:17:44

Question 49Not yet
answeredMarked out of
1.00[Flag question](#)

The nurse would increase the comfort of the patient with appendicitis by:

- ☐ a. Turning the patient onto his or her left side
- ☐ b. Sitting the patient upright in a chair
- ☐ c. Having the patient lie prone
- ☐ d. Flexing the patient's right knee

[Previous page](#)[Next](#)



MoEEP

Back

Time left 1:10:08

Hide

Question 59

Not yet
answered

Marked out of
1.00

Flag question

Administering or providing opioids to a dying patient who is experiencing moderate to severe pain:

- ☐ a. Will likely hasten the person's death.
- ☐ b. Is an appropriate nursing action.
- ☐ c. May cause addiction.
- ☐ d. Will probably be ineffective.

Previous page

Next page

(Comprehensive)

Time left 1:30:37

Which of the following terms best describes the synthesis of major features of decreasing risk of transmission of blood borne pathogens and body substance isolation?

- ☐ a. Universal Precautions
- ☐ b. Medical Asepsis
- ☐ c. Standard Precautions
- ☐ d. Protective isolation

Next page



MoEEP

[Back](#)

Time left 1:07:13

[Hide](#)**Question 66**Not yet
answeredMarked out of
1.00[Flag question](#)

A patient has recently been diagnosed with early stages of breast cancer. Which of the following is most appropriate intervention for the nurse to focus on?

- ☐ a. Discussing the patient's past experiences with her grandmother's cancer
- ☐ b. Preparing a will and advance directives
- ☐ c. Discussing replacement child care for patient's children
- ☐ d. Maintaining patient's hope

[Previous page](#)[Next page](#)

Exam na

1

2

10

11

19

20

28

29

37

38

46

47

55

56

64

65

73

74

82

83

91

92



MoEEP

Back

Time left 1:16:14

Hide

Question 51

Not yet
answered

Marked out of
1.00

Flag question

In planning nursing interventions to increase bladder control in the patient with urinary incontinence, the nurse includes:


- ☐ a. Teaching the patient biofeedback mechanisms to suppress the urge to void
- ☐ b. Counseling the patient concerning choice of incontinence containment device
- ☐ c. Teaching the patient to use Kegel exercises
- ☐ d. Clamping and releasing a catheter to increase bladder tone

Previous page

Next page

Exam na

1	2
10	11
19	20
28	29
37	38
46	47



Back

Time left 1:11:32

Hide

Question 57

Not yet
answered

Marked out of
1.00

Flag question

The factor related to cerebral blood flow that most often determines the extent of cerebral damage from a stroke is the:

- ☐ a. Oxygen content of the blood
- ☐ b. Level of carbon dioxide in the blood
- ☐ c. Amount of cardiac output
- ☐ d. Degree of collateral circulation

Previous page

Next page

Exam nav

1

2

10

11

19

20

28

29

37

38

46

47

55

64

73

82



MoEEP

Back

Question 43

Not yet
answered

Marked out of
1.00

Flag question

A patient with psoriasis tells the nurse that she has quit her job as a receptionist because she feels her appearance is disgusting to customers. The nursing diagnosis that best describes this patient response is

- ☐ a. Social isolation related to fear of rejection secondary to presence of lesions
- ☐ b. Ineffective coping related to lack of social support.
- ☐ c. Anxiety related to lack of knowledge of the disease process.
- ☐ d. Impaired skin integrity related to presence of lesions.

Time left 1:21:59

Hide

Previous page

Next page

Exam navigation

1	2	3	4	5	6	7
10	11	12	13	14	15	16
19	20	21	22	23	24	25
28	29	30	31	32	33	34
37	38	39	40	41	42	43
46	47	48	49	50	51	52
55	56	57	58	59	60	61
64	65	66	67	68	69	70
73	74	75	76	77	78	79
82	83	84	85	86	87	88
91	92	93	94	95	96	97



MoEEP

Back

Time left 1:35:08

Hide

Question 34

Not yet
answered

Marked out of
1.00

Flag question

Which of the following describes an odds ratio of 1?

- ☐ a. The exposure is not associated with the outcome
- ☐ b. There is no association between the exposure and the outcome
- ☐ c. The exposure is associated with a decreased risk of the outcome
- ☐ d. The exposure is associated with an increased risk of the outcome

Previous page

Next page

Exam n

1

2

10

19

21

3



MoEEP

[Back](#)

Time left 1:19:54

[Hide](#)**Question 47**Not yet
answeredMarked out of
1.00[Flag question](#)

A patient who has fallen and injured his wrist carefully cradles it with the other hand. The patient is demonstrating which of the following responses to pain?

- ☐ a. Physiologic
- ☐ b. Affective
- ☐ c. Behavioral
- ☐ d. Involuntary

[Previous page](#)[Next page](#)

Exam navig

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75
82	83	84
91	92	93



MoEEP

Back

Question 58

Not yet
answered

Marked out of
1.00

Flag question

Bladder training interventions in a male patient who has urinary incontinence after a stroke includes:

- ☐ a. Keeping a urinal in place at all times
- ☐ b. Catheterizing the patient every 4 hours
- ☐ c. Limiting fluid intake
- ☐ d. Assisting the patient to stand to void

Time left 1:10:59

Hide

Previous page

Next page

Exam navigation

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75

Back

Question 33

Not yet
answered

Marked out of
1.00

Flag question

Time left 1:36:01

Hide

Which of the following is defined as death of a woman while pregnant, or within 42 days of termination of pregnancy, from any cause related to, or aggravated by, the pregnancy or its management, but not from accidental or incidental death?

- ☐ a. Direct maternal death
- ☐ b. Maternal death
- ☐ c. Indirect maternal death
- ☐ d. Maternal mortality rate

Previous page

Next page



Back

Time left 1:15:24

Hide

Question 52

Not yet
answered

Marked out of
1.00

Flag question

When language differences exist between the nurse and client, the nurse should do which of the following first:

- ☐ a. Use a bilingual family member to facilitate assessment/care
- ☐ b. Act in the client's best interest
- ☐ c. Obtain a key informant to navigate patient's cultural cue
- ☐ d. Establish whether interpreter is needed

Previous page

Next page

Exam navigation

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74
82	83
91	92
100	

[Back](#)

Time left 1:17:44

Question 49Not yet
answeredMarked out of
1.00[Flag question](#)

The nurse would increase the comfort of the patient with appendicitis by:

- ☐ a. Turning the patient onto his or her left side
- ☐ b. Sitting the patient upright in a chair
- ☐ c. Having the patient lie prone
- ☐ d. Flexing the patient's right knee

[Previous page](#)[Next](#)

Question 42

Not yet
answeredMarked out of
1.00

Flag question

Captopril may be administered to a client with HF because it acts as a:

- ☐ a. Vasodilator
- ☐ b. Vasopressor
- ☐ c. Potassium-sparing diuretics
- ☐ d. Volume expander

Previous page

Next page



MoEEP

Back

Question 56

Not yet
answered

Marked out of
1.00

Flag question

When planning care for the infant diagnosed with cleft lip and palate, which action would the nurse take in relation to the priority nursing diagnosis for this child?

- ☐ a. Burp the baby well throughout feedings
- ☐ b. Temporarily refrain from having the baby breastfeed
- ☐ c. Encourage mother to use false palate covering when feeding baby
- ☐ d. Prevent the baby from vigorously crying

Time left 1:12:47

Hide

Previous page

Next page

Exam navigation

1	2	3	4
10	11	12	13
19	20	21	22
28	29	30	31
37	38	39	40
46	47	48	49
55	56	57	58
64	65	66	67
73	74	75	76
82	83	84	85

ing (Comprehensive)

Time left 2:38:52

Which one of the following is the most important for the nurse to assess the client's with anxiety beginning treatment with lorazepam?

- ☐ a. Use of coping mechanisms.
- ☐ b. Family and social support.
- ☐ c. Motivation for treatment.
- ☐ d. Use of alcohol.



MoEEP

Back

Time left 1:16:14

Hide

Question 51

Not yet
answered

Marked out of
1.00

Flag question

In planning nursing interventions to increase bladder control in the patient with urinary incontinence, the nurse includes:

- ☐ a. Teaching the patient biofeedback mechanisms to suppress the urge to void
- ☐ b. Counseling the patient concerning choice of incontinence containment device
- ☐ c. Teaching the patient to use Kegel exercises
- ☐ d. Clamping and releasing a catheter to increase bladder tone

Previous page

Next page

Exam na

1	2
10	11
19	20
28	29
37	38
46	47

Time left 1:25:30

Hide


When explaining the action of a hypertonic solution enema, the nurse incorporates which of the following as the basis for action?

- ☐ a. Softening of fecal contents
- ☐ b. Diffusion of water out of colon
- ☐ c. Bowel mucosa irritation
- ☐ d. Osmosis of water into colon

Next page

Exam navigation

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75
82	83	84
91	92	93



MoEEP

Back

Time left 1:23:50

Hide

Question 40

Not yet
answeredMarked out of
1.00

Flag question

One of your clients with mania begins dancing in the ward. When she twisted her skirt in front of the male clients, it was obvious she had no underwear on. You as a nurse distract her and take her to her room to put on underwear. Which of the following best describe your action?

- ☐ a. Avoid embarrassing the male clients who are watching.
- ☐ b. Minimize the client's embarrassment about her present behavior.
- ☐ c. Teach her about proper attire and hygiene.
- ☐ d. Keep her from dancing with other clients.

Previous page

Next page

Exam navigation

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50
55	56	57	58	59
64	65	66	67	68
73	74	75	76	77
82	83	84	85	86
91	92	93	94	95

Time left 2:25:12

Hide

Question 3

Not yet
answered

Marked out of

Flag question

The physician has ordered an indwelling catheter inserted in a hospitalized male patient. The nurse is aware of which of the following considerations?

- ☐ a. In the hospital, a clean technique is used for catheter insertion
- ☐ b. The male urethra is more vulnerable to injury during insertion
- ☐ c. The catheter is inserted 2 to 3 cm. into the meatus
- ☐ d. Since it uses a closed system, the risk for urinary tract infection is absent

Previous page

Next page

Exam navigation

	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50

[Back](#)**Question 41**Not yet
answeredMarked out of
1.00[Flag question](#)

A 28 years mother has a 4 weeks baby on exclusive breast feeding. The mother is reactive to HIV antibody test. The baby also tested for HIV status and the result is Positive antibody test, but DNA PCR is negative. Which of the following is the best classification of this neonate?


- ☐ a. HIV exposed
- ☐ b. HIV status unknown
- ☐ c. HIV infection unlikely
- ☐ d. HIV infected

Time left 1:23:20

[Hide](#)**Exam navigation**

1	2	3	4
10	11	12	13
19	20	21	22
28	29	30	31
37	38	39	40
46	47	48	49
55	56	57	58
64	65	66	67
73	74	75	76
82	83	84	85
91	92	93	94

[Next page](#)[Previous page](#)



Nursing (Comprehensive)

Back

Time left 0:30

Question 100

Not yet
answered

Marked out of
1.00

Flag question

The most common response of the stroke patient to the change in body image is:

- ☒ a. Denial
- ☐ b. Intellectualization
- ☐ c. Disassociation
- ☐ d. Depression

Clear my choice

Previous page

Finish



MoEEP

Back

Time left 1:21:27

Hide

Question 44

Not yet
answered

Marked out of
1.00

Flag question

Which shistosoma species known to cause Urinary Tract Infection that leads to blood in the urine?

- ☐ a. Shistosoma Japonicum
- ☐ b. Shistosoma Mansoni
- ☐ c. Shistosoma Intercalatum
- ☐ d. Shistosoma Heamatobium

Previous page

Next page



Exam n

1 2


10 1

19

28

37

4



Nursing (Comprehensive)

Back

Question 88

Not yet answered

Marked out of 1.00

Flag question

The nurse would increase the comfort of the patient with appendicitis by:

- ☐ a. Flexing the patient's right knee
- ☐ b. Having the patient lie prone
- ☐ c. Sitting the patient upright in a chair
- ☐ d. Turning the patient onto his or her left side

Previous page



Which one of the following is priority nursing intervention for HIV exposed neonate?

- ☐ a. Give Co-trimoxazole from 4 weeks
- ☐ b. Give Co-trimoxazole from 5 weeks
- ☐ c. Link only neonate to ART clinic and start ART
- ☒ d. Assess feeding and counsel