

# Sequence (Lecture- 4)

Engineering Calculus



School of Engineering and Applied Sciences  
Department of Mathematics  
Bennett University

## Theorem

Let  $\{a_n\}_1^\infty$  and  $\{b_n\}_1^\infty$  be two sequences such that  $\lim_{n \rightarrow \infty} a_n = L$  and  $\lim_{n \rightarrow \infty} b_n = M$ . Then

- (i)  $\lim_{n \rightarrow \infty} (a_n + b_n) = L + M$ .
- (ii)  $\lim_{n \rightarrow \infty} (a_n - b_n) = L - M$ .
- (iii)  $\lim_{n \rightarrow \infty} (ca_n) = cL, \quad c \in \mathbb{R}$ .
- (iv)  $\lim_{n \rightarrow \infty} (a_nb_n) = LM$ .
- (v)  $\lim_{n \rightarrow \infty} \left( \frac{a_n}{b_n} \right) = \frac{L}{M}$  if  $M \neq 0$ .

## Examples

Find the limit of the following sequences:

- (i)  $\left\{ \frac{5}{n^2} \right\}_1^\infty$ , (ii)  $\left\{ \frac{3n^2 - 6n}{5n^2 + 4} \right\}_1^\infty$ , (iii)  $\lim_{n \rightarrow \infty} \left( \frac{n-1}{n} \right)$ .

**Solution:** (i)

$$\lim_{n \rightarrow \infty} \frac{5}{n^2} = \lim_{n \rightarrow \infty} 5 \cdot \frac{1}{n} \cdot \frac{1}{n} = 5 \cdot \lim_{n \rightarrow \infty} \frac{1}{n} \cdot \lim_{n \rightarrow \infty} \frac{1}{n} = 5 \cdot 0 \cdot 0 = 0.$$

(ii) Notice that

$$\frac{3n^2 - 6n}{5n^2 + 4} = \frac{3 - 6/n}{5 + 4/n^2}.$$

Now

$$\lim_{n \rightarrow \infty} (3 - 6/n) = 3 - 6 \lim_{n \rightarrow \infty} 1/n = 3 - 6 \cdot 0 = 3$$

and

$$\lim_{n \rightarrow \infty} (5 + 4/n^2) = 5 + 4 \lim_{n \rightarrow \infty} 1/n^2 = 5 + 4 \cdot 0 = 5.$$

Therefore,

$$\lim_{n \rightarrow \infty} \frac{3n^2 - 6n}{5n^2 + 4} = \lim_{n \rightarrow \infty} \frac{3 - 6/n}{5 + 4/n^2} = \frac{3}{5}.$$

(iii)

$$\lim_{n \rightarrow \infty} \left( \frac{n-1}{n} \right) = \lim_{n \rightarrow \infty} \frac{1 - 1/n}{1} = 1 - \lim_{n \rightarrow \infty} \frac{1}{n} = 1 - 0 = 1.$$

## Sandwich theorem for sequences

Let  $\{a_n\}$ ,  $\{b_n\}$  and  $\{c_n\}$  be three sequences such that  $a_n \leq b_n \leq c_n$  for all  $n \in \mathbb{N}$ . If  $\lim_{n \rightarrow \infty} a_n = L$  and  $\lim_{n \rightarrow \infty} c_n = L$ , then  $\lim_{n \rightarrow \infty} b_n = L$ .

**Proof:** Let  $\epsilon > 0$  be given. As  $\lim_{n \rightarrow \infty} a_n = L$ , there exists  $N_1 \in \mathbb{N}$  such that

$$n \geq N_1 \implies |a_n - L| < \epsilon. \quad (1)$$

Similarly as  $\lim_{n \rightarrow \infty} c_n = L$ , there exists  $N_2 \in \mathbb{N}$

$$n \geq N_2 \implies |c_n - L| < \epsilon. \quad (2)$$

Let  $N = \max\{N_1, N_2\}$ . Then,  $L - \epsilon < a_n$  (from (1)) and  $c_n < L + \epsilon$  (from (2)). Thus

$$L - \epsilon < a_n \leq b_n \leq c_n < L + \epsilon.$$

Thus  $|b_n - L| < \epsilon$  for all  $n \geq N$ . Hence the proof.

## Examples

Using Sandwich theorem, prove the following:

(i)  $\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$

(ii)  $\lim_{n \rightarrow \infty} \frac{1}{2^n} = 0.$

(iii)  $\lim_{n \rightarrow \infty} (-1)^n \frac{1}{n} = 0.$

(iv) If  $0 < b < 1$ , then  $\lim_{n \rightarrow \infty} b^n = 0.$

(v)  $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1.$

**Solution:** (i) Consider the sequence  $\left\{ \frac{\cos n}{n} \right\}_{n=1}^{\infty}$ . Then  $\frac{-1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}$ . Hence by Sandwich theorem,  $\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$

(ii) As  $0 < \frac{1}{2^n} < \frac{1}{n}$  and  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ ,  $\frac{1}{2^n}$  also converges to 0 by Sandwich theorem.

(iii) As  $\frac{-1}{n} \leq (-1)^n \frac{1}{n} \leq \frac{1}{n}$  for all  $n \geq 1$  and  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ ,  $(-1)^n \frac{1}{n}$  also converges to 0 by Sandwich theorem.

(iv) Since  $0 < b < 1$ , we can write  $b = \frac{1}{1+a}$ , where  $a := \frac{1}{b} - 1$  so that  $a > 0$ . Also we have  $(1+a)^n \geq 1+na$ . Hence

$$0 < b^n = \frac{1}{(1+a)^n} \leq \frac{1}{1+na} < \frac{1}{na}.$$

So, by sandwich Theorem, we conclude that  $\lim_{n \rightarrow \infty} b^n = 0$ .

(v) Let  $a_n = n^{\frac{1}{n}} - 1$ . Then  $0 \leq a_n < 1$  for all  $n \in \mathbb{N}$ . Further,

$$n = (1+a_n)^n \geq \frac{n(n-1)}{2} a_n^2.$$

Thus  $0 \leq a_n \leq \sqrt{\frac{2}{(n-1)}}$  ( $n \geq 2$ ). As  $\sqrt{\frac{2}{(n-1)}} \rightarrow 0$  as  $n \rightarrow \infty$ , by Sandwich theorem,  $a_n \rightarrow 0$ , i.e.,  $n^{\frac{1}{n}} \rightarrow 1$  as  $n \rightarrow \infty$ .

*Thank  
You*