

# AI1103 : Assignment 2

Amaan - EP20BTECH11003

Download all python codes from

<https://github.com/amaan28/Assignment2/blob/main/Assignment2/codes/Assignment2.py>

and latex-tikz codes from

<https://github.com/amaan28/Assignment2/blob/main/Assignment2/Assignment2.tex>

$$P = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 0.5 & 0 & 0.5 \\ 0.5 & 0 & 0.5 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix} \quad (47.2)$$

Now, it can be seen that the states 1, 2 are transient, while 3, 4 are absorbing. The standard form of a state transition matrix is,

$$P = \begin{matrix} & \begin{matrix} A & N \end{matrix} \\ \begin{matrix} A \\ N \end{matrix} & \begin{bmatrix} I & O \\ R & Q \end{bmatrix} \end{matrix} \quad (47.3)$$

where,

TABLE 2: Notations and their meanings

Notation	Meaning
$A$	All absorbing states
$N$	All non-absorbing states
$I$	Identity matrix
$O$	Zero matrix
$R, Q$	Other submatrices

- 1)  $\frac{1}{3}$
- 2)  $\frac{1}{5}$
- 3)  $\frac{2}{5}$
- 4)  $\frac{3}{4}$

## GATE 2012 EE Q.47 - SOLUTION

Given, a fair coin is tossed till heads turns up.

$$p = \frac{1}{2}, q = \frac{1}{2} \quad (47.1)$$

Let us define a Markov chain  $\{X_0, X_1, X_2, \dots\}$ , where  $X_n \in S = \{1, 2, 3, 4\}$  where  $n \in \{0, 1, 2, \dots\}$ , such that the state transition matrix for the Markov chain is,

TABLE 1: States and their notations

Notation	State
$S = 1$	Even numbered toss
$S = 2$	Odd numbered toss
$S = 3$	Head appears in odd number of tosses
$S = 4$	Head appears in even number of tosses

Converting (47.2) to standard form, we get

$$P = \begin{matrix} & \begin{matrix} 3 & 4 & 1 & 2 \end{matrix} \\ \begin{matrix} 3 \\ 4 \\ 1 \\ 2 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0.5 & 0 & 0.5 \\ 0.5 & 0 & 0.5 & 0 \end{bmatrix} \end{matrix} \quad (47.4)$$

From (47.4),

$$R = \begin{bmatrix} 0 & 0.5 \\ 0.5 & 0 \end{bmatrix}, Q = \begin{bmatrix} 0 & 0.5 \\ 0.5 & 0 \end{bmatrix} \quad (47.5)$$

The limiting matrix for absorbing Markov chain is,

$$\bar{P} = \begin{bmatrix} I & O \\ FR & O \end{bmatrix} \quad (47.6)$$

where,

$$F = (I - Q)^{-1} \quad (47.7)$$

is called the fundamental matrix of  $P$ .

On solving, we get

$$\bar{P} = \begin{matrix} & \begin{matrix} 3 & 4 & 1 & 2 \end{matrix} \\ \begin{matrix} 3 \\ 4 \\ 1 \\ 2 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \frac{1}{3} & \frac{2}{3} & 0 & 0 \\ \frac{2}{3} & \frac{1}{3} & 0 & 0 \end{bmatrix} \end{matrix} \quad (47.8)$$

A element  $\bar{p}_{ij}$  of  $\bar{P}$  denotes the absorption probability in state  $j$ , starting from state  $i$ . Hence, the required probability is,

$$P = \bar{p}_{14} = \frac{2}{3} \quad (47.9)$$

Therefore, options 3) is correct.

