

6

Iterative Methods

Methods

The goal is to solve

$$Ax = b.$$

Jacobi Method

You write $A = D + R$ where

$$D = \begin{bmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix} \text{ and } R = A - D = \begin{bmatrix} 0 & a_{12} & \cdots & a_{1n} \\ a_{21} & 0 & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & 0 \end{bmatrix}$$

The algorithm is

$$Dx^{k+1} + Rx^k = b \implies x^{k+1} = D^{-1}(b - Rx^k)$$
$$x_i^{k+1} = \frac{1}{a_{ii}} \left(b_i - \sum_{j \neq i} a_{ij} x_j^k \right)$$

Gauss-Seidel Method

You write $A = L + U$ where

$$L = \begin{bmatrix} a_{11} & 0 & \cdots & 0 \\ a_{21} & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \text{ and } U = \begin{bmatrix} 0 & a_{12} & \cdots & a_{1n} \\ 0 & 0 & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}$$

The algorithm is

$$Lx^{k+1} + Ux^k = b \implies x^{k+1} = L^{-1}(b - Ux^k)$$
$$x_i^{k+1} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{k+1} - \sum_{j=i+1}^n a_{ij} x_j^k \right)$$

Exercise

Exercise 1: Write a function that solve the system $Ax = b$ using the Jacobi method. Then test your functions with a test case, and explain why it works.

Jacobi converge if $\sum_{j \neq i} |a_{ij}| < |a_{ii}|$. For example you can pick

```
A = np.array([[10, 1, 3,], [2, 30, 1], [3, 5, 25]])
```

Exercise 2: Write a function that solve the system $Ax = b$ using the Gauss-Seidel method. Then test your functions with a test case, and explain why it works.

Exercise 3: Solve the following system using Jacobi and Gauss-Seidel methods. Specify how many iterations were needed.

$$\begin{cases} 2x_1 + x_2 &= 0 \\ -x_1 + 3x_2 + x_3 &= -4 \\ -x_2 + 3x_3 + x_4 &= 12 \\ -x_3 + 4x_4 + x_5 &= 6 \\ -x_4 + 2x_5 &= 9 \end{cases}$$

Exercise 4: Solve the tridiagonal system $Ax = f$ (using Jacobi and Gauss-Seidel methods) with

$$A_{ii} = 4, \quad A_{i,i-1} = A_{i,i+1} = 1$$

for all i . Let the order of the system be $n = 100$, and let

$$f = [1, 1, \dots, 1]^T$$

Exercise 5: By using an iterative method solve the linear system $x = b + Mx$ with

$$M_{ij} = \frac{1}{2n} \left[\frac{t_i^3}{1 + t_j} + 1 \right], \quad b_j = \frac{1}{4} + t_i - \frac{1}{2} t_i^3,$$

and $t_i = (2i - 1)/2n$.

The true solution is

$$x_i = 1 + t_i \quad 1 \leq i \leq n$$

Solve this system for several values of n , ($n = 25, 50, 75, 100$). Then calculate/plot the error $x - x^{(k)}$, and the ratios with which they decrease (this means the convergence rate).