

Comps Practice

Aniruddha Madhava

January 8, 2026

Contents

Comps Lemma	2
Problem Comps Lemma	2
January 2025	2
Problem 2025-J-I-1 (Algebra)	2
Problem 2025-J-I-2 (Real Analysis)	2
Problem 2025-J-I-3 (Geometry/Topology)	3
Problem 2025-J-II-3 (Algebra)	3
Problem 2025-J-II-4 (Topology)	3
Problem 2025-J-II-5 (Analysis)	3
August 2025	4
Problem 2025-A-I-1 (Geometry/Topology)	4
Problem 2025-A-I-6 (Algebra)	4
Problem 2025-A-II-2 (Geometry/Topology)	4
January 2024	5
Problem 2024-J-I-1 (Algebra)	5
Problem 2024-J-I-2 (Geometry/Topology)	5
Problem 2024-J-I-4 (Algebra)	6
Problem 2024-J-I-5 (Geometry/Topology)	6
Problem 2024-J-II-3 (Geometry/Topology)	7
Problem 2024-J-II-6 (Geometry/Topology)	7
August 2023	7
Problem 2023-A-II-1 (Algebra)	7
January 2023	8
Problem 2023-J-II-4 (Geometry/Topology)	8
Problem 2023-A-I-2 (Geometry/Topology)	8
Problem 2023-A-II-5 (Geometry/Topology)	9
Problem 2023-A-I-1 (Algebra)	9
Problem 2023-A-II-6 (Complex Analysis)	10

Comps Lemma

Problem Comps Lemma. Let M, N be smooth, connected, n -manifolds, and $f : M \rightarrow N$ a (smooth) immersion. If M is compact and nonempty, then N is compact and f is a (smooth) covering map.

Let M, N be smooth, connected n -manifolds and $f : M \rightarrow N$ an immersion. Assume that M is compact and nonempty. Since $\dim N = n$ and f is an immersion, $\text{rank } df_p = n$ at every $p \in M$. Hence, by the Inverse Function Theorem, f is a local diffeomorphism. Since local diffeomorphisms are open maps, $f(M)$ is open in N . On the other hand, since the continuous image of compact sets is compact, $f(M)$ is compact in N . Since N is Hausdorff, $f(M)$ is closed in N . Since N is connected, $f(M) = N$. Therefore, N is compact.

Now, let $q \in N$, and consider $f^{-1}(q) \subset M$. For each $x \in f^{-1}(q)$, let U_x be an open neighborhood of M containing x . Since M is Hausdorff, we can shrink each U_x so that these neighborhoods are pairwise disjoint. This means that each $x \in f^{-1}(q)$ is isolated, and hence $f^{-1}(q)$ is discrete. Since M is compact, we conclude that $f^{-1}(q)$ must be finite; let $f^{-1}(q) = \{x_1, \dots, x_s\}$. As noted above, for each $j = 1, \dots, s$, let U_j be a neighborhood of x_j such that $f|_{U_j} : U_j \rightarrow V_j \subset N$ is a diffeomorphism. Then by the Hausdorff condition on M , shrink each U_j so that $U_i \cap U_j = \emptyset$ for all $i \neq j$; f remains a diffeomorphism on these shrunken neighborhoods. Setting $V = \bigcap_1^s f(U_j)$ and taking $\tilde{U}_j = f^{-1}(V) \cap U_j$ gives us an evenly covered neighborhood of q in N .

January 2025

Problem 2025-J-I-1 (Algebra). Let R be a UFD (unique factorization domain). Let F be its quotient field. Let $p(x) = x^n + b_{n-1}x^{n-1} + \dots + b_0 \in F[x]$ be a monic polynomial with coefficients in R admitting a root $a \in F$. Prove that $a \in R$.

Let R be a UFD, and F its quotient field. Let $p(x) = x^n + b_{n-1}x^{n-1} + \dots + b_0 \in F[x]$ be a monic polynomial with coefficients in R admitting a root $a \in F$. Let $a = c/d$, where $c, d \in R \setminus \{0\}$ so that $\gcd(c, d) = 1$. By definition of a root, we must have

$$0 = p(a) = \left(\frac{c}{d}\right)^n + b_{n-1}\left(\frac{c}{d}\right)^{n-1} + \dots + b_0. \quad (1)$$

Multiplying both sides by d^n ,

$$c^n + d(b_{n-1}c^{n-1} + b_{n-2}c^{n-2}d + \dots + b_0d^{n-1}) = 0 \implies c^n = -d(b_{n-1}c^{n-1} + \dots + b_0d^{n-1}). \quad (2)$$

From this, we observe that $d \mid c^n$. If d is not a unit in R , then every nonidentity irreducible divisor of d is an irreducible divisor of c^n , and hence an irreducible divisor of c . But this contradicts our hypothesis that $\gcd(c, d) = 1$. Hence, d has to be a unit of R . If $v \in R \setminus \{0\}$ such that $dv = vd = 1$, then

$$a = \frac{c}{d} = \frac{c}{d} \cdot \frac{v}{v} = cv \in R. \quad (3)$$

Hence, this concludes the proof.

Problem 2025-J-I-2 (Real Analysis). Let $\{f_n\}_{n \geq 1}$ be a sequence of Lebesgue-measurable functions on $[0, 1]$. Suppose that

$$\int_0^1 f_n^2 dm \leq \frac{1}{n^2} \quad \text{for all } n \geq 1.$$

Prove that f_n converges to 0 a.e. on $[0, 1]$.

Let $\{f_n\}_{n \geq 1}$ be a sequence of Lebesgue-measurable functions on $[0, 1]$ so that

$$\int_0^1 f_n^2 dm \leq \frac{1}{n^2} \quad \text{for all } n \geq 1. \quad (4)$$

Consider the sequence $\{\sum_1^m f_n^2\}$, which is increasing and converges a.e. to $\sum_1^\infty f_n^2$. Hence, by the Monotone Convergence Theorem,

$$\sum_1^\infty \int_0^1 f_n^2 = \lim_{m \rightarrow \infty} \sum_1^m \int_0^1 f_n^2 = \lim_{m \rightarrow \infty} \int_0^1 \sum_1^m f_n^2 = \int_0^1 \sum_1^\infty f_n^2 \leq \int_0^1 \sum_1^\infty \frac{1}{n^2} < \infty. \quad (5)$$

Therefore, $\sum_1^\infty f_n^2 \in L^1(\mathbb{R})$, which means that $\sum_1^\infty f_n^2 < \infty$ a.e. on $[0, 1]$. Hence, $\sum_{n=1}^\infty f_n^2$ converges a.e. on $[0, 1]$. This implies that $f_n^2 \rightarrow 0$ a.e. on $[0, 1]$, and hence $f_n \rightarrow 0$ a.e. on $[0, 1]$.

Problem 2025-J-I-3 (Geometry/Topology). Let M be an orientable, connected, and compact smooth n -manifold with boundary. Show that there is no (smooth) retraction to the boundary, that is, there does not exist a smooth map $f : M \rightarrow \partial M$ such that $f(x) = x$ when $x \in \partial M$.

Let M be an orientable, connected, and compact smooth n -manifold with boundary. Assume to the contrary that there exists a smooth map $f : M \rightarrow \partial M$ such that $f(x) = x$ when $x \in \partial M$. Let $\omega \in \Omega^{n-1}(\partial M)$ be a volume form for the boundary of M . Since volume forms are closed (hence, ω is closed), we have by Stokes's theorem

$$0 = \int_M f^* d\omega = \int_M \partial(f^* \omega) = \int_{\partial M} f^* \omega = \int_{\partial M} \omega > 0, \quad (6)$$

which is a contradiction. Hence, by contradiction, there cannot exist a smooth retraction to the boundary.

Problem 2025-J-II-3 (Algebra). Let V be a vector space of dimension n over \mathbb{Q} . Let $T : V \rightarrow V$ be a linear transformation with minimal polynomial $x^4 - x^2 - 2$ over \mathbb{Q} . Show that n must be even.

Consider V as a module over the ring $\mathbb{Q}[x]$ by letting a polynomial $f(x) \in \mathbb{Q}[x]$ act as the linear operator $f(T)$. Since $\dim V = n$, this module is finitely generated. By the structure theorem for finitely generated modules over principal ideal domains, V is isomorphic to a direct sum of modules of the form $\mathbb{Q}[x]/(p(x))^e$, where $p(x) \in \mathbb{Q}[x]$ is irreducible. Moreover, each $p(x)$ must divide the minimal polynomial of T . We note that over \mathbb{Q} ,

$$x^4 - x^2 - 2 = (x^2 - 2)(x^2 + 1), \quad (7)$$

where both factors are irreducible over \mathbb{Q} . Therefore, the only choices for $p(x)$ are $x^2 - 2$ and $x^2 + 1$. Therefore, $\mathbb{Q}[x]/(p(x))^e$ has dimension $\deg p \cdot e = 2e$ for each choice of p . Since 2 divides these dimensions, we conclude that 2 must divide n . Hence, n is even.

Problem 2025-J-II-4 (Topology). Let Σ_2 be a compact oriented surface of genus 2. Is there a submersion $f : \Sigma_2 \rightarrow S^1 \times S^1$, where S^1 denotes the unit circle?

Assume to the contrary that there exists a submersion $f : \Sigma_2 \rightarrow S^1 \times S^1$, where S^1 denotes the unit circle. Since $\dim \Sigma_2 = \dim S^1 \times S^1 = 2$, df_p must have constant rank 2 at every $p \in \Sigma_2$. Hence, f is a local diffeomorphism. Since f is a local diffeomorphism, $f(\Sigma_2)$ is compact in $S^1 \times S^1$; since $S^1 \times S^1$ is Hausdorff, $f(\Sigma_2)$ must be closed in $S^1 \times S^1$. On the other hand, since local diffeomorphisms are open maps, $f(\Sigma_2)$ is open in $S^1 \times S^1$. Therefore, since $S^1 \times S^1$ is connected, $f(\Sigma_2) = S^1 \times S^1$; i.e., f is surjective. Therefore, f is a covering map. This means that the induced homomorphism, $f_* : \pi_1(\Sigma_2) \rightarrow \pi_1(S^1 \times S^1)$ is injective, and so $f_*(\pi_1(\Sigma_2)) \cong \text{img } f_* \leq \pi_1(S^1 \times S^1)$. However, $\pi_1(S^1 \times S^1) \cong \mathbb{Z} \times \mathbb{Z}$ is an abelian group and cannot have any nonabelian subgroups, whereas $\pi_1(\Sigma_2)$ is nonabelian. Hence, by contradiction, f cannot be a submersion.

Problem 2025-J-II-5 (Analysis). Let V be a topological vector space whose topology is Hausdorff. Let X_1 and X_2 be two Banach spaces, and assume there exist continuous linear bijections $F_1 : X_1 \rightarrow V$ and $F_2 : X_2 \rightarrow V$. Show that there is a continuous linear bijection $G : X_1 \rightarrow X_2$.

Assume the given hypotheses. Let $G = F_2^{-1} \circ F_1$. Since F_1, F_2 are bijections, we conclude that G is a bijection. Likewise, since F_1, F_2 are linear, G must also be linear. It suffices to prove that G is continuous. By the Closed Graph Theorem, continuity of G is equivalent to the graph of G being a closed subspace of $X_1 \times X_2$. Let $\{x_n\} \subset X_1$ be a sequence in X_1 such that $x_n \rightarrow x$ and $y_n = Gx_n \rightarrow y$. We need to show that $y = Gx$. By continuity of F_1 , $F_1x_n \rightarrow F_1x$. By continuity of F_2 ,

$$F_2y = \lim F_2y_n = \lim F_2Gx_n = \lim F_1x_n = F_1x. \quad (8)$$

Since F_2 is bijective, $y = F_2^{-1}F_1x = Gx$. Hence, the graph of G is closed, which implies that G is continuous.

August 2025

Problem 2025-A-I-1 (Geometry/Topology). Let S be a closed orientable surface of genus 4 and C be an embedded circle that partitions S into two subsurfaces of genus 2. Does S retract to C ?

We claim that the answer is no; assume to the contrary that there exists a retraction $r : S \rightarrow C$. Let $i : C \hookrightarrow S$ be the inclusion map so that $r \circ i = \text{id}_C$. Now since C is an embedded circle, $H_1(C)$ (i.e., the first homology) is isomorphic to $H_1(S^1) = \mathbb{Z}$. On the other hand, since C is separating in S , its homology class in $H_1(S)$ is the zero element. Hence, the induced map $i_* : H_1(C) \rightarrow H_1(S)$ is the zero map. But this is impossible since if i_* is the zero map,

$$0 = r_* \circ i_* = (r \circ i)_* = \text{id}_{H_1(C)}, \quad (9)$$

which is a contradiction. Hence, no such retraction can exist.

Problem 2025-A-I-6 (Algebra). Let $f(x)$ be an irreducible polynomial of degree n over a field F , and let $g(x)$ be any polynomial in $F[x]$. Prove that every irreducible factor of the composition $f(g(x))$ has degree divisible by n .

Let $h(x)$ be an irreducible factor of $f(g(x))$ in $F[x]$ and let α be the root of $h(x)$ in some algebraic closure of F . Since h is irreducible and α is a root, the minimum polynomial of α over F is h . Therefore,

$$\deg h = [F(\alpha) : F]. \quad (10)$$

Now since α is a root of $h(x) = f(g(x))$, $f(g(\alpha)) = 0$. In particular, $g(\alpha)$ is a root of f . Since f is irreducible of degree n over F , the minimal polynomial of $g(\alpha)$ over F is f . Hence,

$$[F(g(\alpha)) : F] = n. \quad (11)$$

Since $F \subset F(g(\alpha)) \subset F(\alpha)$, by the Tower Law,

$$\deg h = [F(\alpha) : F] = [F(\alpha) : F(g(\alpha))] \cdot [F(g(\alpha)) : F] = n[F(\alpha) : F(g(\alpha))], \quad (12)$$

so that $n \mid \deg h$. Hence, this concludes the proof.

Problem 2025-A-II-2 (Geometry/Topology). Consider the plane distribution in \mathbb{R}^3 spanned by two vector fields

$$V = \partial_x + 2xy\partial_z, \quad W = x\partial_x + \partial_y + (2x^2y + x^2 - 2y)\partial_z. \quad (13)$$

- (i) Show that this distribution is integrable.
- (ii) Does the pair of vector fields V and W generate a coordinate system on integral surfaces? If not, find a pair that can play this role for the local integral surfaces passing through points $(0, 0, z_0)$.

- (i) Let D be the plane distribution in \mathbb{R}^3 spanned by the two vector fields V and W given above. Then by the Frobenius Theorem, D is integrable if and only if D is involutive, which is true if and only if the Lie Bracket of V and W is a smooth section of D at each $p \in \mathbb{R}^3$. We observe that:

$$\begin{aligned} V(W) &= (\partial_x + 2xy\partial_z)(x\partial_x + \partial_y + (2x^2y + x^2 - 2y)\partial_z) \\ &= \partial_x + (4xy + 2x)\partial_z. \\ W(V) &= (x\partial_x + \partial_y + (2x^2y + x^2 - 2y)\partial_z)(\partial_x + 2xy\partial_z) \\ &= 2xy\partial_z + 2x\partial_z. \end{aligned} \tag{14}$$

Therefore, for any $p \in \mathbb{R}^3$,

$$[V, W] = V(W) - W(V) = \partial_x + 2xy\partial_z = V. \tag{15}$$

Since V is a smooth section of D , we conclude that D is involutive, and hence integrable.

- (ii) Let \mathcal{S} be an integral surface, and assume there are coordinates (u, v) on \mathcal{S} such that $V|_{\mathcal{S}} = \partial_u$ and $W|_{\mathcal{S}} = \partial_v$. Then we observe that $[V|_{\mathcal{S}}, W|_{\mathcal{S}}] = \partial_u(\partial_v) - \partial_v(\partial_u) = 0$. On the other hand,

$$[V|_{\mathcal{S}}, W|_{\mathcal{S}}] = ([V, W])|_{\mathcal{S}} = V|_{\mathcal{S}} \neq 0, \tag{16}$$

which is a contradiction. Therefore, V and W cannot generate a coordinate system on integral surfaces. However, consider the fields $\tilde{V} = V$ and $\tilde{W} = W - xV$ on \mathbb{R}^3 . Then since

$$[\tilde{V}, \tilde{W}] = V(W - xV) - (W - xV)(V) = VW - xVV - W(V) + xVV = 0, \tag{17}$$

and so this pair generates a coordinate system on all integral surfaces.

January 2024

Problem 2024-J-I-1 (Algebra). For distinct odd primes p and q , prove that every finite group of order $2pq$ is a semidirect product of a normal subgroup of order pq and a subgroup of order 2.

Let G be a group of order $2pq$, where p, q are distinct odd primes. Without loss of generality, assume $q > p$. By Sylow's Theorem,

$$n_q \in \{1, 2, p, 2p\} \cap \{1, q+1, \dots\} = 1, \tag{18}$$

since $q > 2$ and $q > p$. Therefore, G has a unique, normal, Sylow q -subgroup, which we denote as Q . Let P be a Sylow p -subgroup of G . By the Second Isomorphism Theorem, we conclude that $N = PQ$ is a subgroup of G of order $|P||Q| = pq$. Since $|G : N| = 2pq/(pq) = 2$, where 2 is the smallest prime dividing $|G|$, we conclude that N is a normal subgroup of G . Next, by Cauchy's Theorem, G contains an element of order 2. Let M be the subgroup generated by this element, which also must have order 2. By Lagrange's Theorem, $N \cap M = \{e\}$. Next,

$$|NM| = \frac{|N||M|}{|N \cap M|} = |N||M| = 2pq = |G|, \tag{19}$$

so that $G = NM$. Therefore, we conclude that $G = N \rtimes M$.

Problem 2024-J-I-2 (Geometry/Topology). Let $p : E \rightarrow B$ be a covering space map, with B and E path connected. Choose a point $e_0 \in E$ and $b_0 \in B$ such that $p(e_0) = b_0$. This gives us a subgroup $H = p_*\pi_1(E, e_0)$ of the fundamental group $G = \pi_1(B, b_0)$. Construct a bijection between the fiber $p^{-1}(b_0)$ and the set of right cosets of H and prove that this is indeed a bijection. Prove that the number of sheets of p equals the index $(G : H)$.

Assume all of the given hypotheses. Let $\phi : \pi_1(B, b_0) \rightarrow p^{-1}(b_0)$ be the lifting correspondence induced by p defined by $\phi([f]) = \tilde{f}(1)$, where \tilde{f} is the lift of f , and let $\Phi : \pi_1(B, b_0)/H \rightarrow p^{-1}(b_0)$ be the map induced by ϕ . It suffices to prove that Φ is a bijection.

- (i) Since E is path connected and $p : E \rightarrow B$ is a covering map, the lifting correspondence ϕ must be surjective. Hence, since Φ is induced by ϕ , it follows that Φ is also surjective.
- (ii) Now we will show that Φ is injective. Let f and g be two paths in B , and \tilde{f}, \tilde{g} their liftings to paths in E that begin at e_0 . We must show that $\tilde{f}(1) = \tilde{g}(1)$ iff $[f] \in H * [g]$.
- (\Leftarrow) Suppose $[f] = [h * g]$, where $h = p \circ \tilde{h}$ for some loop \tilde{h} in E based at e_0 . Since \tilde{g} is a path in E that begins at e_0 , the product $\tilde{h} * \tilde{g}$ is well-defined. Since $[f] = [h * g]$, it follows that \tilde{f} and $\tilde{h} * \tilde{g}$ must end at the same point. Hence, \tilde{f} and \tilde{g} end at the same point. Therefore, $\phi([f]) = \phi([g])$.
- (\Rightarrow) Suppose $\phi([f]) = \phi([g])$, which means that $\tilde{f}(1) = \tilde{g}(1)$. Then the product of \tilde{f} with the reverse of \tilde{g} is well-defined and is a loop \tilde{h} in E based at e_0 . By direct computation, $[\tilde{h} * \tilde{g}] = [\tilde{f}]$. If \tilde{F} is a path homotopy between $\tilde{h} * \tilde{g}$ and \tilde{f} , then $p \circ \tilde{F}$ is a path homotopy between $h * g$ and f , which means that $[f] \in H * [g]$. Hence, this concludes the proof that Φ is injective.

Hence, $|p^{-1}(b_0)| = |G/H| = (G : H)$.

Problem 2024-J-I-4 (Algebra). For each field K , prove that the polynomial ring $K[x, y]$ in two variables is not a principal ideal domain.

Let K be a field, and consider the polynomial ring $K[x, y]$. Let (x, y) be the proper ideal of $K[x, y]$ generated by the monomials x and y . Assume to the contrary that $(x, y) = (f(x, y))$ where $f(x, y) \in K[x, y]$ is not a unit of the polynomial ring. Since $x \in (f(x, y))$, $f(x, y) \mid x$. By our assumption that f is not a unit, it follows that $f(x, y)$ is an associate of x . Likewise, $f(x, y)$ must be an associate of y . But this is impossible since x and y are not associates of each other. This forces $f(x, y)$ to be a unit, which means that $(f(x, y)) = K[x, y]$. But this contradicts the fact that $(x, y) = (f(x, y))$ is a proper ideal. Hence, by contradiction, (x, y) is not a principal ideal, and so $K[x, y]$ is not a principal ideal domain.

Problem 2024-J-I-5 (Geometry/Topology). Let α be a closed 1-form on $\mathbb{R}P^n$, $n > 1$. Show that if $f : [0, 1] \rightarrow \mathbb{R}P^n$ is a smooth function such that $f(0) = f(1)$, then

$$\int_{[0,1]} f^* \alpha = 0.$$

Include all calculations that are relevant to your solution.


We recall that $H^k(\mathbb{R}P^n) = 0$ for all $0 < k < n$ so that $H^1(\mathbb{R}P^n) = 0$ if $n > 1$. In particular, this means that α is also an exact 1-form on $\mathbb{R}P^n$. Let g be a smooth function on $\mathbb{R}P^n$ so that $\alpha = dg$. Then

$$\int_0^1 f^* \alpha = \int_0^1 f^* dg = \int_0^1 d(f^* g) = g(f(1)) - g(f(0)) = 0, \quad (20)$$

where the last equality follows from the fact that $f(1) = f(0)$. Hence, the proof concludes.

Problem 2024-J-II-3 (Geometry/Topology). Let Σ be a genus 2 surface embedded in \mathbb{R}^3 as shown in the picture. Let M be the closure of the *unbounded* component of $\mathbb{R}^3 \setminus \Sigma$; in other words, M is the part of \mathbb{R}^3 which is *not* enclosed by Σ .

- (a) Compute $\pi_1(M)$.
 (b) Is Σ a retract of M ?



(a)

Problem 2024-J-II-6 (Geometry/Topology). Let M be a smooth n -manifold, and let φ be a differential k -form on M which is closed, in the sense that $d\varphi = 0$. At each point $p \in M$, define

$$D_p = \{v \in T_p M : v \lrcorner \varphi = 0\}, \quad (21)$$

where \lrcorner denotes the interior product. Assume $\ell := \dim D_p$, so that $D \subset TM$ is a rank- ℓ vector subbundle of the tangent bundle of M . Prove that D is an integrable distribution of ℓ -planes, in the sense of the Frobenius Theorem.

By the Frobenius Theorem, it suffices to prove that D is involutive, which is to say that if X, Y are smooth sections of D , then $[X, Y]$ is also a smooth section of D . Indeed, let X, Y be smooth sections of D , which means that $X \lrcorner \varphi, Y \lrcorner \varphi = 0$. Observe that,

$$[X, Y] \lrcorner \varphi = \mathcal{L}_X(Y \lrcorner \varphi) - Y \lrcorner (\mathcal{L}_X \varphi). \quad (22)$$

By hypothesis, $Y \lrcorner \varphi = 0$ so that $\mathcal{L}_X(Y \lrcorner \varphi) = 0$. On the other hand, by Cartan's Formula,

$$\mathcal{L}_X \varphi = d(X \lrcorner \varphi) + X \lrcorner d\varphi = 0, \quad (23)$$

by the hypotheses. Hence, this shows that $[X, Y] \lrcorner \varphi = 0$, and so $[X, Y]$ is a smooth section of D . Therefore, D is involutive, which means that it is Frobenius integrable.

August 2023

Problem 2023-A-II-1 (Algebra). A field extension K/L is called algebraic, if every element in K satisfies a polynomial equation with coefficients in L . Let F, K, L be fields such that $F \supset K \supset L$, and F/K and K/L are algebraic extensions. Prove that F/L is also an algebraic extension.

Since subfields of subfields is a subfield, L is a subfield of F . Hence, it suffices to show that every element in F satisfies a polynomial equation with coefficients in L . Let $a \in F$, and let

$$k(x) = k_n x^n + k_{n-1} x^{n-1} + \cdots + k_0 \in K[x] \quad (24)$$

such that $k(a) = 0$; this follows since F/K is an algebraic extension. Each $k_j \in K$, and hence is algebraic over L . Therefore, $L' = L(k_0, \dots, k_n)$ is a finite extension of L . Since $k(a) = 0$ and $k(x)$ now has its coefficients in L' , it follows that a is algebraic over L' so that $L'(a)$ is a finite extension of L . Then since

$$[L(a) : L] = [L(a) : L'] [L' : L], \quad (25)$$

it follows that $L(a)$ is a finite extension of L . Therefore, a is algebraic over L . Since a was arbitrary, F/L is an algebraic extension.

January 2023

Problem 2023-J-II-4 (Geometry/Topology). Prove that $S^2 \times S^2$ is not diffeomorphic to $M_1 \times M_2 \times M_3$, where M_1, M_2, M_3 are smooth manifolds of nonzero dimension.

We begin with a technical lemma, that we will use to prove the desired result.

(Comps Lemma) Let M, N be smooth, connected n -manifolds and $f : M \rightarrow N$ a (smooth) immersion. If M is compact and nonempty, then N is compact and f is a (smooth) covering map.

Proof. Let M, N be smooth connected n -manifolds, $f : M \rightarrow N$ an immersion, and M compact and nonempty. Since $\dim N = n$ everywhere and f is an immersion, $df_p : T_p M \rightarrow T_{f(p)} N$ has constant rank n everywhere. Hence, by the Inverse Function Theorem, f is a local diffeomorphism. Since local diffeomorphisms are open maps, $f(M)$ is open in N . Next since the continuous image of compact sets is compact, $f(M)$ is compact in N . Since N is Hausdorff, $f(M)$ must be closed in N . Therefore, since N is connected, we conclude that $f(M) = N$. This means that N is compact and f is surjective. All that remains is to show that f is a covering map.

Let $q \in N$, and consider $f^{-1}(q)$, which is closed in M . For each $x \in f^{-1}(q)$, there exists a neighborhood U_x of x such that $f|_{U_x}$ is a diffeomorphism. Since M is Hausdorff, we may shrink these neighborhoods so that they are pairwise disjoint. This means that each $x \in f^{-1}(q)$ is isolated. Hence, $f^{-1}(q)$ is discrete in M . Since discrete subspaces of compact spaces must be finite, it follows that $f^{-1}(q)$ is finite; let $f^{-1}(q) = \{x_1, \dots, x_s\}$. As stated above, for each $j = 1, \dots, s$, we can find a neighborhood U_j of x_j such that $f|_{U_j} : U_j \rightarrow V_j \subset N$ is a diffeomorphism. Since M is Hausdorff, we may shrink these neighborhoods so that $U_i \cap U_j = \emptyset$ for all $i \neq j$; f restricted to each of these new U_j 's remains a diffeomorphism. Set $V = \bigcap_1^s f(U_j)$, and define $\tilde{U}_j = f^{-1}(V) \cap U_j$. For each j , $f : \tilde{U}_j \rightarrow V$ is a diffeomorphism and $V = \bigsqcup_1^s f(U_j)$. Hence, V is an evenly covered neighborhood of q , so that f is a covering map. \square

Now, assume to the contrary that $f : S^2 \times S^2 \rightarrow M_1 \times M_2 \times M_3$ is a diffeomorphism; since diffeomorphisms preserve dimensions and M_1, M_2, M_3 have nonzero dimensions, it follows, without loss of generality, that M_1, M_2 are 1-dimensional and M_3 is 2-dimensional. Since diffeomorphisms of manifolds are immersions, by the Comps Lemma, $M_1 \times M_2 \times M_3$ must be compact and connected; by projecting onto each manifold, M_1, M_2, M_3 must be compact and connected. Moreover, the induced group homomorphism $f_* : \pi_1(S^2 \times S^2) \rightarrow \pi_1(M_1 \times M_2 \times M_3) = \pi_1(M_1) \times \pi_1(M_2) \times \pi_1(M_3)$ must be an isomorphism. Since S^2 is simply connected,

$$\pi_1(S^2 \times S^2) = \pi_1(S^2) \times \pi_1(S^2) = \{0\}. \quad (26)$$

On the other hand, since the only compact connected 1-manifold, up to diffeomorphism, is the unit circle S^1 , and $\pi_1(S^1) \cong \mathbb{Z}$ is not trivial, $\pi_1(M_1 \times M_2 \times M_3)$ is not trivial. But this contradicts our claim that f_* is an isomorphism. Hence, by contradiction, f cannot be a diffeomorphism.

Problem 2023-J-II-3 (Geometry/Topology). Consider the form $\omega = (x^2 + x + y)dy \wedge dz$ on \mathbb{R}^3 . Let $S^2 \subset \mathbb{R}^3$ be the unit sphere, and $i : S^2 \rightarrow \mathbb{R}^3$ be the inclusion map.

- Calculate $\int_{S^2} i^* \omega$.
- Construct a closed form α on \mathbb{R}^3 such that $i^* \alpha = i^* \omega$, or show that such a form α does not exist.

Problem 2023-A-I-2 (Geometry/Topology). Let $f : T^2 \rightarrow S^2$ be a smooth map from the 2-torus to the 2-sphere. Can f be an immersion? If the answer is yes, given an explicit example. If the answer is no, then give a proof.

There cannot be an immersion $f : T^2 \rightarrow S^2$. To prove our answer, we will state and proof a technical lemma.

(Comps Lemma) Let M, N be smooth, connected, n -manifolds and $f : M \rightarrow N$ a (smooth) immersion. If M is compact and nonempty, then f is a (smooth) covering map.

Proof. Let M, N be smooth connected n -manifolds, M compact, and $f : M \rightarrow N$ an immersion. Since $\dim N = n$ everywhere and f is an immersion, $df_p : T_p M \rightarrow T_{f(p)} N$ has constant rank n everywhere. Hence, by the Inverse Function Theorem, f is a local diffeomorphism. Let $q \in N$ so that $f^{-1}(q) \subset M$ is closed. For each $x \in f^{-1}(q)$, there exists a neighborhood U_x such that $f|_{U_x} : U_x \rightarrow V_x \subset N$ is a diffeomorphism. Since M is Hausdorff, we can shrink these neighborhoods so that they are pairwise disjoint. Since every $x \in f^{-1}(q)$ is now isolated, it follows that $f^{-1}(q)$ is discrete. Since M is compact, we conclude that $f^{-1}(q)$ must be finite; let $f^{-1}(q) = \{x_1, \dots, x_s\}$. As stated above, for each $j = 1, \dots, s$, we can find a neighborhood U_j of x_j so that $f|_{U_j} : U_j \rightarrow V_j \subset N$ is a diffeomorphism. Again, since M is Hausdorff, we can shrink these neighborhoods so that $U_i \cap U_j = \emptyset$ for all $i \neq j$; f restricted to each of these shrunken neighborhoods remains a diffeomorphism. Now set $V = \bigcap_1^s f(U_j)$, and define $\tilde{U}_j \subset M$ by $\tilde{U}_j = f^{-1}(V) \cap U_j$ for each $j = 1, \dots, s$. Hence, V is an evenly covered neighborhood of $q \in N$, which means f is a covering map. That f is surjective comes from recognizing that $f(M) = N$ due to connectedness of N . \square

Now, assume $f : T^2 \rightarrow S^2$ is an immersion. Since T^2, S^2 are smooth, connected 2-manifolds, and T^2 is compact and nonempty, by the Comps Lemma, f is a covering map. Hence, the induced homomorphism $f_* : \pi_1(T^2) \rightarrow \pi_1(S^2)$ is injective. Since S^2 is simply connected, $\pi_1(S^2) \cong \{0\}$. On the other hand, $\pi_1(T^2) \cong \mathbb{Z} \times \mathbb{Z}$. Since the order of $\pi_1(T^2)$ is more than one, f_* cannot be injective. Hence, f cannot be an immersion.

Problem 2023-A-II-5 (Geometry/Topology). Let (t, x, y, z) be the standard coordinate system on \mathbb{R}^4 , and let ϕ be the non-zero smooth 1-form on \mathbb{R}^4 defined by

$$\phi = dt + ydx + zdy.$$

Let D be the 3-plane field on \mathbb{R}^4 that consists of tangent vectors V such that $\phi(V) = 0$. Is D Frobenius integrable? Support your answer with a proof.

Let D be the 3-plane field on \mathbb{R}^4 defined as follows: for each $p \in \mathbb{R}^4$,

$$D_p = \{v \in T_p \mathbb{R}^4 : \phi(v) = 0\} = \ker \phi_p. \quad (27)$$

Hence, by the Frobenius Theorem, D is Frobenius integrable if and only if $\phi \wedge d\phi = 0$. We compute:

$$d\phi = d(dt + ydx + zdy) = d^2t + dy \wedge dx + dz \wedge dy = dy \wedge dx + dz \wedge dy. \quad (28)$$

Therefore,

$$\phi \wedge d\phi = dt \wedge dy \wedge dx + dt \wedge dz \wedge dy + ydx \wedge dz \wedge dy. \quad (29)$$

Since $\phi \wedge d\phi$ is nowhere vanishing on \mathbb{R}^4 , D is not Frobenius integrable.

Problem 2023-A-I-1 (Algebra). Let V be a n -dimensional vector space over a field F . An element $A \in \text{End } V$ is called *nilpotent*, if $A^k = 0$ for some $k > 1$. Prove that A is nilpotent if and only if

$$\text{Tr}(\Lambda^i A) = 0, \quad i = 1, \dots, n, \quad (30)$$

where $\Lambda^i A$ denotes the induced action of A on the wedge product $\Lambda^i V$ for each i .

Let V be a n -dimensional vector space over a field F , and let $A \in \text{End } V$. Recall that $\Lambda^i A$, the induced action of A on the wedge product $\Lambda^i V$, is defined to be

$$(\Lambda^i A)(v_1 \wedge \cdots \wedge v_i) = Av_1 \wedge \cdots \wedge Av_i, \quad v_j \in V \text{ for all } j = 1, \dots, i. \quad (31)$$

Over an algebraic closure of F , A has eigenvalues $\lambda_1, \dots, \lambda_n$. Suppose A is diagonalizable, with the set of eigenvectors given by $\{v_1, \dots, v_n\}$. Then for each $i = 1, \dots, n$, since the collection

$$\{v_{j_1} \wedge \cdots \wedge v_{j_i} : 1 \leq j_1 < \cdots < j_i \leq n\}$$

is a basis of $\Lambda^i V$, and for each i -tuple, $\Lambda^i A(v_{j_1} \wedge \cdots \wedge v_{j_i}) = Av_{j_1} \wedge \cdots \wedge Av_{j_i} = (\lambda_{j_1} \cdots \lambda_{j_i})(v_{j_1} \wedge \cdots \wedge v_{j_i})$, it follows that the eigenvalues of $\Lambda^i A$ are the set of all products of the form $\lambda_{j_1} \cdots \lambda_{j_i}$ for $1 \leq j_1 < \cdots < j_i \leq n$, counting for multiplicity. Hence,

$$\text{Tr}(\Lambda^i A) = \sum_{1 \leq j_1 < \cdots < j_i \leq n} \lambda_{j_1} \cdots \lambda_{j_i}. \quad (32)$$

If A is not diagonalizable, since the eigenvalues of $\Lambda^i A$ depend only on the eigenvalues of A , we may assume A is in Jordan normal form. Indeed, if $A = PJP^{-1}$, then

$$\Lambda^i(A) = \Lambda^i(PJP^{-1}) = \Lambda^i(P)\Lambda^i(J)\Lambda^i(P^{-1}), \quad (33)$$

so $\Lambda^i A$ and $\Lambda^i J$ are similar and therefore have the same eigenvalues. Thus it suffices to compute the eigenvalues of $\Lambda^i J$, which are exactly the products $\lambda_{j_1} \cdots \lambda_{j_i}$ of the eigenvalues of A .

If A is nilpotent so that $A^k = 0$ for some $k > 1$, then since $0 = A^k v = \lambda^k v$ for all eigenvectors v of A , it follows that every eigenvalue of A is zero. Therefore, the above expression implies that $\text{Tr}(\Lambda^i A) = 0$ for all $i = 1, \dots, n$. On the other hand, expanding the characteristic polynomial for A is given by:

$$p_A(t) = \det(tI - A) = t^n - \text{Tr}(\Lambda^1 A)t^{n-1} + \cdots + (-1)^n \text{Tr}(\Lambda^n A). \quad (34)$$

If $\text{Tr}(\Lambda^i A) = 0$ for all $i = 1, \dots, n$, then we conclude that the characteristic polynomial of A is precisely t^n . Therefore, A 's eigenvalues are all zero. Hence, the minimal polynomial of A is of the form t^k for some $k \leq n$. This implies that $A^k = 0$, and so A is nilpotent.

Problem 2023-A-II-6 (Complex Analysis). Find the number of solutions (counting multiplicity) to $z^8 - 5z^6 + 2z^3 - z - 1 = 0$ that lie inside the unit disk.

Recall Rouché's Formula, which states that

For any two complex-valued functions f and g holomorphic inside some region K with closed and simple contour ∂K , if $|g(z)| < |f(z)|$ on ∂K , then f and $f+g$ have the same number of zeros inside K , where each zero is counted as many times as its multiplicity.

Pick $f(z) = 5z^6$ and set $h(z) = z^8 + 2z^3 - z - 1$ so that $p(z) = z^8 - 5z^6 + 2z^3 - z - 1 = h(z) - f(z)$. On the unit disk $\partial \mathbb{D}^1$, we observe that

$$\begin{aligned} |f(z)| &= |5z^6| = 5 \\ &= 1 + 2 + 1 + 1 \\ &= |z^8| + 2|z^3| + |z| + |1| \\ &\geq |h(z)|. \end{aligned} \quad (35)$$

Hence, $p(z) = h(z) - f(z)$ has the same number of zeros, counting multiplicity, as $f(z)$. Since $f(z)$ has six zeros in the unit disk, we conclude that $p(z)$ must also have six zeros inside the unit disk.