

# Om Algebra Crash Course

Aniruddha Madhava

December 31, 2025

---

## Contents

<b>1</b>	<b>Quotient Groups and Homomorphisms</b>	<b>2</b>
1.1	The Isomorphism Theorems . . . . .	2
1.2	Exercises . . . . .	3
1.3	Composition Series and the Hölder Program . . . . .	4
1.4	Exercises . . . . .	5
<b>2</b>	<b>Group Actions</b>	<b>6</b>
<b>3</b>	<b>Direct and Semidirect Products and Abelian Groups</b>	<b>7</b>

---

# 1 Quotient Groups and Homomorphisms

## 1.1 The Isomorphism Theorems

- **Thm. 16. (First Isomorphism Theorem)** If  $\varphi : G \rightarrow H$  is a group homomorphism, then  $\ker \varphi \trianglelefteq G$  and  $G/\ker \varphi \cong \varphi(G)$ .

Let  $\varphi : G \rightarrow H$  be a group homomorphism with kernel  $K$ . First, we will prove that the kernel is a normal subgroup. Let  $g \in G$  and  $k \in \ker \varphi$ . Then

$$\varphi(gkg^{-1}) = \varphi(g)\varphi(k)\varphi(g)^{-1} = \varphi(g)\varphi(g)^{-1} = 1 \implies gkg^{-1} \in K. \quad (1)$$

This implies that  $gKg^{-1} \subseteq K$  for all  $g \in G$ , and so  $K$  is a normal subgroup of  $G$ . Now let  $\tilde{\varphi} : G/\ker \varphi \rightarrow \varphi(G)$  as follows:  $\tilde{\varphi}(gK) = \varphi(g)$ . First, we start by showing that  $\tilde{\varphi}$  is well-defined. Suppose  $g_1K = g_2K$ , which means that  $g_1g_2^{-1} \in K$ . Therefore,

$$\varphi(g_1g_2^{-1}) = 1 \implies \varphi(g_1) = \varphi(g_2) \implies \tilde{\varphi}(g_1K) = \tilde{\varphi}(g_2K). \quad (2)$$

We need to show that  $\tilde{\varphi}$  is an isomorphism. Suppose  $\tilde{\varphi}(g) = \tilde{\varphi}(h)$ . Then  $\varphi(g) = \varphi(h) \iff gh^{-1} \in K \iff gK = hK$ . This proves injectivity. Now let  $\varphi(g) \in \varphi(G)$ . Hence clearly  $gK \mapsto g$  so that  $\tilde{\varphi}$  is surjective. Finally, if  $g_1K, g_2K \in G/K$ , then

$$\tilde{\varphi}(g_1K \cdot g_2K) = \tilde{\varphi}(g_1g_2K) = \varphi(g_1g_2) = \varphi(g_1)\varphi(g_2) = \tilde{\varphi}(g_1K)\tilde{\varphi}(g_2K). \quad (3)$$

Hence,  $\tilde{\varphi}$  is indeed an isomorphism of groups.

- **Thm. 18. (Diamond Isomorphism Theorem)** Let  $G$  be a group,  $A$  and  $B$  be subgroups of  $G$ , and assume that  $A \leq N_G(B)$ . Then (1)  $AB$  is a subgroup of  $G$ , (2)  $B \trianglelefteq AB$ , (3)  $A \cap B \trianglelefteq A$ , and (4)  $AB/B \cong A/A \cap B$ .

(1) Since  $A \leq N_G(B)$ , it automatically follows that  $AB$  is a subgroup of  $G$ . (2) Since  $A \leq N_G(B)$  and  $B \leq N_G(B)$ , then  $AB \leq N_G(B)$ , which is to say that  $B$  is a normal subgroup of  $AB$ . (3 - 4) Consider the map  $\varphi : A \rightarrow AB/B$  defined by  $\varphi(a) = aB$ . It is straightforward to see that  $\varphi$  is a surjective group homomorphism, which means that  $\varphi(A) = AB/B$ . Now we will determine the kernel:

$$A \ni a \in \ker \varphi \iff aB = 1B \iff a \in B. \quad (4)$$

Hence,  $\ker \varphi = A \cap B$ . By the first Isomorphism Theorem,  $A \cap B \trianglelefteq A$  and  $A/A \cap B \cong AB/B$ .

- **Thm. 19. (Third Isomorphism Theorem)** Let  $G$  be a group and let  $H$  and  $K$  be normal subgroups of  $G$  with  $H \leq K$ . Then  $K/H \trianglelefteq G/H$  and

$$(G/H)/(K/H) \cong G/K. \quad (5)$$

First, we will show that  $K/H \trianglelefteq G/H$ . Define the map  $\varphi : G/H \rightarrow G/K$  by  $\varphi(gH) = gK$ . First, we need to show that this map is well-defined. Suppose  $g_1H = g_2H$ . Then  $g_1 = g_2h$  for some  $h \in H$ . Since  $H \leq K$ ,  $h \in K$ , which shows that  $g_1K = g_2K$ . Hence,  $\varphi$  is well-defined. It is straightforward to see that  $\varphi$  is a surjective homomorphism. Finally, we can easily show that  $\ker \varphi = K/H$ . This means that (1)  $K/H$  is a normal subgroup of  $G/H$ , and (2) by the First Isomorphism Theorem,  $(G/H)/(K/H) \cong G/K$ .

- **Thm. 20. (Lattice Isomorphism Theorem)** Let  $G$  be a group and  $N$  a normal subgroup of  $G$ . Every subgroup of  $G/N$  is of the form  $A/N$ , where  $A$  is a subgroup of  $G$  containing  $N$ . Moreover, for all  $A, B \leq G$ , with  $N \leq A$  and  $N \leq B$ ,

- (1)  $A \leq B$  if and only if  $A/N \leq B/N$ .
- (2) If  $A \leq B$ , then  $|B : A| = |B/N : A/N|$ .
- (3)  $\langle A, B \rangle / N = \langle A/N, B/N \rangle$ .
- (4)  $(A \cap B)/N = (A/N) \cap (B/N)$ .

(5)  $A \trianglelefteq G$  if and only if  $A/N \trianglelefteq G/N$ .

- **Def. (Factoring Through)** In some of the above proofs of the isomorphism theorems, we have had to define a map  $\varphi$  on quotient groups  $G/N$  defined by giving the value of  $\varphi$  on the coset  $gN$  in terms of the representative  $g$  alone. In essence, this defines a homomorphism  $\Phi$  on  $G$ , itself, by specifying the value of  $\varphi$  at  $g$ . Hence, a map on a quotient group  $G/N$  is well-defined if and only if  $N \leq \ker \Phi$ . In this case, we say that the homomorphism  $\Phi$  *factors through*  $N$  and  $\varphi$  is the induced homomorphism on  $G/N$ . Pictorially,

$$\begin{array}{ccc} G & \xrightarrow{\pi} & G/N \\ & \searrow \Phi & \downarrow \varphi \\ & & H \end{array}$$

## 1.2 Exercises

**Exercise 3.** Prove that if  $H$  is a normal subgroup of  $G$  of prime index  $p$  then for all  $K \leq G$  either

- (i)  $K \leq H$  or
- (ii)  $G = HK$  and  $|K : K \cap H| = p$ .

Let  $H$  be a normal subgroup of  $G$  of prime index  $p$ , and let  $K$  be an arbitrary fixed *nontrivial* subgroup of  $G$ . If  $K \leq H$ , we are done; so assume that  $K$  is not a subgroup of  $H$ . Since  $K \leq G = N_G(H)$ , we conclude by the Second Isomorphism Theorem that  $HK$  is a subgroup of  $G$  and that  $K \cap H$  is a normal subgroup of  $K$ . Hence, we have the chain  $H \leq HK \leq G$ . Therefore,

$$p = |G : H| = |G : HK| \cdot |HK : H|. \quad (6)$$

Since  $p$  is a prime, either  $|G : HK| = 1$  (in which case  $G = HK$ ), or  $|HK : H| = 1 \implies HK = H \implies K \leq H$ . By our hypothesis, the latter is not possible. Therefore,  $G = HK$ . From this, we observe that

$$1 = \frac{|HK|}{|G|} = \frac{|H||K|}{|G||K \cap H|} = \frac{p^{-1}|K|}{|K \cap H|} \implies p = \frac{|K|}{|K \cap H|} = |K : K \cap H|. \quad (7)$$

**Exercise 4.** Let  $C$  be a normal subgroup of the group  $A$  and let  $D$  be a normal subgroup of the group  $B$ . Prove that  $(C \times D) \trianglelefteq (A \times B)$  and  $(A \times B)/(C \times D) \cong (A/C) \times (B/D)$ .

Let  $C$  be a normal subgroup of the group  $A$  and  $D$  be a normal subgroup of the group  $B$ . Define the map,

$$\begin{aligned} \varphi : A \times B &\longrightarrow C \times D \\ (a, b) &\longmapsto (aC, bD). \end{aligned}$$

First, we will show that  $\varphi$  is a group homomorphism. Let  $a_1, a_2 \in A$  and  $b_1, b_2 \in B$ . Then

$$\begin{aligned} \varphi(a_1 a_2, b_1 b_2) &= (a_1 a_2 C, b_1 b_2 D) \\ &= (a_1 C, b_1 D) \cdot (a_2 C, b_2 D) \\ &= \varphi(a_1, b_1) \cdot \varphi(a_2, b_2). \end{aligned} \quad (8)$$

This confirms that  $\varphi$  is a group homomorphism;  $\varphi$  is clearly surjective since for any  $(aC, bD) \in (A/C) \times (B/D)$ ,  $\varphi : (a, b) \mapsto (aC, bD)$ . Now we identify the kernel of this map:

$$\begin{aligned} \ker \varphi &= \{(a, b) \in A \times B : aC = 1C \text{ and } bD = 1D\} \\ &= \{(a, b) \in A \times B : a \in C \text{ and } b \in D\} \\ &= C \times D. \end{aligned} \quad (9)$$

Hence, the conclusion proceeds from the First Isomorphism Theorem.

**Exercise 9.** Let  $p$  be a prime and let  $G$  be a group of order  $p^a m$ , where  $p$  does not divide  $m$ . Assume  $P$  is a subgroup of  $G$  of order  $p^a$  and  $N$  is a normal subgroup of  $G$  of order  $p^b n$ , where  $p$  does not divide  $n$ . Prove that  $|P \cap N| = p^b$  and  $|PN/N| = p^{a-b}$ . (The subgroup  $P$  of  $G$  is called a Sylow  $p$ -subgroup of  $G$ . This exercise shows that the intersection of any Sylow  $p$ -subgroup of  $G$  with a normal subgroup  $N$  is a Sylow  $p$ -subgroup of  $N$ .)

Assume all of the given hypotheses. We have the following results:

- (i) since  $P \leq G = N_G(N)$ ,  $PN \leq G$  by the Diamond Isomorphism Theorem;
- (ii)  $P \cap N \leq P$ , which means that  $|P \cap N| = p^j$  for some nonnegative integer  $j \leq a$ ;
- (iii)  $PN \leq G$  implies, by Lagrange's Theorem, that there exists a positive integer  $k$  such that

$$|G| = p^a m = k \cdot |PN| = k \cdot \frac{|P||N|}{|P \cap N|} = k \cdot \frac{p^a p^b n}{p^j} \implies m = k \cdot p^{b-j} n. \quad (10)$$

This shows that  $p^{b-j} \mid m$ . Since  $p \nmid m$ , we must necessarily have  $p^{b-j} = 1 \implies b-j = 0 \implies j = b$ . Therefore,  $|P \cap N| = p^b$ . Next, by the Diamond Isomorphism Theorem, since  $P/(P \cap N) \cong PN/N$ ,  $|PN/N| = |P|/|P \cap N| = p^{a-b}$ .

### 1.3 Composition Series and the Hölder Program

- **Prop. 21. (Element of Prime Order)** If  $G$  is a finite abelian group and  $p$  is a prime dividing  $|G|$ , then  $G$  contains an element of order  $p$ .
- **Def. (Simple Group)** A (finite or infinite) group  $G$  is called *simple* if  $|G| > 1$  and the only normal subgroups of  $G$  are 1 and  $G$ .
- **Def. (Composition Series)** In a group  $G$  a sequence of subgroups

$$1 = N_0 \leq N_1 \leq N_2 \leq \cdots \leq N_{k-1} \leq N_k = G \quad (11)$$

is called a *composition series* if  $N_i \triangleleft N_{i+1}$  and  $N_{i+1}/N_i$  is a simple group for all  $0 \leq i \leq k-1$ . The quotient groups  $N_{i+1}/N_i$  are called *composition factors* of  $G$ .

- **Thm. 22. (Jordan-Hölder)** Let  $G$  be a finite group with  $G \neq 1$ . Then
  - (1)  $G$  has a composition series and
  - (2) the composition factors in a composition series are unique, namely, if  $1 = N_0 \leq N_1 \leq \cdots \leq N_r = G$  and  $1 = M_0 \leq M_1 \leq \cdots \leq M_s = G$  are two composition series for  $G$ , then  $r = s$  and there is some permutation,  $\pi$ , of  $\{1, 2, \dots, r\}$  such that

$$M_{\pi(i)}/M_{\pi(i)-1} \cong N_i/N_{i-1}, \quad 1 \leq i \leq r. \quad (12)$$

- **Thm. (Feit-Thompson)** If  $G$  is a simple group of odd order, then  $G \cong Z_p$  for some prime  $p$ .
- **Def. (Solvable Group)** A group  $G$  is *solvable* if there is a chain of subgroups

$$1 = G_0 \trianglelefteq G_1 \trianglelefteq G_2 \trianglelefteq \cdots \trianglelefteq G_s = G \quad (13)$$

such that  $G_{i+1}/G_i$  is abelian for  $i = 0, 1, \dots, s-1$ .

- **Obs. (Solvability of Groups in Terms of Subgroups)** Let  $G$  be a group and  $N$  a normal subgroup of  $G$ . If  $N$  and  $G/N$  are solvable, then  $G$  is solvable.

Let  $\overline{G} = G/N$ ,  $1 = N_0 \trianglelefteq N_1 \trianglelefteq \cdots \trianglelefteq N_n = N$  be a chain of subgroups of  $N$  such that  $N_{i+1}/N_i$  is abelian for all  $0 \leq i \leq n-1$  and let  $\overline{1} = \overline{G}_0 \trianglelefteq \overline{G}_1 \trianglelefteq \cdots \trianglelefteq \overline{G}_m = \overline{G}$  be a chain of subgroups such that  $\overline{G}_{i+1}/\overline{G}_i$  is abelian for  $0 \leq i \leq m-1$ . By the Lattice Isomorphism Theorem, there are

subgroups  $G_i$  of  $G$  with  $N \leq G_i$  such that  $G_i/N = \overline{G}_i$  and  $G_i \trianglelefteq G_{i+1}$ ,  $0 \leq i \leq m-1$ . By the Third Isomorphism Theorem,

$$\overline{G}_{i+1}/\overline{G}_i = (G_{i+1}/N)/(G_i/N) \cong G_{i+1}/G_i. \quad (14)$$

Hence, the chain

$$1 = N_0 \trianglelefteq N_1 \trianglelefteq \cdots \trianglelefteq N_n = N = G_0 \trianglelefteq \cdots \trianglelefteq G_m = G \quad (15)$$

is a composition series for  $G$ , which proves that  $G$  is solvable.

## 1.4 Exercises

**Exercise 1.** Prove that if  $G$  is an abelian simple group then  $G \cong \mathbb{Z}_p$  for some prime  $p$  (do not assume  $G$  is a finite group).

Let  $G$  be a nontrivial, abelian, simple group. Since  $G$  is nontrivial, it must contain some nonidentity element  $x \in G$ . Consider the subgroup  $\langle x \rangle$  generated by this element. Since  $G$  is abelian,  $\langle x \rangle$  is a normal subgroup of  $G$ . And since  $G$  is simple,  $\langle x \rangle = G$ . Therefore,  $G$  is a cyclic group.

Suppose  $G$  is an infinite group. Then  $G \cong \mathbb{Z}$  via the isomorphism  $\varphi : \mathbb{Z} \rightarrow G$  that maps  $n \mapsto x^n$ . However,  $\mathbb{Z}$  is not a simple group, since for example, the subgroup  $4\mathbb{Z}$  is a proper normal subgroup of  $\mathbb{Z}$ . Hence, by contradiction,  $G$  must be a finite group.

Assume  $|G| = pm$  for some prime  $p$ . By Cauchy's Theorem,  $G$  contains an element  $y$  of order  $p$ ; since  $G$  is abelian, the subgroup  $\langle y \rangle$  of index  $m$  generated by this element is proper unless  $m = 1$ . But if  $m = 1$ ,  $G$  is a cyclic group of prime order  $p$ . Then it is easily shown that the map  $\varphi : \mathbb{Z}/p\mathbb{Z} \rightarrow G$  defined by  $\varphi(n) = x^n$  is an isomorphism. Therefore,  $G$  is isomorphic to  $\mathbb{Z}_p$  for some prime  $p$ .

**Exercise 4.** Use Cauchy's Theorem and induction to show that a finite abelian group has a subgroup of order  $n$  for each positive divisor  $n$  of its order.

Let  $G$  be a finite abelian group. We will proceed by complete induction; assume that the result holds true for all groups of order less than  $|G|$ .

## 2 Group Actions

### **3 Direct and Semidirect Products and Abelian Groups**