

Comps Mega Review

Aniruddha Madhava

June 4, 2026

Contents

1	About	6
2	Differential Geometry and Topology	7
2.1	Frobenius Theorem	7
	Problem 2025-A-II-2	7
	Problem 2024-A-II-2	7
	Problem 2024-J-II-6	7
	Problem 2023-A-II-5	8
	Problem 2019-J-II-6	8
	Problem 2015-J-II-3	9
	Problem 2013-A-II-4	9
	Problem 2012-J-I-4	10
	Problem 2011-A-II-5	10
	Problem 2010-J-I-2	10
	Problem 1997-J-I-4	10
2.2	Stokes' Theorem	12
	Problem 2025-J-I-3	12
	Problem 2023-J-II-3	12
	Problem 2016-J-I-5	12
	Problem 2009-J-II-3	13
	Problem 2004-J-II-1	13
	Problem 2002-J-I-6	13
	Problem 1999-A-I-4	13
	Problem 1998-J-I-5	14
2.3	Submersions and Immersions	15
	Problem Comps Lemma (Submersion)	15
	Problem Comps Lemma (Local Homeomorphism)	15
	Problem 2025-J-II-4	15
	Problem 2024-A-I-5	15
	Problem 2024-A-II-1	16
	Problem 2023-A-I-2	16
	Problem 2023-A-I-5	16
	Problem 2019-A-I-4	16
	Problem 2017-A-I-1	17
	Problem 2017-J-I-1	17
	Problem 2015-A-II-3	17
	Problem 2011-J-I-3	17
	Problem 2010-J-I-3	17
	Problem 2002-A-I-1	18
	Problem 2002-J-II-1	18
2.4	de Rham Cohomology	19
	Problem 2026-J-II-6	19
	Problem 2026-J-II-6 (Modified)	19
	Problem 2024-A-I-1	20
	Problem 2024-J-I-5	20
	Problem 2021-A-II-1	20

Problem 2020-J-II-2	20
Problem 2019-A-II-4	21
Problem 2019-J-I-3	21
Problem 2015-J-I-5	22
Problem 2013-A-II-6	22
Problem 2013-J-II-6	23
Problem 2002-J-II-5	23
Problem 1997-A-I-5	23
Problem 2001-A-II-3	24
Problem 2007-A-II-3	24
2.5 Fundamental Groups	25
Problem 2023-J-II-4	25
Problem 2003-J-II-5	25
2.6 Vector Fields	26
Problem 2020-J-I-4	26
Problem 2017-J-II-1	26
2.7 Covering Spaces	26
Problem 2021-A-I-6	26
Problem 2016-J-I-2	27
Problem 2013-J-II-5	27
Problem 2013-J-II-5 (Modified)	28
Problem 2010-A-II-4	28
Problem 2004-A-I-5	28
3 Algebra	30
3.1 Linear Algebra	30
Problem 2022-A-I-6	30
Problem 2019-J-I-1	30
3.2 Abelian Groups	31
Problem 2020-A-I-4	31
Problem 2020-J-I-1	31
Problem 2012-J-I-6	31
3.3 Ring Theory	31
Problem 2025-J-I-1	31
3.4 Galois Theory	32
Problem Galois Lemma I	32
Problem 2025-J-II-6	32
Problem 2024-A-II-5	32
Problem 2024-J-II-4	32
Problem 2023-A-II-3	33
Problem 2023-J-I-5	33
Problem 2022-A-I-2	34
Problem 2021-A-I-4	34
Problem 2021-J-I-5	35
Problem 2019-A-I-6	35
Problem 2017-A-II-3	35
Problem 2017-J-I-3	36
Problem 2016-J-II-2	36
Problem 2015-A-II-5	36
Problem 2014-A-II-1	37
Problem 2014-J-I-5	38
Problem 2013-A-I-2	38
Problem 2011-A-II-1	38
Problem 2010-A-II-2	39

Problem 2010-J-I-5	39
Problem 2005-J-I-4	39
3.5 Sylow's Theorems	40
Problem 2022-J-I-3	40
Problem 2022-A-II-4	40
Problem 2018-J-II-5	40
Problem 2017-J-II-6	41
Problem 2016-J-I-1	41
Problem 2013-J-I-2	41
Problem 2009-J-I-5	42
Problem 2006-A-II-5	42
Problem 2001-J-I-3	42
Problem 1999-A-I-1	42
Problem D&F-4.5-13	42
Problem D&F-4.5-14	42
Problem D&F-4.5-15	43
Problem D&F-4.5-17	43
Problem D&F-4.5-32	44
Problem D&F-4.5-33	44
Problem D&F-4.5-34	44
Problem D&F-4.5-36	44
3.6 Classification of Groups	46
Problem 2024-J-I	46
Problem 2023-J-II-6	46
Problem 2010-J-II-5	47
Problem 2008-J-I-3	47
Problem 2004-A-I-6	47
Problem 2003-A-II-6	48
Problem 2003-J-I-6	48
Problem D&F-5.2-1	48
3.7 Module Theory	50
Problem 2025-J-I-5	50
Problem 2025-J-II-3 (Modified)	50
Problem 2025-J-II-3	50
Problem 2021-A-I-5	50
Problem 2005-J-II-4	51
Problem D&F-10.5-1	51
Problem D&F-10.5-2	52
Problem D&F-10.5-3	52
Problem D&F-10.5-4	53
Problem D&F-10.5-5	53
Problem D&F-10.5-6	53
Problem D&F-10.5-7	54
Problem D&F-10.5-9	54
Problem D&F-10.5-10	54
Problem D&F-10.5-12	55
Problem D&F-10.5-20	56
Problem FR-1-1	56
Problem D&F-17.1-6	57
Problem D&F-17.1-7	58
Problem D&F-17.1-10	58
Problem D&F-17.1-12	60
Problem D&F-17.1-13	60
Problem D&F-17.1-14	61

Problem D&F-17.1-16	61
Problem D&F-17.1-17	62
Problem D&F-17.1-19	62
Problem D&F-17.1-21	63
Problem D&F-17.1-22	64
Problem D&F-17.1-23	65
4 Real Analysis	66
4.1 Convergence of Functions	66
Problem 2026-J-II-2	66
Problem 2025-J-I-2	66
Problem 2004-A-II-1	66
Problem 2021-J-II-2	67
Problem 2019-A-I-3	68
Problem 2015-J-II-5	68
4.2 L^p Spaces	69
Problem 2025-A-II-4	69
Problem 2017-J-II-2	69
Problem F-6-1-2	69
Problem F-6-1-3	71
Problem F-6-1-7	71
Problem F-6-1-10	72
Problem F-6-1-14	72
Problem F-6-1-4	73
4.3 Distribution Theory	75
Problem F-9-1-1	75
Problem F-9-1-2	75
Problem F-9-1-3	76
Problem F-9-1-5	76
Problem F-9-1-6	77
5 Complex Analysis	78
5.1 Holomorphic Functions	78
Problem 2026-J-II-4	78
5.2 Rouché's Theorem	78
Problem 2023-A-II-6	78
Problem 2020-A-II-1	78
Problem 2018-A-I-3	79
Problem 2015-A-I-3	79
Problem 2012-A-I-4	79
6 Partial Differential Equations	80
6.1 Introduction	80
Problem E-1-1	80
Problem E-1-2	80
Problem E-1-4	81
7 Notes	82
7.1 Differential Geometry and Topology	82
7.1.1 De Rham Cohomology	82
Problem LSM-17-5	86
Problem LSM-17-6	87
Problem LSM-17-7	88
7.2 Partial Differential Equations	90
7.2.1 Introduction	90

7.2.2	Four Important Linear Partial Differential Equations	90
7.2.3	Sobolev Spaces	105
7.3	Semidirect Products	107

1 About

The following document contains questions that I have solved as preparation for the August 2026 comprehensive exams given by the Department of Mathematics at Stony Brook University. The problems are divided by subjects. The problem labeling scheme is as follows:

- (1) Any problem that has the format “{20--}{ }-{ }-{ }” or “{19--}{ }-{ }-{ }” are questions from past comprehensive exams. The labels indicate (1) in what year, (2) in what month, (3) in which part, and (4) in what the order the problem was given, respectively.
- (2) Any problem that has the format “LSM-{ }-{ }” was taken from Lee’s *Introduction to Smooth Manifolds* (2nd Ed.)
- (3) Any problem that the format “D&F-{ }-{ }” was taken from Dummit& Foote’s *Abstract Algebra* (3rd ed.)
- (4) Any problem that has the format “F-{ }-{ }-{ }” was taken from Folland’s *Real Analysis*.

Some resources I found indispensable while studying for this exam are:

- (1) Math StackExchange
- (2) Wikipedia
- (3) The SBU Past Comps website

2 Differential Geometry and Topology

2.1 Frobenius Theorem

Problem 2025-A-II-2. Consider the plane distribution in $\mathbb{R}^3$ spanned by the two vector fields

$$V = \partial_x + 2xy\partial_z, \quad W = x\partial_x + \partial_y + (2x^2y + x^2 - 2y)\partial_z.$$

- (i) Show that this distribution is integrable.
- (ii) Does the pair of vector fields V and W generate a coordinate system on integral surfaces? If not, find a pair that can play this role for the local integral surfaces passing through the point $(0, 0, z_0)$.

- (i) Let D denote the distribution spanned by the two vector fields V and W . By the Frobenius Theorem, D is integrable if and only if $[V, W]$ is a smooth section of D . Given the vector fields V and W , we first compute

$$V(W) = \partial_x + (4xy + 2x)\partial_z.$$

$$W(V) = (2xy + 2x)\partial_z.$$

Thus $[V, W] = V(W) - W(V) = \partial_x + 2xy\partial_z = V$. Since V is a smooth section of D , we conclude that D is integrable.

- (ii) We recall that a pair of vector fields generates a coordinate system on integral surfaces if and only if the Lie bracket of the vector fields is identically zero. However, in (i), we showed that the Lie bracket of V and W is not identically zero. Thus, V and W cannot generate a coordinate system on integral surfaces. However, we can easily show that the pair of vector fields $(\tilde{V}, \tilde{W})$ given by $\tilde{V} = V$ and $\tilde{W} = W - xV$ has identically zero Lie bracket. Moreover, at the point $(0, 0, z_0)$, $\tilde{V}$ and $\tilde{W}$ are linearly independent. Thus, this pair of vector fields generates a coordinate system on local integral surfaces passing through the point $(0, 0, z_0)$.

Problem 2024-A-II-2. On $\mathbb{R}^5$, equipped with standard coordinates (v, w, x, y, z) , consider the 1-form

$$\theta = dz + v dx + w dy.$$

Are there two smooth functions $f, g : \mathbb{R}^5 \rightarrow \mathbb{R}$ such that $\theta = f dg$? Justify your answer by means of concrete calculations.

No. Assume to the contrary that there exist two smooth functions $f, g : \mathbb{R}^5 \rightarrow \mathbb{R}$ so that $\theta = f dg$. Then we compute,

$$\theta \wedge d\theta = (f dg) \wedge (df \wedge dg) = 0.$$

On the other hand,

$$\begin{aligned} (dz + v dx + w dy) \wedge d(dz + v dx + w dy) &= (dz + v dx + w dy) \wedge (dv \wedge dx + dw \wedge dy) \\ &= dz \wedge dv \wedge dx + dz \wedge dw \wedge dy + v dx \wedge dw \wedge dy \\ &\quad + w dy \wedge dv \wedge dx, \end{aligned}$$

which is not identically zero because of the first two terms. Thus, the proof concludes by contradiction.

Problem 2024-J-II-6. Let M be a smooth n -manifold, and let φ be a differential k -form on M which is closed, in the sense that $d\varphi = 0$. At each point $p \in M$, define

$$D_p = \{v \in T_p M : v \lrcorner \varphi = 0\},$$

where $\lrcorner$ denotes the interior product. Assume $\ell := \dim D_p$ is independent of p , so that $D \subset TM$ is a rank- ℓ vector subbundle of the tangent bundle of M . Prove that D is an integrable distribution

$T_x M = D_x$ for all $x \in M$, it suffices to show that D is an integrable distribution. To do so, we will show that if V, W are smooth sections of D , then their Lie bracket $[V, W]$ is also a smooth section of D . Then

$$\mathcal{L}_{[V, W]}\omega = \mathcal{L}_V(\mathcal{L}_W\omega) - \mathcal{L}_W(\mathcal{L}_V\omega).$$

Cartan's formula tells us that

$$\begin{aligned}\mathcal{L}_V\omega &= V \lrcorner d\omega + d(V \lrcorner \omega) \\ &= 0 + d(0) = 0, \\ \mathcal{L}_W\omega &= W \lrcorner d\omega + d(W \lrcorner \omega) \\ &= 0 + d(0) = 0,\end{aligned}$$

where in both lines, the first terms on the right sides are zero since ω is closed, and the second terms are zero since V and W are smooth sections of D . Thus, we conclude that $\mathcal{L}_{[V, W]}\omega = 0$. But this means that $[V, W]$ must be a smooth section of D . Thus, by the Frobenius Theorem, we conclude that D is integrable, which means that there exist m -dimensional submanifolds $M \subseteq N$ so that $T_x M = D_x$ for all $x \in M$.

where the final equality follows from the fact that $d\omega = 0$ since ω is a top form on ∂M . On the other hand, since Φ is the identity map on ∂M ,

$$\int_{\partial M} \Phi^* \omega = \int_{\partial M} \omega > 0.$$

This produces a contradiction. Therefore, we conclude that no such retraction Φ can exist.

Problem 1998-J-I-5. Show that a nowhere vanishing 2-form on a compact orientable 2-dimensional manifold cannot be exact.

Let M denote a compact orientable 2-dimensional manifold without boundary, and let ω be a nowhere vanishing 2-form on M . Assume to the contrary that ω is exact; i.e., assume there exists some 1-form α so that $\omega = d\alpha$. Since ω is nowhere vanishing, it is a volume form on M so that if M is equipped with the orientation induced

$$\int_M \omega > 0.$$

On the other hand, since M is without boundary, by Stokes' Theorem,

$$\int_M \omega = \int_M d\alpha = 0,$$

which is a contradiction. Thus, by contradiction, we conclude that ω cannot be exact.

Problem 2017-A-I-1. Let M be a smooth compact connected n -manifold (without boundary), and let $F : M \rightarrow \mathbb{R}^n$ be a smooth map. Does F necessarily have a critical point?

Let M be a smooth compact connected n -manifold (without boundary), and let $F : M \rightarrow \mathbb{R}^n$ be a smooth map. Then F necessarily has a critical point. Recall that a point $p \in M$ is said to be a **regular point** of F if and only if $dF_p : T_p M \rightarrow T_{F(p)} \mathbb{R}^n$ is surjective, and **critical** otherwise. Let $F = (f_1, \dots, f_n)$, where for each $1 \leq j \leq n$, $f_j : M \rightarrow \mathbb{R}$ is a smooth function. Since M is compact, by the extreme value theorem, each f_j has an extrema at some point $p_j \in M$, which means that $df_j|_{p_j} = 0$. But this means that dF_{p_j} (which can be represented by a square $n \times n$ matrix) does not always have full rank n , and therefore, is not surjective. Thus, each p_j is a critical point of F .

Problem 2017-J-I-1. Let Σ_1 be a torus and let Σ_2 be a genus 2 surface. Show that there is no submersion from Σ_2 to Σ_1 .

Let Σ_1 be a torus and let Σ_2 be a genus 2 surface. Assume to the contrary that there exists a submersion from $f : \Sigma_2 \rightarrow \Sigma_1$. Then by the Comps Lemma, f must be a k -sheeted covering map for some finite integer $k > 0$. Thus, if $\chi(\Sigma_j)$ denotes the Euler characteristic of Σ_j , we must have $\chi(\Sigma_2) = k \cdot \chi(\Sigma_1)$. We compute,

$$\chi(\Sigma_1) = 2 - 2(1) = 0 \quad \text{and} \quad \chi(\Sigma_2) = 2 - 2(2) = 2 - 4 = -2.$$

Since there exists no positive integer k such that $-2 = 0(k) = 0$, we have arrived at a contradiction. Thus, there exist no submersions from Σ_2 to Σ_1 .

Problem 2015-A-II-3. Suppose $f : \Sigma_2 \rightarrow \Sigma_1$ is a continuous map from a compact orientable surface Σ_2 of genus 2 to a torus Σ_1 (a compact orientable surface of genus 1). Show that f is not a local homeomorphism (i.e., for some $x \in \Sigma_2$, there exists no neighborhood on which f is a homeomorphism onto its image).

Let $f : \Sigma_2 \rightarrow \Sigma_1$ be a continuous map. Assume to the contrary that f is a local homeomorphism. Then by the Comps Lemma, f is a k -sheeted covering map for some finite positive integer k . Since f is a covering map, we must have $\chi(\Sigma_2) = k \cdot \chi(\Sigma_1)$. But

$$\chi(\Sigma_2) = 2 - 2(2) = 2 - 4 = -2 \quad \text{and} \quad \chi(\Sigma_1) = 2 - 2(1) = 0.$$

Since $k \cdot 0 = 0 \neq -2$, we have reached a contradiction. Thus by contradiction, there cannot be a local homeomorphism from Σ_2 to Σ_1 .

Problem 2011-J-I-3. Let Σ be the double torus (i.e., a compact oriented surface of genus 2) and $T = S^1 \times S^1$ the torus, where S^1 is the circle. Suppose that $f : \Sigma \rightarrow T$ is continuous. Show that f is not a local homeomorphism.

Let $f : \Sigma_2 \rightarrow T$ be a continuous map. Assume to the contrary that f is a local homeomorphism. Then by the Comps Lemma, f is a k -sheeted covering map for some finite positive integer k . Since f is a covering map, we must have $\chi(\Sigma_2) = k \cdot \chi(T)$. But

$$\chi(\Sigma_2) = 2 - 2(2) = 2 - 4 = -2 \quad \text{and} \quad \chi(T) = 2 - 2(1) = 0.$$

Since $k \cdot 0 = 0 \neq -2$, we have reached a contradiction. Thus by contradiction, there cannot be a local homeomorphism from Σ_2 to T .

Problem 2010-J-I-3. Let $f : S^1 \times S^1 \rightarrow \mathbb{RP}^2$ be a smooth map from the 2-torus to the projective plane. Prove that f cannot be an immersion.

Let $f : S^1 \times S^1 \rightarrow \mathbb{RP}^2$ be a smooth map from the 2-torus to the projective plane. Assume to the contrary that f is an immersion. Then from the Comps Lemma, f must be a covering map, so that the induced map $f_* : \pi_1(S^1 \times S^1) \rightarrow \pi_1(\mathbb{RP}^2)$ is injective, which means that the first fundamental group of the two torus would have to be isomorphic to a subgroup of the first fundamental group of the projective plane. But $\pi_1(S^1 \times S^1) \cong \mathbb{Z} \times \mathbb{Z}$ (which is an infinite group), while $\pi_1(\mathbb{RP}^2) \cong \mathbb{Z}/2\mathbb{Z}$ (a finite group). Thus, the proof concludes by contradiction.

Problem 2002-A-I-1. Let $f : S^3 \rightarrow \mathbb{R}^3$ be a C^∞ -map. Show that there must exist at least one point where f does not locally have a C^∞ inverse.

Let $f : S^3 \rightarrow \mathbb{R}^3$ be a smooth map, and assume to the contrary that f has a local smooth inverse at every $p \in S^3$. By definition, it follows that f is a local diffeomorphism. Since diffeomorphisms are open maps, $f(S^3)$ is a nonempty open subset of the manifold $\mathbb{R}^3$. On the other hand, since the image of compact spaces under continuous maps is compact, $f(S^3)$ is compact in $\mathbb{R}^3$. Since $\mathbb{R}^3$ is Hausdorff and compact spaces of Hausdorff spaces are closed, $f(S^3)$ is closed in $\mathbb{R}^3$. Thus, $f(S^3)$ is a nonempty closed and open subset of $\mathbb{R}^3$. Since $\mathbb{R}^3$ is connected, it follows that $f(S^3) = \mathbb{R}^3$. But then, $\mathbb{R}^3$ must be a compact subspace of itself, which is a contradiction. Thus, by contradiction, there must exist some point $p \in S^3$ so that f does not have a smooth inverse in some neighborhood of p .

Problem 2002-J-II-1. Let M and N be smooth, connected, n -dimensional manifolds, and let $f : M \rightarrow N$ be an immersion (i.e., assume that the derivative of f always sends nonzero tangent vectors to non-zero tangent vectors). If M is compact and nonempty, show that N is compact, and that f is a covering map.

Note that this is essentially the same as the Comps Lemma from above. Let M, N be smooth, connected, n -dimensional manifolds, and let $f : M \rightarrow N$ be an immersion. Since the image of compact spaces under continuous maps is compact, $f(M)$ is a nonempty, compact subset of N ; since N is Hausdorff, and compact subspaces of Hausdorff spaces are closed, $f(M)$ is a nonempty closed subset of N . Now since f is an immersion, $df_p : T_p M \rightarrow T_{f(p)} N$ is injective for all $p \in M$. Since $\dim T_p M = \dim M = \dim N = \dim T_{f(p)} N$, it follows that df_p must also be surjective. Thus df_p is bijective. By the Inverse Function Theorem, it follows that f is a local diffeomorphism. Since local diffeomorphisms are open maps, $f(M)$ is an open subset of N . Thus $f(M)$ is a nonempty open and closed subset of N . Since N is connected, it follows that $f(M) = N$. And since $f(M)$ was shown to be compact in N , N is compact.

Now let $q \in N$, and consider $f^{-1}(q)$. Since $\{q\}$ is closed in N and f is continuous, $f^{-1}(q)$ is closed in M . Furthermore, we claim that $f^{-1}(q) \subset M$ is a discrete set. To see this, let $x \in f^{-1}(q)$ and assume to the contrary that $f^{-1}(q)$ is not discrete. Since f is a local diffeomorphism, there exists some neighborhood U of x in M so that $f : U \rightarrow f(U)$ is a diffeomorphism (in particular, is injective). Since x is not isolated, there exists some distinct $y \in f^{-1}(q) \cap U$ such that $f(y) = q$. But then $f(x) = q = f(y)$ and $x \neq y$, which contradicts the claim that f is injective. Thus by contradiction, x must be isolated. Since x was arbitrary, we conclude that every point in $f^{-1}(q)$ is isolated so that $f^{-1}(q)$ is discrete. Since every closed discrete subset of a compact space is finite, we conclude that $f^{-1}(q)$ is finite. Let $q \in N$ be fixed, and assume $\{x_1, x_2, \dots, x_s\} = f^{-1}(q)$. For each $1 \leq j \leq s$, let $\tilde{U}_j$ be a neighborhood of x_j in M so that $f : \tilde{U}_j \rightarrow f(\tilde{U}_j)$ is a diffeomorphism. Without loss of generality, we can assume that the $\tilde{U}_j$ are disjoint. If they are not, then we can shrink each U_j so that they are disjoint and f remains a diffeomorphism on each shrunken neighborhood. Let $V = \bigcap_1^s f(\tilde{U}_j)$, and define $U_j = f^{-1}(V) \cap \tilde{U}_j$. Thus V is an evenly covered neighborhood of q in N . This concludes the proof that f is a covering map.

2.4 de Rham Cohomology

Problem 2026-J-II-6. Let M be the closed 3-manifold obtained from the 3-torus by removing two small disjoint 3-balls and identifying their boundary 2-spheres by an orientation-reversing diffeomorphism. Compute the de Rham cohomology of M .

Let M be the closed 3-manifold obtained by the construction given above. We recall that this procedure is called the *autosum* of M and denotes the connected sum of the 3-torus, T^3 , with the handle $S^2 \times S^1$. Thus, for every p , $H_{\text{dR}}^p(M) \cong H_{\text{dR}}^p(T^3 \# (S^2 \times S^1))$. Since $\dim M = 3$, $H_{\text{dR}}^k(M) \cong 0$ for all $k \geq 4$. Since M is compact connected and orientable, $H_{\text{dR}}^3(M) \cong \mathbb{R}$. Likewise, since M is connected, $H_{\text{dR}}^0(M) \cong \mathbb{R}$. Since T^3 and $S^2 \times S^1$ are connected and orientable, by Problem LSM-17-7, one has $H_{\text{dR}}^p(M) \cong H_{\text{dR}}^p(T^3) \oplus H_{\text{dR}}^p(S^2 \times S^1)$ for $p = 1, 2$. We begin by computing the following:

$$\begin{aligned}
 H_{\text{dR}}^1(T^3) &= H_{\text{dR}}^1(S^1 \times T^2) = \bigoplus_{i+j=1} H_{\text{dR}}^i(S^1) \otimes H_{\text{dR}}^j(T^2) \\
 &= H_{\text{dR}}^0(S^1) \otimes H_{\text{dR}}^1(T^2) \oplus H_{\text{dR}}^1(S^1) \otimes H_{\text{dR}}^0(T^2) \\
 &= \mathbb{R} \otimes \mathbb{R}^2 \oplus \mathbb{R} \otimes \mathbb{R} \cong \mathbb{R}^2 \oplus \mathbb{R} \cong \mathbb{R}^3. \\
 H_{\text{dR}}^1(S^2 \times S^1) &= \bigoplus_{i+j=1} H_{\text{dR}}^i(S^2) \otimes H_{\text{dR}}^j(S^1) \\
 &= H_{\text{dR}}^0(S^2) \otimes H_{\text{dR}}^1(S^1) \oplus H_{\text{dR}}^1(S^2) \otimes H_{\text{dR}}^0(S^1) \\
 &= \mathbb{R} \otimes \mathbb{R} \oplus 0 \otimes \mathbb{R} \cong \mathbb{R} \oplus 0 \cong \mathbb{R}. \\
 H_{\text{dR}}^2(T^3) &= H_{\text{dR}}^2(S^1 \times T^2) = \bigoplus_{i+j=2} H_{\text{dR}}^i(S^1) \otimes H_{\text{dR}}^j(T^2) \\
 &= H_{\text{dR}}^0(S^1) \otimes H_{\text{dR}}^2(T^2) \oplus H_{\text{dR}}^1(S^1) \otimes H_{\text{dR}}^1(T^2) \oplus H_{\text{dR}}^2(S^1) \otimes H_{\text{dR}}^0(T^2) \\
 &= \mathbb{R} \otimes \mathbb{R} \oplus \mathbb{R} \otimes \mathbb{R}^2 \oplus 0 \otimes \mathbb{R} \\
 &= \mathbb{R} \oplus \mathbb{R}^2 \cong \mathbb{R}^3. \\
 H_{\text{dR}}^2(S^2 \times S^1) &= \bigoplus_{i+j=2} H_{\text{dR}}^i(S^2) \otimes H_{\text{dR}}^j(S^1) \\
 &= H_{\text{dR}}^0(S^2) \otimes H_{\text{dR}}^2(S^1) \oplus H_{\text{dR}}^1(S^2) \otimes H_{\text{dR}}^1(S^1) \oplus H_{\text{dR}}^2(S^2) \otimes H_{\text{dR}}^0(S^1) \\
 &= \mathbb{R} \otimes 0 \oplus 0 \otimes \mathbb{R} \oplus \mathbb{R} \otimes \mathbb{R} \cong \mathbb{R}.
 \end{aligned}$$

Therefore, using the above information, we conclude that

$$H_{\text{dR}}^k(M) = \begin{cases} \mathbb{R}, & k = 0, 3 \\ \mathbb{R}^4, & k = 1, 2 \\ 0, & k \geq 4 \end{cases}$$

Problem 2026-J-II-6 (Modified). Let M be the closed 3-manifold obtained by removing two small 3-balls from two 3-tori and identifying their boundary 2-spheres by an orientation-reversing diffeomorphism. Compute the de Rham cohomology groups $H_{\text{dR}}^k(M, \mathbb{R})$.

Let M be the closed 3-manifold obtained from the 3-torus by removing two small disjoint 3-balls from 2 3-tori and identifying their boundary 2-spheres by an orientation-reversing diffeomorphism. I.e., let $M = T^3 \# T^3$ be the connected sum of two 3-tori. Since M is connected, $H_{\text{dR}}^0(M) \cong \mathbb{R}$. Since $\dim M = 3$, $H_{\text{dR}}^k(M) \cong 0$ for all $k \geq 4$. On the other hand, since T^3 is compact and orientable, by Problem LSM-17-7,

$$H_{\text{dR}}^1(M) \cong H_{\text{dR}}^1(T^3) \oplus H_{\text{dR}}^1(T^3) \cong \mathbb{R}^3 \oplus \mathbb{R}^3 \cong \mathbb{R}^6.$$

Likewise, by Problem LSM-17-7,

$$H_{\text{dR}}^2(M) \cong H_{\text{dR}}^2(T^3) \oplus H_{\text{dR}}^2(T^3) \cong \mathbb{R}^3 \oplus \mathbb{R}^3 \cong \mathbb{R}^6.$$

Therefore, we conclude that

$$H_{\text{dR}}^k(M) = \begin{cases} \mathbb{R}, & k = 0, 3 \\ \mathbb{R}^6, & k = 1, 2 \\ 0, & k \geq 4. \end{cases}$$

Problem 2024-A-I-1. Let M be a smooth compact manifold without boundary, and let φ be a smooth closed 1-form on M that has the property that $\varphi \neq 0$ at every point of M . Prove that the first de Rham cohomology $H_{\text{dR}}^1(M)$ of the given manifold is non-zero.

Let M be a smooth compact manifold without boundary, and let φ be a smooth closed 1-form on M so that $\varphi \neq 0$ at every point of M . Assume to the contrary that $H_{\text{dR}}^1(M) = 0$, which means that φ must be a closed 1-form; i.e., there exists a smooth function $f : M \rightarrow \mathbb{R}$ so that $\varphi = df$. Since M is compact and f is a smooth function, f has an extrema at some point $p \in M$. But this necessarily means that p is a critical point of f , which implies $df_p = 0$. I.e., φ is zero at $p \in M$, which is a contradiction. Thus, by contradiction, φ is not exact, which means that $H_{\text{dR}}^1(M) \neq 0$ since there exists a closed 1-form that is not exact.

Problem 2024-J-I-5. Let α be a closed 1-form on $\mathbb{R}P^n$, $n > 1$. Show that if $f : [0, 1] \rightarrow \mathbb{R}P^n$ is a smooth function such that $f(0) = f(1)$, then

$$\int_{[0,1]} f^* \alpha = 0.$$

Include all calculations that are relevant to your solution.

Let α be a closed 1-form on $\mathbb{R}P^n$, $n > 1$, and let $f : [0, 1] \rightarrow \mathbb{R}P^n$ be a smooth function so that $f(0) = f(1)$. To calculate the integral $\int_{[0,1]} f^* \alpha$, we need to know the de Rham cohomology groups of $\mathbb{R}P^n$; in particular, we must know the first de Rham cohomology group of $\mathbb{R}P^n$. **!!! Show explicit computation of the cohomology groups !!!** Since $H_{\text{dR}}^1(\mathbb{R}P^n) = 0$, α must also be an exact 1-form on $\mathbb{R}P^n$. I.e., there exists a smooth function $\beta : \mathbb{R}P^n \rightarrow \mathbb{R}$ so that $\alpha = d\beta$. Then one has

$$\int_{[0,1]} f^* \alpha = \int_{[0,1]} f^* d\beta = \beta(f(1)) - \beta(f(0)) = \beta(f(0)) - \beta(f(0)) = 0.$$

Problem 2021-A-II-1. Let M be a compact manifold (without boundary) and $\pi : M \rightarrow S^1$ a submersion onto the circle. Show that the de Rham group $H_{\text{dR}}^1(M) \neq 0$.

Let M be a compact smooth manifold without boundary, and let $\pi : M \rightarrow S^1$ be a submersion onto the circle. Assume to the contrary that $H_{\text{dR}}^1(M) = 0$, which means that every closed form on M is an exact form. The cohomology class of the closed 1-form $d\theta$ is a generator of $H_{\text{dR}}^1(S^1) \cong \mathbb{R}$. Then $\pi^* d\theta$ is a closed 1-form on M . Since $H_{\text{dR}}^1(M) = 0$, there exists a smooth function $f : M \rightarrow \mathbb{R}$ so that $\pi^* d\theta = df$. In particular, by the extreme value theorem, there exists some $p \in M$ so that f has an extrema at p . But this means that $df_p = 0$, so that $\pi^* d\theta$ vanishes at p . We will show that this is a contradiction. Since π is a submersion, its differential $d\pi_x$ is surjective at every $x \in M$. Then since $d\theta$ is nowhere vanishing, for all $x \in M$, $d\theta_{\pi(x)} \neq 0$. But then $(\pi^* d\theta)_x := d\theta_{\pi(x)} \circ d\pi_x \neq 0$, directly contradicting our above assumption. Thus by contradiction, we conclude that $H_{\text{dR}}^1(M) \neq 0$.

Problem 2020-J-II-2. Let φ be a smooth closed 4-form on a compact 12-manifold M with smooth boundary. If $\varphi \wedge \varphi \wedge \varphi \neq 0$ everywhere on M , prove that the de Rham cohomologies $H_{\text{dR}}^4(M)$ and $H_{\text{dR}}^8(M)$ are both non-zero.

Let φ be a smooth closed 4-form on a compact 12-manifold M with smooth boundary. Assume also that $\varphi \wedge \varphi \wedge \varphi =: \varphi^3$ is nowhere vanishing on M ; i.e., φ^3 defines an orientation form on M . Then $\int_M \varphi^3 \neq 0$.

Now assume to the contrary that $H_{\text{dR}}^4(M) = 0$. Then φ must be an exact form so that $\varphi = d\eta$ for some smooth 3-form η . Because $d\varphi = 0$, one can write

$$\varphi \wedge \varphi \wedge \varphi = d(\eta \wedge \varphi \wedge \varphi).$$

But then, by Stokes' Theorem,

$$\int_M \varphi^3 = \int_M d(\eta \wedge \varphi \wedge \varphi) = \int_{\partial M} \eta \wedge \varphi \wedge \varphi = 0,$$

since $\partial M = \emptyset$. This is a contradiction. Thus, $H_{\text{dR}}^4(M) \neq 0$. Likewise, assume $H_{\text{dR}}^8(M) = 0$. This means that φ^2 is exact; there exists some smooth 7-form η on M so that $\varphi^2 = d\eta$. Thus since $d\varphi = 0$, we can write $\varphi^3 = d(\eta \wedge \varphi)$. Then by Stokes' Theorem,

$$0 \neq \int_M \varphi^3 = \int_M d(\eta \wedge \varphi) = \int_{\partial M} \eta \wedge \varphi = 0,$$

since $\partial M = \emptyset$. By contradiction, we conclude that $H_{\text{dR}}^8(M) \neq 0$.

Problem 2019-A-II-4. Let M^n be a compact, connected, oriented n -manifold, and assume that n is odd. Compute the Euler characteristic $\chi(M^n \setminus \{\text{pt}\})$.

Let M^n be a compact, connected, oriented n -manifold, and assume that n is odd. Let us fix some point $p \in M^n$, and let B be a small n -ball in M centered at p . Then B deformation retracts onto $\{p\}$, and $M \setminus \{p\}$ deformation retracts onto $M \setminus B$. Moreover, $B \cap (M \setminus B) = S^{n-1}$. Thus by additivity of the Euler characteristic, one has

$$\chi(M) = \chi(M \setminus B) + \chi(B) - \chi(S^{n-1}).$$

Since n is odd, $n-1$ is even so that $\chi(S^{n-1}) = 2$. On the other hand, by Poincaré duality, one has $\chi(M) = 0$, and $\chi(B) = 1$. Thus, we conclude that $\chi(M \setminus B) = \chi(M \setminus \{p\}) = 1$.

Problem 2019-J-I-3. Let M and N be connected, compact, oriented n -manifolds. Prove that if $H_{\text{dR}}^k(N) \neq 0$ and $H_{\text{dR}}^k(M) = 0$ for some $k \neq 0$, then any smooth map $f : M \rightarrow N$ has degree 0.

Let M and N be connected, compact, oriented n -manifolds so that $H_{\text{dR}}^k(N) \neq 0$ and $H_{\text{dR}}^k(M) = 0$ for some $k \neq 0$. Further, let $f : M \rightarrow N$ be a smooth map. The map f induces a smooth pullback map $f^* : H_{\text{dR}}^k(N) \rightarrow H_{\text{dR}}^k(M)$. But since $H_{\text{dR}}^k(M) = 0$, f^* must be the zero map. Let α be a closed k -form on N so that $[\alpha] \neq 0$. Since f^* is the zero map, one must have $[f^*\alpha] = 0 \in H_{\text{dR}}^k(M)$ so that $f^*\alpha$ is an exact form. Let β be a $(k-1)$ -form on M so that $f^*\alpha = d\beta$. Then by the Poincaré duality for de Rham cohomology, there exists a closed $(n-k)$ -form γ on N so that

$$\int_N \alpha \wedge \gamma \neq 0.$$

But this implies that

$$\int_M f^*(\alpha \wedge \gamma) = \int_M f^*\alpha \wedge f^*\gamma = \int_M d\beta \wedge f^*\gamma = \int_M d(\beta \wedge f^*\gamma),$$

where in the final equality, we used the fact that $f^*\gamma$ is closed. Since M is without boundary, Stokes' Theorem tells us that the above integral must be zero. Thus,

$$\int_M f^*(\alpha \wedge \gamma) = 0 \cdot \int_N (\alpha \wedge \gamma).$$

By the definition of the degree of a smooth map, we conclude that $\deg f = 0$.

Problem 2015-J-I-5. Let X be a connected smooth manifold, and let U and V be connected open subsets such that $X = U \cup V$. Show that if $U \cap V$ is not connected, then the de Rham cohomology group $H_{\text{dR}}^1(X, \mathbb{R})$ must be nonzero.

Let X be a connected smooth manifold, and let U and V be connected open subsets such that $X = U \cup V$. Further assume that $U \cap V$ is not connected; let n denote the number of connected components of $U \cap V$. Then, by the Mayer-Vietoris theorem, we have the exact sequence,

$$0 \rightarrow H_{\text{dR}}^0(X) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^0(U) \oplus H_{\text{dR}}^0(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^0(U \cap V) \xrightarrow{\delta} H_{\text{dR}}^1(X) \\ \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^1(U \cap V) \xrightarrow{\delta} \dots,$$

where $i : U \cap V \hookrightarrow U$, $j : U \cap V \hookrightarrow V$, $k : U \hookrightarrow X$, and $l : V \hookrightarrow X$. Since X, U, V are connected and $U \cap V$ has n connected components, the exact sequence becomes,

$$0 \rightarrow \mathbb{R} \xrightarrow{k^* \oplus l^*} \mathbb{R}^2 \xrightarrow{i^* - j^*} \mathbb{R}^n \xrightarrow{\delta} H_{\text{dR}}^1(X) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^1(U \cap V) \xrightarrow{\delta} \dots,$$

To show that $H_{\text{dR}}^1(X)$ is nonzero, it suffices to show that the map δ is nonzero. We recall that each equivalence class in $H_{\text{dR}}^0(U)$ and $H_{\text{dR}}^0(V)$ is represented by a constant function on U and V , respectively. Since each such function restricts to a constant on each connected component W_j of $U \cap V$, we have that on each connected component W_j of $U \cap V$, $(i^* - j^*)|_{W_j}(c, d) = c - d$. Thus under the identification $H_{\text{dR}}^0(U \cap V) \cong \mathbb{R}^n$, $i^* - j^*$ is given by the map $(c, d) \mapsto (c - d, \dots, c - d)$. The image of this map is the set $\{(t, \dots, t) \in \mathbb{R}^n : t \in \mathbb{R}\} \cong \mathbb{R}$ so that $\text{rank}(i^* - j^*) = 1$. By exactness, $\text{nullity}(\delta) = 1$. Thus by the rank-nullity theorem, we must have

$$\text{rank}(\delta) + \text{nullity}(\delta) = \dim(\mathbb{R}^n) = n \implies \text{rank}(\delta) = n - 1.$$

Since we assumed $U \cap V$ was not connected, $n > 1$ (thus, $n - 1 > 0$). Therefore, since the rank of δ is positive, δ cannot be the zero map. Hence, $H_{\text{dR}}^1(X)$ cannot be zero. The proof concludes.

Problem 2013-A-II-6. Let M be a smooth compact oriented n -manifold with boundary. Assume that ∂M is connected, and let $j : \partial M \hookrightarrow M$ denote the inclusion map. Prove that the induced map $j^* : H_{\text{dR}}^{n-1}(M) \rightarrow H_{\text{dR}}^{n-1}(\partial M)$ of de Rham cohomology is zero. Then use this to show there is no smooth retraction $F : M \rightarrow \partial M$.

Let M be a smooth compact oriented n -manifold with boundary, assume that ∂M is connected, and let $j : \partial M \hookrightarrow M$ denote the inclusion map. Consider the induced map $j^* : H_{\text{dR}}^{n-1}(M) \rightarrow H_{\text{dR}}^{n-1}(\partial M)$ of de Rham cohomology groups. Let $[\beta] \in H_{\text{dR}}^{n-1}(M) \setminus \{0\}$ (i.e., let β be a closed, non-exact $(n-1)$ -form on M). Assume to the contrary that $j^*\beta$ is not an exact $(n-1)$ -form on N . Then since ∂M is also compact and oriented, one must have

$$\int_{\partial M} j^*\beta \neq 0.$$

But by Stokes' Theorem,

$$\int_{\partial M} j^*\beta = \int_M d\beta = 0,$$

since $d\beta$ is an exact n -form on M and M is compact. This is a contradiction. Thus, by contradiction $j^*\beta$ is exact. Since β was arbitrary, j^* maps every closed $(n-1)$ -form on M to an exact $(n-1)$ -form on ∂M so that j^* is the zero map on de Rham cohomology groups. Now assume to the contrary that there exists a smooth retraction $F : M \twoheadrightarrow \partial M$. Since ∂M is oriented, let ω be a smooth nowhere vanishing $(n-1)$ -form on ∂M (i.e., a volume form). Let ∂M be equipped with the orientation induced by ω so that

$$\int_{\partial M} \omega > 0.$$

Note that ω is a closed differential form. Then, $F^*\omega$ is a closed $(n-1)$ -form on M . By our above work, $j^*F^*\omega$ must be an exact $(n-1)$ -form on ∂M so that since ∂M is compact, one has $\int_{\partial M} j^*F^*\omega = 0$. But $(F \circ j) = \text{id}_{\partial M}$ so that $j^* \circ F^* = (F \circ j)^* = \text{id}_{\partial M}^*$. This means that

$$\int_{\partial M} j^*F^*\omega = \int_{\partial M} \omega > 0,$$

which contradicts our previous finding. Thus, by contradiction, there cannot exist a smooth retraction $F : M \rightarrow \partial M$.

Problem 2013-J-II-6. Let M be a smooth compact manifold, and suppose that there is a smooth map $F : M \rightarrow S^1$ whose derivative is non-zero at every point. Prove that the de Rham cohomology space $H_{\text{dR}}^1(M)$ is non-zero.

Let M be a smooth compact manifold, and suppose there is a smooth map $F : M \rightarrow S^1$ whose derivative is non-zero at every point $p \in M$. Assume to the contrary that $H_{\text{dR}}^1(M) = 0$. Since $H_{\text{dR}}^1(S^1) \cong \mathbb{R}$, this cohomology space is generated by the cohomology class of the nowhere vanishing closed 1-form $d\theta$ on S^1 . Then $F^*d\theta$ is a closed 1-form on M . Since $H_{\text{dR}}^1(M) = 0$, there exists a smooth function $f : M \rightarrow \mathbb{R}$ so that $F^*d\theta = df$. In particular, by the extreme value theorem, there exists some $p \in M$ so that f has an extrema at p . But this means that $df_p = 0$, so that $F^*d\theta$ vanishes at p . We will show that this is a contradiction. For any $p \in M$, $d\theta_{F(p)} \neq 0$. Moreover, by the given hypothesis, one has $dF_p \neq 0$. But this implies that

$$(F^*d\theta)_p := d\theta_{F(p)} \circ dF_p \neq 0,$$

directly contradicting our result from above. Thus by contradiction, we conclude that $H_{\text{dR}}^1(M) \neq 0$.

Problem 2002-J-II-5. Let θ be a closed 1-form on a compact manifold M without boundary. Further suppose that $\theta \neq 0$ at each point of M . Prove that $H_{\text{dR}}^1(M) \neq 0$.

Let θ be a closed 1-form on a compact manifold M without boundary, and suppose that $\theta \neq 0$ at each point of M . Assume to the contrary that $H_{\text{dR}}^1(M) = 0$. This means that θ is an exact 1-form, so that there exists some smooth 0-form f satisfying $\theta = df$ (i.e., $f : M \rightarrow \mathbb{R}$ is a smooth function). But since M is compact, f has an extrema at some point $p \in M$. This means that $df_p = 0$, and thus by extension, $\theta_p = 0$. But this contradicts our hypothesis on θ . Thus, by contradiction, θ cannot be an exact 1-form. Since we have shown the existence of a closed 1-form that is not exact, $H_{\text{dR}}^1(M) \neq 0$.

Problem 1997-A-I-5. For which values of p is the de Rham cohomology $H_{\text{dR}}^p(S^1 \times S^2)$ non-trivial? Explain.

First since $S^1 \times S^2$ is a smooth 3-manifold, $H_{\text{dR}}^p(S^1 \times S^2) = 0$ for all $p \geq 4$. Moreover, since $S^1 \times S^2$ is connected, $H_{\text{dR}}^0(S^1 \times S^2) \cong \mathbb{R}$ is non-trivial. Thus, it suffices to examine the case for $p = 1, 2, 3$. We recall Künneth's Formula, which states that for smooth manifolds M and N ,

$$H_{\text{dR}}^k(M \times N) \cong \bigoplus_{i+j=k} H_{\text{dR}}^i(M) \otimes H_{\text{dR}}^j(N).$$

Thus, we have

$$H_{\text{dR}}^1(S^1 \times S^2) \cong (H_{\text{dR}}^0(S^1) \otimes H_{\text{dR}}^1(S^2)) \oplus (H_{\text{dR}}^1(S^1) \otimes H_{\text{dR}}^0(S^2)) \cong \mathbb{R} \otimes 0 \oplus \mathbb{R} \otimes \mathbb{R} \cong 0 \oplus \mathbb{R} \cong \mathbb{R}.$$

Next, we have

$$\begin{aligned} H_{\text{dR}}^2(S^1 \times S^2) &\cong (H_{\text{dR}}^2(S^1) \otimes H_{\text{dR}}^0(S^2)) \oplus (H_{\text{dR}}^1(S^1) \otimes H_{\text{dR}}^1(S^2)) \oplus (H_{\text{dR}}^0(S^1) \otimes H_{\text{dR}}^2(S^2)) \\ &\cong (0 \otimes \mathbb{R}) \oplus (\mathbb{R} \otimes 0) \oplus (\mathbb{R} \otimes \mathbb{R}) \cong \mathbb{R}. \end{aligned}$$

Finally,

$$H_{\text{dR}}^3(S^1 \times S^2) \cong (H_{\text{dR}}^1(S^1) \otimes H_{\text{dR}}^2(S^2)) \cong \mathbb{R}.$$

Thus, $H_{\text{dR}}^p(S^1 \times S^2)$ is non-trivial for $p = 0, 1, 2, 3$.

Problem 2001-A-II-3. We say that a map $f : M \rightarrow N$ between two n -dimensional orientable manifolds is of “degree 1” if it induces an isomorphism $f_* : H_{\text{dR}}^n(N) \rightarrow H_{\text{dR}}^n(M)$. Prove that there is no map of degree 1 from the 2-sphere S^2 to the 2-dimensional torus T^2 .

Assume to the contrary that there exists a map $f : S^2 \rightarrow T^2$ of degree 1, so that it induces an isomorphism $f_* : H_{\text{dR}}^n(T^2) \rightarrow H_{\text{dR}}^n(S^2)$ of de Rham cohomology groups. [!! Complete Later !!]

Problem 2007-A-II-3. Calculate the de Rham cohomology groups $H_{\text{dR}}^k(\mathbb{T}^2 \# \mathbb{P}^2, \mathbb{R})$, where $\mathbb{T}^2$ is the torus and $\mathbb{P}^2$ is the projective plane.

Let $\mathbb{T}^2$ and $\mathbb{P}^2$ denote 2-torus and the projective plane, respectively. Since both of these are connected, $\mathbb{T}^2 \# \mathbb{P}^2$ is also connected. Thus,

$$H_{\text{dR}}^0(\mathbb{T}^2 \# \mathbb{P}^2) \cong \mathbb{R}.$$

Since $\mathbb{T}^2, \mathbb{P}^2$ are both compact, $\mathbb{T}^2 \# \mathbb{P}^2$ is compact. However, since $\mathbb{P}^2$ is nonorientable, $\mathbb{T}^2 \# \mathbb{P}^2$ is nonorientable. Thus since the connected sum is a smooth compact, non-orientable, and connected 2-manifold, $H_{\text{dR}}^2(\mathbb{T}^2 \# \mathbb{P}^2) = 0$; $H_{\text{dR}}^k(\mathbb{T}^2 \# \mathbb{P}^2) \cong 0$ for all $k \geq 3$ since the connected sum is a 2-manifold. Thus, we just need to compute the first de Rham cohomology group. We can solve this using the Euler characteristic. We recall that

$$\chi(M) = \sum_0^{\dim M} (-1)^i H_{\text{dR}}^i(M).$$

On the other hand, $\chi(M_1 \# M_2) = \chi(M_1) + \chi(M_2) - 2$. Therefore,

$$-1 = 0 + 1 - 2 = 0 - \dim H_{\text{dR}}^1(\mathbb{T}^2 \# \mathbb{P}^2) + 1 = 1 - \dim H_{\text{dR}}^1(\mathbb{T}^2 \# \mathbb{P}^2).$$

Thus, $\dim H_{\text{dR}}^1(\mathbb{T}^2 \# \mathbb{P}^2) = 2$ so that $H_{\text{dR}}^1(\mathbb{T}^2 \# \mathbb{P}^2) = \mathbb{R}^2$. Therefore, we have the following:

$$H_{\text{dR}}^k(\mathbb{T}^2 \# \mathbb{P}^2) = \begin{cases} \mathbb{R}, & k = 0 \\ \mathbb{R}^2, & k = 1 \\ 0, & k \geq 2 \end{cases}.$$

2.5 Fundamental Groups

Problem 2023-J-II-4. Prove that $S^2 \times S^2$ is not diffeomorphic to $M_1 \times M_2 \times M_3$, where M_1, M_2, M_3 are smooth manifolds of nonzero dimension.

Assume to the contrary that $f : S^2 \times S^2 \rightarrow M_1 \times M_2 \times M_3$ is a diffeomorphism. Since $\dim(M_1 \times M_2 \times M_3) = \sum \dim(M_j) = 4$, we have (up to relabeling of indices), $\dim M_1, \dim M_2 = 1$, and $\dim M_3 = 2$. Since $S^2 \times S^2$ is a compact manifold, f is continuous, and surjective, it follows that $M_1 \times M_2 \times M_3$ is compact. Thus, each M_j is compact. By the classification of manifolds of dimension 1, there exists exactly one smooth compact 1-manifold (up to diffeomorphism), which is the unit circle S^1 . Thus, $M_1, M_2 \cong S^1$. This means that $\pi_1(M_1 \times M_2 \times M_3) \cong \mathbb{Z} \times \mathbb{Z}$ is nontrivial. But this is a contradiction: one has $\pi_1(S^2 \times S^2) \cong \pi_1(S^2) \times \pi_1(S^2) \cong 0$, and $f_* : \pi_1(S^2 \times S^2) \rightarrow \pi_1(M_1 \times M_2 \times M_3)$ is an isomorphism since f is a diffeomorphism implies that $0 \cong \mathbb{Z} \times \mathbb{Z}$. Thus, by contradiction, $S^2 \times S^2$ is not diffeomorphic to $M_1 \times M_2 \times M_3$.

Problem 2003-J-II-5. Is the unit sphere S^3 diffeomorphic to the product of two other smooth manifolds of dimension > 0 ?

Assume that the unit sphere S^3 is diffeomorphic to the product of two other smooth manifolds, M_1, M_2 , both of nonzero dimension. Since diffeomorphisms preserve the total dimension, $\dim S^3 = \dim M_1 \times M_2 = \dim M_1 + \dim M_2$. Thus, up to relabeling of indices, $\dim M_1 = 1$ and $\dim M_2 = 2$. Moreover, since S^3 is compact, $M_1 \times M_2$ must also be compact. Since the product of manifolds is compact if and only if each factor is compact, M_1, M_2 are both compact. Since there exists only one compact smooth 1-manifold up to diffeomorphism, namely the unit circle S^1 , $M_1 \cong S^1$. Diffeomorphic manifolds have isomorphic fundamental groups so that $\pi_1(S^3) \cong \pi_1(M_1 \times M_2) \cong \pi_1(M_1) \times \pi_1(M_2) \cong \pi_1(S^1) \times \pi_1(M_2)$. Since S^3 is simply connected, $\pi_1(S^3)$ is trivial. On the other hand, since $\pi_1(S^1) \cong \mathbb{Z}$, $\pi_1(M_1 \times M_2)$ must be nontrivial, which is a contradiction. Thus by contradiction, the unit sphere S^3 cannot be diffeomorphic to the product of two other smooth manifolds of nonzero dimensions.

we must have $\chi_{\tilde{M}} = k \cdot \chi_M$, where χ denotes the Euler characteristic. Since $\chi_M = 2 - 2(1) = 0$, $\chi_{\tilde{M}} = 2 - 2(2) = -2$, and there exists no finite positive integer k so that $-2 = k \cdot 0$, we conclude that M has no finitely-sheeted covering spaces homeomorphic to a surface 2 genus surface. On the other hand, any infinitely-sheeted covering space of M has to be noncompact, and thus cannot be homeomorphic to a (compact) surface of genus 2.

- (b) Now let M denote a surface of genus 2. As above, it suffices to check if there exists a finitely-sheeted covering space of M with genus 3. By looking at the Euler characteristics, *if* M has a finitely-sheeted covering space of genus 3, then it must be a 2-sheeted covering space. But by replicating the work from Problem [2021-A-I-6](#), we see that by Galois correspondence of covering spaces, M must have a 2-sheeted covering space. Thus, we conclude that M has a covering space which is homeomorphic to a surface of genus 2.

3 Algebra

3.1 Linear Algebra

Problem 2022-A-I-6. Denote by $\text{Mat}(m, \mathbb{C})$ and $\text{Mat}(n, m, \mathbb{C})$ the space of $m \times m$ and $n \times m$ matrices with complex coefficients. Let $A \in \text{Mat}(n, \mathbb{C})$, $B \in \text{Mat}(m, \mathbb{C})$ be given and suppose A and B have no common eigenvalues. Show that the only solution of

$$AX = XB, \quad X \in \text{Mat}(n, m, \mathbb{C}),$$

is the trivial solution $X = 0$.

Assume all of the given hypotheses. Observe that $A^2X = A(AX) = (AX)B = XB^2$. Hence, by induction, it follows that for all k ,

$$A^kX = XB^k.$$

In particular, $p(A)X = Xp(B)$ for any $p \in \mathbb{C}[z]$. Let p be the characteristic polynomial $p_B(z)$ of B . By the Cayley-Hamilton theorem, one obtains,

$$p_B(A)X = Xp_B(B) = 0.$$

We can write

$$p_B(z) = \prod_{i=1}^k (z - \lambda_k)^{m_k},$$

where each λ_k is an eigenvalue of B , and m_k is its multiplicity. By the hypothesis, λ_k is not an eigenvalue of A . Hence, $(A - I\lambda_k)^{m_k}$ is invertible. Therefore, since each factor is invertible, $p_B(A)$ is invertible. Hence,

$$p_B(A)X = 0 \implies p_B^{-1}(A)p_B(A)X = 0 \implies X = 0.$$

Problem 2019-J-I-1. Let A and B be $n \times n$ invertible matrices over complex numbers, satisfying

$$AB = \lambda BA \quad \text{for some } \lambda \in \mathbb{C}.$$

Prove that A^n and B commute.

Let A and B be $n \times n$ invertible matrices over the complex numbers, and $\lambda \in \mathbb{C}$ so that $AB = \lambda BA$. First we will show that λ^n is a root of unity. Taking the determinant on both sides and using the fact that $\det : \text{GL}(n, \mathbb{C}) \rightarrow \mathbb{C}^\times$ is a group homomorphism, one gets,

$$\det A \det B = \det AB = \det(\lambda BA) = \lambda^n \det A \det B.$$

Since A, B are invertible, $\det A, \det B \neq 0$, so that dividing through by these quantities gets us $\lambda^n = 1$. Now we observe the following:

$$AB = \lambda BA.$$

$$A^2B = \lambda(AB)A = \lambda^2BA^2.$$

$$A^3B = \lambda^2(AB)A^2 = \lambda^3BA^3.$$

Repeating this procedure inductively, one finally obtains $A^nB = \lambda^nBA^n$. But we showed above that $\lambda^n = 1$. Thus, one gets $A^nB = BA^n$, which concludes the proof that A^n and B commute.

From the first equation, one gets $a = -c$. Thus making this substitution, one obtains the following system of equations:

$$\begin{cases} b + d - c^2 = -5 \\ c(b - d) = 0 \\ bd = 5. \end{cases}$$

Since $\mathbb{Q}$ is an integral domain, either $c = 0$ or $b - d = 0$. If $c = 0$, then one has $b + d = -5$ and $bd = 5$. From this, one obtains $b = 5/d$. Plugging this into the first equation, one gets

$$-5 = b + d = \frac{5}{d} + d \implies -5d = 5 + d^2 \implies d^2 + 5d + 5 = 0.$$

But there exist no rational solutions to $d^2 + 5d + 5 = 0$. Now suppose $b - d = 0$ so that $b = d$. Then one has the system of equations

$$\begin{aligned} 2b - c^2 &= -5 \\ b^2 &= 5. \end{aligned}$$

Then second equation implies $b = \pm\sqrt{5}$ so that

$$\pm 2\sqrt{5} - c^2 = -5 \implies c^2 = \pm 2\sqrt{5} + 5 \implies c = \{\pm\sqrt{2\sqrt{5} + 5}, \pm\sqrt{-2\sqrt{5} + 5}\},$$

neither of which are rational. Thus, we conclude that $p(x)$ is irreducible over $\mathbb{Q}$. To find the splitting field and the Galois group, we will compute the roots of $p(x)$. Setting $u = x^2$, one has $0 = p(u) = u^2 - 5u + 5$ so that

$$u = \frac{5 \pm \sqrt{25 - 20}}{2} = \frac{5 \pm \sqrt{5}}{2} \implies x = \pm\alpha, \pm\beta,$$

where

$$\alpha := \sqrt{\frac{5 + \sqrt{5}}{2}}, \beta := \sqrt{\frac{5 - \sqrt{5}}{2}}.$$

Let K denote the splitting field of $p(x)$ over $\mathbb{Q}$. Then K must contain both α and β (and thus the field extension $\mathbb{Q}(\alpha, \sqrt{5})$ since $\alpha\beta = \sqrt{5}$). On the other hand, $\mathbb{Q}(\alpha, \sqrt{5})$ contains all of the roots of $p(x)$. Then the splitting field is $\mathbb{Q}(\alpha, \sqrt{5})$. Since in fact, $\sqrt{5} \in \mathbb{Q}(\alpha)$, we conclude that the splitting field is actually just $\mathbb{Q}(\alpha)$. Since $\mathbb{Q}(\alpha)$ is the splitting field of an irreducible monic polynomial, $\mathbb{Q}(\alpha)/\mathbb{Q}$ is Galois. The Galois group of the polynomial consists of the automorphisms on $\mathbb{Q}(\alpha)$ that preserve the base field $\mathbb{Q}$ and the algebraic relations between the roots. Thus we have the four automorphisms:

$$\begin{aligned} \alpha &\mapsto \alpha & \alpha &\mapsto -\alpha \\ \alpha &\mapsto \beta & \alpha &\mapsto -\beta. \end{aligned}$$

The first automorphism is trivial; the second automorphism has order 2, the third automorphism has order 4, and the fourth automorphism has order 4. Thus, we conclude that the Galois group is isomorphic to $\mathbb{Z}/4\mathbb{Z}$.

Problem 2014-A-II-1. Determine the Galois group of the polynomial $x^4 + 4$ over $\mathbb{Q}$.

Let $p(x) = x^4 + 4$. First, we begin by checking if $p(x)$ is irreducible over $\mathbb{Q}$. It is straightforward to see that p has no linear factors since it has no rational roots. Instead, we look for a decomposition of the form $(x^2 + ax + b)(x^2 + cx + d)$. Doing so, we obtain the decomposition,

$$x^4 + 4 = (x^2 - 2x + 2)(x^2 + 2x + 2).$$

Each of these quadratic are irreducible as they have no rational (nor real) roots. Solving these quadratics, we find the four roots of $p(x)$ to be $1 \pm i, -1 \pm i$. Thus, it is clear that the splitting field of $p(x)$ over $\mathbb{Q}$ must be $\mathbb{Q}(i)$. Since the degree of this field extension is 2, we conclude that the Galois group must be $\mathbb{Z}/2\mathbb{Z}$.

If $n_5 = 1$, then we are done. So assume $n_5 = 16$. By the prime factorization of $|G|$, each Sylow 5-subgroup of G must be of order 5, and consequently, must intersect trivially (i.e., if P, Q are distinct Sylow 5-subgroups, then $P \cap Q = \{1\}$). Each Sylow 5-subgroup contains 4 non-identity elements. This means that G contains 64 elements whose order is exactly 5. This leaves exactly 15 elements of G to have an order of the form 2^α . Since each Sylow 2-subgroup contains exactly 15 nonidentity elements of G with order equal to some power of 2, all of the aforementioned elements must be contained in exactly one Sylow 2-subgroup (i.e., $n_2 = 1$). Thus, $n_2 = 1$, and so this subgroup is normal in G . In any case, G contains a normal subgroup.

Problem 2017-J-II-6. Without using the classification of simple groups, show that every simple group of order 60 has exactly 10 subgroups of order 3. Hint: Consider the action of this group on subgroups of order 3 by conjugation.

Let G denote a simple group of order $60 = 2^2 \cdot 3 \cdot 5$, and let n_2, n_3, n_5 denote the number of Sylow 2-, 3-, and 5-subgroups. By Sylow's Theorems, one has $n_3 \in \{1, 2, 4, 5, 10, 20\} \cap \{1, 4, 7, 10, \dots, 19, 22, \dots\} = \{1, 4, 10\}$. Moreover, it follows that every subgroup of G of order 3 is precisely a Sylow 3-subgroup, and every Sylow-3 subgroup of G is a subgroup of order 3. Since G is simple, it has no normal subgroups, and therefore, $n_3 \neq 1$. Assume to the contrary that G has exactly 4 subgroups of order 3. Then the group action of G on the set $\text{Syl}_3(G)$ of Sylow 3-subgroups by conjugation induces a homomorphism $G \rightarrow S_4$. If the kernel of this action were trivial, then this homomorphism is injective. I.e., G is isomorphic to a subgroup of S_4 . But $|S_4| = 24$, while $|G| = 60 > 24$, which is a contradiction. Thus, the kernel must be nontrivial. If the kernel is all of G , then the action is trivial, which forces $n_3 = 1$ (a contradiction). Thus we conclude that $n_3 = 10$.

Problem 2016-J-I-1. Let G be a finite group of order 80. Prove that G is not simple.

Let G be a group order $80 = 2^4 \cdot 5$. Let n_2 and n_5 denote the number of Sylow 2- and Sylow 5-subgroups of G , respectively. By Sylow's Theorems, n_5 must divide 2^4 and be of the form $\equiv 1 \pmod{5}$. Thus, we must have

$$n_5 \in \{1, 2, 4, 8, 16\} \cap \{1, 6, 11, 16, \dots\} = \{1, 16\}.$$

If $n_5 = 1$, then we are done. So assume $n_5 = 16$. By the prime factorization of $|G|$, each Sylow 5-subgroup of G must be of order 5, and consequently, must intersect trivially (i.e., if P, Q are distinct Sylow 5-subgroups, then $P \cap Q = \{1\}$). Each Sylow 5-subgroup contains 4 non-identity elements. This means that G contains 64 elements whose order is exactly 5. This leaves exactly 15 elements of G to have an order of the form 2^α . Since each Sylow 2-subgroup contains exactly 15 nonidentity elements of G with order equal to some power of 2, all of the aforementioned elements must be contained in exactly one Sylow 2-subgroup (i.e., $n_2 = 1$). Thus, $n_2 = 1$, and so this subgroup is normal in G . In any case, G contains a normal subgroup, and hence is not simple.

Problem 2013-J-I-2. Prove that every group of order 72 is non-simple.

Let G denote a group of order $72 = 2^3 \cdot 3^2$, and let n_2, n_3 denote the number of Sylow 2- and 3-subgroups, respectively. By Sylow's Theorems,

$$n_3 \in \{1, 2, 4, 8\} \cap \{1, 4, 7, \dots\} = \{1, 4\}.$$

If $n_3 = 1$, then we are done since a unique Sylow subgroup must necessarily be normal. So assume $n_3 = 4$. Let G act on the set $\text{Syl}_3(G)$ of Sylow 3-subgroups by conjugation. Then this action induces a homomorphism $\varphi : G \rightarrow S_4$. Let K denote the kernel of this homomorphism. If $K = \{e\}$, then $\varphi : G \hookrightarrow S_4$ is an embedding, which is impossible since $|G| = 72 > |S_4| = 24$. So K is nontrivial. If $K = G$, then the group action is trivial, which means that for every $g \in G$ and every $P \in \text{Syl}_3(G)$, $gPg^{-1} = P$. But since every Sylow 3-subgroup has to be conjugate of each other, this constraint forces $n_3 = 1$, contradicting our assumption that $n_3 = 4$. This, $\ker \varphi$ is a nontrivial, proper normal subgroup of G which means that G cannot be simple.

Problem 2009-J-I-5. Let G be a group of order 63. Prove that G is not simple (i.e., that G contains a nontrivial normal subgroup).

Let G be a group of order $63 = 3^2 \cdot 7$, and let n_7 denote the number of distinct Sylow 7-subgroups of G . By Sylow's Theorems, $n_7 \in \{1, 3, 9\} \cap \{1, 8, 15, \dots\} = \{1\}$. Thus since there exists a single Sylow 7-subgroup, by Sylow's Theorems, this subgroup must be normal. Thus since this subgroup has order $0 < 7 < 63$, we conclude that G contains a nontrivial normal subgroup, and hence cannot be simple.

Problem 2006-A-II-5. Show that a group of order 55 is not simple.

Let G denote a group of order $55 = 5 \cdot 11$, and let n_{11} denote the number of distinct Sylow 11-subgroups of G . By Sylow's theorems $n_{11} \in \{1, 5\} \cap \{1, 12, \dots\} = \{1\}$ so that G contains a single Sylow-11 subgroup. By Sylow's theorems, this subgroup must be normal. Since this subgroup has order $0 < 11 < 55$, we conclude that G contains a nontrivial normal subgroup, and hence is not simple.

Problem 2001-J-I-3. Let G be a finite group and $P \subset G$ a normal Sylow p -subgroup. Show that for any group homomorphism $\varphi : G \rightarrow G$, $\varphi(P) \subset P$.

Let G be a finite group, and $P \subset G$ a normal Sylow p -subgroup. This necessarily implies that P is the unique Sylow p -subgroup of G . Therefore, every element of G with order equal to some power of p must be contained in P . Now let $\varphi : G \rightarrow G$ be a group homomorphism. Then for any $x \in P$, $|\varphi(x)| \mid |x| = p^{\alpha_x}$ for some positive integer α_x . So since $\varphi(x)$ has an order equal to some power of p , $\varphi(x) \in P$. Therefore, since $x \in P$ was arbitrary, we must have $\varphi(P) \subset P$.

Problem 1999-A-I-1. Let P be a Sylow p -subgroup of G , and let H be a p -subgroup of G contained in $N(P)$. Here, $N(P)$ is the normalizer of P in G . Prove H is contained in P .

Let P be a Sylow p -subgroup of G , and let H be a p -subgroup of G contained in $N(P)$, the normalizer of P in G . First, this implies that PH is a subgroup of G of order

$$|PH| = \frac{|P||H|}{|P \cap H|}.$$

In particular, it follows that PH is also a p -subgroup of G . Assume to the contrary that H is not entirely contained in P . This means that $|H|/|P \cap H| > 1$. But then $|PH| > |P|$. But this contradicts the fact that P is a Sylow p -subgroup of G , and therefore has maximal order amongst p -subgroups of G . Thus, by contradiction, H is entirely contained in P .

Problem D&F-4.5-13. Prove that a group of order 56 has a normal Sylow p -subgroup for some prime p dividing its order.

Let G denote a group of order $56 = 2^3 \cdot 7$, and let n_2, n_7 denote the number of Sylow 2- and 7-subgroups. By Sylow's Theorems, $n_7 \in \{1, 2, 4, 8\} \cap \{1, 8, \dots\} = \{1, 8\}$. Likewise, $n_2 \in \{1, 7\} \cap \{1, 3, 5, 7, \dots\} = \{1, 7\}$. If $n_7 = 1$, we are done since by Sylow's Theorems, a unique Sylow subgroup must necessarily be normal. So suppose $n_7 = 8$. Since each Sylow 7-subgroup has prime order 7, each pair of distinct Sylow 7-subgroups intersect trivially. This means that G contains 48 elements of order 7. If $n_2 = 7$, then each pair of distinct Sylow 2-subgroups can share at most 4 elements, so that G contains at least $8 + 6(4) = 32$ elements whose order is of the form 2^α for some positive integer α . But this means that G contains at least $32 + 48 + 1 = 81 > 56$ elements, which is a contradiction. Thus, by contradiction, $n_2 = 1$. In this case, G contains a normal Sylow 2-subgroup. Since this subgroup is of order $8 < 56$, this subgroup must be proper. The proof concludes.

Problem D&F-4.5-14. Prove that a group of order 312 has a normal Sylow p -subgroup for some prime p dividing its order.

Let G denote a group of order $312 = 2^3 \cdot 3 \cdot 13$, and let n_{13} denote the number of Sylow 13-subgroups. By Sylow's Theorems,

$$n_{13} \in \{1, 2, 3, 4, 6, 8, 12, 24\} \cap \{1, 14, 27, \dots\} = \{1\}.$$

Thus G contains a unique Sylow 13-subgroup, which must then be normal in G .

Problem D&F-4.5-15. Prove that a group of order 351 has a normal Sylow p -subgroup for some prime p dividing its order.

Let G denote a group of order $351 = 3^3 \cdot 13$, and let n_3, n_{13} denote the number of Sylow 3- and 13-subgroups in G . By Sylow's Theorems,

$$n_3 \in \{1, 13\} \cap \{1, 4, \dots\} = \{1, 13\}, \quad n_{13} \in \{1, 27\}.$$

If n_{13} or n_3 is equal to one, then we are done. So assume $n_{13} = 27$ and $n_3 = 13$. Since each Sylow 13-subgroup is of prime order 13, each distinct Sylow 13-subgroup intersects trivially so that G contains exactly $12(27) = 324$ elements of order 13. On the other hand, each pair of distinct Sylow 3-subgroups may share at most 9 elements, so that the two subgroups alone contain at least 45 unique elements whose order is some power of 3. But this is impossible since $324 + 45 > |G|$. Thus G must contain either a normal Sylow 13-subgroup or a normal Sylow 3-subgroup.

Problem D&F-4.5-17. Prove that if $|G| = 105$ then G has a normal Sylow 5-subgroup and a normal Sylow 7-subgroup.

Let G denote a group of order $105 = 3 \cdot 5 \cdot 7$, and let n_3, n_5, n_7 denote the number of Sylow 3-, 5-, and 7-subgroups, respectively. By Sylow's Theorems,

$$\begin{aligned} n_3 &\in \{1, 5, 7, 35\} \cap \{1, 4, 7, \dots\} = \{1, 7\}. \\ n_5 &\in \{1, 3, 7, 21\} \cap \{1, 6, 11, 16, 21, \dots\} = \{1, 21\}. \\ n_7 &\in \{1, 3, 5, 15\} \cap \{1, 8, 15, \dots\} = \{1, 15\}. \end{aligned}$$

First, assume to the contrary that G has 15 Sylow 7-subgroups. Since each Sylow 7-subgroup has prime order, each pair of distinct Sylow 7-subgroups intersects trivially so that G contains exactly 90 elements of order 7. Since Sylow p -subgroups intersect trivially with Sylow q -subgroups for $p \neq q$, this leaves exactly 14 elements of G to have an order of either 3 or 5. This is a contradiction: for each of the possible values of (n_3, n_5) we calculate the total number of elements of G with order either 3 or 5 to be:

(n_3, n_5)	(# Elements with Order 3, # Elements with Order 5)	Total
(1, 1)	(2, 4)	$2 + 4 = 6$
(1, 21)	(2, 84)	$2 + 84 = 86$
(7, 1)	(14, 4)	$14 + 4 = 18$
(7, 21)	(14, 84)	$14 + 84 = 98$

Thus, we must have $n_7 = 1$. Likewise, assume $n_5 = 21$. This means that G contains exactly 84 elements of order 5. Therefore, G must contain exactly 20 elements of order either 3 or 7. Since G already is known to have exactly one Sylow 7-subgroup (and thus 6 elements of order 7), G must have exactly 14 elements of order 3. Altogether, we've identified the order of every element of G : 14 elements of order 3, 6 elements of order 7, 84 elements of order 5, and the identity. This means that there exist no elements of G of composite orders. Denote by P the Sylow 7-subgroup and Q a Sylow 5-subgroup. Since $Q \leq N_G(P)$, PQ is a subgroup of G of order,

$$|PQ| = \frac{|P||Q|}{|P \cap Q|} = 7 \cdot 5 = 35.$$

Since every group of order 35 is cyclic², it follows that $PQ = \langle x \rangle$, where $x \in G$ is an element of order 35. But this is a contradiction. Thus, we must have $n_5 = 1$. The proof concludes.

²Use Sylow's Theorems again to quickly prove this!

Problem D&F-4.5-32. Let P be a Sylow p -subgroup of H and let H be a subgroup of K . If $P \trianglelefteq H$ and $H \trianglelefteq K$, prove that P is normal in K . Deduce that if $P \in \text{Syl}_p(G)$ and $H = N_G(P)$, then $N_G(H) = H$ (in words: *normalizers of Sylow p -subgroups are self-normalizing*).

Let G be a finite group, and $P \trianglelefteq H \trianglelefteq K \leq G$. Our first goal is to show that $kPk^{-1} = P$ for all $k \in K$. Since H is a normal subgroup of K , we must have that $kPk^{-1} \leq H$ for all k at the very least. But kPk^{-1} is another Sylow p -subgroup of H . Since P is a normal Sylow p -subgroup of H , it must be the unique Sylow p -subgroup so that $P = kPk^{-1}$ for all k . Thus, P is normal in K .

Now assume that $P \in \text{Syl}_p(G)$ and $H = N_G(P)$. By definition, we have $P \trianglelefteq H \trianglelefteq N_G(H) \leq G$. By our above work, one must have $P \trianglelefteq N_G(H)$. But by the definition of a normalizer of a subgroup, this constraint forces $N_G(H) \leq H$. Thus, we must have $N_G(H) = H$.

Problem D&F-4.5-33. Let P be a normal Sylow p -subgroup of G and let H be any subgroup of G . Prove that $P \cap H$ is the unique Sylow p -subgroup of H .

Let P be a normal (i.e., unique) Sylow p -subgroup of G , and let H be a fixed arbitrary subgroup of G . First, it is clear that $P \cap H$ is a p -subgroup of H : since $P \cap H \leq P$, $|P \cap H| \mid |P| = p^\alpha$, where α is some positive integer so that p^α is the largest power of p dividing $|G|$, so that $|P \cap H|$ is also a power of p . Let $|P \cap H| = p^\beta$. First, we must show that $P \cap H$ is a Sylow p -subgroup of H . Suppose p^γ is the largest power of p that divides $|H|$, where $\gamma \not\geq \beta$. Then since $H \leq N_G(P)$, PH is a subgroup of G of order

$$|PH| = \frac{|P||H|}{|P \cap H|} = p^{\alpha+\gamma-\beta} m > p^\alpha = |P|,$$

which contradicts our hypothesis that P is the Sylow p -subgroup of G . Thus, we must have $\gamma = \beta$. So since $P \cap H$ is a subgroup of H with order $p^\gamma = p^\beta$, it must be a Sylow p -subgroup of H . Now, we will show that it is normal in H ; i.e., we wish to show that for any $h \in H$, $h(P \cap H)h^{-1} = P \cap H$. One direction is trivial: since P is normal in G , for any $h \in H$, $hPh^{-1} = P$, and $hHh^{-1} = H$ so that $h(P \cap H)h^{-1} \leq P \cap H$. Now let $g \in P \cap H$, and choose $g' = h^{-1}gh$ for some fixed $h \in H$. Then $g' \in h^{-1}(P \cap H)h$. By the previous inclusion, it follows that $g' \in P \cap H$. Thus, since $g = hg'h^{-1} \in h(P \cap H)h^{-1}$, $P \cap H \leq h(P \cap H)h^{-1}$. Since $h \in H$ was arbitrary, this concludes the proof that $P \cap H$ is normal in H . Therefore, $P \cap H$ is the unique Sylow p -subgroup of H .

Problem D&F-4.5-34. Let $P \in \text{Syl}_p(G)$ and assume $N \trianglelefteq G$. Use the conjugacy part of Sylow's Theorem to prove that $P \cap N$ is a Sylow p -subgroup of N . Deduce that PN/N is a Sylow p -subgroup of G/N (note that this may also be done by the Second Isomorphism Theorem — cf. Exercise 9, Section 3.3).

Let G be a finite group of order $p^\alpha M$, where $p \nmid M$ (so that p^α is the largest power of the prime integer p that divides $|G|$), $P \in \text{Syl}_p(G)$ a Sylow p -subgroup, and N a normal subgroup of G . Assume $|N| = p^\gamma m$, where $p \nmid m$ for some nonnegative integer γ . Consider the subgroup $P \cap N$. Since $P \cap N \leq P$, it follows that $P \cap N$ is a p -subgroup of N . Suppose $|P \cap N| = p^\beta$ for some positive integer β , and assume to the contrary that $\gamma > \beta$. Then since PN is a subgroup of G (because N is normal in G), $|PN| = |P||N|/|P \cap N| = p^{\alpha+\gamma-\beta} m > p^\alpha m > p^\alpha = |P|$. But this is a contradiction since by virtue of P being a Sylow p -subgroup of G , p^α is the largest power of p that divides $|G|$. Thus, $\gamma = \beta$, and we have $P \cap N$ is a Sylow p -subgroup of N . Now, by the Second Isomorphism Theorem, one has that $G/N \geq PN/N \cong P/(P \cap N)$. By our work from above, since $P \cap N$ is a p -subgroup as is P , $P/(P \cap N)$ is a Sylow p -subgroup of G/N . In particular, $|G/N| = p^\alpha M/p^\gamma m = p^{\alpha-\gamma}(M/m)$, where $M/m \in \mathbb{N}$ by Lagrange's Theorem and $p^{\alpha-\gamma} \nmid (M/m)$. Thus, $p^{\alpha-\gamma}$ is the greatest power of p that divides $|G/N|$. But $|PN/N| = p^\alpha m/p^\gamma m = p^{\alpha-\gamma}$. Thus, PN/N must be a Sylow p -subgroup of G/N .

Problem D&F-4.5-36. Prove that if N is a normal subgroup of G then $n_p(G/N) \leq n_p(G)$.

Let N be a normal subgroup of G , so that the quotient G/N is well-defined. Let G have the prime factorization,

$$G = p_1^{\alpha_1} \cdots p_m^{\alpha_m}.$$

Likewise, N has the prime factorization,

$$N = p^{\tilde{\alpha}} p_1^{\tilde{\alpha}_1} \cdots p_m^{\tilde{\alpha}_m},$$

where $0 \leq \tilde{\alpha}, \tilde{\alpha}_j \leq \alpha, \alpha_j$ for each $1 \leq j \leq m$. Note that since $N \leq G$, by Lagrange's Theorem, the prime factorization of N cannot contain any primes not contained in the prime factorization of G . **!!! Complete Later !!!**

3.6 Classification of Groups

Problem 2024-J-I. For distinct odd primes p and q , prove that every finite group of order $2pq$ is a semidirect product of a normal subgroup of order pq and a subgroup of order 2.

Let p, q be distinct odd primes, and let G be a group of order $2pq$. Without loss of generality, assume that $p < q$. **!! Complete Later !!**

Problem 2023-J-II-6. Classify (up to isomorphism) all groups of order 20.

Let G be a group of order $20 = 2^2 \cdot 5$. By the Sylow Theorems, G contains exactly one Sylow 5-subgroup H so that H is normal in G . Let K be a Sylow 2-subgroup of G (with order 4). By Lagrange's theorem, $H \cap K = \{1\}$, and $HK = G$ since

$$|HK| = \frac{|H||K|}{|H \cap K|} = |H||K| = |G|.$$

Therefore, it follows that $G \cong H \rtimes_{\varphi} K$ for some group homomorphism $\varphi : K \rightarrow \text{Aut}(H)$. Since H has prime order, it is a cyclic group; in particular, $H \cong \mathbb{Z}_5$. Therefore, $\text{Aut}(H) \cong (\mathbb{Z}_5)^{\times} \cong \mathbb{Z}_4$. On the other hand, by the classification of groups of order 4, $K \cong \mathbb{Z}_4$ or $K \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. We will consider each case independently.

(Case 1) Assume that $K \cong \mathbb{Z}_4$. Each homomorphism $\varphi : K \rightarrow \text{Aut}(H) \cong \mathbb{Z}_4$ is uniquely determined by $\varphi(1)$, with the caveat that the order of $\varphi(1)$ divides the order of 1. Since the order of 1 is 4 in $\mathbb{Z}_4$, $\varphi(1) \in \{0, 1, 2, 3\}$, making for four total group homomorphisms.

- (i) Consider the homomorphism $\varphi_0 : 1 \mapsto 0$. Then since φ_0 maps every element of K to the element $0 \in \mathbb{Z}_4$, one must have that φ_0 is the trivial homomorphism. Therefore, $H \rtimes_{\varphi_0} K$ is isomorphic to the abelian direct product $H \times K \cong \mathbb{Z}_5 \times \mathbb{Z}_4 \cong \mathbb{Z}_{20}$, with the final isomorphism following from the Chinese Remainder Theorem.
- (ii) Consider the map $\alpha : \mathbb{Z}_4 \rightarrow \mathbb{Z}_4$ defined by $\alpha(x) = 3x$. Then α is easily seen to be an automorphism on $\mathbb{Z}_4$ with $\alpha : 1 \mapsto 3, 3 \mapsto 1$. This means that if $\varphi_1 : K \rightarrow \text{Aut}(H)$ and $\varphi_3 : K \rightarrow \text{Aut}(H)$ are the maps mapping $1 \mapsto 1$ and $1 \mapsto 3$, respectively, then $\varphi_3 = \varphi_1 \circ \alpha$. Since these two homomorphisms differ by an automorphism on the domain group K , we conclude that the nonabelian semidirect product groups $\mathbb{Z}_5 \rtimes_{\varphi_1} \mathbb{Z}_4$ and $\mathbb{Z}_5 \rtimes_{\varphi_3} \mathbb{Z}_4$ are isomorphic.
- (iii) Now consider the homomorphism $\varphi_2 : 1 \mapsto 2$. Since the order of 2 in $\mathbb{Z}_4$ is 2, there cannot exist any automorphisms on $\mathbb{Z}_4$ that maps 2 to any other element of $\mathbb{Z}_4$. Moreover, the image of φ_2 has order 2, while none of the other homomorphisms have image with order 2. Thus, the nonabelian semidirect product group $H \rtimes_{\varphi_2} K$ cannot be isomorphic to $\mathbb{Z}_{20}$ nor to the nonabelian group $\mathbb{Z}_5 \rtimes_{\varphi_1} \mathbb{Z}_4$.

Therefore, for the case with $K \cong \mathbb{Z}_4$, we have found three distinct isomorphism classes of groups of order 20, one of which is abelian.

(Case 2) Assume that $K \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Each homomorphism $\varphi : K \rightarrow \text{Aut}(H) \cong \mathbb{Z}_4$ is uniquely determined by $\varphi((1, 0))$ and $\varphi((0, 1))$ (where $(1, 0), (0, 1)$ are the two generators of $\mathbb{Z}_2 \times \mathbb{Z}_2$, with the caveat that the order of $\varphi(\cdot)$ divides the order of $\cdot$). Since the order of $(1, 0)$ and $(0, 1)$ is 2, $\varphi((1, 0)), \varphi((0, 1)) \in \{0, 2\}$, making for four total group homomorphisms.

- (i) Consider the homomorphism $\varphi_0 : (0, 1), (1, 0) \mapsto 0$. Then φ_0 is the trivial homomorphism. Thus, $\mathbb{Z}_5 \rtimes_{\varphi_0} (\mathbb{Z}_2 \times \mathbb{Z}_2)$ is isomorphic to the abelian direct product $\mathbb{Z}_{10} \times \mathbb{Z}_2$.
- (ii) On the other hand, since $\text{Aut}(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}) \cong GL_2(\mathbb{F}_2)$ acts transitively on the nonzero elements of K , all three of the remaining homomorphisms differ only by a precomposition by an automorphism of K . Thus, the nonabelian semidirect product groups corresponding to each such homomorphism are isomorphic to each other.

Altogether, we conclude that there are five distinct isomorphism classes of groups of order 20, 2 of which are abelian.

Problem 2010-J-II-5. Classify (up to isomorphism) all groups of order 45.

Let G denote a group of order $45 = 3^2 \cdot 5$. By the Sylow theorems, if n_3 and n_5 denote the number of Sylow 3- and Sylow 5-subgroups, then

$$n_3 \in \{1, 5\} \cap \{1, 4, 7, \dots\} = \{1\}.$$

$$n_5 \in \{1, 3\} \cap \{1, 6, 11, \dots\} = \{1\}.$$

Thus G has a normal Sylow 3-subgroup H and a normal Sylow 5-subgroup K . Since by Lagrange's theorem, $H \cap K = \{1\}$ and $HK = G$, it follows that G is isomorphic to the abelian direct product $H \times K$. Since K has prime order 5, $K \cong \mathbb{Z}_5$. On the other hand, by classification of groups of order p^2 , $H \cong \mathbb{Z}_9$ or $\mathbb{Z}_3 \times \mathbb{Z}_3$. Thus, we conclude that (up to isomorphism) there are precisely 2 groups of order 45: (1) $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_5 \cong \mathbb{Z}_3 \times \mathbb{Z}_{15}$ (by the Chinese Remainder Theorem), and (2) $\mathbb{Z}_9 \times \mathbb{Z}_5 \cong \mathbb{Z}_{45}$ (by the Chinese Remainder Theorem).

Problem 2008-J-I-3. Classify (up to isomorphism) all groups of order 28.

Let G denote a group of order $28 = 2^2 \cdot 7$. By Sylow's theorems, G must contain a normal Sylow 7-subgroup N of order 7. Let K denote a Sylow 2-subgroup of order 4. By Lagrange's theorem, $N \cap K = \{1\}$. Therefore, since

$$|NK| = |N||K|/|N \cap K| = |N||K| = |G|,$$

one has $NK = G$. Hence, we must have $G \cong N \rtimes_{\varphi} K$ for some group homomorphism $\varphi : K \rightarrow \text{Aut}(N)$. Since N has prime order, it is cyclic of order 7 so that $\text{Aut}(N) \cong (\mathbb{Z}/7\mathbb{Z})^{\times} \cong \mathbb{Z}/6\mathbb{Z}$. On the other hand, by the classification of groups of order 4, either $K \cong \mathbb{Z}_4$ or $K \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Therefore, we will consider two cases.

(Case 1) Suppose $K \cong \mathbb{Z}_4 = \langle 1 \rangle$. Any homomorphism $\varphi \in \text{Hom}_{\text{Grp}}(K, \text{Aut}(N))$ is uniquely determined by $\varphi(1)$ under the caveat that the order of $\varphi(1) \in \mathbb{Z}_6 \cong \text{Aut}(N)$ divides the order of $1 \in \mathbb{Z}_4$ (which is 4). Therefore, $\varphi(1) \in \{0, 3\}$. The homomorphism $\varphi_0 : 1 \mapsto 0$ is the trivial homomorphism, and corresponds to the abelian direct product $G \cong N \times K \cong \mathbb{Z}_7 \times \mathbb{Z}_4 \cong \mathbb{Z}_{28}$ (with the last isomorphism following from the Chinese Remainder Theorem). On the other hand, the nontrivial isomorphism $\varphi_1 : 1 \mapsto 3$ corresponds to a nonabelian semidirect product group $N \rtimes_{\varphi_1} K$, which has the group presentation

$$\langle x, y \mid x^7 = x^4 = 1, yxy^{-1} = x^{-1} \rangle.$$

(Case 2) Suppose $K \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, with the two generators $(0, 1)$ and $(1, 0)$. Each of these have order 2 in K . Thus any automorphism $\varphi : K \rightarrow \mathbb{Z}/6\mathbb{Z}$ is uniquely determined by $\varphi((0, 1))$ and $\varphi((1, 0))$ with the caveat that the images of these generators divide 2. Hence, $\varphi((0, 1)), \varphi((1, 0)) \in \{0, 3\}$. If φ_0 maps both generators to 0, then φ_0 is the trivial homomorphism and corresponds to the abelian direct product $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7 \cong \mathbb{Z}_2 \times \mathbb{Z}_{14}$ (by the Chinese Remainder Theorem). On the other hand, since $\text{Aut}(\mathbb{Z}_2 \times \mathbb{Z}_2) \cong \text{GL}_2(\mathbb{F}_2)$ acts transitively on the nonzero elements of K , all three of the remaining (nontrivial) homomorphisms differ by only a precomposition by an automorphism of K , and thus the nonabelian semidirect product groups corresponding to these homomorphisms are all isomorphic, and has the group presentation

$$\langle x, y, z \mid x^7 = y^2 = z^2 = 1, yz = zy, yxy^{-1} = x^{-1}, zyz^{-1} = z \rangle.$$

Thus altogether, we conclude that up to isomorphism, there exist precisely 4 groups of order 28, two which are abelian.

Problem 2004-A-I-6. Let G and H be groups of order 42 and 55, respectively. Show that the only group homomorphism from G to H is the constant homomorphism.

Let G and H be groups of order 42 and 55, respectively. Let $\varphi : G \rightarrow H$ be a group homomorphism, and let $g \in G$ be an arbitrary element. By Lagrange's theorem,

$$|g| \in \{1, 2, 3, 6, 7, 14, 21, 42\} =: A.$$

On the other hand, one must have $|\varphi(g)| \mid |g|$ and

$$|\varphi(g)| \in \{1, 5, 11, 55\} =: B.$$

But the non-identity elements of B are coprime to the non-identity elements in A so that $|\varphi(g)| = 1$. Therefore, for every $g \in G$, $\varphi(g) = 1$, which means that φ is the constant (i.e., trivial) homomorphism.

Problem 2003-A-II-6. Show that Σ_7 , the group of permutations of 7 elements, does not contain any subgroup with 15 elements.

Let Σ_7 denote the group of permutations of 7 elements, and assume to the contrary that Σ_7 contains a subgroup H with $15 = 3 \cdot 5$ elements. By Sylow's theorems, H must contain a unique Sylow 5-subgroup of order 5, and a unique Sylow 3-subgroup of order 3. Thus, H contains precisely $15 - (1 + 4 + 2) = 8$ elements of order 15. In particular, Σ_7 must contain some element τ whose order is 15. Let τ be expanded as the product $\alpha_1 \alpha_2 \cdots \alpha_n$ of disjoint cycles. Then since Σ_7 is the permutation group on 7 letters, $\sum_1^n \ell_i = 7$, where for each i , ℓ_i denotes the length of α_i . On the other hand, since $|\tau| = 15$, we must have $\text{lcm}_i \alpha_i = 15$. Looking at the partitions of 7, we find there is no disjoint cycle decomposition of τ that satisfies both criteria. This suggests that τ cannot have order 15, which is a contradiction! Thus, by contradiction, Σ_7 cannot contain a subgroup of order 15.

Problem 2003-J-I-6.

- (a) Prove that a group of order p^2 , p a prime number, is abelian.
 (b) Classify groups of order p^2 up to isomorphism.

- (a) Let G be a group of order p^2 , where p is prime, and let $Z(G)$ denote the center of G . By Lagrange's theorem, $|Z(G)| \in \{1, p, p^2\}$. If $|Z(G)| = p^2$, then we are done. Now assume to the contrary that $|Z(G)| = p$. Then $|G/Z(G)| = |G|/|Z(G)| = p^2/p = p$. By classification of prime order groups, it follows that $G/Z(G)$ is cyclic, and therefore, G is abelian. Now suppose $|Z(G)| = 1$. By the class equation, one has

$$|G| = |Z(G)| + \sum_{i=1} [G : C_G(g_i)].$$

Each $[G : C_G(g_i)]$ must be a multiple of p . Thus taking the modulus of both sides of the above equation with respect to p , one obtains $0 \equiv 1 \pmod{p}$, which is a contradiction. Thus by contradiction, we conclude that $|Z(G)| = p^2$, and so G is abelian.

- (b) Now we know that every nonidentity element of G must have order p or p^2 . If G contains an element y of order p^2 , then since $|\langle y \rangle| = |G|$, G must be a cyclic abelian group of order p^2 ; i.e., $G \cong \mathbb{Z}_{p^2}$. Now suppose every nonidentity element of G has order exactly p . Then let $x \in G$ be an element of order p , and let $y \in G \setminus \langle x \rangle$. Then $|\langle y \rangle| = p$ and $\langle x \rangle \cap \langle y \rangle = 1$. Thus since $\langle x \rangle \langle y \rangle = G$, one has that $G \cong \langle x \rangle \times \langle y \rangle \cong \mathbb{Z}_p \times \mathbb{Z}_p$.

Problem D&F-5.2-1. In each of parts (a) to (e) give the number of nonisomorphic abelian groups of the specified order — do not list the groups: (a) order 100, (b) order 576, (c) order 1155, (d) order 42875, (e) order 2704.

- (a) We begin by noting that $100 = 2^2 \cdot 5^2$. The number of partitions of 2 is 2. Thus, we conclude that there are exactly 4 nonisomorphic abelian groups of order 100.
 (b) We begin by noting that $576 = 2^6 \cdot 3^2$. As noted above, the number of partitions of 2 is 2, and now, the number of partitions of 6 is 11. Thus, the total number of nonisomorphic abelian groups of order 576 is 22.

- (c) We begin by noting that $1155 = 3 \cdot 5 \cdot 7 \cdot 11$. Since 1155 is a squarefree integer, we conclude that there exists exactly one nonisomorphic abelian group of the specified order.
- (d) We start by noting that $42875 = 5^3 \cdot 7^3$. The number of partitions of 3 is 3. Thus, there exist exactly 9 nonisomorphic abelian groups of order 42875.
- (e) We start by noting that $2704 = 2^4 \cdot 13^2$. The number of partitions of 4 is 5. Thus, there exist exactly 10 nonisomorphic abelian groups of order 2704.

3.7 Module Theory

Problem 2025-J-I-5. Let R be a nonzero commutative ring with multiplicative identity. Let m and n be distinct positive integers. Prove that R^n and R^m , with their natural structure of R -modules, are not isomorphic R -modules.

Let R be a nonzero commutative ring with multiplicative identity, and assume that $R^m \cong R^n$. Let I denote a maximal ideal of R , and consider the quotient $F = R/I$, which must be a field since I is maximal. Then since $R^m \cong R^n$, tensoring on the left by F gives the isomorphism $F \otimes_R R^m \cong F \otimes_R R^n$. But as R -modules, for any k , $F \otimes_R R^k \cong F^k$. Thus we have $F^m \cong F^n$ as vector spaces over F . This means that $m = n$. Therefore, by the contrapositive, if $n \neq m$, then $R^m \not\cong R^n$.

Problem 2025-J-II-3 (Modified) . Let V be a vector space of dimension n over $\mathbb{Q}$. Let $T : V \rightarrow V$ be a linear transformation with minimal polynomial $x^4 - x - 2$ over $\mathbb{Q}$. Show that n must be even.

Let V be a vector space of dimension n over $\mathbb{Q}$, and let $T : V \rightarrow V$ be a linear transformation with minimal polynomial $x^4 - x - 2$ over $\mathbb{Q}$. We first note that V is a finitely generated module over the PID $\mathbb{Q}[x]$, where x acts by the linear operator T . Thus by the Structure Theorem for Finitely Generated Modules over PIDs, we have the decomposition,

$$V \cong \bigoplus_{i=1}^N \frac{\mathbb{Q}[x]}{(\alpha_i(x))},$$

where each $\alpha_i(x)$ is an invariant factor of T , $\alpha_1(x) \mid \alpha_2(x) \mid \cdots \mid \alpha_N(x)$, and $\alpha_N(x)$ is the minimal polynomial of T . It is straightforward to show that $\alpha_N(x)$ is irreducible over $\mathbb{Q}$. This means that each nonconstant $\alpha_i(x)$ is an associate of $\alpha_N(x)$ (and thus has degree 4). The constant $\alpha_i(x)$ all have degree 0. Thus, since

$$n = \dim V = \sum_{i=1}^N \deg \alpha_i(x)$$

is divisible by 4 (and hence by 2), we conclude that V is an even-dimensional vector space.

Problem 2025-J-II-3. Let V be a vector space of dimension n over $\mathbb{Q}$. Let $T : V \rightarrow V$ be a linear transformation with minimal polynomial $x^4 - x^2 - 2$ over $\mathbb{Q}$. Show that n must be even.

Let V be a vector space of dimension n over $\mathbb{Q}$, and let $T : V \rightarrow V$ be a linear transformation with minimal polynomial $x^4 - x^2 - 2$. We first note that V is a finitely generated module over the PID $\mathbb{Q}[x]$, where x acts by the linear operator T . Thus by the Structure Theorem for Finitely Generated Modules over PIDs, we have the decomposition,

$$V \cong \bigoplus_{i=1}^N \frac{\mathbb{Q}[x]}{(\alpha_i(x))},$$

where each $\alpha_i(x)$ is an invariant factor of T , $\alpha_1(x) \mid \alpha_2(x) \mid \cdots \mid \alpha_N(x)$, and $\alpha_N(x)$ is the minimal polynomial of T . We note that $\alpha_N(x) = x^4 - x^2 - 2$ is reducible over $\mathbb{Q}$ as the product $(x^2 + 1)(x^2 - 2)$ of irreducible quadratics. Thus, since

$$n = \dim V = \sum_{i=1}^N \deg \alpha_i(x),$$

and the degree of each direct summand is even, we conclude that n is even.

Problem 2021-A-I-5. Let R be an integral domain with field of fractions Q . Show that $Q/R \otimes_R Q/R = 0$.

Let R be an integral domain with field of fractions Q . We note that every tensor in $Q/R \otimes_R Q/R$ is a finite sum of the tensors of the form,

$$\left(\frac{a}{b} + R\right) \otimes_R \left(\frac{c}{d} + R\right),$$

where $a, b, c, d \in R$ and $b, d \neq 0$. We compute,

$$\frac{a}{b} \otimes_R \frac{c}{d} = d \left(\frac{a}{db} \otimes_R \frac{c}{d} \right) = \frac{a}{db} \otimes_R c.$$

But since $c \in R$, its representative class in Q/R is the zero class. Thus, $(a/db) \otimes c = 0$. Since every such simple tensor is identically zero, we conclude that every tensor in $Q/R \otimes_R Q/R$ is identically zero so that $Q/R \otimes_R Q/R = 0$.

Problem 2005-J-II-4. Let $A : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be a linear map such that the characteristic polynomial of A is $\chi_A(t) = t^4 + 1$. How many A -invariant subspaces are there in $\mathbb{R}^4$? (A subspace W is called A -invariant if $AW \subset W$.)

Let $A : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be a linear map such that the characteristic polynomial of A is $\chi_A(t) = t^4 + 1$. Since $\mathbb{R}^4$ is a finitely generated vector space, we note that $\mathbb{R}^4$ can be viewed as a finitely generated module over $\mathbb{R}[x]$, where x acts by the linear operator A . Thus by the Structure Theorem for Finitely Generated Modules over PIDs, we have the decomposition,

$$V \cong \bigoplus_{i=1}^N \frac{\mathbb{R}[x]}{(\alpha_i(x))},$$

where each $\alpha_i(x)$ is an invariant factor of A , $\alpha_1(x) \mid \alpha_2(x) \mid \cdots \mid \alpha_N(x)$, and $\alpha_1(x)\alpha_2(x)\cdots\alpha_N(x) = \chi_A(x) = x^4 + 1$. It is immediately straightforward to see that $\chi_A(x)$ has no real roots so that it cannot be decomposed as a product containing a linear factor. So, we will check if $\chi_A(x)$ is equal to the product of two irreducible quadratics. Doing so, we conclude that

$$(x^4 + 1) = (x^2 + \sqrt{2}x + 1)(x^2 - \sqrt{2}x + 1).$$

Each A -invariant subspace of $\mathbb{R}^4$ corresponds to one of the direct summands above. Thus since $\chi_A(t)$ has four divisors $(1, x^2 \pm \sqrt{2}x + 1, \text{ and } x^4 + 1)$, we conclude that there are exactly 4 A -invariant subspaces of $\mathbb{R}^4$.

Problem D&F-10.5-1. Suppose that

$$\begin{array}{ccccc} A & \xrightarrow{\psi} & B & \xrightarrow{\varphi} & C \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ A' & \xrightarrow{\psi'} & B' & \xrightarrow{\varphi'} & C' \end{array}$$

is a commutative diagram of groups and that the rows are exact. Prove that

- (a) if φ and α are surjective, and β is injective then γ is injective.
- (b) if ψ' , α , and γ are injective, then β is injective.
- (c) If φ , α , and γ are surjective, then β is surjective.
- (d) if β is injective, α and γ are surjective, then γ is injective.

- (a) Let $c \in C$ so that $\gamma(c) = 0$. Since φ is surjective, there exists some $b \in B$ so that $\varphi(b) = c$. By commutativity of the diagram, $\varphi'(\beta(b)) = 0$ so that $\beta(b) \in \ker \varphi'$. By exactness of the rows, $\beta(b) \in \text{img } \psi'$ so that there exists $a' \in A'$ with $\psi'(a') = \beta(b)$. Since α is surjective, there exists $a \in A$ so that $\psi'(\alpha(a)) = \beta(b)$. By commutativity of the diagram, $\beta(\psi(a)) = \beta(b)$; by injectivity of β , $b = \psi(a)$. Since $b \in \text{img } \psi$, by exactness, $b \in \ker \varphi$. This means that $c = \varphi(b) = 0$. Hence, γ is injective.

- (b) Let $b \in B$ so that $\beta(b) = 0$. This means that $\varphi'(\beta(b)) = 0$; i.e., $\beta(b) \in \ker \varphi'$. By commutativity of the right square, $\gamma(\varphi(b)) = 0$; since γ is injective, $\varphi(b) = 0$. I.e., $b \in \ker \varphi$; by exactness, there exists $a \in A$ such that $\psi(a) = b$. Then by commutativity of the diagram, $\psi'(\alpha(a)) = \beta(b) = 0$; since ψ' is injective, $\alpha(a) = 0$. Since α is injective, $a = 0$. Therefore, $b = \psi(a) = 0$. Hence, this concludes the proof that β is injective.
- (c) Let $b' \in B'$. Since γ is surjective, there exists a $c \in C$ so that $\gamma(c) = \varphi'(b')$. Since φ is surjective, there exists a $b \in B$ such that $c = \varphi(b) \implies \gamma(\varphi(b)) = \varphi'(b')$. By commutativity of the diagram, $\varphi'(\beta(b)) = \varphi'(b')$. So consider the element $\beta(b) - b' \in B'$. By the previous observation, $\beta(b) - b' \in \ker \varphi'$ so that by exactness, there exists $a' \in A'$ with $\psi'(a') = \beta(b) - b'$. And by surjectiveness of α , there exists $a \in A$ with $\beta(b) - b' = \psi'(\alpha(a))$. By commutativity of the right square, $\psi'(\alpha(a)) = \beta(\psi(a))$ so that in fact, $\beta(b) - b' = \beta(\psi(a)) \implies b' = \beta(b - \psi(a))$, where $b - \psi(a) \in B$. Hence, β is surjective.
- (d) Let $c \in C$ so that $\gamma(c) = 0$.

Problem D&F-10.5-2. Suppose that

$$\begin{array}{ccccccc} A & \xrightarrow{\psi} & B & \xrightarrow{\varphi} & C & \xrightarrow{\phi} & D \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta \\ A' & \xrightarrow{\psi'} & B' & \xrightarrow{\varphi'} & C' & \xrightarrow{\phi'} & D' \end{array}$$

is a commutative diagram of groups and that the rows are exact. Prove that

- (a) if α is surjective, and β, δ are injective, then γ is injective.
(b) if δ is injective, and α, γ are surjective, then β is surjective.

- (a) Let α be surjective, and β, δ be injective. Let $c \in C$ such that $\gamma(c) = 0$. Then $\phi'(\gamma(c)) = 0$. By commutativity of the right square, it then follows that $\delta(\phi(c)) = 0$; since δ is injective, $\phi(c) = 0$, which means $c \in \ker \phi$. By exactness of the first row, $c \in \text{im } \varphi$. I.e., there exists some $b \in B$ so that $\varphi(b) = c$. But then, by commutativity of the diagram, $\varphi'(\beta(b)) = 0$ so that $\beta(b) \in \ker \varphi'$. By exactness of the second row, $\beta(b) \in \text{im } \psi'$, which means there exists $a' \in A'$ with $\psi'(a') = \beta(b)$. Since α is surjective, there exists $a \in A$ so that $\psi'(a') = \psi'(\alpha(a)) = \beta(b)$. By commutativity of the diagram, $\beta(\psi(a)) = \beta(b)$ so that $\psi(a) = b$ (by injectivity of β). This implies $b \in \text{im } \psi$. By exactness of the first row, $b \in \ker \varphi$. Hence, $c = \varphi(b) = 0$, which proves that γ is injective.
- (b) Let δ be injective, and α, γ be surjective. Let $b' \in B'$. By surjectivity of γ , there exists some $c \in C$ so that $\gamma(c) = \varphi'(b')$. By exactness of the second row, $\varphi'(b') \in \ker \phi'$. By commutativity of the diagram, $\delta(\phi(c)) = 0$. Since δ is injective, $\phi(c) = 0$. By exactness of the first row, there exists some $b \in B$ so that $c = \varphi(b)$. This implies $\gamma(\varphi(b)) = \varphi'(b')$. By commutativity of the middle square, $\varphi'(\beta(b) - b') = 0$ so that, by exactness, $\beta(b) - b' \in \text{im } \psi'$. Then by surjectivity of α , there exists some $a \in A$ so that $\psi'(\alpha(a)) = \beta(b) - b'$. By commutativity of the left square, $\beta(\psi(a)) = \psi'(\alpha(a)) = \beta(b) - b'$ so that $b' = \beta(b - \psi(a))$. Hence, since $b - \psi(a) \in B$, we have proven that β is surjective.

Problem D&F-10.5-3. Let P_1 and P_2 be R -modules. Prove that $P_1 \oplus P_2$ is a projective R -module if and only if both P_1 and P_2 are projective.

($\Leftarrow$) Let P_1 and P_2 be projective modules. Fix arbitrary left R -modules N, M and let $f : N \twoheadrightarrow M$ be a surjection. Let $g : P_1 \oplus P_2 \rightarrow M$ be a module homomorphism. Then by the universal property for direct sums, g is uniquely determined by the maps $(g_i)_{i=1,2}$, where for each i , $g_i = g \circ \iota_i : P_i \rightarrow M$ is a module homomorphism and $\iota_i : P_i \rightarrow P_1 \oplus P_2$ is the canonical inclusion $P_i \hookrightarrow P_1 \oplus P_2$. Since each P_i is a projective R -module, there exist module homomorphisms $h_i : P_i \rightarrow N$ such that $fh_i = g_i$. By the universal property of direct sums, there exists a unique module homomorphism $h : P_1 \oplus P_2 \rightarrow N$ such that for each i , $h \circ \iota_i = h_i$, so that in fact, $fh \circ \iota_i = g_i = g \circ \iota_i$ for each i . But by uniqueness from the universal property, it follows that $fh = g$. Hence, by definition, $P_1 \oplus P_2$ is projective.

($\Rightarrow$) Assume $P_1 \oplus P_2$ is projective. Again fix arbitrary left R -modules N and M , and let $f : N \rightarrow M$ be a surjective module homomorphism. Let $g_i : P_i \rightarrow M$ be a module homomorphism for each $i = 1, 2$. By the universal property for direct sums, $(g_i)_{i=1,2}$ determine a unique homomorphism $g : P_1 \oplus P_2 \rightarrow M$; by definition of a projective module, there exists a module homomorphism $h : P_1 \oplus P_2 \rightarrow N$ such that $fh = g$. But by the universal property for direct sums, h is uniquely determined by the family of maps $(h_i)_{i=1,2}$, where for each i , $h_i = h \circ \iota_i : P_i \rightarrow N$ (where $\iota_i : P_i \hookrightarrow P_1 \oplus P_2$ is the canonical inclusion) and $fh = g$. Then we compute $f(h \circ \iota_i) = (g \circ \iota_i) \implies fh_i = g_i$. Hence, by definition, each P_i is projective.

Problem D&F-10.5-4. Let Q_1 and Q_2 be R -modules. Prove that $Q_1 \oplus Q_2$ is an injective R -module if and only if both Q_1 and Q_2 are injective.

Let Q_1 and Q_2 be R -modules. Fix arbitrary R -modules A and B , and an injection $f : A \hookrightarrow B$. Assume that Q_1 and Q_2 are injective. Let $g : A \rightarrow Q_1 \oplus Q_2$ be a module homomorphism. By the universal property for direct sums, g is uniquely determined by the family of maps $(g_i)_{i \in I}$, where for each i , $g_i = \pi_i \circ g : A \rightarrow Q_i$ and $\pi_i : Q_1 \oplus Q_2 \rightarrow Q_i$ are the canonical projection maps. Since each Q_i is injective, there exist morphisms $h_i : B \rightarrow Q_i$ such that $h_i f = g_i$. But by the universal property for direct sums, the family of maps $(h_i)_{i \in I}$ uniquely determines a map $h : B \rightarrow Q_1 \oplus Q_2$ so that for each i , $h_i = \pi_i h$. But then we observe that

$$\pi_i g = g_i = h_i f = (\pi_i h) f = \pi_i(hf), \quad i = 1, 2.$$

By uniqueness from the universal property for direct sums, we conclude that $hf = g$, and so $Q_1 \oplus Q_2$ is injective. Now suppose $Q_1 \oplus Q_2$ are injective, and let $g_i : A \rightarrow Q_i$ be morphisms. By the universal property for direct sums, these maps uniquely determine a map $g : A \rightarrow Q_1 \oplus Q_2$ such that for each i , $g_i = \pi_i \circ g$. Since $Q_1 \oplus Q_2$ is injective, there exists a morphism $h : B \rightarrow Q_1 \oplus Q_2$ so that $hf = g$. But then, h is uniquely determined by maps $h_i : B \rightarrow Q_i$, where for each i , $h_i = \pi_i \circ h$. We then note that

$$h_i f = (\pi_i h) f = \pi_i(hf) = \pi_i g = g_i.$$

Hence, we conclude that Q_1 and Q_2 are injective.

Problem D&F-10.5-5. [!! Complete Later !!]

Problem D&F-10.5-6. Prove that the following are equivalent for a ring R :

- (i) Every R -module is projective.
- (ii) Every R -module is injective.

Let R be a ring. We will first prove (i) $\Rightarrow$ (ii). Let P be an R -module, A and B be arbitrary R -modules, $f : A \hookrightarrow B$ an injection, and $g : A \rightarrow P$ be an R -module homomorphism. It suffices to show there exists a homomorphism $h : B \rightarrow P$ such that $hf = g$. Consider the quotient module B/A ; taking $\pi : B \rightarrow B/A$ to be the canonical projection map, we obtain the short exact sequence

$$0 \rightarrow A \xrightarrow{f} B \xrightarrow{\pi} B/A \rightarrow 0.$$

Since every R -module is projective by hypothesis, B/A is a projective R -module. Therefore, the short exact sequence splits, which means that $B \cong A \oplus (B/A)$. Now let $k : B/A \rightarrow P$ be any R -module homomorphism and define the map $h : B \rightarrow P$ as follows: $h(a, c) = g(a) + k(c)$ for any $(a, c) \in A \oplus (B/A)$. We observe that

$$hf(a) = h(a, 0) = g(a) + k(0) = g(a), \quad \forall a \in A,$$

so that $hf = g$. Hence, P is an injective R -module.

Now we will show that (ii) $\Rightarrow$ (i). Let Q be an R -module, A and B be arbitrary fixed R -modules, $f : A \rightarrow B$ a surjection, and $g : Q \rightarrow B$ be an R -module homomorphism. It suffices to prove that there exists an R -module homomorphism $h : Q \rightarrow A$ such that $fh = g$. Let $K = \ker f$, and consider the short exact sequence

$$0 \rightarrow K \xrightarrow{i} A \xrightarrow{f} B \rightarrow 0,$$

where $i : K \hookrightarrow A$ is an injective mapping. Since every R -module is injective by hypothesis (i.e., K is injective), the short exact sequence splits, which means that $A \cong B \oplus K$. Since the sequence splits, there also exists a morphism $s : B \rightarrow A$ such that $f \circ s = \text{id}_B$. Let $h = sg$. Then we observe that $fh = (fs)g = \text{id}_B g = g$.

Problem D&F-10.5-7. Let A be a nonzero finite abelian group.

- (a) Prove that A is not a projective $\mathbb{Z}$ -module.
- (b) Prove that A is not an injective $\mathbb{Z}$ -module.

Let A be a nonzero finite abelian group.

- (a) Assume to the contrary that A is a projective $\mathbb{Z}$ -module. This implies that A must be the direct summand of a free $\mathbb{Z}$ -module. But $\mathbb{Z}$ -modules are precisely the abelian groups. Hence, A must be the direct summand of a free abelian group. Since subgroups of free abelian groups are free abelian, A must be a free abelian group. But there exists exactly one free abelian group of finite cardinality, which is the trivial group $\{0\}$. This contradicts our hypothesis that A is nonzero. Hence, A cannot be a projective $\mathbb{Z}$ -module.
- (b) Now assume to the contrary that A is an injective $\mathbb{Z}$ -module. By Proposition 36, A must necessarily be a divisible $\mathbb{Z}$ -module. But the only finite divisible $\mathbb{Z}$ -module is the trivial module $\{0\}$. This contradicts our hypothesis that A is nonzero. Hence, A cannot be an injective $\mathbb{Z}$ -module.

Problem D&F-10.5-9. Assume R is commutative with 1.

- (a) Prove that the tensor product of two free R -modules is free. [Use the fact that tensor products commute with direct sums.]
- (b) Use (a) to prove that the tensor product of two projective R -modules is projective.

Assume R is commutative with 1.

- (a) Let M_1, M_2 be fixed arbitrary free R -modules so that $M_1 \cong R^{(I)}$ and $M_2 \cong R^{(J)}$ for some index sets I, J . Since the tensor product commutes with direct sums,

$$M_1 \otimes_R M_2 \cong R^{(I)} \otimes_R R^{(J)} = \bigoplus_{i \in I} R^{(i)} \otimes_R \bigoplus_{j \in J} R^{(j)} = \bigoplus_{(i,j) \in I \times J} R^{(i,j)} = R^{(I \times J)}.$$

Hence, we conclude that $M_1 \otimes_R M_2$ is a free R -module.

- (b) Now let M_1, M_2 be projective R -modules. Since every projective R -module is the direct summand of a free module, suppose there exist R -modules N_1, N_2 so that $M_1 \oplus N_1 \cong R^{(I)}$ and $M_2 \oplus N_2 \cong R^{(J)}$ for some index sets I, J . Then

$$R^{(I \times J)} \cong R^{(I)} \otimes_R R^{(J)} \cong (M_1 \oplus N_1) \otimes_R (M_2 \oplus N_2) = (M_1 \otimes_R M_2) \oplus (M_1 \otimes_R N_2) \oplus \cdots \oplus (N_1 \otimes_R N_2).$$

Since $M_1 \otimes_R M_2$ is the direct summand of a free R -module, we conclude that $M_1 \otimes_R M_2$ is a projective R -module. This concludes the claim.

Problem D&F-10.5-10. Let R and S be rings with 1 and let M and N be left R -modules. Assume also that M is an (R, S) -bimodule.

- (a) For $s \in S$ and for $\varphi \in \text{Hom}_R(M, N)$, define $(s\varphi) : M \rightarrow N$ by $(s\varphi)(m) = \varphi(ms)$. Prove that $s\varphi$ is a homomorphism of left R -modules, and that this action of S on $\text{Hom}_R(M, N)$ makes it into a left S -module.

- (a) First, we will show that $s\varphi : M \rightarrow N$ is a left R -module homomorphism. Let $m_1, m_2 \in M$, and $r \in R$. Then

$$\begin{aligned} (s\varphi)(m_1 + m_2) &= \varphi((m_1 + m_2)s) \\ &= \varphi(m_1s + m_2s) \\ &= \varphi(m_1s) + \varphi(m_2s) = (s\varphi)(m_1) + (s\varphi)(m_2). \\ (s\varphi)(rm_1) &= \varphi(rm_1s) \\ &= \varphi(r(m_1s)) = r\varphi(m_1s) = r(s\varphi)(m_1). \end{aligned}$$

Above, we are using the fact that M is a *right* S -module when multiplying m_i by s on the right. Hence, we conclude that $(s\varphi)$ is a left R -module homomorphism. Now, we will show that this action of S on $\text{Hom}_R(M, N)$ makes it a left S -module. We already have, by definition of the hom set, that $\text{Hom}_R(M, N)$ is an abelian additive group. Now fix an arbitrary $x \in M$, let $s, \tilde{s} \in S$, and $\varphi, \phi \in \text{Hom}_R(M, N)$. Then we observe the following:

$$\begin{aligned}
 (s(\varphi + \phi))(x) &= (\varphi + \phi)(xs) \\
 &= \varphi(xs) + \phi(xs) \\
 &= (s\varphi)(x) + (s\phi)(x). \\
 \implies (s \cdot (\varphi + \phi)) &= (s\varphi) + (s\phi). \\
 (s + \tilde{s}) \cdot \varphi(x) &= \varphi(x \cdot (s + \tilde{s})) \\
 &= \varphi(xs + x\tilde{s}) \quad (\text{since } M \text{ is a right } S\text{-module}) \\
 &= \varphi(xs) + \varphi(x\tilde{s}) = (s\varphi)(x) + (\tilde{s}\varphi)(x) \\
 \implies (s + \tilde{s}) \cdot \varphi &= (s\varphi) + (\tilde{s}\varphi). \\
 (s\tilde{s})(\varphi)(x) &= \varphi(xs\tilde{s}) = \varphi((xs)\tilde{s}) \\
 &= \tilde{s} \cdot \varphi(xs) \\
 &= s(\tilde{s}\varphi)(x).
 \end{aligned}$$

That the operation satisfies the unit condition is straightforward to verify. Hence, $\text{Hom}_R(M, N)$ is a left S -module.

Problem D&F-10.5-12. Let A be an R -module, let I be a nonempty index set and for each $i \in I$, let B_i be an R -module. Prove the following isomorphisms of abelian groups; when R is commutative also that these are R -module isomorphisms. (Arbitrary direct sums and direct products of modules are introduced in Exercise 20 of Section 3.)

- (a) $\text{Hom}_R(\bigoplus_{i \in I} B_i, A) \cong \prod_{i \in I} \text{Hom}_R(B_i, A)$.
 (b) $\text{Hom}_R(A, \prod_{i \in I} B_i) \cong \prod_{i \in I} \text{Hom}_R(A, B_i)$.

- (a) Let $g \in \text{Hom}_R(\bigoplus_{i \in I} B_i, A)$; by the universal property for direct sums, g is uniquely determined by the family of maps $(g_i)_{i \in I}$, where for each i , $g_i = g \circ \iota_i$ and $g_i : B_i \rightarrow A$. Hence, define the map $\mathcal{F} : \text{Hom}_R(\bigoplus_{i \in I} B_i, A) \rightarrow \prod_{i \in I} \text{Hom}_R(B_i, A)$ as follows:

$$\mathcal{F}(\varphi) = (\varphi_i)_{i \in I} \implies \mathcal{F}(\varphi)((b_i)_{i \in I}) = ((\varphi \circ \iota_i)(b_i))_{i \in I} =: (\varphi_i(b_i))_{i \in I},$$

where $(b_i)_{i \in I} \in \bigoplus_{i \in I} B_i$. First, we will show that $\mathcal{F}$ is a group homomorphism. Let $\varphi, \phi \in \text{Hom}_R(\bigoplus_{i \in I} B_i, A)$, and fix some arbitrary $(b_i)_{i \in I} \in \bigoplus_{i \in I} B_i$. Then we observe that

$$\begin{aligned}
 \mathcal{F}(\varphi + \phi)((b_i)) &= ((\varphi + \phi)_i(b_i))_{i \in I} \\
 &= (\varphi_i(b_i) + \phi_i(b_i))_{i \in I} \\
 &= (\varphi_i(b_i))_{i \in I} + (\phi_i(b_i))_{i \in I} \\
 &= \mathcal{F}(\varphi)((b_i)) + \mathcal{F}(\phi)((b_i)).
 \end{aligned}$$

Hence, $\mathcal{F}$ is a group homomorphism. Now let $(\varphi_i)_{i \in I} \in \prod_{i \in I} \text{Hom}_R(B_i, A)$. By the universal property for direct sums, there exists a unique morphism $\varphi : \bigoplus_{i \in I} B_i \rightarrow A$ such that for each i , $\varphi_i = \varphi \circ \iota_i : B_i \rightarrow A$. Hence, $\mathcal{F} : \varphi \mapsto (\varphi_i)_{i \in I}$ so that it is surjective. Finally, we will show that $\mathcal{F}$ is injective. Let $\varphi, \phi \in \text{Hom}_R(\bigoplus_{i \in I} B_i, A)$ and assume that $\mathcal{F}(\varphi) = \mathcal{F}(\phi)$. Then for any $b \in \bigoplus_{i \in I} B_i$, we note that

$$0 = \mathcal{F}(\varphi)(b) - \mathcal{F}(\phi)(b) = \mathcal{F}(\varphi - \phi)(b),$$

so that $\varphi - \phi$ must be the trivial (i.e., zero) homomorphism. Hence, $\varphi = \phi$ so that $\mathcal{F}$ is injective. Altogether, we conclude that $\mathcal{F}$ is an isomorphism of groups.

- (b) Now let $g \in \text{Hom}_R(A, \prod_{i \in I} B_i)$. By the universal property for direct products, g is uniquely determined by a family of maps $(g_i)_{i \in I} : A \rightarrow B_i$ where for each i , $g_i = \pi_i \circ g$. Therefore, define the map $\mathcal{F} : \text{Hom}_R(A, \prod_{i \in I} B_i) \rightarrow \prod_{i \in I} \text{Hom}_R(A, B_i)$ as follows:

$$\text{for each } \varphi \in \text{Hom}_R(A, \prod_{i \in I} B_i). \quad \mathcal{F}(\varphi) = (\pi_i \circ \varphi)_{i \in I}.$$

First, we will show that $\mathcal{F}$ is a group homomorphism. Let $\varphi, \phi \in \text{Hom}_R(A, \prod_{i \in I} B_i)$. Then

$$\begin{aligned} \mathcal{F}(\varphi + \phi) &= (\pi_i \circ (\varphi + \phi))_{i \in I} \\ &= (\pi_i \circ \varphi + \pi_i \circ \phi)_{i \in I} \\ &= (\pi_i \circ \varphi)_{i \in I} + (\pi_i \circ \phi)_{i \in I} = \mathcal{F}(\varphi) + \mathcal{F}(\phi). \end{aligned}$$

Now let $(\varphi_i)_{i \in I}$ be a family of maps, where for each i , $\varphi_i : A \rightarrow B_i$. Then by the universal property for direct products, there exists a unique homomorphism $\varphi : A \rightarrow \prod_{i \in I} B_i$ such that for each i , $\varphi_i = \pi_i \circ \varphi$. Hence, $\mathcal{F} : \varphi \mapsto (\varphi_i)_{i \in I}$, so that this map is surjective. Finally, generalizing the argument from (a), we see that $\mathcal{F}$ is injective. Therefore, $\text{Hom}_R(A, \prod_{i \in I} B_i) \cong \prod_{i \in I} \text{Hom}_R(A, B_i)$.

Problem D&F-10.5-20. Prove that the polynomial ring $R[x]$ in the indeterminate x over the commutative ring R is a flat R -module.

Let R be a commutative ring and $R[x]$ the polynomial ring over R in the indeterminate x . Consider the set $\{x^n : n \geq 0\} \subset R[x]$. It is straightforward to see that this set is linearly independent since each element in this collection has a distinct degree. Moreover, since any $p(x) \in R[x]$ can be written as the finite sum

$$a_0 + a_1x + \cdots + a_nx^n$$

for some nonnegative integer n and $a_0, a_1, \dots, a_n \in R$,

$$\text{span}\{x^n : n \geq 0\} = R[x].$$

Hence, this set forms a basis for $R[x]$. This means that $R[x]$ is a free R -module:

$$R[x] = \bigoplus_{n=0}^{\infty} R \cdot x^n.$$

Since free R -modules are flat (Corollary 42), we conclude that $R[x]$ is flat.

Problem FR-1-1. Let R be a commutative ring with 1. Let I and J be ideals in R . Let M be an R -module.

- Describe $R/I \otimes_R M$ in terms of M , its R -submodules, and its quotient R -modules.
- Describe $R/I \otimes_R R/J$ as a quotient of R .
- Using the following short exact sequence of R -modules, describe $\text{Tor}_1^R(R/I, R/J)$ in terms of ideals coming from I and J .

$$0 \longrightarrow J \longrightarrow R \longrightarrow R/J \longrightarrow 0.$$

- Describe the finitely generated abelian group $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/9\mathbb{Z}, \mathbb{Z}/27\mathbb{Z})$ in the “standard form” $\mathbb{Z}/a_1\mathbb{Z} \times \cdots \times \mathbb{Z}/a_m\mathbb{Z}$ with $a_1 \mid a_2 \mid \cdots \mid a_m$, i.e., give the integer m and the sequence of elementary divisors $a_1, \dots, a_m$.

- We claim that $R/I \otimes_R M \cong M/IM$.

- We have $R/I \otimes_R R/J \cong R/(I+J)$. To prove this result, construct the map $\theta : R \rightarrow R/I \otimes_R R/J$, defined by $\theta(r) = \bar{1} \otimes \bar{r}$. It follows that this map is surjective: given any $R/I \otimes_R R/J \ni \bar{a} \otimes \bar{b} = \bar{1} \otimes \overline{ab}$. Hence, θ maps ab to this desired tensor product. Now we will compute the kernel of this map. Let $i \in I, j \in J$. Then

$$\theta(i+j) = \bar{1} \otimes (\bar{i} + \bar{j}) = \bar{1} \otimes \bar{i} + \bar{1} \otimes \bar{j} = 0,$$

since $\bar{i} = 0 \in R/I$ and $\bar{j} = 0 \in R/J$. Thus, $I + J \subseteq \ker \theta$. To show that $\ker(\theta) \subseteq I + J$, define the map

$$\psi : R/I \otimes_R R/J \rightarrow R/(I + J)$$

by

$$\psi(\bar{a} \otimes \bar{b}) = \overline{ab}.$$

This map is well defined: if $a - a' \in I$, then

$$ab - a'b = (a - a')b \in I \subseteq I + J,$$

so $\overline{ab} = \overline{a'b}$ in $R/(I + J)$. Similarly, if $b - b' \in J$, then

$$ab - ab' = a(b - b') \in J \subseteq I + J.$$

Now let $r \in \ker(\theta)$. Then

$$0 = \theta(r) = \bar{1} \otimes \bar{r}.$$

Applying ψ , we obtain

$$0 = \psi(\bar{1} \otimes \bar{r}) = \bar{r},$$

which implies that $r \in I + J$. Therefore,

$$\ker(\theta) \subseteq I + J.$$

Therefore, by the First Isomorphism Theorem, we conclude that $R/(I + J) \cong R/I \otimes_R R/J$.

(c) Consider the short exact sequence

$$0 \longrightarrow J \longrightarrow R \longrightarrow R/J \longrightarrow 0.$$

Then we have the Tor long exact sequence:

$$\cdots \rightarrow \operatorname{Tor}_1^R(R/I, R) \rightarrow \operatorname{Tor}_1^R(R/I, R/J) \rightarrow R/I \otimes_R J \xrightarrow{1 \otimes \beta} R/I \otimes_R R \xrightarrow{1 \otimes \gamma} R/I \otimes_R R/J \rightarrow 0.$$

We know that $\operatorname{Tor}_1^R(R/I, R) = 0$. Hence, we must have

$$\operatorname{Tor}_1^R(R/I, R/J) \cong \ker(R/I \otimes_R J \longrightarrow R/I \otimes_R R).$$

By (a), $R/I \otimes_R R \cong R/I$, and $R/I \otimes_R J \cong J/IJ$. We are interested in the kernel of the map from $J/IJ \rightarrow R/I$. But this kernel is precisely $(I \cap J)/IJ$, and so $\operatorname{Tor}_1^R(R/I, R/J) \cong (I \cap J)/IJ$.

(d) By Part (c), we then have

$$\operatorname{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/9\mathbb{Z}, \mathbb{Z}/27\mathbb{Z}) \cong \frac{(9) \cap (27)}{(243)} \cong \frac{(27)}{(243)} \cong \mathbb{Z}/9\mathbb{Z}.$$

Problem D&F-17.1-6. Let $0 \rightarrow \mathcal{A} \xrightarrow{\alpha} \mathcal{B} \xrightarrow{\beta} \mathcal{C} \rightarrow 0$ be a short exact sequence of cochain complexes. Prove that if any two of $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are exact, then so is the third. [Use Theorem 2.]

Let $0 \rightarrow \mathcal{A} \xrightarrow{\alpha} \mathcal{B} \xrightarrow{\beta} \mathcal{C} \rightarrow 0$ be a short exact sequence of cochain complexes. Then there exists a corresponding long exact sequence in cohomology:

$$0 \rightarrow H^0(\mathcal{A}) \rightarrow H^0(\mathcal{B}) \rightarrow H^0(\mathcal{C}) \xrightarrow{\delta_1} H^1(\mathcal{A}) \rightarrow H^1(\mathcal{B}) \rightarrow H^1(\mathcal{C}) \xrightarrow{\delta_1} H^2(\mathcal{A}) \rightarrow \cdots.$$

First assume $\mathcal{A}$ and $\mathcal{C}$ are exact. Then $H^\bullet(\mathcal{A})$ and $H^\bullet(\mathcal{C})$ are zero. So, our long exact sequence in cohomology collapses to the series of exact segments:

$$0 \rightarrow 0 \rightarrow H^\bullet(\mathcal{B}) \rightarrow 0 \xrightarrow{\delta_1} 0.$$

Hence, we conclude that each $H^\bullet(\mathcal{B}) = 0$ so that $\mathcal{B}$ is exact. In fact, repeating this argument for the other possible combinations concludes the proof.

Problem D&F-17.1-7. Prove that a finitely generated abelian group A is free if and only if

$$\text{Ext}^1(A; \mathbb{Z}) = 0.$$

Let A be a finitely generated abelian group; note that every finitely generated abelian group can be considered as a $\mathbb{Z}$ -module. First assume that A is free; since every free module is projective, A is a projective $\mathbb{Z}$ -module. So then by Proposition 11, $\text{Ext}_{\mathbb{Z}}^n(A, \mathbb{Z}) = 0$ for all $n \geq 1$. Now assume that $\text{Ext}_{\mathbb{Z}}^1(A, \mathbb{Z}) = 0$. By the Fundamental Theorem for Modules Generated over a PID, one has

$$A \cong \mathbb{Z}^r \oplus \bigoplus_{i=1}^s \mathbb{Z}/n_i\mathbb{Z}.$$

Since $\mathbb{Z}^r$ is a projective $\mathbb{Z}$ -module, one has

$$0 = \text{Ext}_{\mathbb{Z}}^1(A, \mathbb{Z}) \cong \bigoplus_{i=1}^s \text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/n_i\mathbb{Z}, \mathbb{Z}) \cong \bigoplus_{i=1}^s \mathbb{Z}/n_i\mathbb{Z}.$$

Since $s < \infty$, each n_i must be 1 so that $A \cong \mathbb{Z}^r$, and is therefore free.

Problem D&F-17.1-10.

- (a) Prove that an arbitrary direct sum $\bigoplus_{i \in I} P_i$ of projective modules P_i is projective and that an arbitrary direct product $\prod_{j \in J} Q_j$ of injective modules is injective.
- (b) Prove that an arbitrary direct sum of projective resolutions is again projective and use this to show $\text{Ext}_n^R(\bigoplus_{i \in I} A_i, B) \cong \prod_{i \in I} \text{Ext}_n^R(A_i, B)$ for any collection of R -modules A_i ($i \in I$). [cf. Exercise 12 in Section 10.5.]
- (c) Prove that an arbitrary direct product of injective resolutions is an injective resolution and use this to show $\text{Ext}_n^R(A, \prod_{j \in J} B_j) \cong \prod_{j \in J} \text{Ext}_n^R(A, B_j)$. [cf. Exercise 12 in Section 10.5.]
- (d) Prove that $\text{Tor}_n^R(A, \bigoplus_{j \in J} B_j) \cong \bigoplus_{j \in J} \text{Tor}_n^R(A, B_j)$.

- (a) Let I be an index set and $\{P_i\}_{i \in I}$ a collection of projective R -modules. Fix arbitrary left R -modules M and N , and let $f : M \twoheadrightarrow N$ be a surjection. Further, let $g : \bigoplus_{i \in I} P_i \rightarrow N$ be a R -module homomorphism. By the universal property of direct sums, g is uniquely determined by a family of maps $(g_i)_{i \in I} : P_i \rightarrow N$. Since each P_i is projective, there exists a family of module homomorphisms $h_i : P_i \rightarrow M$ such that $fh_i = g_i$ for each i . By the universal property for direct sums, there exists a unique module homomorphism $h : \bigoplus_{i \in I} P_i \rightarrow M$ such that for each i , $h_i = h \circ \iota_i$, where $\iota_i : P_i \hookrightarrow \bigoplus_{i \in I} P_i$. But then we observe that

$$g_i = fh_i \iff g \circ \iota_i = (fh) \circ \iota_i.$$

By the uniqueness property of morphisms given by the universal property, $fh = g$. Hence, $\bigoplus_{i \in I} P_i$ is a projective R -module. Now let J be an index set and $\{Q_j\}_{j \in J}$ a collection of injective R -modules. Fix arbitrary R -modules N and M , and let $f : N \hookrightarrow M$ be an injective homomorphism. Let $g : M \rightarrow \prod_{j \in J} Q_j$ be a module homomorphism. By the universal property of direct products, g is uniquely determined by a family $(g_j)_{j \in J}$ of morphisms $g_j : M \rightarrow Q_j$ for each j , where $g_j = \pi_j \circ g$, with $\pi_j : \prod_{j \in J} Q_j \twoheadrightarrow Q_j$ being the canonical projection map. Since each Q_j is an injective R -module, there exists a morphism $h_j : N \rightarrow Q_j$ so that $h_j f = g_j$ for each j . Again invoking the universal property for direct products, the family $(h_j)_{j \in J}$ uniquely determines a map $h : N \rightarrow \prod_{j \in J} Q_j$ so that for each j , $h_j = \pi_j \circ h$. But then we observe that

$$\pi_j \circ g = g_j = h_j f = \pi_j \circ (hf).$$

By the uniqueness property of morphisms given by the universal property, we conclude that $hf = g$. Hence, we conclude that $\prod_{j \in J} Q_j$ is an injective R -module.

- (b) Let I be an arbitrary index set, A any R -module, and $\{(P_\bullet^{(i)}, d_\bullet^{(i)})\}_{i \in I}$ an arbitrary collection of projective resolutions of A . Then the direct sum of these projective resolutions is defined to be the sequence $(\bigoplus_{i \in I} P_\bullet^{(i)}, \bigoplus_{i \in I} d_\bullet^{(i)})$. Since the arbitrary direct sum of projective modules is projective (by part (a)), for each $j \in \mathbb{N}_0$, $\bigoplus_{i \in I} P_j^{(i)}$ is projective. It suffices to now show that the sequence

$$\cdots \rightarrow \bigoplus_{i \in I} P_n^{(i)} \xrightarrow{\bigoplus_{i \in I} d_n^{(i)}} \bigoplus_{i \in I} P_{n-1}^{(i)} \rightarrow \cdots \xrightarrow{\bigoplus_{i \in I} d_1^{(i)}} \bigoplus_{i \in I} P_0^{(i)} \xrightarrow{\bigoplus_{i \in I} \varepsilon^{(i)}} \bigoplus_{i \in I} A \rightarrow 0$$

is exact. But this follows immediately due to the following: for each $j \in \mathbb{N}_0$,

$$\operatorname{im} \bigoplus_{i \in I} d_{j+1}^{(i)} = \bigoplus_{i \in I} \operatorname{im} d_{j+1}^{(i)} = \bigoplus_{i \in I} \ker d_j^{(i)} = \ker \bigoplus_{i \in I} d_j^{(i)}$$

Hence, the arbitrary direct sum of projective resolutions is a projective resolution. Now, we will show that $\operatorname{Ext}_R^n(\bigoplus_{i \in I} A_i, B) \cong \prod_{i \in I} \operatorname{Ext}_R^n(A_i, B)$. Let $\{(P_\bullet^{(i)}, d_\bullet^{(i)})\}_{i \in I}$ be a collection of projective resolutions for each A_i . By our work above, $(\bigoplus_{i \in I} P_\bullet^{(i)}, \bigoplus_{i \in I} d_\bullet^{(i)})$ is a projective resolution for $\bigoplus_{i \in I} A_i$. Then by definition,

$$\begin{aligned} \operatorname{Ext}_R^n\left(\bigoplus_{i \in I} A_i, B\right) &:= H^n\left(\operatorname{Hom}_R\left(\bigoplus_{i \in I} P_\bullet^{(i)}, B\right)\right) \cong H^n\left(\prod_{i \in I} \operatorname{Hom}_R(P_\bullet^{(i)}, B)\right) \\ &\cong \prod_{i \in I} H^n\left(\operatorname{Hom}_R(P_\bullet^{(i)}, B)\right) \cong \prod_{i \in I} \operatorname{Ext}_R^n(A_i, B), \end{aligned}$$

where in the first isomorphism, we used Exercise 12 from Section 10.5, and the second isomorphism follows from the fact that cohomology commutes with products since both kernels and images commute with products.

- (c) Let I be an arbitrary index set, A any R -module, and $\{(Q_\bullet^{(i)}, d_\bullet^{(i)})\}_{i \in I}$ an arbitrary collection of injective resolutions of A . Then the direct product of these injective resolutions is defined to be the sequence $(\prod_{i \in I} Q_\bullet^{(i)}, \prod_{i \in I} d_\bullet^{(i)})$. Since the arbitrary direct product of injective modules is injective (by part (a)), for each $j \in \mathbb{N}_0$, $\prod_{i \in I} Q_j^{(i)}$ is injective. It suffices to now show that the sequence

$$0 \rightarrow \prod_{i \in I} A \xrightarrow{\prod_{i \in I} \varepsilon} \prod_{i \in I} Q_0^{(i)} \xrightarrow{\prod_{i \in I} d_1^{(i)}} \cdots \xrightarrow{\prod_{i \in I} d_{n-1}^{(i)}} \prod_{i \in I} Q_n^{(i)} \xrightarrow{\prod_{i \in I} d_n^{(i)}} \cdots$$

is exact. But since kernels and images commute with products, by injectivity of each resolution, we observe that

$$\ker \prod_{i \in I} d_n^{(i)} \cong \prod_{i \in I} \ker d_n^{(i)} \cong \prod_{i \in I} \operatorname{image} d_{n+1}^{(i)} \cong \operatorname{image} \prod_{i \in I} d_{n+1}^{(i)},$$

and so the sequence is indeed exact. Therefore, the arbitrary direct product of injective resolutions is an injective resolution. Now, we will show that $\operatorname{Ext}_R^n(A, \prod_{j \in J} B_j) \cong \prod_{j \in J} \operatorname{Ext}_R^n(A, B_j)$. Let $\{(Q_\bullet^{(j)}, d_\bullet^{(j)})\}_{j \in J}$ be a collection of injective resolutions for each B_j . By our work above,

$$\left(\prod_{j \in J} Q_\bullet^{(j)}, \prod_{j \in J} d_\bullet^{(j)}\right)$$

is an injective resolution for $\prod_{j \in J} B_j$. Then by definition

$$\begin{aligned} \operatorname{Ext}_R^n\left(A, \prod_{j \in J} B_j\right) &:= H^n\left(\operatorname{Hom}_R\left(A, \prod_{j \in J} Q_\bullet^{(j)}\right)\right) \cong H^n\left(\prod_{j \in J} \operatorname{Hom}_R(A, Q_\bullet^{(j)})\right) \\ &\cong \prod_{j \in J} H^n\left(\operatorname{Hom}_R(A, Q_\bullet^{(j)})\right) =: \prod_{j \in J} \operatorname{Ext}_R^n(A, B_j), \end{aligned}$$

where the first isomorphism follows from Exercise 12 in Section 10.5.

- (d) Let A be an arbitrary right R -module and $\{B_j\}_{j \in J}$ left R -modules. Let $(P_\bullet, d_\bullet)$ be a projective resolution for A . Then

$$\begin{aligned} \operatorname{Tor}_n^R\left(A, \bigoplus_{j \in J} B_j\right) &\cong H_n\left(P_\bullet \otimes_R \bigoplus_{j \in J} B_j\right) \cong H_n\left(\bigoplus_{j \in J} (P_\bullet \otimes_R B_j)\right) \cong \bigoplus_{j \in J} H_n(P_\bullet \otimes_R B_j) \\ &\cong \bigoplus_{j \in J} \operatorname{Tor}_n^R(A, B_j), \end{aligned}$$

where in the second isomorphism, we used the fact that tensor products commute with arbitrary direct sums, and in the third isomorphism, we used the fact that homology commutes with arbitrary direct sums.

Problem D&F-17.1-12. Prove Proposition 13: $\operatorname{Tor}_0^R(D, A) \cong D \otimes_R A$. [Follow the proof of Proposition 3.]

Let D be a fixed arbitrary right R -module, A a fixed arbitrary left R -module, and let $(P_\bullet, d_\bullet)$ be an arbitrary projective resolution for A . Then we have the exact sequence

$$P_1 \xrightarrow{d_1} P_0 \xrightarrow{\varepsilon} A \rightarrow 0.$$

Since tensor products is right exact, tensoring on the left by D , we obtain the exact sequence

$$D \otimes_R P_1 \xrightarrow{1 \otimes d_1} D \otimes_R P_0 \xrightarrow{1 \otimes \varepsilon} D \otimes_R A \rightarrow 0.$$

By exactness, we must have $(1 \otimes \varepsilon)$ surjective, and $\operatorname{im}(1 \otimes d_1) = \ker(1 \otimes \varepsilon)$. By definition of Tor ,

$$\operatorname{Tor}_0^R(D, A) \cong (D \otimes_R P_0) / \operatorname{im}(1 \otimes d_1) = (D \otimes_R P_0) / \ker(1 \otimes \varepsilon) \cong D \otimes_R A,$$

where the final isomorphism comes from the First Isomorphism Theorem for Modules.

Problem D&F-17.1-13. Prove Proposition 16 characterizing flat modules.

Proposition 16 states the following:

Proposition 16. For a right R -module D the following are equivalent:

- (a) D is a flat R -module.
- (b) $\operatorname{Tor}_1^R(D, B) = 0$ for all left R -modules B , and
- (c) $\operatorname{Tor}_n^R(D, B) = 0$ for all left R -modules B and all $n \geq 1$.

Clearly (c) implies (b). Therefore, we will show that (b) implies (a) and (a) implies (c). We start with the latter. Let D be a flat R -module, and fix an arbitrary left R -module B . Let $(P_\bullet, d_\bullet)$ be a projective resolution for B . Then tensoring by D on the right, we get the *exact* (since flat R -modules preserve exact sequences) chain complexes

$$\cdots \rightarrow D \otimes_R P_2 \xrightarrow{1 \otimes d_2} D \otimes_R P_1 \xrightarrow{1 \otimes d_1} D \otimes_R P_0 \xrightarrow{1 \otimes \varepsilon} D \otimes_R B \rightarrow 0.$$

Since this complex is exact, for each $n \geq 1$,

$$\operatorname{Tor}_n^R(D, B) = \ker(1 \otimes d_n) / \operatorname{image}(1 \otimes d_{n+1}) = 0.$$

Now, we will prove that (b) implies (a). Suppose that $\operatorname{Tor}_1^R(D, B) = 0$ for all left R -modules. Consider the short exact sequence of R -modules, $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$. Then there exists the Tor long exact sequence

$$\begin{aligned} \cdots \rightarrow \operatorname{Tor}_2^R(D, N) \xrightarrow{\delta_1} \operatorname{Tor}_1^R(D, L) \rightarrow \operatorname{Tor}_1^R(D, M) \rightarrow \operatorname{Tor}_1^R(D, N) \\ \xrightarrow{\delta_0} D \otimes_R L \rightarrow D \otimes_R M \rightarrow D \otimes_R N \rightarrow 0. \end{aligned}$$

Applying the hypothesis, we obtain the short exact sequence

$$0 \rightarrow D \otimes_R L \rightarrow D \otimes_R M \rightarrow D \otimes_R N \rightarrow 0.$$

Hence, since for any short exact sequence of R -modules, applying the functor $D \otimes_R _$ preserves short exactness, it follows that D is a flat R -module.

Problem D&F-17.1-14. Suppose $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is a short exact sequence of R -modules. Prove that if C is a flat R -module, then A is flat if and only if B is also flat. [Use the Tor long exact sequence.] Given an example to show that if A and B are flat then C need not be flat.

Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a short exact sequence of R -modules, and let C be a flat R -module. First assume that B is also a flat module. Then for any R -module M , there exists a long exact sequence

$$\begin{aligned} \cdots \rightarrow \operatorname{Tor}_2^R(M, C) \xrightarrow{\delta_1} \operatorname{Tor}_1^R(M, A) \rightarrow \operatorname{Tor}_1^R(M, B) \\ \rightarrow \operatorname{Tor}_1^R(M, C) \rightarrow M \otimes_R A \rightarrow M \otimes_R B \rightarrow M \otimes_R C \rightarrow 0. \end{aligned}$$

Since B and C are flat, $\operatorname{Tor}_i^R(M, B)$ and $\operatorname{Tor}_i^R(M, C)$ are zero for all $i \geq 1$. This means that we get the exact sequence $0 \rightarrow \operatorname{Tor}_1^R(M, A) \rightarrow 0$, and so $\operatorname{Tor}_1^R(M, A) = 0$. Since M was an arbitrary R -module, we conclude that A is flat. Now suppose A is a flat R -module so that for any left R -module M , $\operatorname{Tor}_1^R(M, A) = 0$. Then the above Tor long exact sequence gives us the exact sequence

$$0 \rightarrow \operatorname{Tor}_1^R(M, B) \rightarrow 0 \quad \text{so that } \operatorname{Tor}_1^R(M, B) = 0.$$

Since this is true for all left R -modules M , we conclude that B is flat.

Problem D&F-17.1-16. Suppose R is a P.I.D. and A and B are R -modules. If $t(B)$ denotes the torsion submodule of B show that $\operatorname{Tor}_1(A, t(B)) \cong \operatorname{Tor}_1^R(t(A), t(B))$ and deduce that $\operatorname{Tor}_1^R(A, B)$ is isomorphic to $\operatorname{Tor}_1^R(t(A), t(B))$. [Use Exercise 26 in Section 10.5 to show that $B/t(B)$ is flat over R , then use the Tor long exact sequence with $D = A$ applied to the short exact sequence $0 \rightarrow t(B) \rightarrow B \rightarrow B/t(B) \rightarrow 0$ and the remarks following Proposition 16.]

Let R be a PID, and A, B arbitrary R -modules. Let $t(B)$ denote the torsion submodule of B . First, we will show that $B/t(B)$ is a torsion-free module. Let $[b] \in B/t(B)$ be nonzero and assume to the contrary there exists some nonzero $r \in R$ such that $r[b] = [rb] = 0$. But then $rb \in t(B)$, which means there exists some nonzero $s \in R$ so that $(sr)b = s(rb) = 0$. Hence, $b \in t(B)$, which contradicts our hypothesis that $b \notin t(B)$. Hence, $B/t(B)$ is torsion-free. By Exercise 26 in Section 10.5, we conclude that $B/t(B)$ is flat over R . So now consider the short exact sequence

$$0 \rightarrow t(B) \rightarrow B \rightarrow B/t(B) \rightarrow 0.$$

Then there exists a long exact sequence

$$\begin{aligned} \cdots \rightarrow \operatorname{Tor}_2^R(A, B/t(B)) \xrightarrow{\delta_1} \operatorname{Tor}_1^R(A, t(B)) \rightarrow \operatorname{Tor}_1^R(A, B) \rightarrow \operatorname{Tor}_1^R(A, B/t(B)) \xrightarrow{\varepsilon} A \otimes_R t(B) \\ \rightarrow A \otimes_R B \rightarrow A \otimes_R B/t(B) \rightarrow 0. \end{aligned}$$

Since $B/t(B)$ is flat, $\operatorname{Tor}_n^R(A, B/t(B)) = 0$ for all $n \geq 1$. And so we obtain the short exact sequence

$$0 \rightarrow \operatorname{Tor}_1^R(A, t(B)) \rightarrow \operatorname{Tor}_1^R(A, B) \rightarrow 0;$$

by exactness, we conclude that $\operatorname{Tor}_1^R(A, t(B)) \cong \operatorname{Tor}_1^R(A, B)$. Repeating this argument for the short exact sequence $0 \rightarrow t(A) \rightarrow A \rightarrow A/t(A) \rightarrow 0$, we obtain the isomorphism $\operatorname{Tor}_1^R(A, B) \cong \operatorname{Tor}_1^R(t(A), B)$. Finally, using the Tor long exact sequence with $D = t(A)$ applied to the short exact sequence $0 \rightarrow t(B) \rightarrow B \rightarrow B/t(B) \rightarrow 0$, we obtain the isomorphism $\operatorname{Tor}_1^R(t(A), t(B)) \cong \operatorname{Tor}_1^R(t(A), B)$. Hence, using our previous results, we conclude that $\operatorname{Tor}_1^R(A, B) \cong \operatorname{Tor}_1^R(t(A), t(B))$.

Problem D&F-17.1-17. Let $A = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z} \oplus \cdots$. Prove that $\text{Ext}^1(A, B) \cong (B/2B) \times (B/3B) \times (B/4B) \times \cdots$ for any abelian group B . [Use Exercise 10.] Prove that $\text{Ext}^1(A, B) = 0$ if and only if B is divisible.

Define the group A as prescribed above, and let B be a fixed arbitrary abelian group. In Exercise 10, we proved that

$$\text{Ext}^1(A, B) = \text{Ext}^1\left(\bigoplus_{j=2}^{\infty} \mathbb{Z}/j\mathbb{Z}, B\right) \cong \prod_{j=2}^{\infty} \text{Ext}^1(\mathbb{Z}/j\mathbb{Z}, B).$$

We shall work out $\text{Ext}^1(\mathbb{Z}/j\mathbb{Z}, B)$ for some specified j first. Consider the projective resolution

$$0 \rightarrow \mathbb{Z} \xrightarrow{j} \mathbb{Z} \rightarrow \mathbb{Z}/j\mathbb{Z} \rightarrow 0.$$

Applying the functor $\text{Hom}(-, B)$, we obtain the chain complex

$$0 \rightarrow \text{Hom}(\mathbb{Z}/j\mathbb{Z}, B) \rightarrow \text{Hom}(\mathbb{Z}, B) \xrightarrow{j^*} \text{Hom}(\mathbb{Z}, B) \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \cdots.$$

We recall the isomorphism $\text{Hom}(\mathbb{Z}, B) \cong B$ so that we have the cochain complex

$$0 \rightarrow {}_jB \rightarrow B \xrightarrow{j} B \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \cdots.$$

Hence,

$$\text{Ext}^1(\mathbb{Z}/j\mathbb{Z}, B) \cong \frac{\ker(B \rightarrow 0)}{\text{image}(B \xrightarrow{j} B)} \cong B/jB.$$

Therefore, we conclude that

$$\text{Ext}^1(A, B) \cong \prod_{j=2}^{\infty} B/jB.$$

Now suppose B is a divisible group, which means that for every positive integer j , $jB = B$. This means that $B/jB = 0$ so that

$$\text{Ext}^1(A, B) \cong \prod_{j=2}^{\infty} B/jB = \prod_{j \geq 2} 0 = 0.$$

Now suppose $\text{Ext}^1(A, B) \cong 0$. This means that

$$\prod_{j \geq 2} B/jB = 0.$$

This forces every $B/jB = 0$ since if not, the product cannot be equal to zero. If $B/jB = 0$, then we immediately conclude that $B/jB = 0$ for all $j \geq 2$. Moreover, it follows trivially that $B = (1)B$. Hence, B is divisible.

Problem D&F-17.1-19. Suppose $r \neq 0$ is not a zero divisor in the commutative ring R .

- (a) Prove that multiplication by r gives a free resolution $0 \rightarrow R \xrightarrow{r} R \rightarrow R/rR \rightarrow 0$ of the quotient R/rR .
- (b) Prove that $\text{Ext}_R^0(R/rR, B) = {}_rB$ is the set of elements $b \in B$ with $rb = 0$, that $\text{Ext}_R^1(R/rR, B) \cong B/rB$, and that $\text{Ext}_R^n(R/rR, B) = 0$ for $n \geq 2$ for every R -module B .
- (c) Prove that $\text{Tor}_0^R(A, R/rR) = A/rA$, that $\text{Tor}_1^R(A, R/rR) = {}_rA$ is the set of elements $a \in A$ with $ra = 0$, and that $\text{Tor}_n^R(A, R/rR) = 0$ for $n \geq 2$ for every R -module A .

Suppose $r \neq 0$ is not a zero divisor in the commutative ring R .

- (a) First, it is straightforward that seen as R -modules, R is a free R -module. Hence, it suffices to show that the sequence $0 \rightarrow R \xrightarrow{r} R \xrightarrow{\pi} R/rR \rightarrow 0$ is exact. The kernel of the map $R/rR \rightarrow 0$ is precisely R/rR . On the other hand, π being the canonical projection is surjective so that $\text{image } \pi = R/rR$. Hence, the sequence is exact at R/rR . Likewise, the kernel of multiplication by r is precisely 0 since if the kernel includes any nonzero elements u , then $ru = 0$, contradicting the fact that r is not a zero divisor. On the other hand, the image of the trivial homomorphism $0 \rightarrow R$ is exactly 0. Hence, the sequence is exact at the first R . On the other hand, the kernel of π is precisely rR , while the image of multiplication of R by r is also rR . Hence, the sequence is exact at the second R . Therefore, we conclude that multiplication by r gives a free resolution $0 \rightarrow R \xrightarrow{r} R \rightarrow R/rR \rightarrow 0$ of the quotient R/rR .
- (b) Since any free module is projective, the above free resolution gives a projective resolution for R/rR . Since the functor $-\otimes_R N$ is right exact, we obtain by tensoring B on the right, the exact sequence

$$R \otimes_R B \xrightarrow{r \otimes 1} R \otimes_R B \rightarrow R/rR \otimes_R B \rightarrow 0.$$

Since $R \otimes_R B \cong B$, we actually obtain the exact sequence

$$B \xrightarrow{r} B \rightarrow R/rR \otimes_R B \rightarrow 0.$$

Hence, it follows that

$$\text{Ext}_R^0(R/rR, B) = \ker(B \xrightarrow{r} B) = \{b \in B : rb = 0\} =: {}_rB.$$

Now we will calculate $\text{Ext}_R^1(R/rR, B)$. Applying the functor $\text{Hom}_R(-, B)$ to the free resolution, we obtain the exact sequence

$$0 \rightarrow \text{Hom}_R(R, B) \xrightarrow{r^*} \text{Hom}_R(R, B) \rightarrow 0.$$

Since $\text{Hom}_R(R, B) \cong B$ (by means of the identification $f \mapsto f(1)$), $\ker r^* = {}_rB$. Hence,

$$\text{Ext}_R^1(R/rR, B) \cong \frac{\ker 0}{\text{image } r} = B/{}_rB.$$

Since the sequence only has two nonzero components, it follows that $\text{Ext}_R^n(R/rR, B) = 0$ for all $n \geq 2$.

- (c) Consider again the free (projective) resolution $0 \rightarrow R \xrightarrow{r} R \rightarrow R/rR \rightarrow 0$. Applying the right-exact functor $A \otimes_R -$, we obtain the exact sequence

$$A \otimes_R R \xrightarrow{1 \otimes r} A \otimes_R R \rightarrow 0 \implies A \xrightarrow{r} A \rightarrow 0.$$

Hence, we conclude from this that

$$\text{Tor}_0^R(A, R/rR) = \text{coker } A \xrightarrow{r} A = A/rA.$$

On the other hand, $\text{Tor}_1^R(A, R/rR) = \ker A \xrightarrow{r} A = {}_rA$. And since the sequence terminates here, $\text{Tor}_n^R(A, R/rR) = 0$ for all $n \geq 2$.

Problem D&F-17.1-21. Let $R = k[x, y]$ where k is a field, and let I be the ideal (x, y) in R .

- (a) Let $\alpha : R \rightarrow R^2$ be the map $\alpha(r) = (yr, -xr)$ and let $\beta : R^2 \rightarrow R$ be the map $\beta((r_1, r_2)) = r_1x + r_2y$. Show that

$$0 \rightarrow R \xrightarrow{\alpha} R^2 \xrightarrow{\beta} R \rightarrow k \rightarrow 0$$

where the map $R \rightarrow R/I = k$ is the canonical projection, gives a free resolution of k as an R -module.

- (b) Use the resolution in (a) to show that $\text{Tor}_2^R(k, k) \cong k$.

Let $R = k[x, y]$, where k is a field, and let I be the ideal (x, y) in R .

- (a) It is straightforward to see that R, R^2, R are free R -modules by definition. Hence, it suffices to show that this sequence is exact. We observe the following

$$\ker \alpha = \{r \in R : xr = 0 \text{ and } yr = 0\} = \{0\}, \quad \ker \beta = \{(r_1, r_2) \in R : r_2y = -r_1x\}$$

$$\ker(R \rightarrow k) = I, \quad \ker(k \rightarrow 0) = k.$$

Hence,

- (i) At the first R ,

$$\text{image}(0 \rightarrow R) = \{0\} = \ker \alpha,$$

so that the sequence is exact at the first R .

- (ii) At R^2 we compute the following: (1) for any $(yr, -xr) \in \text{image } \alpha$,

$$\beta((yr, -xr)) = (yr)x - (xr)y = (yrx) - (yrx) = 0.$$

So $\text{image } \alpha \subset \ker \beta$; (2) for any $(r_1, r_2) \in \ker \beta$, $r_1x + r_2y = 0$ so that $x \mid r_2$ and $y \mid r_1$. I.e., there exists some $r \in R$ so that $r_2 = -xr$ and $r_1 = yr$. Hence, $\ker \beta \subseteq \text{image } \alpha$. Altogether, this proves exactness at R^2 .

- (iii) Likewise, we can keep going to show that the whole sequence is exact. Hence, this is a free resolution of k as an R -module.

- (b) From the provided free (hence, projective) resolution for k , applying the functor $k \otimes_R -$ yields the exact sequence

$$k \xrightarrow{\alpha} k^2 \xrightarrow{\beta} k \rightarrow 0$$

since $k \otimes_R R \cong k$ and $k \otimes_R R^2 \cong k^2$. So therefore, $\text{Tor}_2^R(k, k) = \ker \alpha = k$.

Problem D&F-17.1-22. (*Flat Base Change for Tor*) Suppose R and S are commutative rings and $f : R \rightarrow S$ is a ring homomorphism making S into an R -module as in Example 6 following Corollary 12 in Section 10.4. Prove that if S is flat as an R -module, then $\text{Tor}_n^R(A, B) \cong \text{Tor}_n^S(S \otimes_R A, B)$ for all R -modules A and all S -modules B . [Show that since S is flat, tensoring an R -module projective resolution for A with S gives an S -module projective resolution of $S \otimes_R A$.]

Let R and S be commutative rings, and $f : R \rightarrow S$ a ring homomorphism that turns S into an R -module. Further assume that S is flat as an R -module. Let A, B be fixed arbitrary R - and S -modules, respectively. Let $(P_\bullet, d_\bullet)$ be a projective resolution for A . Since S is flat, the sequence $(S \otimes_R P_\bullet, 1 \otimes d_\bullet)$ is an exact sequence. Now we will show that each $S \otimes_R P_j$ is a projective S -module. Let N, M be S -modules (hence R -modules by restriction of scalars), $f : N \twoheadrightarrow M$ a surjection (hence an R -module surjection by restriction of scalars) and $g : S \otimes_R P_j \rightarrow M$ a S -module homomorphism. By the universal property for tensor products, g is uniquely determined by an R -linear map $g_2 : P_j \rightarrow M$. Since P_j is a projective R -module, there exists a morphism $h_2 : P_j \rightarrow N$ so that $fh_2 = g_2$. So define the map $h : S \otimes_R P_j \rightarrow M$ as follows: $h(s \otimes p) = sh_2(p)$. Then we check that

$$fh(s \otimes p) = f(sh_2(p)) = sf(h_2(p)) = sg_2(p) = g(s \otimes p),$$

where the final equality follows from the fact that g is S -linear. Hence, $fh = g$, which proves that $S \otimes_R P_j$ is a projective S -module for all nonnegative integers j . This means that $(S \otimes_R P_\bullet, d_\bullet)$ is a projective resolution of $S \otimes_R A$. So since $(S \otimes_R P_\bullet) \otimes_S B \cong S \otimes_R (P_\bullet \otimes_R B)$, we conclude that

$$\begin{aligned} \text{Tor}_n^S(S \otimes_R A, B) &:= H_n((S \otimes_R P_\bullet) \otimes_S B) = H_n(S \otimes_R (P_\bullet \otimes_R B)) = H_n((S \otimes_R P_\bullet) \otimes_R B) \\ &=: \text{Tor}_n^R(S \otimes_R A, B). \end{aligned}$$

Problem D&F-17.1-23. (Localization and Tor) Let $D^{-1}R$ be the localization of the commutative ring R with respect to the multiplicative subset D of R . Prove that localization commutes with Tor, i.e.,

$D^{-1}\mathrm{Tor}_n^R(A, B) \cong \mathrm{Tor}_n^{D^{-1}R}(D^{-1}A, D^{-1}B)$. [Use the previous exercise and the fact that $D^{-1}R$ is flat over R , cf. Proposition 42(6) in Section 15.4.]

Let R be a commutative ring, D a multiplicative subset of R , and $D^{-1}R$ the localization of R . Let $(P_\bullet, d_\bullet)$ be a projective resolution for A so that

$$\mathrm{Tor}_n^R(A, B) := H_n(P_\bullet \otimes_R B).$$

Since localizations are exact and preserve projective modules, $D^{-1}P_\bullet$ is a projective resolution for $D^{-1}A$. This means that

$$\mathrm{Tor}_n^{D^{-1}R}(D^{-1}A, D^{-1}B) = H_n(D^{-1}P_\bullet \otimes_{D^{-1}R} D^{-1}B).$$

Here, we note that since localization commutes with tensor products,

$$D^{-1}P_\bullet \otimes_R D^{-1}B \cong D^{-1}(P_\bullet \otimes_R B).$$

Then since localizations commute with homology, $H_n(D^{-1}(P_\bullet \otimes_R B)) \cong D^{-1}H_n(P_\bullet \otimes_R B)$. Hence,

$$\mathrm{Tor}_n^{D^{-1}R}(D^{-1}A, D^{-1}B) = H_n(D^{-1}P_\bullet \otimes_{D^{-1}R} D^{-1}B) \cong D^{-1}H_n(P_\bullet \otimes_R B) \cong D^{-1}\mathrm{Tor}_n^R(A, B).$$

This concludes the proof that localization commutes with Tor.

4 Real Analysis

4.1 Convergence of Functions

Problem 2026-J-II-2. Let c be a positive real number, and let $\{f_n(x)\}_{n \in \mathbb{N}}$ be a sequence of integrable functions from $[0, 1]$ to $[-c, c]$. For the following functions on $[0, 1]$,

$$g_n(x) := \int_0^x f_n(t) dt,$$

prove that some subsequence $(g_{n_m})_{m \in \mathbb{N}}$ converges uniformly to a c -Lipschitz function.

Define the sequence $\{g_n\}$ of functions on $[0, 1]$ as given above. We begin by showing that each g_n is c -Lipschitz. Fix $\varepsilon > 0$, and let $\delta = \varepsilon/c > 0$. For any $x, y \in [0, 1]$ with $x > y$ and $|x - y| < \delta$, one computes,

$$|g_n(x) - g_n(y)| = \left| \int_y^x f_n(t) dt \right| \leq \int_y^x c dt = c|x - y| < \varepsilon.$$

Thus we have that each g_n is c -Lipschitz, and the sequence $\{g_j\}$ is equicontinuous. Thus by the Arzelà-Ascoli theorem, it follows that there exists a subsequence $\{g_{n_m}\}_{m=1}^\infty$ that converges uniformly to a function g on $[0, 1]$. We will now show that this limit is c -Lipschitz. Fix $\varepsilon > 0$, and take $x, y \in [0, 1]$ with $x > y$. Then

$$\begin{aligned} |g(x) - g(y)| &= |g(x) - g_{n_m}(x) + g_{n_m}(x) - g_{n_m}(y) + g_{n_m}(y) - g(y)| \\ &\leq |g(x) - g_{n_m}(x)| + |g_{n_m}(x) - g_{n_m}(y)| + |g_{n_m}(y) - g(y)| \\ &\leq |g(x) - g_{n_m}(x)| + c|x - y| + |g_{n_m}(y) - g(y)|. \end{aligned}$$

By uniform convergence of the subsequence, in the limit $m \rightarrow \infty$, one has $|g(x) - g_{n_m}(x)|, |g(y) - g_{n_m}(y)| \rightarrow 0$. Thus, we conclude that $|g(x) - g(y)| \leq c|x - y|$ so that g is c -Lipschitz.

Problem 2025-J-I-2. Let $\{f_n\}_{n \geq 1}$ be a sequence of Lebesgue measurable functions on $[0, 1]$. Suppose that

$$\int_0^1 f_n^2 dm \leq \frac{1}{n^2}$$

for all $n \geq 1$. Prove that f_n converges to 0 a.e. on $[0, 1]$.

Let $\{f_n\}_{n \geq 1}$ be a sequence of Lebesgue measurable functions on $[0, 1]$ such that for each $n \geq 1$,

$$\int_0^1 f_n^2 dm \leq \frac{1}{n^2}.$$

To show that $f_n \rightarrow 0$ a.e., it suffices to show that $f_n^2 \rightarrow 0$ a.e., and thus $\sum_1^\infty f_n^2$ converges a.e. We note that $\sum_1^N f_n^2 \rightarrow \sum_1^\infty f_n^2$ a.e., and the sequence of such partial sums is increasing. By the Monotone Convergence Theorem, one has the following:

$$\int_0^1 \sum_1^\infty f_n^2 dm = \int_0^1 \lim_{N \rightarrow \infty} \sum_1^N f_n^2 dm = \lim_{N \rightarrow \infty} \int_0^1 \sum_1^N f_n^2 dm = \sum_1^\infty \int_0^1 f_n^2 dm \leq \sum_1^\infty \frac{1}{n^2} < \infty.$$

Thus, $\sum_1^\infty f_n^2 < \infty$ a.e. on $[0, 1]$ so that the sum converges a.e. Since the tail of a convergent series must tend to zero a.e., we must have $f_n^2 \rightarrow 0$ a.e., and thus $f_n \rightarrow 0$ a.e.

Problem 2004-A-II-1. Let $\{f_n\}_{n=1}^\infty$ be a sequence of real valued absolutely continuous functions defined on $[0, 1]$. Assume there exist constants c_0 and c_1 such that

$$\sup_{0 \leq x \leq 1} |f_n(x)| \leq c_0 \quad \text{and} \quad \int_0^1 |f_n'(x)|^2 dx \leq c_1$$

for $n = 1, 2, 3, \dots$. Prove that $\{f_n\}_{n=1}^\infty$ has a subsequence which converges uniformly on $[0, 1]$.

Let $\{f_n\}_{n=1}^\infty$ be a sequence of real-valued absolutely continuous functions defined on $[0, 1]$ with the above properties. Since $[0, 1]$ is compact, by the Arzelà-Ascoli theorem, it suffices to show that the sequence is uniformly bounded and equicontinuous; the first condition on the sequence already proves uniform boundedness. Therefore, we will prove that the sequence is equicontinuous. Let $\varepsilon > 0$, and let $\delta = \varepsilon^2/c_1$. Then for any $x, y \in [0, 1]$ with $x > y$ and all $j \geq 1$, one has

$$\begin{aligned} |f_j(x) - f_j(y)| &= \left| \int_y^x f'_j(x) dx \right| \leq \int_y^x |f'_j(x)| dx \\ &\leq \left(\int_y^x |f'_j(x)|^2 dx \right)^{1/2} \left(\int_y^x dx \right)^{1/2} \quad (\text{by Cauchy-Schwarz Inequality}) \\ &\leq c_1^{1/2} \cdot |x - y|^{1/2} < \varepsilon. \end{aligned}$$

Thus it follows that $\{f_j\}_{j \geq 1}$ is uniformly continuous. By Arzelà-Ascoli, we conclude that the sequence has a subsequence which converges uniformly on $[0, 1]$.

Problem 2021-J-II-2. A real-valued function $f : [0, 1] \rightarrow \mathbb{R}$ is said to be Hölder continuous of order α if there is a constant $C > 0$ such that $|f(x) - f(y)| \leq C|x - y|^\alpha$ for all $x, y \in [0, 1]$. Define

$$\|f\|_\alpha = \max_{x \in [0, 1]} |f(x)| + \sup_{x, y \in [0, 1]} \frac{|f(y) - f(x)|}{|x - y|^\alpha}.$$

Show that for $\alpha \in (0, 1)$, the set of functions

$$S_\alpha = \{f \in C([0, 1]) : \|f\|_\alpha \leq 1\}$$

is a compact subset of $C([0, 1])$ in the topology of uniform convergence.

Fix some order $\alpha \in (0, 1)$ and define the set of functions,

$$S_\alpha = \{f \in C([0, 1]) : \|f\|_\alpha \leq 1\}.$$

By the Arzelà-Ascoli theorem, S_α is a compact subset of $C([0, 1])$ if and only if it is closed, bounded, and equicontinuous under the topology of uniform convergence. Boundedness is immediately clear: for all $f \in S_\alpha$, one has

$$\|f\|_\infty = \max_{x \in [0, 1]} |f(x)| \leq \|f\|_\alpha = 1,$$

where the upper bound of 1 is independent of the individual function f . Now we will show that S_α is closed under the supremum norm; we must show that if a sequence $\{f_n\} \subset S_\alpha$ converges uniformly to a function f , then $\|f\|_\alpha \leq 1$. We begin by showing first that f is Hölder continuous of exponent α . For any $x, y \in [0, 1]$, one has

$$\begin{aligned} |f(x) - f(y)| &\leq |f(x) - f_n(x)| + |f_n(x) - f_n(y)| + |f_n(y) - f(y)| \\ &\leq |f(x) - f_n(x)| + C|x - y|^\alpha + |f(y) - f_n(y)| \quad (\text{since each } f_n \text{ is Hölder continuous}) \\ &\xrightarrow{n \rightarrow \infty} C|x - y|^\alpha \quad (\text{by uniform convergence}). \end{aligned}$$

Now, we must show that $\|f\|_\alpha \leq 1$. We will do this by showing that the Hölder norm is lower semicontinuous. For each distinct pair $x, y \in [0, 1]$, the functional $\rho_{x,y} : C([0, 1]) \rightarrow [0, \infty)$ given by

$$\rho_{x,y}(g) := \frac{|g(x) - g(y)|}{|x - y|^\alpha}$$

is continuous under the topology of uniform convergence (hence lower semicontinuous). Since the supremum of lower semicontinuous functions is lower semicontinuous, the Hölder seminorm $[\cdot]_\alpha$

must be lower semicontinuous. Thus since $\|\cdot\|_\infty$ is continuous (hence, lower semicontinuous), one has that the Hölder norm is lower semicontinuous. This means that

$$\|f\|_\alpha \leq \liminf_{n \rightarrow \infty} \|f_n\|_\alpha \leq 1.$$

Thus, S_α is closed. Finally, we will show equicontinuity. Let $\varepsilon > 0$. and take $\delta = \varepsilon^{1/\alpha}$. Then for any $x, y \in [0, 1]$ with $|x - y| < \delta$, one has that for all n ,

$$|f_n(x) - f_n(y)| \leq |x - y|^\alpha < \varepsilon,$$

since $[f_n]_\alpha \leq 1$ (which means we can take $C = 1$) for all n .

Problem 2019-A-I-3. For every positive integer n , let $T_n : X \rightarrow Y$ be a linear transformation with X, Y normed vector spaces. Suppose that $\|T_n\| \leq 1$. Let $\{x_\alpha\}$ be a dense subset of X . Suppose that for any α ,

$$\lim_{n \rightarrow \infty} T_n(x_\alpha) = y_\alpha.$$

for some $y_\alpha \in Y$. Suppose further that if $x = \lim_i x_{\alpha_i}$, then y_{α_i} converges to some $y_x \in Y$. Show that

$$\lim_{n \rightarrow \infty} T_n(x) = y_x.$$

Fix some arbitrary $x \in X$, and let $\varepsilon > 0$. By the density of $\{x_\alpha\}$ in X , there exists a subsequence $\{x_{\alpha_i}\}_{i=1}^\infty$ such that the following holds: there exists a positive integer I such that for all $i \geq I$, $\|x_{\alpha_i} - x\|_X \leq \varepsilon/3$ (i.e., $x_{\alpha_i} \rightarrow x$). By the hypotheses, for each $i \geq 1$, there exists a corresponding $y_{\alpha_i} \in Y$ and a positive integer N_i so that for all $n \geq N_i$, $\|T_n x_{\alpha_i} - y_{\alpha_i}\|_Y \leq \varepsilon/3$. By the final hypothesis, there exists some $y_x \in Y$ and some positive integer J such that for all $i \geq J$, $\|y_{\alpha_i} - y_x\|_Y \leq \varepsilon/3$. Set $M = \max\{I, J\}$. Then for all $n \geq N_M$, one has

$$\begin{aligned} \|T_n x - y_x\|_Y &= \|T_n x - T_n x_{\alpha_M} + T_n x_{\alpha_M} - y_{\alpha_M} + y_{\alpha_M} - y_x\|_Y \\ &\leq \|T_n(x - x_{\alpha_M})\|_Y + \|T_n x_{\alpha_M} - y_{\alpha_M}\|_Y + \|y_{\alpha_M} - y_x\|_Y \\ &\leq \|x - x_{\alpha_M}\|_Y + \|T_n x_{\alpha_M} - y_{\alpha_M}\|_Y + \|y_{\alpha_M} - y_x\|_Y \quad (\text{since } \|T_n\| \leq 1) \\ &\leq 3 \cdot \frac{\varepsilon}{3} = \varepsilon. \end{aligned}$$

Thus, since ε was arbitrary, we conclude that $T_n x \rightarrow y_x$ under the norm on Y .

Problem 2015-J-II-5. Let E be a measurable subset of the real numbers whose Lebesgue measure is finite. Let $\chi_E : \mathbb{R} \rightarrow \{0, 1\}$ be the characteristic function of E (which equals 1 on the set E , and 0 otherwise). Show that $\chi_E(x - t)$ converges to $\chi_E(x)$ in L^1 as $t \rightarrow 0$.

Let E be a finite Lebesgue measurable subset of $\mathbb{R}$, and let χ_E denote the characteristic function of E . We wish to show that

$$\int_{\mathbb{R}} |\chi_E(x) - \chi_E(x - t)| dx \xrightarrow{t \rightarrow 0} 0.$$

We begin by studying the function $\chi_E(x) - \chi_E(x - t)$. We note the following:

$$\chi_E(x) - \chi_E(x - t) = \begin{cases} 1, & x \in E \cap (E + t)^c \\ 0, & x \in E \cap (E + t) \text{ or } x \in E^c \cap (E + t)^c \\ -1, & x \in (E + t) \cap E^c \end{cases}$$

Therefore,

$$\int_{\mathbb{R}} |\chi_E(x) - \chi_E(x - t)| dx = \int_{E \cap (E + t)^c} dx + \int_{(E + t) \cap E^c} dx = \mu(E \cap (E + t)^c) + \mu((E + t) \cap E^c).$$

Hence, we need to show that each of the terms on the right side converges to zero as $t \rightarrow 0$. **!!! Complete Later !!!**

4.2 L^p Spaces**Problem 2025-A-II-4.**

- (a) Characterize L^1 -functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f * f = f$.
 (b) The same question for L^2 -functions.

Problem 2017-J-II-2. Suppose $f : [0, 1] \rightarrow \mathbb{R}$ is measurable. Suppose further that for all $g \in L^2([0, 1])$ we have that $fg \in L^2([0, 1])$. Show f is in $L^\infty([0, 1])$.

Suppose $f : [0, 1] \rightarrow \mathbb{R}$ is measurable. Define the functional $T : L^2([0, 1]) \rightarrow L^2([0, 1])$ by $Tg := fg$ for any $g \in L^2([0, 1])$. It is immediately clear that this operator is linear. We will now show that T is a closed linear operator. Let $\{g_j\}_{j=1}^\infty \subset L^2([0, 1])$ be a sequence of square-integrable functions so that $g_j \rightarrow 0$ under the L^2 norm. Further assume that Tg_j converges to some $y \in L^2([0, 1])$ under the L^2 -norm. Since convergence under the L^2 norm implies pointwise a.e. convergence of a subsequence, there exists a subsequence $\{g_{j_k}\}_{k=1}^\infty$ so that $g_{j_k} \rightarrow 0$ pointwise a.e. Since $Tg_{j_k} \rightarrow y$ under the L^2 norm, there exists a further subsequence $\{g_{j_{k_l}}\}_{l=1}^\infty$ so that $Tg_{j_{k_l}}$ converges to y pointwise a.e. and by construction, $g_{j_{k_l}}$ converges to 0 pointwise a.e. By definition of T , it then follows that $f(x)g_{j_{k_l}}(x)$ converges to 0 pointwise a.e. so that $y = 0$ a.e. Thus in fact, we have shown that the graph of T , $\Gamma(T)$, is closed. By the closed graph theorem, it follows that T is continuous, and thus bounded. Therefore, we conclude that f must be bounded, and hence, must be in $L^\infty([0, 1])$.

Problem F-6-1-2. Prove Theorem 6.8.

Theorem 6.8 states the following

6.8 Theorem.

- If f and g are measurable functions on X , then $\|fg\|_1 \leq \|f\|_1 \|g\|_\infty$. If $f \in L^1$ and $g \in L^\infty$, $\|fg\|_1 = \|f\|_1 \|g\|_\infty$ iff $|g(x)| = \|g\|_\infty$ a.e. on the set where $f(x) \neq 0$.
- $\|\cdot\|_\infty$ is a norm on L^∞ .
- $\|f_n - f\|_\infty \rightarrow 0$ iff there exists $E \in \mathcal{M}$ such that $\mu(E^c) = 0$ and $f_n \rightarrow f$ uniformly on E .
- L^∞ is a Banach space.
- Simple functions are dense in L^∞ .

- a. Let f, g be measurable functions on X . Since $|g(x)| \leq \|g\|_\infty$ a.e., one has

$$\|fg\|_1 = \int_X |fg| d\mu \leq \int_X |f| \|g\|_\infty d\mu = \|f\|_1 \|g\|_\infty.$$

Now suppose $|g(x)| = \|g\|_\infty$ a.e. on the set $E = \{x : f(x) \neq 0\}$. Then

$$\|fg\|_1 = \int_X |fg| d\mu = \int_E |f| \|g\|_\infty d\mu = \|f\|_1 \|g\|_\infty.$$

Now suppose $\|fg\|_1 = \|f\|_1 \|g\|_\infty$ on E . I.e., one has

$$\int_X |fg| d\mu = \int_E |f| \|g\|_\infty d\mu = \int_E |f| \|g\|_\infty d\mu \implies \int_E |f| (|g| - \|g\|_\infty) d\mu = 0.$$

We note that on E , $|f| \not\equiv 0$. Moreover, $|g| - \|g\|_\infty \leq 0$ on E . Thus, the integral is zero if and only if $|g| - \|g\|_\infty = 0$ a.e. on E so that $|g(x)| = \|g\|_\infty$ a.e. on E .

- b. Recall that we defined the following

$$\|f\|_\infty = \inf\{a \geq 0 : |f(x)| \leq a \text{ a.e.}\}.$$

Thus it follows immediately that for any $f \in L^\infty$, $\|f\|_\infty \geq 0$. Now suppose $f(x) = 0$ a.e. Then $|f(x)| = 0$ a.e. so that $\|f\|_\infty = 0$. On the other hand, if $\|f\|_\infty = 0$, then $0 \leq |f(x)| \leq 0$ for a.e. x so that

$f(x) = 0$ a.e. Now we will show that the essential supremum is homogeneous. If $\lambda = 0$, the proof is immediate. So suppose $\lambda \neq 0$. Let $\mathcal{A} = \{a \geq 0 : |f(x)| \leq a \text{ a.e.}\}$, and let $\mathcal{B} = \{b \geq 0 : |\lambda f(x)| \leq b \text{ a.e.}\}$. Suppose $|f(x)| \leq a$ a.e. Then it follows that for any $\lambda \in \mathbb{C} \setminus \{0\}$, $|\lambda f(x)| = |\lambda| |f(x)| \leq |\lambda| a$ a.e. This shows that $|\lambda| \mathcal{A} \subset \mathcal{B}$ (which means that $|\lambda| \|f\|_\infty = |\lambda| \inf \mathcal{A} \geq \|\lambda f\|_\infty = \inf \mathcal{B}$). Now suppose $|\lambda f(x)| \leq b$ a.e. so that $|f(x)| \leq b/|\lambda|$ a.e. This means that $(1/|\lambda|) \mathcal{B} \subset \mathcal{A}$ so that $(1/|\lambda|) \|\lambda f\|_\infty \geq \|f\|_\infty \iff \|\lambda f\|_\infty \geq |\lambda| \|f\|_\infty$. Thus, one has $\|\lambda f\|_\infty = |\lambda| \|f\|_\infty$. Now we will show that the triangle inequality holds. Let $f, g \in L^\infty$. Further define the sets $\mathcal{A} = \{a \geq 0 : |f(x) + g(x)| \leq a \text{ a.e.}\}$, $\mathcal{F} = \{b \geq 0 : |f(x)| \leq b \text{ a.e.}\}$, and $\mathcal{G} = \{c \geq 0 : |g(x)| \leq c \text{ a.e.}\}$. Let $b \in \mathcal{F}, c \in \mathcal{G}$. Then a.e. x ,

$$|f(x) + g(x)| \leq |f(x)| + |g(x)| \leq b + c,$$

so that $\mathcal{F} + \mathcal{G} \subset \mathcal{A}$. Taking the infimum of both sides, one has $\|f + g\|_\infty = \inf \mathcal{A} \leq \inf \mathcal{F} + \inf \mathcal{G} = \|f\|_\infty + \|g\|_\infty$. Thus, the proof that $\|\cdot\|_\infty$ is a norm concludes.

- c. Suppose there exists $E \in \mathcal{M}$ so that $\mu(E^c) = 0$ and $f_n \rightarrow f$ uniformly on E . I.e., we have that $f_n \rightarrow f$ uniformly a.e. This means that for every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ large enough so that for all $n \geq N$ and almost all $x \in X$, $|f_n(x) - f(x)| \leq \varepsilon$. By definition of the essential supremum, it follows that for all $n \geq N$, $\|f_n - f\|_\infty \leq \varepsilon$. Thus we conclude that $\|f_n - f\|_\infty \rightarrow 0$. Now suppose that $\|f_n - f\|_\infty \rightarrow 0$. This means that for every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ large enough so that for all $n \geq N$, $\|f_n - f\|_\infty \leq \varepsilon$. For each fixed $n \geq N$, let Z_n denote the set $\{x : |f_n(x) - f(x)| > \|f_n - f\|_\infty\}$; note that $\mu(Z_n) = 0$. Let $E_n = X \setminus Z_n$. One has that $f_n \rightarrow f$ uniformly on $E := \bigcap_{n \geq N} E_n$. On the other hand,

$$\mu\left(\left(\bigcap_{n \geq N} E_n\right)^c\right) = \mu\left(\bigcup_{n \geq N} Z_n\right) = 0,$$

by countable subadditivity.

- d. Let $\{f_n\}_{n \geq 1} \subseteq L^\infty(X)$ be a Cauchy sequence. It suffices to show that $\{f_n\}$ converges to some $f \in L^\infty$. by definition of Cauchy sequences, for every $\varepsilon > 0$, there exists a positive integer N so that for all $n, m \geq N$, $\|f_n - f_m\|_\infty \leq \varepsilon$. For each pair $(n, m) \geq (N, N)$, let $E_{n,m}$ denote the set $\{x : |f_n(x) - f_m(x)| > \|f_n - f_m\|_\infty\}$, and let $E = \bigcup_{n,m=1}^\infty E_{n,m}$. By countable subadditivity, one has $\mu(E) = 0$. On the other hand, for every $x \in E^c$, the sequence $\{f_n(x)\}_{n=1}^\infty$ is a Cauchy sequence in $\mathbb{C}$ and thus converges to some value. Define the function $f : X \rightarrow \mathbb{C}$ as follows

$$f(x) = \begin{cases} \lim_{n \rightarrow \infty} f_n(x), & x \in E^c \\ 0, & x \in E. \end{cases}$$

First, we must show that $f \in L^\infty$. Since $\{f_n\}$ is a Cauchy sequence, it follows that $\{\|f_n\|_\infty\}_{n \geq 1}$ is a Cauchy sequence in $\mathbb{R}$, and thus must be bounded above by some finite constant $M > 0$. Therefore, for every $x \in E^c$, one has

$$|f(x)| = \lim_{n \rightarrow \infty} |f_n(x)| \leq \lim_{n \rightarrow \infty} \|f_n\|_\infty = M,$$

so that $\|f\|_\infty \leq M < \infty$. Thus, $f \in L^\infty$. Now we need to show that $f_n \rightarrow f$ under the L^∞ norm. Let $\varepsilon > 0$. Let $N \in \mathbb{N}$ so that for all $n, m \geq N$, $\|f_n - f_m\|_\infty < \varepsilon$. Then for every $x \in E^c$,

$$|f_n(x) - f(x)| = \lim_{m \rightarrow \infty} |f_n(x) - f_m(x)| \leq \varepsilon.$$

Since this inequality holds for all $n \geq N$, $x \in E^c$, we conclude that $\|f_n - f\|_\infty \leq \varepsilon$, and thus $f_n \rightarrow f$ under the L^∞ -norm.

- e. First, we will start by showing that simple functions are contained in L^∞ . To do so, we will look at *characteristic functions*. Since for every $x \in X$, $|\chi_E(x)| \leq 1$ for any measurable subset $E \subseteq X$, it follows that χ_E is bounded everywhere on X , and hence, is contained in L^∞ . Thus, it follows that every simple function is contained in L^∞ . Now we will show that the set of simple functions in L^∞ is dense in this space. Fix an arbitrary $f \in L^\infty$. By Theorem 2.10, there exists a sequence $\{\phi_n\}$ of simple functions such that $0 \leq |\phi_1| \leq |\phi_2| \leq \dots \leq |f|$, $\phi_n \rightarrow f$ pointwise, and $\phi_n \rightarrow f$ uniformly on

any set on which f is bounded. Since f is essentially bounded, there exists some measurable set $E \in \mathcal{M}$ so that $\mu(E^c) = 0$ and f is bounded on E . Thus, $\phi_n \rightarrow f$ uniformly on E . But by part (c), it follows that $\|\phi_n - f\|_\infty \rightarrow 0$; i.e., for every $\varepsilon > 0$, there exists some positive integer N so that for all $n \geq N$, $\|\phi_n - f\|_\infty \leq \varepsilon$. Thus, this proves density of simple functions in L^∞ .

Problem F-6-1-3. If $1 \leq p < r \leq \infty$, $L^p \cap L^r$ is a Banach space with norm $\|f\| = \|f\|_p + \|f\|_r$, and if $p < q < r$, the inclusion map $L^p \cap L^r \rightarrow L^q$ is continuous.

Fix $1 \leq p < r \leq \infty$, and define the map $\|\cdot\| : L^p \cap L^r \rightarrow [0, \infty)$ as given above. First, we will begin by showing that this map is a well-defined norm. Let $f \in L^p \cap L^r$, and assume $\|f\| = \|f\|_p + \|f\|_r = 0$. Then since $\|f\|_p, \|f\|_r \geq 0$, one must have $\|f\|_p = 0$ and $\|f\|_r = 0$. But by positive definiteness of the L^p norms, one must have $f = 0$ a.e. Now we will show homogeneity. Let $\lambda \in \mathbb{C}$. Then

$$\begin{aligned} \|\lambda f\| &= \|\lambda f\|_p + \|\lambda f\|_r \\ &= |\lambda|(\|f\|_p + \|f\|_r) \\ &= |\lambda| \|f\|. \end{aligned}$$

Finally, we will show that the triangle inequality is satisfied. Let $f, g \in L^p \cap L^r$. Then

$$\begin{aligned} \|f + g\| &= \|f + g\|_p + \|f + g\|_r \\ &\leq \|f\|_p + \|g\|_p + \|f\|_r + \|g\|_r \\ &= \|f\| + \|g\|. \end{aligned}$$

Thus $\|\cdot\|$ is a norm on $L^p \cap L^r$. Now we will show that $L^p \cap L^r$ is a Banach space under this norm. Let $\{f_n\}_{n \geq 1} \subset L^p \cap L^r$ be a Cauchy sequence. Then by construction of $\|\cdot\|$, one has that $\{f_n\}$ is a Cauchy sequence in L^p and in L^r . Thus since each of these spaces are Banach, $f_n \rightarrow f$ in L^p and $f_n \rightarrow g$ in L^r . Our first task is to show that $f = g$ a.e. Since $\{f_n\}$ converges to f in L^p , there exists a subsequence $\{f_{n_k}\}$ that converges to f pointwise a.e. But since $\{f_{n_k}\} \subset \{f_n\}$ and the latter converges to g in L^r , there exists a further subsequence $\{f_{n_{k_j}}\}$ that converges pointwise to g a.e. But since $\{f_{n_{k_j}}\} \rightarrow f$ pointwise a.e., every subsequence must also converge to f pointwise a.e. Thus by uniqueness of limits, we must have $f = g$ a.e. This shows that $f, g \in L^p \cap L^r$. Thus,

$$\|f_n - f\| = \|f_n - f\|_p + \|f_n - f\|_r \xrightarrow{n \rightarrow \infty} 0.$$

This concludes the proof that $(L^p \cap L^r, \|\cdot\|)$ is Banach. Now let $p < q < r$. By Proposition 6.10, one has $L^p \cap L^r \subset L^q$ so that the inclusion $\iota : L^p \cap L^r \hookrightarrow L^q$ is well-defined. To show that ι is continuous, it suffices to show that it is bounded. By Proposition 6.10 and Young's inequality (in that order), one has

$$\|\iota(f)\|_q \leq \|f\|_p^\lambda \|f\|_r^{1-\lambda} \leq \lambda \|f\|_p + (1-\lambda) \|f\|_r \leq \|f\|_p + \|f\|_r = \|f\|,$$

where λ is defined by the relation $q^{-1} = \lambda p^{-1} + (1-\lambda)r^{-1}$ and $\lambda \in (0, 1)$. Thus, since $f \in L^p \cap L^r$ was arbitrary, ι is bounded, and thus continuous.

Problem F-6-1-7. If $f \in L^p \cap L^\infty$ for some $p < \infty$, so that $f \in L^q$ for all $q > p$, then $\|f\|_\infty = \lim_{q \rightarrow \infty} \|f\|_q$.

Fix some $f \in L^p \cap L^\infty$ for some $p < \infty$. By Proposition 6.10, one has that $f \in L^q$ for all $q > p$. Let $\varepsilon > 0$, and let $A = \{x : |f(x)| > \|f\|_\infty - \varepsilon\}$; by definition of the essential supremum, A has positive measure; note that since $f \in L^q$, A must have finite measure. Then one has

$$\|f\|_q^q = \int_X |f(x)|^q d\mu \geq \int_A |f(x)|^q d\mu \geq \mu(A)(\|f\|_\infty - \varepsilon)^q.$$

Since both sides are finite, one has $\|f\|_q \geq \mu(A)^{1/q}(\|f\|_\infty - \varepsilon)$. Since $\varepsilon > 0$ was arbitrary, one has $\|f\|_q \geq \mu(A)^{1/q} \|f\|_\infty$. Thus taking the limit $q \rightarrow \infty$, one has

$$\lim_{q \rightarrow \infty} \|f\|_q \geq \|f\|_\infty.$$

On the other hand, set $\lambda \in (0, 1)$ so that $q^{-1} = \lambda p^{-1}$. Then by Proposition 6.10, one has

$$\|f\|_q \leq \|f\|_p^\lambda \|f\|_\infty^{1-\lambda} = \|f\|_p^{p/q} \|f\|_\infty^{1-p/q}.$$

Thus, taking the limit $q \rightarrow \infty$ on both sides yields,

$$\lim_{q \rightarrow \infty} \|f\|_q \leq \lim_{q \rightarrow \infty} \|f\|_p^{p/q} \|f\|_\infty^{1-p/q} = \|f\|_\infty.$$

Thus the proof concludes.

Problem F-6-1-10. Suppose $1 \leq p < \infty$. If $f_n, f \in L^p$, and $f_n \rightarrow f$ a.e., then $\|f_n - f\|_p \rightarrow 0$ iff $\|f_n\|_p \rightarrow \|f\|_p$. (Use Exercise 20 in §2.3.)

Suppose $1 \leq p < \infty$, $f_n, f \in L^p$, and $f_n \rightarrow f$ a.e. Exercise 20 in §2.3 states the following

(Generalized Dominated Convergence Theorem) If $f_n, g_n, f, g \in L^1$, $f_n \rightarrow f$ and $g_n \rightarrow g$ a.e., $|f_n| \leq g_n$, and $\int g_n \rightarrow \int g$, then $\int f_n \rightarrow \int f$.

First assume that $\|f_n - f\|_p \rightarrow 0$. Then one has

$$\|f_n\|_p \leq \|f_n - f\|_p + \|f\|_p.$$

Thus taking the limit as $n \rightarrow \infty$ on both sides, one has

$$\lim_{n \rightarrow \infty} \|f_n\|_p = \lim_{n \rightarrow \infty} \|f_n - f\|_p + \|f\|_p = \|f\|_p.$$

Now suppose $\|f_n\|_p \rightarrow \|f\|_p$. We will use Exercise 20 from §2.3. Since $f_n, f \in L^p$, one has that $|f_n - f|^p, 0, 2^{p-1}(|f_n|^p + |f|^p), 2^p|f|^p \in L^1$, $|f_n - f|^p \leq 2^{p-1}(|f_n|^p + |f|^p)$ for all n , and $\int 2^{p-1}(|f_n|^p + |f|^p) \rightarrow \int 2^p|f|^p$ (by the hypothesis). Thus by the Generalized Dominated Convergence Theorem, we conclude that

$$\|f_n - f\|_p^p = \int |f_n - f|^p \rightarrow \int 0 = 0 \implies \|f_n - f\|_p \rightarrow 0.$$

Problem F-6-1-14. If $g \in L^\infty$, the operator T defined by $Tf = fg$ is bounded on L^p for $1 \leq p \leq \infty$. Its operator norm is at most $\|g\|_\infty$, with equality if μ is semifinite.

Fix an arbitrary $g \in L^\infty$, $1 \leq p \leq \infty$, and define the operator T on L^p by $Tf := fg$. We begin by showing that for any $f \in L^p$, $fg \in L^p$. We compute (for $1 \leq p < \infty$),

$$\|fg\|_p^p = \int_X |f|^p |g|^p d\mu \leq \int_X \|g\|_\infty^p |f|^p d\mu = \|g\|_\infty^p \|f\|_p^p < \infty \implies \|fg\|_p < \infty.$$

The case for $p = \infty$ can be shown trivially. Thus, one has that for any $f \in L^p$,

$$\|Tf\|_p = \|fg\|_p \leq \|g\|_\infty \|f\|_p,$$

so that T is a bounded operator. Indeed we see that the operator norm of T is at most $\|g\|_\infty$. Now assume that μ is semifinite. Let $\varepsilon > 0$, and first assume that $1 \leq p < \infty$. By semifiniteness, there exists a measurable set $E \in \mathcal{M}$ of finite positive measure so that $|g(x)| > \|g\|_\infty - \varepsilon$ on E . Then one has that for all $f \in L^p$,

$$\|fg\|_p^p = \int_X |f|^p |g|^p d\mu \geq \int_E |f|^p (\|g\|_\infty - \varepsilon)^p d\mu.$$

Let

$$f = \frac{\chi_E \operatorname{sgn}(g)}{\mu(E)^{1/p}} \implies \|f\|_p = 1.$$

For this particular choice of f , one has

$$\|Tf\|_p^p \geq (\|g\|_\infty - \varepsilon)^p \implies \|Tf\|_p \geq \|g\|_\infty - \varepsilon.$$

Since this is true for all $\varepsilon > 0$, $\|Tf\|_p \geq \|g\|_\infty$. Thus in fact, one has

$$\|T\| = \sup_{\|h\|_p \leq 1} \|Th\| \geq \|Tf\|_p \geq \|g\|_\infty.$$

Therefore, in this case, we conclude that the operator norm of T is exactly $\|g\|_\infty$. Now consider the case $p = \infty$, and let $f = \chi_E \operatorname{sgn}(g)$. Then, one has

$$|f(x)g(x)| = |g(x)| > \|g\|_\infty - \varepsilon \implies \|fg\|_\infty \geq \|g\|_\infty - \varepsilon.$$

Therefore, one has

$$\|T\| = \sup_{\|h\|_\infty \leq 1} \|Th\|_\infty \geq \|fg\|_\infty \geq \|g\|_\infty - \varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, we conclude that $\|T\| \geq \|g\|_\infty$. Thus the proof concludes in both cases by using the upper bound we proved above.

Problem F-6-1-4. If $1 \leq p < r \leq \infty$, $L^p + L^r$ is a Banach space with norm $\|f\| = \inf\{\|g\|_p + \|h\|_r : f = g + h\}$, and if $p < q < r$, the inclusion map $L^q \hookrightarrow L^p + L^r$ is continuous.

Let $1 \leq p < r \leq \infty$, and let $\|\cdot\| : L^p + L^r \rightarrow [0, \infty)$ be the map defined by

$$\|\cdot\| := \inf\{\|g\|_p + \|h\|_r : \cdot = g + h\}.$$

First, we will show that the above map is a norm. We begin by showing positive definiteness. Suppose $f \in L^p + L^r$ with $f = 0$ a.e. Then the sum $0 + 0$ is a decomposition of f into a sum of an L^p function with an L^r function; i.e., $f = 0 + 0$ a.e. But then

$$\|f\| = \inf\{\|g\|_p + \|h\|_r : f = g + h\} \leq \|0\|_p + \|0\|_r = 0 + 0 = 0.$$

Now assume $\|f\| = 0$. By definition of infimum, for every $j \geq 1$, there exists a decomposition $g_j + h_j$ of f so that $\|g_j\|_p + \|h_j\|_r \leq j^{-1}$. In particular, one has $\|g_j\|_p \leq j^{-1}$ and $\|h_j\|_r \leq j^{-1}$. Consider the sequence $\{g_j\}_{j=1}^\infty$. By the above reasoning, it follows that the sequence converges to $0 = \|0\|_p$. Thus since $\{g_j\}$ converges to 0 under the L^p norm, it follows that there exists a subsequence $\{g_{j_k}\}_{k=1}^\infty$ that converges to 0 pointwise a.e. Let $\{h_{j_k}\}_{k=1}^\infty$ be the corresponding subsequence. Then since $\{h_j\} \rightarrow 0$ under the L^r norm, $\{h_{j_k}\}_{k=1}^\infty$ converges to 0 under the L^r norm. Thus there exists a subsequence $\{h_{j_{k_m}}\}_{m=1}^\infty$ that converges pointwise to 0 a.e. By construction, $\{g_{j_{k_m}}\}_{m=1}^\infty$ also converges pointwise to 0 a.e. Thus, since $f = g_{j_{k_m}} + h_{j_{k_m}}$, taking the limit $m \rightarrow \infty$ yields that $f = 0$ a.e. Thus, we conclude that $\|\cdot\|$ is positive definite. Now we will show that $\|\cdot\|$ is homogeneous. If $\lambda \in \mathbb{C}$ is zero, the proof follows trivially. Thus let $\lambda \in \mathbb{C} \setminus \{0\}$, $\mathcal{A} = \{\|g\|_p + \|h\|_r : f = g + h\}$ and $\mathcal{B} = \{\|i\|_p + \|j\|_r : \lambda f = i + j\}$. If $g + h$ is a decomposition of f , then $\lambda g + \lambda h$ is a decomposition of λf . Thus, we have $|\lambda|\mathcal{A} \subseteq \mathcal{B}$ so that $\|\lambda f\| = \inf \mathcal{B} \leq |\lambda| \inf \mathcal{A} = |\lambda| \|f\|$. On the other hand, suppose $i + j$ is a decomposition of λf . Then $(1/\lambda)(i + j)$ is a decomposition of f so that $1/|\lambda| \mathcal{B} \subseteq \mathcal{A}$, which implies $\|f\| \leq \frac{1}{|\lambda|} \inf \mathcal{B} = \frac{1}{|\lambda|} \|\lambda f\| \implies \|\lambda f\| \geq |\lambda| \|f\|$. Thus we conclude that $\|\lambda f\| = |\lambda| \|f\|$ so that $\|\cdot\|$ is homogeneous. Finally, we will show that the triangle inequality is satisfied. Let $f_1, f_2 \in L^p + L^r$, and define the following sets

$$\begin{aligned} \mathcal{A} &= \{\|g_1\|_p + \|h_1\|_r : f_1 = g_1 + h_1\} \\ \mathcal{B} &= \{\|g_2\|_p + \|h_2\|_r : f_2 = g_2 + h_2\} \\ \mathcal{C} &= \{\|g_3\|_p + \|h_3\|_r : f_1 + f_2 = g_3 + h_3\}. \end{aligned}$$

Let $g_1 + h_1$ and $g_2 + h_2$ be decompositions of f_1 and f_2 , respectively. Then $(g_1 + g_2) + (h_1 + h_2)$ is a decomposition of $f + g$ so that $\mathcal{A} + \mathcal{B} \subset \mathcal{C}$. This implies that

$$\|f + g\| = \inf \mathcal{C} \leq \inf(\mathcal{A} + \mathcal{B}) = \inf \mathcal{A} + \inf \mathcal{B} = \|f\| + \|g\|.$$

Thus, we conclude that $\|\cdot\|$ is indeed a valid norm on $L^p + L^r$. Now we need to show that $(L^p + L^r, \|\cdot\|)$ is Banach. Let $\{f_n\}_{n=1}^\infty$ be a Cauchy sequence under this norm.

4.3 Distribution Theory

Problem F-9-1-1. Suppose that $f_1, f_2, \dots$, and f are in $L^1_{\text{loc}}(U)$. The conditions in (a) and (b) below imply that $f_n \rightarrow f$ in $\mathcal{D}'(U)$, but the condition in (c) does not.

- a. $f_n \in L^p(U)$ ($1 \leq p \leq \infty$) and $f_n \rightarrow f$ in the L^p norm or weakly in L^p .
- b. For all n , $|f_n| \leq g$ for some $g \in L^1_{\text{loc}}(U)$, and $f_n \rightarrow f$ a.e.
- c. $f_n \rightarrow f$ pointwise.

- a. First assume that $f_n \in L^p(U)$, and further assume that $f_n \rightarrow f$ in the L^p norm. Fix an arbitrary test function $\phi \in C_c^\infty(U)$; let K be a compact subset of U containing the support of ϕ . Then one has

$$\begin{aligned} \left| \int_U f_n \phi - \int_U f \phi \right| &= \left| \int_K (f_n - f) \phi \right| \\ &\leq \int_K |f_n - f| |\phi| \\ &\leq \|f_n - f\|_p \|\phi\|_q \quad (\text{by Hölder's Inequality}) \xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

so that by arbitrariness of ϕ , $f_n \rightarrow f$ in $\mathcal{D}'(U)$. Now suppose that $f_n \rightarrow f$ weakly in L^p for $1 \leq p < \infty$. I.e., if q is the conjugate exponent of p , then $\int_U f_n g \rightarrow \int_U f g$ for all $g \in L^q$. We know that $C_c^\infty(U) \subset L^q(U)$. Thus it follows immediately that for every $\phi \in C_c^\infty(U)$, $\int_U f_n \phi \rightarrow \int_U f \phi$ so that $f_n \rightarrow f$ in $\mathcal{D}'(U)$.

- b. Assume that $f_n \rightarrow f$ pointwise a.e. and assume there exists some $g \in L^1_{\text{loc}}(U)$ so that $|f_n| \leq g$ for all n . Fix some test function $\phi \in C_c^\infty(U)$ and let $K \subset U$ be a compact subset that contains the support of ϕ . Then since each $f_n, g \in L^1(K)$, one has by the dominated convergence theorem that

$$\lim_{n \rightarrow \infty} \int_U f_n \phi = \lim_{n \rightarrow \infty} \int_K f_n \phi = \int_K \lim_{n \rightarrow \infty} f_n \phi = \int_K f \phi = \int_U f \phi.$$

Thus, $f_n \rightarrow f$ in $\mathcal{D}'(U)$.

- c. Consider the sequence $\{n\chi_{(0,1/n)}\}_{n \in \mathbb{N}}$. Pointwise, the sequence converges to the zero distribution a.e., but as distributions, the sequence converges to the Dirac delta distribution supported at the origin, which is not identically zero. Thus $f_n \rightarrow f$ pointwise does not guarantee convergence on the level of distributions.

Problem F-9-1-2. The product rule for derivatives is valid for products of smooth functions and distributions.

Let $U \subseteq \mathbb{R}^n$ be an open domain, $\ell \in \mathcal{D}'(U)$, and fix arbitrary $f, \phi \in C_c^\infty(U) = \mathcal{D}(U)$. We will solve this problem in two steps.

(Step 1): Let $|\alpha| = 1$; there exists some $i \in \{1, 2, \dots, n\}$ so that $D^\alpha = \partial_i$. In particular, we must have

$$\begin{aligned} \langle \partial_i(f\ell), \phi \rangle &= -\langle f\ell, \partial_i \phi \rangle = -\langle \ell, f \partial_i \phi \rangle = -\langle \ell, \partial_i(f\phi) \rangle + \langle \ell, \partial_i(f)\phi \rangle \\ &= \langle f \partial_i \ell, \phi \rangle + \langle \partial_i(f)\ell, \phi \rangle = \langle f \partial_i \ell + \partial_i f \ell, \phi \rangle. \end{aligned} \tag{1}$$

(Step 2): So now consider the case when $|\alpha| = m$ for some $m > 1$. Then there exist indices $\{i_j\}_{j=1}^m \subset$

$\{1, \dots, n\}$ so that $D^\alpha = \partial_{i_1} \cdots \partial_{i_m}$. Then,

$$\begin{aligned}
 \langle \partial_{i_1} \cdots \partial_{i_m} (f\ell), \phi \rangle &= (-1)^m \langle f\ell, \partial_{i_1} \cdots \partial_{i_m} \phi \rangle \\
 &= (-1)^m \langle \ell, f \partial_{i_1} \cdots \partial_{i_m} \phi \rangle \\
 &= (-1)^m \langle \ell, f \partial_{i_1} \psi \rangle & (\psi &:= \partial_{i_2} \cdots \partial_{i_m} \phi) \\
 &= (-1)^{m+1} \langle f \partial_{i_1} \ell + \partial_{i_1} f \ell, \partial_{i_2} \phi \rangle & (\phi &:= \partial_{i_3} \cdots \partial_{i_m} \phi) \\
 &= (-1)^{m+1} [\langle \partial_{i_1} \ell, f \partial_{i_2} \phi \rangle + \langle \ell, (\partial_{i_1} f) \partial_{i_2} \phi \rangle] \\
 &= (-1)^{m+2} [\langle f \partial_{i_1} \partial_{i_2} \ell, \partial_{i_3} \phi \rangle + \langle (\partial_{i_2} f) (\partial_{i_1} \ell), \partial_{i_3} \phi \rangle \\
 &\quad + \langle (\partial_{i_1} f) (\partial_{i_2} \ell), \partial_{i_3} \phi \rangle + \langle (\partial_{i_1} \partial_{i_2} f) \ell, \partial_{i_3} \phi \rangle] \\
 &\vdots \\
 &= \sum_{|\beta| \leq m} \binom{\alpha}{\beta} \langle D^\beta (f) D^{\alpha-\beta} \ell, \phi \rangle,
 \end{aligned} \tag{2}$$

where $\Phi := \partial_{i_4} \cdots \partial_{i_m} \phi$, and we repeated the same types of calculation to get to the last line. Hence, since ϕ was arbitrary, we conclude that

$$D^\alpha (f\ell) = \sum_{|\beta| \leq m} \binom{\alpha}{\beta} (D^\beta f) (D^{\alpha-\beta} \ell)$$

Problem F-9-1-3. On $\mathbb{R}$, if $\psi \in C^\infty$ then $\psi \delta^{(k)} = \sum_0^k (-1)^j \binom{k}{j} \psi^{(j)}(0) \delta^{(k-j)}$, where the superscripts denote derivatives.

Let $\psi \in C^\infty(\mathbb{R})$, and let $\delta := \delta_0 \in \mathcal{D}'(\mathbb{R})$ be the Dirac delta function. Let us fix arbitrary $\phi \in C_c^\infty(\mathbb{R})$. For some $k \geq 1$, we compute

$$\begin{aligned}
 \langle \psi \delta^k, \phi \rangle &= (-1)^k \langle \phi, \sum_{j=0}^k \binom{k}{j} \psi^{(j)} \phi^{(k-j)} \rangle \\
 &= (-1)^k \sum_{j=0}^k \binom{k}{j} \psi^{(j)}(0) \phi^{(k-j)}(0) \\
 &= \sum_{j=0}^k \binom{k}{j} \psi^{(j)}(0) (-1)^k (-1)^{k-j} \langle \delta^{(k-j)}, \phi \rangle \\
 &= \sum_{j=0}^k (-1)^j \binom{k}{j} \psi^{(j)}(0) \langle \delta^{(k-j)}, \phi \rangle.
 \end{aligned} \tag{3}$$

Since ϕ was arbitrary, we conclude that

$$\psi \delta^{(k)} = \sum_{j=0}^k (-1)^j \binom{k}{j} \psi^{(j)}(0) \delta^{(k-j)}.$$

Problem F-9-1-5. Suppose that f is continuously differentiable on $\mathbb{R}$ except at $x_1, \dots, x_m$, where f has jump discontinuities, and that its pointwise derivative df/dx (defined except at the x_j 's) is in $L^1_{\text{loc}}(\mathbb{R})$. Then the distribution derivative f' of f is given by $f' = (df/dx) + \sum_1^m [f(x_{j+}) - f(x_{j-})] \tau_{x_j} \delta$.

Suppose that f is continuously differentiable on $\mathbb{R}$, except at the points $x_1, \dots, x_m$, where f has jump discontinuities. Let df/dx , defined except at the x_j 's denote its pointwise derivative. Let f' denote the distribution derivative of f . Fix an arbitrary test function $\phi \in C_c^\infty(\mathbb{R})$, and let $[x_0, x_{m+1}]$ be a compact

set so that $\text{supp } \phi \not\subseteq K$, where $x_0 < x_1 < x_2 < \dots < x_{m+1}$. We begin by looking at $\langle f, \phi' \rangle := \int_{\mathbb{R}} f(x) \phi'(x) dx$. We have the following:

$$\begin{aligned} \langle f, \phi' \rangle &= \int_{\mathbb{R}} f(x) \phi'(x) dx = \int_{x_0}^{x_{m+1}} f(x) \phi'(x) dx \\ &= \sum_{j=0}^m \int_{x_j}^{x_{j+1}} f(x) \phi'(x) dx \\ &= \sum_{j=0}^m \left[f(x) \phi(x) \Big|_{x_j}^{x_{j+1}} - \int_{x_j}^{x_{j+1}} \frac{df}{dx} \phi(x) dx \right] \quad (\text{by integration by parts}) \\ &= \sum_{j=0}^m [f(x_{j+1}-) \phi(x_{j+1}) - f(x_j+) \phi(x_j)] - \sum_{j=0}^m \int_{x_j}^{x_{j+1}} \frac{df}{dx} \phi(x) dx. \end{aligned}$$

Telescoping the sum and using the fact that $\phi(x_{m+1}) = \phi(x_0) = 0$ since $\text{supp } \phi \not\subseteq [x_0, x_{m+1}]$, one has

$$\begin{aligned} \langle f, \phi' \rangle &= - \left\langle \sum_{j=1}^m [f(x_j+) - f(x_j-)] \tau_{x_j} \delta, \phi \right\rangle - \left\langle \frac{df}{dx}, \phi \right\rangle \\ &= - \left\langle \frac{df}{dx} + \sum_{j=1}^m [f(x_j+) - f(x_j-)] \tau_{x_j} \delta, \phi \right\rangle. \end{aligned}$$

By definition of the distribution derivative, we conclude that the distribution derivative of f is given by,

$$f' = \frac{df}{dx} + \sum_{j=1}^m [f(x_j+) - f(x_j-)] \tau_{x_j} \delta.$$

Problem F-9-1-6. If f is absolutely continuous on compact subsets of an interval $U \subset \mathbb{R}$, the distribution derivative $f' \in \mathcal{D}'(U)$ coincides with the pointwise (a.e.-defined) derivative of f .

Let f be absolutely continuous on compact subsets of an interval $U \subset \mathbb{R}$ (in particular, since f is continuous on compact subsets of U , one has $f \in L^1_{\text{loc}}(U)$). Let f' denote the distribution derivative of f , and let df/dx be the pointwise (a.e.-defined) derivative of f . Fix an arbitrary test function $\phi \in C_c^\infty$ and let $K = [a, b]$ be a compact subset of U so that $\text{supp } \phi \not\subseteq [a, b]$. Then one has

$$\langle f, \phi' \rangle = \int_a^b f(x) \phi'(x) dx = f(x) \phi(x) \Big|_a^b - \int_a^b \frac{df}{dx} \phi(x) dx,$$

where the second equality follows from integration by parts. Since $\text{supp } \phi \not\subseteq [a, b]$, one has $\phi(a) = \phi(b) = 0$. Thus the first term vanishes. It remains to be seen that df/dx is locally integrable. This follows immediately: since f is absolutely continuous, its pointwise a.e.-defined derivative must be integrable, and hence, must be contained in $L^1_{\text{loc}}(U)$ so that $df/dx \in \mathcal{D}'(U)$. Therefore, by definition of the distribution derivative, for every test function $\phi \in C_c^\infty(U)$, one has

$$\langle f', \phi \rangle = - \langle f, \phi' \rangle = \left\langle \frac{df}{dx}, \phi \right\rangle,$$

whence $f' = df/dx$ a.e.

5 Complex Analysis

5.1 Holomorphic Functions

Problem 2026-J-II-4. Let U be an open subset of $\mathbb{C}$. Let $(f_j(z))$ be a collection of holomorphic functions on U that converges uniformly on every compact subset. Carefully prove that the pointwise limit is holomorphic.

We will use Morera's theorem for this proof. Recall the statement of Morera's Theorem:

Theorem. (Morera) A continuous, complex-valued function f defined on an open set D in the complex plane that satisfies

$$\oint_{\gamma} f(z) dz = 0$$

for every closed piecewise differentiable curve γ in D is holomorphic on D .

Assume the hypotheses, and define f on U to be $f(z) = \lim_{j \rightarrow \infty} f_j(z)$ for all $z \in U$. Since each f_j is holomorphic (and thus continuous), and the uniform limit of continuous functions is continuous, f is continuous. Then for any closed piecewise differentiable curve γ in U ,

$$\oint_{\gamma} f(z) dz = \oint_{\gamma} \lim_{j \rightarrow \infty} f_j(z) dz = \lim_{j \rightarrow \infty} \oint_{\gamma} f_j(z) dz = \lim_{j \rightarrow \infty} 0 = 0,$$

where the limit comes out of the integral due to uniform on compacta convergence of the f_j , and holomorphicity of each f_j implies $\oint_{\gamma} f_j(z) dz = 0$. Thus, we conclude by Morera's Theorem that f is holomorphic in U .

5.2 Rouché's Theorem

Problem 2023-A-II-6. Find the number of solutions (counting multiplicity) to $z^8 - 5z^6 + 2z^3 - z - 1 = 0$ that lie inside the unit disk.

Let $p(z) = z^8 - 5z^7 + 2z^3 - z - 1$, and let $g(z) = -5z^6 + 2z^3$, $\Delta := \{z \in \mathbb{C} : |z| \leq 1\}$. It is straightforward to check that neither $p(z)$ nor $g(z)$ have any zeros on $\partial\Delta$. Indeed, on $\partial\Delta$, one has

$$\begin{aligned} |p(z) - g(z)| &= |z^8 - z - 1| \leq |z|^8 + |z| + |1| \\ &= 3 < 7 = |g(z)|. \end{aligned}$$

By Rouché's Theorem, it follows that $p(z)$ has the same number of zeros as $g(z)$ in Δ . We can write $g(z) = z^3(-5z^3 + 2)$. Thus, since $g(z)$ has six roots (counting multiplicity) inside the unit disk, we conclude that $p(z)$ also has only six roots inside the unit disk.

Problem 2020-A-II-1. How many roots (counted with multiplicity) does the function

$$g(z) = 6z^3 + e^z + 1$$

have in the unit disk $|z| < 1$?

Let $g(z) = 6z^3 + e^z + 1$, $p(z) = 6z^3$, and $\Delta := \{z \in \mathbb{C} : |z| \leq 1\}$. Then it is straightforward to see that neither $p(z)$ nor $g(z)$ has any roots in $\partial\Delta$. Thus, since on $\partial\Delta$ one has

$$|g(z) - p(z)| = |e^z + 1| \leq |e^z| + 1 \leq e + 1 < 6 = |p(z)|.$$

Thus, by Rouché's Theorem, $g(z)$ has the same number of roots as $p(z)$ inside the unit disk (counting multiplicity). Thus, since $p(z)$ has 3 roots inside Δ , we conclude that $g(z)$ also has 3 roots inside the unit disk.

Problem 2018-A-I-3. Show that if $c > 1$, then the function

$$f(z) = ze^{c-z} - 1$$

has precisely one root in $\Delta = \{|z| < 1\}$, and this root is real and positive.

Let $c > 1$, and define $f(z) = ze^{c-z} - 1$. Define the function $g(z) = ze^{c-z}$. Then on $\partial\Delta$, one has

$$|f(z) - g(z)| = 1 < e^{c-1} = |g(z)|,$$

so that by Rouché's Theorem, $f(z)$ has the same number of zeros as $g(z)$ inside Δ . Since the exponential function is never zero, and the polynomial z is zero exactly at the origin (with multiplicity one), $g(z)$ has exactly one zero inside the unit disk. Thus, $f(z)$ also has only one zero inside the unit disk. Now consider the restriction $f|_{[0,1]}$ given by $f(x) = xe^{c-x} - 1$, for $x \in [0, 1]$. By elementary calculus, $f(x)$ is continuous on $[0, 1]$, and $f(0) = -1$, $f(1) = e^{c-1} - 1 > 0$ since $c > 1$. Thus, by the Intermediate Value Theorem, there exists some $x_0 \in (0, 1)$ so that $f(x_0) = 0$. Thus, x_0 is a root of f in Δ . Since f has exactly one root inside the unit disk, and $x_0 > 0$, the proof concludes.

Problem 2015-A-I-3. Let f be a holomorphic function defined on a neighborhood of the closed unit disk $\overline{\mathbb{D}} := \{z \in \mathbb{C} : |z| \leq 1\}$, and assume that $|f(z)| < 1$ for all $|z| = 1$. Determine the number of fixed points of f in $\overline{\mathbb{D}}$.

Let $f(z)$ be a holomorphic function defined on a neighborhood of the closed unit disk $\overline{\mathbb{D}}$ with the aforementioned properties. The number of fixed points of f is equivalent to the number of roots of the holomorphic function $f(z) - z$ in $\overline{\mathbb{D}}$. Indeed, by Rouché's theorem, since for all $z \in \partial\overline{\mathbb{D}}$,

$$|(f(z) - z) - (-z)| = |f(z)| < 1 = |-z|,$$

$f(z) - z$ has the same number of roots as $-z$ inside the closed unit disk. Since the latter has exactly one zero inside the closed unit disk, $f(z) - z$ has exactly one zero inside the closed unit disk. Thus, $f(z)$ has exactly one fixed point inside $\overline{\mathbb{D}}$.

Problem 2012-A-I-4. Find, with proof, the precise number of zeros of the complex polynomial $p(z) = z^9 - 2z^6 + z^2 - 8z - 2$ inside the annulus $1 < |z| < 2$.

Let $p(z) = z^9 - 2z^6 + z^2 - 8z - 2$. By Rouché's theorem, the number of zeros inside the annulus is equal to the difference of the number of zeros contained in the outer disk $|z| < 2$ and the number of zeros contained in the inner disk $|z| < 1$ (on a quick note, it is important and straightforward to check that $p(z)$ has no roots in $|z| = 1$ nor $|z| = 2$).

FINDING NUMBER OF ROOTS IN $|z| < 2$:

Let $g(z) = z^9$. Then on $|z| = 2$,

$$\begin{aligned} |-2z^6 + z^2 - 8z + 2| &\leq 2|z|^6 + \dots + 2 \\ &= 2^7 + 2^2 + 2^4 + 2 < 2^9 = |g(z)|. \end{aligned}$$

Thus, by Rouché's theorem, $p(z)$ has the same number of zeros inside $|z| < 2$ as does $g(z)$ (which is 9).

FINDING NUMBER OF ROOTS IN $|z| < 1$: Let $g(z) = -8z$. Then on $|z| = 1$,

$$\begin{aligned} |p(z) - g(z)| &\leq |z|^9 + 2|z|^6 + |z|^2 + 2 \\ &= 1 + 2 + 1 + 2 = 6 < 8 = |g(z)|. \end{aligned}$$

Thus, by Rouché's Theorem, $p(z)$ has the same number of zeros inside $|z| < 1$ as does $g(z)$ (which is 1).

Therefore, we conclude that $p(z)$ has only eight zeros inside the annulus.

6 Partial Differential Equations

6.1 Introduction

Problem E-1-1. Classify each of the partial differential equations in §1.2 as follows:

- (a) Is the PDE linear, semilinear, quasilinear or fully nonlinear?
 (b) What is the order of the PDE?

Name	Equation	Classification	Order
Laplace's Equation	$\Delta u = 0$	Linear	2
Helmholtz's (Eigenvalue) Equation)	$-\Delta u = \lambda u$	Linear	2
Linear Transport Equation	$u_t + \sum_{i=1}^n b^i u_{x_i} = 0$	Linear	1
Liouville's Equation	$u_t - \sum_{i=1}^n (b^i u)_{x_i} = 0$	Linear	1
Heat (or diffusion) Equation	$u_t - \Delta u = 0$	Linear	2
Schrödinger's Equation	$i u_t + \Delta u = 0$	Linear	2
Kolmogorov's Equation	$u_t - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} = 0$	Linear	2
Fokker-Planck Equation	$u_t - \sum_{i,j=1}^n (a^{ij} u)_{x_i x_j} - \sum_{i=1}^n (b^i u)_{x_i} = 0$	Linear	2
Wave Equation	$u_{tt} - \Delta u = 0$	Linear	2
Klein-Gordon Equation	$u_{tt} - \Delta u + m^2 u = 0$	Linear	2
Telegraph Equation	$u_{tt} + 2d u_t - u_{xx} = 0$	Linear	2
General Wave Equation	$u_{tt} - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} = 0$	Linear	2
Airy's Equation	$u_t + u_{xxx} = 0$	Linear	3
Beam Equation	$u_{tt} + u_{xxxx} = 0$	Linear	4

Name	Equation	Classification	Order
Eikonal Equation	$ Du = 1$	Fully Nonlinear	1
Nonlinear Poisson Equation	$-\Delta u = f(u)$	Semilinear	2
p -Laplacian Equation	$\operatorname{div}(Du ^{p-2} Du) = 0$	Quasilinear	2
Minimal Surface Equation	$\operatorname{div}\left(\frac{Du}{(1 + Du ^2)^{1/2}}\right) = 0$	Quasilinear	2
Monge-Ampère Equation	$\det(D^2 u) = f$	Fully Nonlinear	2
Hamilton-Jacobi Equation	$u_t + H(Du, x) = 0$	Quasilinear	1
Scalar Conservation Law	$u_t + \operatorname{div} F(u) = 0$	Fully Nonlinear	1
Inviscid Burger's Equation	$u_t + uu_x = 0$	Quasilinear	1
Scalar Reaction-Diffusion Equation	$u_t - \Delta u = f(u)$	Semilinear	2
Porous Medium Equation	$u_t - \Delta(u^p) = 0$	Quasilinear	2
Nonlinear Wave Equation	$u_{tt} - \Delta u + f(u) = 0$	Semilinear	2
Korteweg-de Vries (KdV) Equation	$u_t + uu_x + u_{xxx} = 0$	Semilinear	3
Nonlinear Schrödinger Equation	$i u_t + \Delta u = f(u ^2)u$	Semilinear	2

Problem E-1-2. Let k be a positive integer. Show that a smooth function defined in $\mathbb{R}^n$ has in

general

$$\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$$

distinct partial derivatives of order k .

(Hint: This is the number of ways of inserting $n-1$ dividers $|$ within a row of k symbols $\circ$: for example $\circ\circ || \circ\circ\circ | \circ | \circ\circ\circ || \circ\circ\circ\circ |$. Explain why each such pattern corresponds to precisely one of the partial derivatives of order k .)

Let $k \geq 1$ be a positive integer, and let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function. We want to compute the total number of distinct partial derivatives of u of the form

$$\frac{\partial^k u}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_k}} \quad (1 \leq i_1, \dots, i_k \leq n).$$

Problem E-1-4. Prove *Leibniz's Formula*:

$$D^\alpha(uv) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} D^\beta u D^{\alpha-\beta} v,$$

where $u, v : \mathbb{R}^n \rightarrow \mathbb{R}$ are smooth, $\binom{\alpha}{\beta} = \frac{\alpha!}{\beta!(\alpha-\beta)!}$, and $\beta \leq \alpha$ means $\beta_i \leq \alpha_i$ ($i = 1, \dots, n$).

We can prove Leibniz's formula using induction. First, we consider the case where $|\alpha| = 1$, and assume that $D^\alpha = \partial_{x_i}$ for some positive integer $1 \leq i \leq n$ (i.e., $\alpha = (0, \dots, 1, \dots, 0)$ where the 1 is in the i^{th} place). We note that

$$\mathcal{A} := \{\beta \in \mathbb{N}^n : \beta \leq \alpha\} = \{(0, \dots, 0), \alpha\}.$$

Then by the usual product rule for one variable, one has

$$D^\alpha(uv) = \frac{\partial}{\partial x_i}(uv) = (u) \left(\frac{\partial v}{\partial x_i} \right) + \left(\frac{\partial u}{\partial x_i} \right) (v) = \sum_{\beta \in \mathcal{A}} \binom{\alpha}{\beta} D^\beta u D^{\alpha-\beta} v.$$

Now assume that the result holds for all $|\alpha| = k$ for some $k \geq 1$. We now consider D^α , where $|\alpha| = k+1$. There exists a $(k+1)$ -tuple $(i_1, \dots, i_{k+1})$ so that $D^\alpha = \partial_{x_{i_1}} \cdots \partial_{x_{i_{k+1}}}$. Let γ be the k -tuple so that $D^\gamma = \partial_{x_{i_2}} \cdots \partial_{x_{i_{k+1}}}$. Then by the induction hypothesis, one has

$$D^\gamma(uv) = \sum_{\beta \leq \gamma} \binom{\gamma}{\beta} D^\beta u D^{\gamma-\beta} v,$$

so that

$$\begin{aligned} D^\alpha(uv) &= \sum_{\beta \leq \gamma} \binom{\gamma}{\beta} \partial_{x_{i_1}} (D^\beta u D^{\gamma-\beta} v) \\ &= \sum_{\beta \leq \gamma} \binom{\gamma}{\beta} [\partial_{x_{i_1}} (D^\beta u) D^{\gamma-\beta} v + D^\beta u \partial_{x_{i_1}} (D^{\gamma-\beta} v)]. \end{aligned}$$

Now we note that $\partial_{x_{i_1}} D^{\gamma-\beta} v = D^{\alpha-\beta} v$. On the other hand, we can write $\partial_{x_{i_1}} (D^\beta u) D^{\gamma-\beta} v$ as

$$\partial_{x_{i_1}} (D^\beta u) D^{\gamma-\beta} v = D^{\beta+e_{i_1}} u \cdot D^{\alpha-(\beta+e_{i_1})} v.$$

7 Notes

7.1 Differential Geometry and Topology

7.1.1 De Rham Cohomology

- **Def. (de Rham Cohomology Group)** Let M be a smooth manifold with or without boundary, and let p be a nonnegative positive integer. Define the following spaces:

$$\begin{aligned}\mathcal{Z}^p(M) &= \ker(d : \Omega^p(M) \rightarrow \Omega^{p+1}(M)) = \{\text{closed } p\text{-forms on } M\}, \\ \mathcal{B}^p(M) &= \text{im}(d : \Omega^{p-1}(M) \rightarrow \Omega^p(M)) = \{\text{exact } p\text{-forms on } M\}.\end{aligned}$$

Then we define the *de Rham cohomology group in degree p* to be the quotient vector space

$$H_{\text{dR}}^p(M) := \frac{\mathcal{Z}^p(M)}{\mathcal{B}^p(M)}.$$

- **Def. (Cohomologous Closed Form)** Let M be a smooth manifold with or without boundary, and let ω, ω' be closed p -forms on M . Then ω and ω' are said to be *cohomologous* if and only if $\omega = \omega' + \beta$, where β is an exact p -form. Then $[\omega] = [\omega']$ in $H_{\text{dR}}^p(M)$.
- **Prop. (Induced Cohomology Groups)** For any smooth map $F : M \rightarrow N$ between smooth manifolds with or without boundary, the pullback $F^* : \Omega^p(N) \rightarrow \Omega^p(M)$ carries $\mathcal{Z}^p(N)$ and $\mathcal{B}^p(N)$ into $\mathcal{Z}^p(M)$ and $\mathcal{B}^p(M)$, respectively. Thus it descends into a linear map from $H_{\text{dR}}^p(N)$ to $H_{\text{dR}}^p(M)$.
- **Prop. (Homotopic Maps and De Rham Cohomology)** Suppose M and N are smooth manifolds with or without boundary, and let $F, G : M \rightarrow N$ be homotopic smooth maps. For every p , the induced cohomology maps $F^*, G^* : H_{\text{dR}}^p(N) \rightarrow H_{\text{dR}}^p(M)$ are equal.
- **Thm. (Homotopy Invariance of De Rham Cohomology)** If M and N are homotopy equivalent smooth manifolds with or without boundary, then $H_{\text{dR}}^p(M) \cong H_{\text{dR}}^p(N)$ for each p . The isomorphisms are induced by any smooth homotopy equivalence $F : M \rightarrow N$.
- **Cor. (Topological Invariance of De Rham Cohomology)** The de Rham cohomology groups are topological invariants: if M and N are homeomorphic smooth manifolds with or without boundary, then their de Rham cohomology groups are isomorphic.
- **Def. (Contractible Topological Space)** A topological space X is said to be *contractible* if the identity map of X is homotopic to a constant map.
- **Thm. (Contractible Manifolds)** If M is a contractible smooth manifold with or without boundary, then $H_{\text{dR}}^p(M) = 0$ for $p \geq 1$.

Proof. Let M be a contractible smooth manifold. Then there exists some $q \in M$ so that id_M is homotopic to the constant map $c_q : M \rightarrow M$ sending all of M to $\{q\}$. Let $\iota_q : \{q\} \hookrightarrow M$ denote the inclusion map. Then since $c_q \circ \iota_q = \text{Id}_M = \iota_q \circ c_q$, ι_q is a homotopy equivalence. This means that $H_{\text{dR}}^p(\{q\}) \cong H_{\text{dR}}^p(M)$ for each p . But $H_{\text{dR}}^p(\{q\}) \cong 0$ for each $p \geq 1$ since $\{q\}$ is a 0-manifold. Thus the proof concludes. $\square$

- **Thm. (Poincaré Lemma)** If U is a star-shaped open subset of $\mathbb{R}^n$ or $\mathbb{H}^n$ then $H_{\text{dR}}^p(U) = 0$ for $p \geq 1$.

Proof. Let U be star-shaped with respect to $c \in U$. Then the straight-line homotopy $H(x, t) = c + t(x - c)$ shows that U is contractible. Thus by the above theorem, $H_{\text{dR}}^p(U) = 0$ for all $p \geq 1$. $\square$

- **Cor. (Local Exactness of Closed Forms)** Let M be a smooth manifold with or without boundary. Each point of M has a neighborhood on which every closed form is exact.

Proof. Every point of M has a neighborhood diffeomorphic to an open ball in $\mathbb{R}^n$ or an open half-ball in $\mathbb{H}^n$, each of which is star-shaped. Thus by the Poincaré Lemma and the diffeomorphism invariance of the de Rham cohomology, the proof concludes. $\square$

- **Thm. (First Cohomology and the Fundamental Group)** Suppose M is a connected smooth manifold. For each $q \in M$, the linear map $\Phi : H_{\text{dR}}^1(M) \rightarrow \text{Hom}(\pi_1(M, q), \mathbb{R})$ is well-defined and injective (in particular, an isomorphism).
- **Cor. (Finite Fundamental Group and First Cohomology)** If M is a connected smooth manifold with finite fundamental group, then $H_{\text{dR}}^1(M) = 0$.

Proof. Let M be a connected smooth manifold with finite fundamental group. By the above theorem, one has that $H_{\text{dR}}^1(M) \cong \text{Hom}(\pi_1(M), \mathbb{R})$. Since there is no nontrivial homomorphism from a finite group to $\mathbb{R}$, it follows that $\text{Hom}(\pi_1(M), \mathbb{R})$ is trivial. Thus the proof concludes. $\square$

Exercise 17.19. A group Γ is called a **torsion group** if for each $g \in \Gamma$, there exists an integer k such that $g^k = 1$. Show that if M is a connected smooth manifold whose fundamental group is a torsion group, then $H_{\text{dR}}^1(M) = 0$.

Let M be connected smooth manifold whose fundamental group $\pi_1(M)$ is a torsion group. By the above theorem, one has $H_{\text{dR}}^1(M) \cong \text{Hom}(\pi_1(M), \mathbb{R})$. Suppose $\varphi : \pi_1(M) \rightarrow \mathbb{R}$ is a homomorphism. Let $g \in \pi_1(M)$ be an arbitrary nonidentity element, and let k be a positive integer so that $g^k = 1$. By the property of group homomorphisms,

$$0 = \varphi(e) = \varphi(g^k) = k \cdot \varphi(g).$$

Since k is a positive integer and $\mathbb{R}$ is an integral domain, we conclude that $\varphi(g) = 0$. But since g was an arbitrary non-identity element of the fundamental group, we conclude that φ is the trivial homomorphism. Thus in fact, $\text{Hom}(\pi_1(M), \mathbb{R}) = 0$ so that $H_{\text{dR}}^1(M) = 0$.

- **Thm. (Mayer-Vietoris)** Let M be a smooth manifold with or without boundary, and let U, V be open subsets of M whose union is M . For each p , there is a linear map $\delta : H_{\text{dR}}^p(U \cap V) \rightarrow H_{\text{dR}}^{p+1}(M)$ such that the following sequence called the **Mayer-Vietoris sequence** for the open cover $\{U, V\}$ is exact:

$$\cdots \xrightarrow{\delta} H_{\text{dR}}^p(M) \xrightarrow{k^* \oplus \ell^*} H_{\text{dR}}^p(U) \oplus H_{\text{dR}}^p(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^p(U \cap V) \xrightarrow{\delta} H_{\text{dR}}^{p+1}(M) \xrightarrow{k^* \oplus \ell^*} \cdots,$$

where $i : U \cap V \hookrightarrow U$, $k : U \hookrightarrow M$, $\ell : V \hookrightarrow M$, and $j : U \cap V \hookrightarrow V$ are inclusions.

- **Thm. (Cohomology of Spheres)** For $n \geq 1$, the de Rham cohomology groups of S^n are

$$H_{\text{dR}}^p(S^n) \cong \begin{cases} \mathbb{R}, & \text{if } p = 0 \text{ or } p = n, \\ 0, & \text{if } 0 < p < n. \end{cases}$$

Proof. Since S^n is connected, $H_{\text{dR}}^0(S^n) \cong \mathbb{R}$. Thus we only need to prove the above result for $p \geq 1$. We will do so by induction. For $n = 1$, since any orientation form on S^1 has nonzero integral, they must all be non-exact. Thus $\dim H_{\text{dR}}^1(S^1) \geq 1$. On the other hand, there is an injective linear map from $H_{\text{dR}}^1(S^1) \rightarrow \text{Hom}(\pi_1(S^1, 1), \mathbb{R})$, which is 1-dimensional. Thus $H_{\text{dR}}^1(S^1)$ has dimension exactly 1, and is spanned by the cohomology class of any orientation form.

Now suppose $n \geq 2$, and assume by induction that the theorem is true for S^{n-1} . Since S^n is simply connected, $H_{\text{dR}}^1(S^n) = 0$. For $p > 1$, we use the Mayer-Vietoris theorem as follows. Let N and S be the north and south poles in S^n , respectively, and let $U = S^n \setminus \{S\}$ and $V = S^n \setminus \{N\}$. By

stereographic projection, both U and V are diffeomorphic to $\mathbb{R}^n$ so that $U \cap V$ is diffeomorphic to $\mathbb{R}^n \setminus \{0\}$. Part of the Mayer-Vietoris sequence for $\{U, V\}$ reads

$$H_{\text{dR}}^{p-1}(U) \oplus H_{\text{dR}}^{p-1}(V) \rightarrow H_{\text{dR}}^{p-1}(U \cap V) \rightarrow H_{\text{dR}}^p(S^n) \rightarrow H_{\text{dR}}^p(U) \oplus H_{\text{dR}}^p(V).$$

Since U and V are diffeomorphic to $\mathbb{R}^n$, the groups on both ends are trivial when $p > 1$, which implies that $H_{\text{dR}}^p(S^n) \cong H_{\text{dR}}^{p-1}(U \cap V)$. Moreover, $U \cap V$ is diffeomorphic to $\mathbb{R}^n \setminus \{0\}$ and therefore homotopy equivalent to $\mathbb{S}^{n-1}$. Thus, we conclude that $H_{\text{dR}}^p(S^n) \cong H_{\text{dR}}^{p-1}(S^{n-1})$ for $p > 1$. Therefore, the above result follows by induction. As in the $n = 1$ case, any smooth orientation form on $\mathbb{S}^n$ determines a nonzero cohomology class, which therefore spans $H_{\text{dR}}^n(S^n)$. $\square$

- **Cor. (Cohomology of Punctured Euclidean Space)** Suppose $n \geq 2$ and $x \in \mathbb{R}^n$. Let $M = \mathbb{R}^n \setminus \{x\}$. The only nontrivial de Rham groups of M are $H_{\text{dR}}^0(M)$ and $H_{\text{dR}}^{n-1}(M)$, both of which are 1-dimensional. A closed $(n-1)$ -form η on M is exact if and only if $\int_S \eta = 0$ for some (and hence every) $(n-1)$ -dimensional sphere $S \subseteq M$ centered at x .

Proof. Let $S \subseteq M$ be any $(n-1)$ -dimensional sphere centered at x . Because inclusion $\iota: S \hookrightarrow M$ is a homotopy equivalence, $\iota^*: H_{\text{dR}}^p(M) \rightarrow H_{\text{dR}}^p(S)$ is an isomorphism for each p , so the assertion about the dimension of $H_{\text{dR}}^p(M)$ follows from the cohomology of spheres. If η is a closed $(n-1)$ -form on M , it follows that η is exact if and only if $\iota^*\eta$ is exact on S , which in turn is true if and only if $\int_S \eta = \int_S \iota^*\eta = 0$ by Exercise 17.22. $\square$

- **Cor. (Punctured Open Set)** Suppose $n \geq 2$, $U \subseteq \mathbb{R}^n$ is any open subset, and $x \in U$. Then $H_{\text{dR}}^{n-1}(U \setminus \{x\}) \neq 0$.
- **Lem. (Poincaré Lemma with Compact Support)** Let $n \geq p \geq 1$, and suppose ω is a compactly supported closed p -form on $\mathbb{R}^n$. If $p = n$, suppose in addition that $\int_{\mathbb{R}^n} \omega = 0$. Then there exists a compactly supported smooth $(p-1)$ -form η on $\mathbb{R}^n$ such that $\omega = d\eta$.

Proof. Let $n \geq p \geq 1$, and suppose ω is a compactly supported closed p -form on $\mathbb{R}^n$. By the Poincaré Lemma, $H_{\text{dR}}^k(\mathbb{R}^n) = 0$ for all $k \geq 1$ so that ω is automatically exact. Thus, it suffices to show that ω is the exterior derivative of a *compactly supported* form. Our proof of this lemma will be split into a few different cases.

CASE 1: Suppose $n = p = 1$. Assume $\int_{\mathbb{R}^n} \omega = 0$. Then we can write $\omega = f dx$ for some smooth, compactly supported function $f \in C_c^\infty(\mathbb{R})$. Define $F: \mathbb{R} \rightarrow \mathbb{R}$ by

$$F(x) = \int_{-\infty}^x f(t) dt.$$

By the fundamental theorem of calculus, $dF = F' dx = f dx = \omega$. Thus it suffices to show that F is compactly supported. Let $R > 0$ so that $\text{supp } f \subset [-R, R]$. When $x < -R$, by construction, $F(x) = 0$. When $x > R$, since $\int_{\mathbb{R}} \omega = 0$, one has

$$F(x) = \int_{-\infty}^x f(t) dt = \int_{-\infty}^{\infty} f(t) dt = 0.$$

Thus, $\text{supp } F \subset [-R, R]$. Since F is compactly supported and $\omega = dF$, the proof concludes.

CASE 2: Now assume $n \geq 2$, and let $B, B' \subseteq \mathbb{R}^n$ be open balls centered at the origin such that $\text{supp } \omega \subseteq B \subseteq \bar{B} \subseteq B'$. By the ordinary Poincaré Lemma, there exists a smooth (but not necessarily compactly supported) $(p-1)$ -form η_0 on $\mathbb{R}^n$ such that $d\eta_0 = \omega$. In particular, $d\eta_0 = 0$ on $\mathbb{R}^n \setminus \bar{B}$. Consider the further three cases:

CASE A: $p = 1$. In this case, η_0 must be a smooth function. Since $\mathbb{R}^n \setminus \bar{B}$ is connected when $n \geq 2$, it follows that η_0 is equal to a constant c on $\mathbb{R}^n \setminus \bar{B}$. Let $\eta = \eta_0 - c$. Then $\eta = \eta_0 - c$ is compactly supported and satisfies $d\eta = \omega$ as claimed.

CASE B: $1 < p < n$. The restriction of η_0 to $\mathbb{R}^n \setminus \bar{B}$ is a closed $(p-1)$ -form. Since $H_{\text{dR}}^{p-1}(\mathbb{R}^n \setminus \bar{B}) = 0$, there is a smooth $(p-2)$ -form γ on $\mathbb{R}^n \setminus \bar{B}$ such that $d\gamma = \eta_0$ there. Let ψ be a smooth bump function that is supported in $\mathbb{R}^n \setminus \bar{B}$ and equal to 1 on $\mathbb{R}^n \setminus B'$. Then $\eta = \eta_0 - d(\psi\gamma)$ is smooth on all of $\mathbb{R}^n$ and satisfies $d\eta = d\eta_0 = \omega$. Because $d(\psi\gamma) = d\gamma = \eta_0$ on $\mathbb{R}^n \setminus B'$, η is compactly supported.

CASE C: $p = n$. In this case, we cannot use the same argument as in Case B since $H_{\text{dR}}^{n-1}(\mathbb{R}^n \setminus \bar{B}) \neq 0$. However, the restriction of η_0 to $\mathbb{R}^n \setminus \bar{B}$ is exact provided its integral is zero over some sphere centered at the origin and contained in $\mathbb{R}^n \setminus \bar{B}$. By Stokes' Theorem,

$$0 = \int_{\mathbb{R}^n} \omega = \int_{\bar{B}'} \omega = \int_{\bar{B}'} d\eta_0 = \int_{\partial B'} \eta_0.$$

Thus η_0 is exact on $\mathbb{R}^n \setminus \bar{B}$, and the proof proceeds exactly as in Case B. □

- **Def. (p th Compactly Supported de Rham Cohomology Group)** Let M be a smooth manifold with or without boundary, and let $\Omega_c^p(M)$ denote the vector space of compactly supported smooth p -forms on M . The p^{th} compactly supported de Rham cohomology group of M is the quotient space

$$H_c^p(M) := \frac{\ker(d : \Omega_c^p(M) \rightarrow \Omega_c^{p+1}(M))}{\text{im}(\Omega_c^{p-1}(M) \rightarrow \Omega_c^p(M))}.$$

- **Thm. (Compactly Supported Cohomology of $\mathbb{R}^n$)** For $n \geq 1$, the compactly supported de Rham cohomology groups of $\mathbb{R}^n$ are

$$H_c^p(\mathbb{R}^n) = \begin{cases} 0, & \text{if } 0 \leq p < n, \\ \mathbb{R}, & \text{if } p = n. \end{cases}$$

Exercise 17.29. Prove this theorem.

Fix some $n \geq 1$, and let $M = \mathbb{R}^n$. For all $1 \leq p < n$, by the Poincaré Lemma with compact support, every compactly supported closed p -form on $\mathbb{R}^n$ is the exterior derivative of some compactly supported smooth $(p-1)$ -form on $\mathbb{R}^n$. Thus for all $1 \leq p < n$, $H_c^p(M) = 0$. Now let $p = 0$, and $\omega \in \Omega_c^0(M)$. Since ω is closed, $d\omega$ is identically zero on $\mathbb{R}^n$. Since $\mathbb{R}^n$ is connected and ω is actually just a smooth function on $\mathbb{R}^n$, ω must be a constant everywhere on $\mathbb{R}^n$. But since ω also has to be compactly supported, $\omega = 0$. Thus since every compactly supported closed smooth 0-form on $\mathbb{R}^n$ is identically zero, $H_c^0(M) = 0$. We have proven the first half of the theorem. Now suppose $p = n$. First assume $n = 1$. Let $\psi(x)$ be a smooth, non-negative bump function supported in some interval $B = [a, b]$ so that $\int_{\mathbb{R}} \psi(x) dx > 0$, and let $\omega = \psi(x) dx$. Let f be a smooth function on $\mathbb{R}$ such that $\omega = df$. In particular, we must then have

$$f(x) = \int_{-\infty}^x \psi(t) dt.$$

For any $x > b$,

$$f(x) = \int_{-\infty}^x \psi(t) dt = \int_{-\infty}^{\infty} \psi(t) dt = C > 0,$$

which means f is not compactly supported. Thus, $\dim H_c^1(\mathbb{R}) \geq 1$ since we have shown the existence of a closed, non-exact, compactly supported 1-form. [!! Complete Later !!]

- **Thm. (Top Cohomology, Orientable Compact Support Case)** If M is a connected oriented smooth n -manifold, then the integration map $I : H_c^n(M) \rightarrow \mathbb{R}$ is an isomorphism so that $H_c^n(M)$ is 1-dimensional.

- **Thm. (Top Cohomology, Orientable Compact Case)** If M is a compact connected orientable smooth n -manifold, then $H_{\text{dR}}^n(M)$ is 1-dimensional, and is spanned by the cohomology class of any smooth orientation form.
- **Thm. (Top Cohomology, Orientable Noncompact Case)** If M is a noncompact connected orientable smooth n -manifold, then $H_{\text{dR}}^n(M) = 0$.
- **Lem. (Nonorientability and Cohomology)** Suppose M is a connected nonorientable smooth manifold and $\hat{\pi} : \hat{M} \rightarrow M$ is its orientation covering. For each p , the induced cohomology maps $\hat{\pi}^* : H_{\text{dR}}^p(M) \rightarrow H_{\text{dR}}^p(\hat{M})$ and $\hat{\pi}^* : H_c^p(M) \rightarrow H_c^p(\hat{M})$ are injective.
- **Thm. (Top Cohomology, Nonorientable Case)** If M is a connected nonorientable smooth n -manifold, then $H_c^n(M) = 0$ and $H_{\text{dR}}^n(M) = 0$.

Problem LSM-17-5. For each $n \geq 1$, compute the de Rham cohomology groups of $\mathbb{R}^n \setminus \{e_1, -e_1\}$; and for each nonzero cohomology group, give specific differential forms whose cohomology classes form a basis.

We will work through this problem through several cases.

CASE 1: First suppose that $n = 1$, and fix $e_1 \in \mathbb{R}$. Then $\mathbb{R} \setminus \{e_1, -e_1\}$ is the disjoint union of three connected components M_1, M_2, M_3 so that $H_{\text{dR}}^1(M) \cong \mathbb{R}^3$. Next, since each connected component is contractible,

$$H_{\text{dR}}^1(M) \cong \prod_{j=1}^3 H_{\text{dR}}^1(M_j) = 0.$$

CASE 2: Now let $n \geq 2$. Since M is connected, $H_{\text{dR}}^0(M) = \mathbb{R}$. Since M is connected, noncompact, and orientable, $H_{\text{dR}}^n(M) = 0$. Thus it suffices to compute the de Rham cohomology groups for $1 \leq p \leq n-1$. We first focus on $1 < p < n-1$. Let $e_1 = (1, 0, \dots, 1)$, and define the open subsets $U = \{x \in \mathbb{R}^n : x_1 > -0.5\} \setminus \{e_1\}$ and $V = \{x \in \mathbb{R}^n : x_1 < 0.5\} \setminus \{e_2\}$. U and V are both homotopy equivalent to S^{n-1} and $U \cap V$ is contractible. By the Mayer-Vietoris Theorem, we have the exact sequence

$$\cdots \rightarrow H_{\text{dR}}^{p-1}(U) \oplus H_{\text{dR}}^{p-1}(V) \rightarrow H_{\text{dR}}^{p-1}(U \cap V) \rightarrow H_{\text{dR}}^p(M) \rightarrow H_{\text{dR}}^p(U) \oplus H_{\text{dR}}^p(V) \rightarrow H_{\text{dR}}^p(U \cap V) \rightarrow \cdots,$$

which then becomes

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow H_{\text{dR}}^p(M) \rightarrow 0 \rightarrow 0 \rightarrow \cdots.$$

Thus for $1 < p < n-1$, one has $H_{\text{dR}}^p(M) = 0$. Now let $p = n-1$. Then Mayer-Vietoris gives us the exact sequence

$$\cdots \rightarrow H_{\text{dR}}^{n-2}(U \cap V) \rightarrow H_{\text{dR}}^{n-1}(M) \rightarrow H_{\text{dR}}^{n-1}(U) \oplus H_{\text{dR}}^{n-1}(V) \rightarrow H_{\text{dR}}^{n-1}(U \cap V) \rightarrow \cdots.$$

For $n > 2$, the above exact sequence becomes

$$\cdots \rightarrow 0 \rightarrow H_{\text{dR}}^{n-1}(M) \rightarrow \mathbb{R}^2 \rightarrow 0 \rightarrow \cdots,$$

so that $H_{\text{dR}}^{n-1}(M) \cong \mathbb{R}^2$. Finally, we consider the case where $p = 1$. By the Mayer-Vietoris Theorem, we have the exact sequence

$$\cdots \rightarrow H_{\text{dR}}^0(U) \oplus H_{\text{dR}}^0(V) \rightarrow H_{\text{dR}}^0(U \cap V) \rightarrow H_{\text{dR}}^1(M) \rightarrow H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V) \rightarrow H_{\text{dR}}^1(U \cap V) \rightarrow \cdots,$$

which becomes

$$\cdots \rightarrow \mathbb{R}^2 \rightarrow \mathbb{R} \rightarrow H_{\text{dR}}^1(M) \rightarrow H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V) \rightarrow 0 \rightarrow \cdots.$$

The map $\mathbb{R}^2 \rightarrow \mathbb{R}$ is defined by $(f, g) \mapsto f|_{U \cap V} - g|_{U \cap V}$. But since $H_{\text{dR}}^0(U), H_{\text{dR}}^0(V) \cong \mathbb{R}$, we can treat the elements in the direct sum $H_{\text{dR}}^0(U) \oplus H_{\text{dR}}^0(V)$ simply as pairs of real numbers (c_1, c_2) . Thus the map is surjective. By the First Isomorphism Theorem,

$$\text{Im}(\mathbb{R} \rightarrow H_{\text{dR}}^1(M)) \cong \mathbb{R} / \ker(\mathbb{R} \rightarrow H_{\text{dR}}^1(M)).$$

But $\ker(\mathbb{R} \rightarrow H_{\text{dR}}^1(M)) = \text{Im}(\mathbb{R}^2 \rightarrow \mathbb{R}) = \mathbb{R}$. Thus, $\text{Im}(\mathbb{R} \rightarrow H_{\text{dR}}^1(M)) = 0$. This means that we have the exact sequence

$$0 \rightarrow H_{\text{dR}}^1(M) \rightarrow H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V) \rightarrow 0,$$

so that $H_{\text{dR}}^1(M) \cong H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V)$. Thus, by using the cohomology of spheres, the computations conclude.

Problem LSM-17-6. Let M be a connected smooth manifold of dimension $n \geq 3$. For any $x \in M$ and $0 \leq p \leq n-2$, prove that the map $H_{\text{dR}}^p(M) \rightarrow H_{\text{dR}}^p(M \setminus \{x\})$ induced by inclusion $M \setminus \{x\} \hookrightarrow M$ is an isomorphism. Prove that the same is true for $p = n-1$ if M is compact and orientable. [Hint: use the Mayer-Vietoris theorem. The cases $p = 0$, $p = 1$, and $p = n-1$ require special handling.]

Let M be a connected smooth manifold of dimension $n \geq 3$. Fix an arbitrary $x \in M$. Let us denote the following: (1) $U = M \setminus \{x\}$, (2) V an open neighborhood of x in M , (3) $i : U \cap V \hookrightarrow U$, (4) $j : U \cap V \hookrightarrow V$, (5) $k : U \hookrightarrow M$, and $l : V \hookrightarrow M$. We will work through the proof through several different cases.

CASE 1: $1 < p \leq n-2$, $n > 3$. For each such p , the Mayer-Vietoris theorem gives us the exact sequence

$$\cdots \rightarrow H_{\text{dR}}^{p-1}(U \cap V) \rightarrow H_{\text{dR}}^p(M) \rightarrow H_{\text{dR}}^p(U) \oplus H_{\text{dR}}^p(V) \rightarrow H_{\text{dR}}^p(U \cap V) \rightarrow \cdots$$

Since $U \cap V$ is homotopy equivalent to S^{n-1} , for $1 < p \leq n-2$ (with $n > 3$), $H_{\text{dR}}^p(U \cap V), H_{\text{dR}}^{p-1}(U \cap V) \cong 0$. Moreover, since V is contractible, $H_{\text{dR}}^p(V) \cong 0$. Thus, we have the short exact segment

$$0 \rightarrow H_{\text{dR}}^p(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^p(U) \oplus 0 \rightarrow 0,$$

so that $H_{\text{dR}}^p(M) \cong H_{\text{dR}}^p(U) \oplus 0$. Thus $k^* \oplus l^*$ is an isomorphism, which then implies that k^* is an isomorphism since l^* is the zero map.

CASE 2: Now we will look at the case $p = 0$ for all $n \geq 3$. The Mayer-Vietoris theorem gives us the following exact sequence

$$0 \rightarrow H_{\text{dR}}^0(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^0(U) \oplus H_{\text{dR}}^0(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^0(U \cap V) \xrightarrow{\delta} H_{\text{dR}}^1(M) \rightarrow \cdots$$

In particular, the above sequence becomes,

$$0 \rightarrow \mathbb{R} \xrightarrow{k^* \oplus l^*} \mathbb{R}^2 \xrightarrow{i^* - j^*} \mathbb{R} \xrightarrow{\delta} H_{\text{dR}}^1(M) \rightarrow \cdots$$

In particular, we already know that $k^* \oplus l^*$ is injective by exactness. On the other hand, $i^* - j^*$ is the map that maps $(a, b) \mapsto a - b$, and therefore is surjective. By exactness, $\ker \delta = \text{Im}(i^* - j^*) = \mathbb{R}$ so that δ is the zero map. So we actually have the short exact sequence

$$0 \rightarrow \mathbb{R} \xrightarrow{k^* \oplus l^*} \mathbb{R}^2 \xrightarrow{i^* - j^*} \mathbb{R} \rightarrow 0.$$

Since $k^* \oplus l^* : c \mapsto (k^*c, l^*c) = (c, c)$, $k^* : c \mapsto c$. Since $H_{\text{dR}}^0(M) \cong H_{\text{dR}}^0(U)$ and k^* maps each $c \in \mathbb{R}$ to itself, k^* is the identity map, and thus is an isomorphism.

CASE 3: Now we will look at the case $p = 1$ (which also includes the case $n = 3$). The Mayer-Vietoris Theorem gives us the following exact sequence

$$\cdots \rightarrow H_{\text{dR}}^0(U \cap V) \xrightarrow{\delta} H_{\text{dR}}^1(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(U) \oplus H_{\text{dR}}^1(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^1(U \cap V) \xrightarrow{\delta} H_{\text{dR}}^2(M) \rightarrow \cdots$$

In particular, one has

$$\cdots \rightarrow \mathbb{R} \xrightarrow{\delta} H_{\text{dR}}^1(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(U) \oplus 0 \xrightarrow{i^* - j^*} 0 \xrightarrow{\delta} H_{\text{dR}}^2(M) \rightarrow \cdots$$

But in our previous work, we showed that δ was the zero map. So in fact, we must have the short exact segment

$$0 \xrightarrow{\delta} H_{\text{dR}}^1(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(U) \oplus 0 \xrightarrow{i^* - j^*} 0,$$

so that $k^* \oplus l^*$ is an isomorphism, and therefore, k^* is an isomorphism.

CASE 4: Now assume that M is compact and orientable. We will look at the case $p = n-1$. The Mayer-Vietoris Theorem gives us the following exact sequence

$$\begin{aligned} \cdots \rightarrow H_{\text{dR}}^{n-2}(U \cap V) \xrightarrow{\delta} H_{\text{dR}}^{n-1}(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^{n-1}(U) \oplus H_{\text{dR}}^{n-1}(V) \xrightarrow{i^* - j^*} H_{\text{dR}}^{n-1}(U \cap V) \\ \xrightarrow{\delta} H_{\text{dR}}^n(M) \rightarrow H_{\text{dR}}^n(U) \oplus H_{\text{dR}}^n(V) \rightarrow \cdots \end{aligned}$$

In particular, using the information we have, the above short exact segment can be written as the following:

$$\cdots \rightarrow 0 \xrightarrow{\delta} H_{\text{dR}}^{n-1}(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^{n-1}(U) \oplus 0 \xrightarrow{i^* - j^*} \mathbb{R} \xrightarrow{\delta} \mathbb{R} \xrightarrow{k^* \oplus l^*} 0,$$

where the penultimate term follows from the compactness, connectedness, and orientability of M , and the last term follows from the fact that U is a noncompact, connected, orientable smooth n -manifold. Thus, we have that $\delta : \mathbb{R} \rightarrow \mathbb{R}$ is surjective. Since δ is a surjection between equal dimensional vector spaces, δ is an isomorphism. Thus, $0 = \ker \delta = \text{Im}(i^* - j^*)$. Therefore, we have the short exact segment $\cdots \rightarrow 0 \rightarrow H_{\text{dR}}^{n-1}(M) \rightarrow H_{\text{dR}}^{n-1}(U) \oplus 0 \rightarrow 0$, whence we conclude that k^* is an isomorphism.

Problem LSM-17-7. Let M_1, M_2 be connected smooth manifolds of dimension $n \geq 3$, and let $M_1 \# M_2$ denote their smooth connected sum (Example 9.31). Prove that $H_{\text{dR}}^p(M_1 \# M_2) \cong H_{\text{dR}}^p(M_1) \oplus H_{\text{dR}}^p(M_2)$ for $0 < p < n - 1$. Prove that the same is true for $p = n - 1$ if M_1 and M_2 are both compact and orientable. [Hint: use Problems 9-12 and 17-6.]

Let M_1, M_2 be connected smooth manifolds of dimension $n \geq 3$, and let M denote their smooth connected sum. Fix $p_1 \in M_1$ and $p_2 \in M_2$. By Problem 9-12, there exist open subsets $\tilde{M}_1, \tilde{M}_2 \subseteq M$ diffeomorphic to $M_1 \setminus \{p_1\}$ and $M_2 \setminus \{p_2\}$, respectively, such that $\tilde{M}_1 \cup \tilde{M}_2 = M$ and $\tilde{M}_1 \cap \tilde{M}_2$ is diffeomorphic to $(-1, 1) \times S^{n-1}$.

Next, for each $0 \leq p < n - 1$, $H_{\text{dR}}^p(\tilde{M}_j) \cong H_{\text{dR}}^p(M_j / \{p_j\}) \cong H_{\text{dR}}^p(M_j)$ for $j = 1, 2$, where the first isomorphism follows by diffeomorphism invariance of the de Rham cohomology group, and the second isomorphism follows from Problem LSM-17-6. Now we will examine a few different cases.

CASE 1: Suppose $1 < p \leq n - 2$, $n > 3$. The Mayer-Vietoris theorem gives us the following exact sequence,

$$\cdots \rightarrow H_{\text{dR}}^{p-1}(\tilde{M}_1 \cap \tilde{M}_2) \rightarrow H_{\text{dR}}^p(M) \rightarrow H_{\text{dR}}^p(\tilde{M}_1) \oplus H_{\text{dR}}^p(\tilde{M}_2) \rightarrow H_{\text{dR}}^p(\tilde{M}_1 \cap \tilde{M}_2) \rightarrow \cdots,$$

which then begets the exact sequence,

$$\cdots \xrightarrow{i^* - j^*} H_{\text{dR}}^{p-1}((-1, 1) \times S^{n-1}) \xrightarrow{\delta} H_{\text{dR}}^p(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^p(M_1) \oplus H_{\text{dR}}^p(M_2) \xrightarrow{i^* - j^*} H_{\text{dR}}^p((-1, 1) \times S^{n-1}) \xrightarrow{\delta} \cdots.$$

Since $(-1, 1) \times S^{n-1}$ is homotopy equivalent to S^{n-1} , for $1 < p \leq n - 2$,

$$H_{\text{dR}}^{p-1}((-1, 1) \times S^{n-1}), H_{\text{dR}}^p((-1, 1) \times S^{n-1}) \cong 0.$$

Thus, we have the short exact sequence

$$0 \xrightarrow{\delta} H_{\text{dR}}^p(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^p(M_1) \oplus H_{\text{dR}}^p(M_2) \xrightarrow{i^* - j^*} 0,$$

CASE 2: Now consider the case $p = 1$ (which also includes the case $n = 3$). From the Mayer-Vietoris theorem, we have the following exact sequence:

$$\begin{aligned} \cdots \rightarrow H_{\text{dR}}^0(M_1) \oplus H_{\text{dR}}^0(M_2) \xrightarrow{i^* - j^*} H_{\text{dR}}^0(\tilde{M}_1 \cap \tilde{M}_2) \xrightarrow{\delta} H_{\text{dR}}^1(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(M_1) \oplus H_{\text{dR}}^1(M_2) \\ \xrightarrow{i^* - j^*} H_{\text{dR}}^1(\tilde{M}_1 \cap \tilde{M}_2) \xrightarrow{\delta} H_{\text{dR}}^2(M) \rightarrow \cdots, \end{aligned}$$

which begets the following exact sequence

$$\cdots \rightarrow \mathbb{R}^2 \xrightarrow{i^* - j^*} \mathbb{R} \xrightarrow{\delta} H_{\text{dR}}^1(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(M_1) \oplus H_{\text{dR}}^1(M_2) \xrightarrow{i^* - j^*} 0 \rightarrow \cdots.$$

Here $i^* - j^* : (c, d) \in \mathbb{R}^2 \mapsto c - d$ so that $i^* - j^*$ is surjective. By exactness, $\ker \delta = \mathbb{R}$ so that δ is the zero map. Thus we actually have the exact sequence

$$0 \rightarrow H_{\text{dR}}^1(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^1(M_1) \oplus H_{\text{dR}}^1(M_2) \xrightarrow{i^* - j^*} 0,$$

CASE 3: Now assume both M_1 and M_2 are compact and orientable, and let $p = n - 1$. Then by Problems LSM-9-12 and LSM-17-6, one has for $j = 1, 2$: $H_{\text{dR}}^{n-1}(\tilde{M}_j) \cong H_{\text{dR}}^{n-1}(M_j)$. Thus the Mayer-Vietoris theorem gives the following exact sequence:

$$\begin{aligned} \cdots \rightarrow H_{\text{dR}}^{n-2}(\tilde{M}_1 \cap \tilde{M}_2) \xrightarrow{\delta} H_{\text{dR}}^{n-1}(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^{n-1}(M_1) \oplus H_{\text{dR}}^{n-1}(M_2) \xrightarrow{i^* - j^*} H_{\text{dR}}^{n-1}(\tilde{M}_1 \cap \tilde{M}_2) \\ \xrightarrow{\delta} H_{\text{dR}}^n(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^n(\tilde{M}_1) \oplus H_{\text{dR}}^n(\tilde{M}_2) \rightarrow \cdots \end{aligned}$$

Using the information we have, we obtain the following exact sequence:

$$\cdots \rightarrow 0 \xrightarrow{\delta} H_{\text{dR}}^{n-1}(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^{n-1}(M_1) \oplus H_{\text{dR}}^{n-1}(M_2) \xrightarrow{i^* - j^*} \mathbb{R} \xrightarrow{\delta} \mathbb{R} \xrightarrow{k^* \oplus l^*} 0,$$

where we used the facts that the top de Rham cohomology of a compact connected orientable manifold is one-dimensional, $M_1 \# M_2$ is compact connected and orientable, and $\tilde{M}_1, \tilde{M}_2$ are both noncompact and orientable so that their top de Rham cohomology groups are trivial. By exactness, δ must be surjective; since it is a surjective linear map between equal dimensional vector spaces, it must also be injective. Thus by exactness, $i^* - j^*$ is the zero map. Thus, we have the exact sequence

$$0 \xrightarrow{\delta} H_{\text{dR}}^{n-1}(M) \xrightarrow{k^* \oplus l^*} H_{\text{dR}}^{n-1}(M_1) \oplus H_{\text{dR}}^{n-1}(M_2) \rightarrow 0,$$

whence $H_{\text{dR}}^{n-1}(M) \cong H_{\text{dR}}^{n-1}(M_1) \oplus H_{\text{dR}}^{n-1}(M_2)$.
The proof concludes.

7.2 Partial Differential Equations

7.2.1 Introduction

- **Def. (k^{th} -Order Partial Differential Equation)** Fix an integer $k \geq 1$, and let U denote an open subset of $\mathbb{R}^n$. An expression of the form

$$F(D^k u(x), D^{k-1} u(x), \dots, Du(x), u(x), x) = 0, \quad (x \in U),$$

where $F : \mathbb{R}^k \times \mathbb{R}^{k-1} \times \dots \times \mathbb{R}^n \times \mathbb{R} \times U \rightarrow \mathbb{R}$ is given and $u : U \rightarrow \mathbb{R}$ is an unknown function, is called a k -th order partial differential equation (PDE).

- **Def. (Classification of PDE Types)** Consider a k -th order PDE of the above form. Then:
 - (i) the PDE is *linear* iff it has the form

$$\sum_{|\alpha| \leq k} a_\alpha(x) D^\alpha u(x) = f(x),$$

for given functions $a_\alpha(x)$ ($|\alpha| \leq k$), f . The linear PDE is *homogeneous* if $f \equiv 0$;

- (ii) the PDE is *semilinear* if it has the form

$$\sum_{|\alpha|=k} a_\alpha(x) D^\alpha u + a_0(D^{k-1} u, \dots, Du, u, x) = 0;$$

- (iii) the PDE is *quasilinear* if it has the form

$$\sum_{|\alpha|=k} a_\alpha(D^{k-1} u, \dots, Du, u, x) D^\alpha u + a_0(D^{k-1} u, \dots, Du, u, x) = 0;$$

- (iv) the PDE is *fully nonlinear* if it depends nonlinearly upon the highest order derivatives.

- **Def. (k -th Order System of PDE)** Fix an integer $k \geq 1$, and let U denote an open subset of $\mathbb{R}^n$. An expression of the form

$$\mathbf{F}(D^k \mathbf{u}(x), D^{k-1} \mathbf{u}(x), \dots, Du(x), \mathbf{u}(x), x) = \mathbf{0}, \quad (x \in U),$$

where $\mathbf{F} : \mathbb{R}^{mn^k} \times \mathbb{R}^{mn^{k-1}} \times \dots \times \mathbb{R}^{mn} \times \mathbb{R}^m \times U \rightarrow \mathbb{R}^m$ is given and $\mathbf{u} : U \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$ is the unknown, is called a k -th order system of PDE.

- **Def. (Well-Posed PDE)** A PDE is said to be *well-posed* iff:
 - (i) the PDE problem has a solution;
 - (ii) the solution is unique;
 - (iii) the solution depends *continuously* on the data given in the problem (i.e., “a small change in the initial data only changes the solution by a small amount”).
- **Def. (Classical Solution to a k -th Order PDE)** A function u is called a *classical solution* to a k -th order partial differential equation problem if and only if u satisfies the PDE and $u \in C^k(U)$. While higher regularity is desirable, this could be too much to ask for, in general (e.g., the scalar conservation law).

7.2.2 Four Important Linear Partial Differential Equations

Transport Equation

- **Def. (Transport Equation with Constant Coefficients)** The transport equation with constant coefficients is the PDE,

$$u_t + b \cdot Du = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

where $b = (b_1, \dots, b_n) \in \mathbb{R}^n$ is fixed, $u = u(x, t) : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$ is the unknown, and $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$ denotes the gradient of u with respect to the spatial variables x .

- **Def. (Initial-Value Problem)** The initial-value problem for the transport equation with constant coefficients is given by

$$\begin{cases} u_t + b \cdot Du = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where $b \in \mathbb{R}^n$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are known, and the problem is to compute the unknown function g .

- **Prop. (Solving Transport Equation with Constant Coefficients)** To determine which functions u solve the transport equation, we begin by assuming we have *some* solution u and try to compute this. First fix some point $(x, t) \in \mathbb{R}^n \times (0, \infty)$ and define

$$z(s) := u(x + sb, t + s) \quad (s \in \mathbb{R}).$$

Then we compute

$$z(s) = Du(x + sb, t + s) \cdot b + u_t(x + sb, t + s) = 0.$$

I.e., $z(\cdot)$ is a constant function of s . Consequently, for each point (x, t) , u is constant on the line through (x, t) with the direction $(b, 1) \in \mathbb{R}^{n+1}$. Therefore, if we know the value of u at any point on each such line we know its value everywhere in $\mathbb{R}^n \times (0, \infty)$. Thus given (x, t) as above, the line through (x, t) with direction $(b, 1)$ is represented parametrically by $(x + sb, t + s)$ ($s \in \mathbb{R}$). This line hits the plane $\Gamma := \mathbb{R}^n \times \{t = 0\}$ when $s = -t$, at the point $(x - tb, 0)$. Since u is constant on this line, and $u(x - tb, 0) = g(x - tb)$, we deduce that

$$u(x, t) = g(x - tb) \quad (x \in \mathbb{R}^n, t \geq 0).$$

So *if* the transport problem with constant coefficients has a sufficiently regular solution u , it must be given by the above equation. Conversely, if $g \in C^1$, then u defined by above is certainly a solution of the transport equation. If $g \notin C^1$, then the transport equation cannot have any C^1 solution. But we will declare $u(x, t) = g(x - tb)$ to be a *weak solution* of the transport equation.

- **Prop. (Solving Nonhomogeneous Transport Equation with Constant Coefficients)** Consider the associated nonhomogeneous problem

$$\begin{cases} u_t + b \cdot Du = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

As before, fix $(x, t) \in \mathbb{R}^{n+1}$, and set $z(s) = u(x + sb, t + s)$ for $s \in \mathbb{R}$. Then

$$z(s) = Du(x + sb, t + s) \cdot b + u_t(x + sb, t + s) = f(x + sb, t + s).$$

Consequently,

$$\begin{aligned} u(x, t) - g(x - tb) &= z(0) - z(-t) = \int_{-t}^0 z(s) \, ds \\ &= \int_{-t}^0 f(x + sb, t + s) \, ds \\ &= \int_0^t f(x + (s - t)b, s) \, ds. \end{aligned}$$

This means that

$$u(x, t) = g(x - tb) + \int_0^t f(x + (s - t)b, s) \, ds \quad (x \in \mathbb{R}^n, t \geq 0)$$

solves the initial-value problem.

Laplace's Equation

- **Def. (Laplace's and Poisson's Equations)** Fix an open subset $U \subseteq \mathbb{R}^n$, and let the unknown function be $u : \bar{U} \rightarrow \mathbb{R}$, $u = u(x)$ ($x \in U$). Also fix some function $f : U \rightarrow \mathbb{R}$. Then:

(1) (LAPLACE'S EQUATION) $\Delta u = 0$;

(2) (POISSON'S EQUATION) $\Delta u = f$;

where $\Delta u := \sum_{i=1}^n u_{x_i x_i}$ is the *Laplacian of u* .

- **Def. (Harmonic Function)** a C^2 function u satisfying Laplace's equation.
- **Prop. (Derivation of Laplace's Equation)** Let $U^{\text{open}} \subseteq \mathbb{R}^n$ be a fixed open subset, V any smooth subregion within U , and let u denote the density of some quantity (e.g., a chemical concentration) in equilibrium. Then the net flux of u through ∂V is zero (equilibrium means net flux is zero):

$$\int_{\partial V} \mathbf{F} \cdot \mathbf{v} \, dS = 0,$$

where $\mathbf{F}$ denotes the flux density and $\mathbf{v}$ denotes the unit outer normal field. By the Gauss-Green Theorem, one has

$$\int_V \operatorname{div} \mathbf{F} \, dx = \int_{\partial V} \mathbf{F} \cdot \mathbf{v} \, ds = 0,$$

so that

$$\operatorname{div} \mathbf{F} = 0$$

since V was arbitrary. Assume $\mathbf{F} = -aDu$ ($a > 0$). Then substituting into the above result, we have that

$$\Delta u = \operatorname{div}(Du) = 0.$$

- **Prop. (Derivation of Fundamental Solution to Laplace's Equation)** Our strategy to study Laplace's equation is to identify explicit solutions, then exploit linearity to assemble more complicated solutions. In particular, we will restrict attention to functions with certain symmetry properties. Thus, we will begin by looking for radial solutions (i.e., functions of $r = |x|$) since the Laplace equation is rotation invariant. Let $u(x) = v(r)$, where $r = |x| = (x_1^2 + \cdots + x_n^2)^{1/2}$ and v is to be selected (if possible) so that $\Delta u = 0$. First, we observe that for $i = 1, \dots, n$,

$$\frac{\partial r}{\partial x_i} = \frac{1}{2}(x_1^2 + \cdots + x_n^2)^{-1/2} 2x_i = \frac{x_i}{r} \quad (x \neq 0).$$

Therefore, we must have

$$u_{x_i} = v'(r) \frac{x_i}{r} \implies u_{x_i x_i} = v''(r) \frac{x_i^2}{r^2} + v'(r) \left(\frac{1}{r} - \frac{x_i^2}{r^3} \right).$$

Therefore,

$$\begin{aligned} \Delta u &= \sum_{i=1}^n u_{x_i x_i} = \frac{v''(r)}{r^2} \sum_{i=1}^n x_i^2 + \frac{v'(r)}{r} \sum_{i=1}^n 1 - \frac{v'(r)}{r^3} \sum_{i=1}^n x_i^2 \\ &= v''(r) + \frac{nv'(r)}{r} - \frac{v'(r)}{r} = v''(r) + \frac{n-1}{r} v'(r). \end{aligned}$$

Thus, $\Delta u = 0$ if and only if

$$v''(r) + \frac{n-1}{r} v'(r) = 0.$$

Suppose $v' \neq 0$. Then we have

$$\log(v')' = \frac{v''}{v'} = -\frac{n-1}{r},$$

so that $v'(r) = ar^{1-n}$ for some constant a . Consequently, if $r > 0$, we have

$$v(r) = \begin{cases} b \log r + c & (n = 2) \\ br^{2-n} + c & (n \geq 3) \end{cases},$$

where b and c are constants.

- **Def. (Fundamental Solutions of Laplace's Equation)** The function

$$\Phi(x) = \begin{cases} -\frac{1}{2\pi} \log |x| & (n = 2) \\ \frac{1}{n(n-2)\alpha(n)} \frac{1}{|x|^{n-2}} & (n \geq 3), \end{cases}$$

defined for $x \in \mathbb{R}^n \setminus \{0\}$, is the fundamental solution of Laplace's equation, where $\alpha(n)$ denotes the volume of the unit ball in $\mathbb{R}^n$. Note that the fundamental solution is radial as it only depends on $|x|$.

- **Prop. (Estimates for the Fundamental Solution)** Let $\Phi(x)$ denote the fundamental solution to Laplace's equation. Then, one has the following estimates:

$$|D\Phi(x)| \leq \frac{C}{|x|^{n-1}} \quad \text{and} \quad |D^2\Phi(x)| \leq \frac{C}{|x|^n} \quad (x \neq 0),$$

for some constant $C > 0$.

- **Obs. (Naive Solution to Poisson's Equation)** By construction, the function $x \mapsto \Phi(x)$ is harmonic for $x \neq 0$. Since shifting the origin to the new point $y \in \mathbb{R}^n$ leaves Laplace's equation intact, the map $x \mapsto \Phi(x - y)$ is also harmonic as a function of x , for $x \neq y$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Then it follows that the mapping $x \mapsto \Phi(x - y)f(y)$ ($x \neq y$) is harmonic for each point $y \in \mathbb{R}^n$, and thus so is the sum of finitely many such expressions built for different points y . Thus *one might think* that the convolution

$$u(x) = \int_{\mathbb{R}^n} \Phi(x - y)f(y) dy = \begin{cases} -\frac{1}{2\pi} \int_{\mathbb{R}^2} \log(|x - y|)f(y) dy & (n = 2) \\ \frac{1}{n(n-2)\alpha(n)} \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n-2}} dy & (n \geq 3) \end{cases}$$

will solve Laplace's equation. But this is not the case. The major obstacle is differentiation under the integral sign. In particular, this is because $D^2\Phi(x - y)$ is not integrable at $y = x$ so that differentiation through the integral sign is unjustified.

- **Prop. (Solving the Poisson Equation)** Let u be defined as

$$u(x) = \int_{\mathbb{R}^n} \Phi(x - y)f(y) dy,$$

where $f \in C_c^2(\mathbb{R}^n)$. Then

- (i) $u \in C^2(\mathbb{R}^n)$;
- (ii) $-\Delta u = f$ in $\mathbb{R}^n$.

Consequently, the above equation does give a formula for a solution of Poisson's equation in $\mathbb{R}^n$.

Proof. The proof of this result will be split into four steps. In the first step, we will establish that $u \in C^2(\mathbb{R}^n)$; in the second, we will compute Δu on some small ball since Φ blows up at the origin; in the third, we will continue working through Δu over small balls; finally, in the fourth step, we will take limits as our balls shrink to the origin.

- (1) Remember that convolutions are symmetric (i.e., $f * g = g * f$). Thus,

$$u(x) = \int_{\mathbb{R}^n} \Phi(x-y)f(y) dy = \int_{\mathbb{R}^n} \Phi(y)f(x-y) dy.$$

From this, if $h \neq 0$ and $e_i = (0, \dots, 1, \dots, 0)$ (the 1 in the i^{th} -slot), then

$$\frac{u(x + he_i) - u(x)}{h} = \int_{\mathbb{R}^n} \Phi(y) \left[\frac{f(x + he_i - y) - f(x - y)}{h} \right] dy.$$

Since $f \in C_c^2(\mathbb{R}^n)$,

$$\frac{f(x + he_i - y) - f(x - y)}{h} \xrightarrow{h \rightarrow 0} f_{x_i}(x - y) \text{ uniformly on } \mathbb{R}^n.$$

Therefore, since the limit can pass inside the integral sign,

$$u_{x_i}(x) = \int_{\mathbb{R}^n} \Phi(y) f_{x_i}(x - y) dy. \quad (i = 1, \dots, n).$$

Likewise,

$$u_{x_i x_j}(x) = \int_{\mathbb{R}^n} \Phi(y) f_{x_i x_j}(x - y) dy \quad (i, j = 1, \dots, n).$$

Since the integral on the right side is continuous in x , we conclude that $u \in C^2(\mathbb{R}^n)$.

- (2) Now we turn to the second result. Since Φ blows up at 0, we will need to isolate this singularity inside a small ball. Fix $\varepsilon > 0$. Then

$$\Delta u(x) = \int_{B(0, \varepsilon)} \Phi(y) \Delta_x f(x - y) dy + \int_{\mathbb{R}^n \setminus B(0, \varepsilon)} \Phi(y) \Delta_x f(x - y) dy =: I_\varepsilon + J_\varepsilon.$$

Let us see what happens with I_ε and J_ε . We first have³,

$$\begin{aligned} |I_\varepsilon| &= \left| \int_{B(0, \varepsilon)} \Phi(y) \Delta_x f(x - y) dy \right| \leq \int_{B(0, \varepsilon)} |\Phi(y)| |\Delta_x f(x - y)| dy \\ &\leq C \|D^2 f\|_{L^\infty(\mathbb{R}^n)} \int_{B(0, \varepsilon)} |\Phi(y)| dy \leq \begin{cases} C\varepsilon^2 |\log \varepsilon| & (n = 2) \\ C\varepsilon^2 & (n \geq 3) \end{cases} \end{aligned}$$

Therefore, as $\varepsilon \rightarrow 0$, $I_\varepsilon \rightarrow 0$. Now we shift to J_ε . Using integration by parts, one has

$$\begin{aligned} J_\varepsilon &= \int_{\mathbb{R}^n \setminus B(0, \varepsilon)} \Phi(y) \Delta_y f(x - y) dy \\ &= \int_{\mathbb{R}^n \setminus B(0, \varepsilon)} D\Phi(y) \cdot D_y f(x - y) dy + \int_{\partial B(0, \varepsilon)} \Phi(y) \frac{\partial f}{\partial \nu}(x - y) dS(y) \\ &=: K_\varepsilon + L_\varepsilon. \end{aligned}$$

where ν denotes the *inward* pointing unit normal along $\partial B(0, \varepsilon)$. We check

$$|L_\varepsilon| \leq \|Df\|_{L^\infty(\mathbb{R}^n)} \int_{\partial B(0, \varepsilon)} |\Phi(y)| dS(y) \leq \begin{cases} C\varepsilon |\log(\varepsilon)| & (n = 2) \\ C\varepsilon & (n \geq 3). \end{cases}$$

Thus, $L_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$.

³Recall that $\|D^2 f\|_{L^\infty} = \max_{1 \leq i, j \leq n} \|\partial_{x_i x_j} f\|_{L^\infty}$. On the other hand, one has (using polar coordinates),

$$\begin{aligned} \int_{B(0, \varepsilon)} |\log |x|| dx &= \int_0^\varepsilon \int_{\partial B(0, r)} r^{n-1} |\log(r)| dS dr = C' \int_0^\varepsilon r^{n-1} |\log(r)| dr \\ &= C' \cdot \varepsilon^n (-n^{-1} \log \varepsilon + n^{-2}) \leq C \cdot \varepsilon^n |\log \varepsilon| \leq C\varepsilon^2 |\log \varepsilon|, \end{aligned}$$

where without loss of generality, we assume that $0 < \varepsilon < 1$. This covers the $n = 2$ case. For $n \geq 3$, one has,

$$\int_{B(0, \varepsilon)} |x|^{2-n} dx = \int_0^\varepsilon \int_{\partial B(0, r)} r^{n-1} r^{2-n} dS dr = C' \cdot \int_0^\varepsilon r dr = C\varepsilon^2.$$

- (3) Now we investigate K_ε . If we can show that $K_\varepsilon \rightarrow -f(x)$ as $\varepsilon \rightarrow 0$, then we will be done. Using integration by parts, one has

$$\begin{aligned} K_\varepsilon &= \int_{\mathbb{R}^n \setminus B(0, \varepsilon)} \Delta \Phi(y) f(x-y) dy - \int_{\partial B(0, \varepsilon)} \frac{\partial \Phi}{\partial \nu}(y) f(x-y) dS(y) \\ &= - \int_{\partial B(0, \varepsilon)} \frac{\partial \Phi}{\partial \nu}(y) f(x-y) dS(y), \end{aligned}$$

since Φ is harmonic away from the origin. By definition of Φ , one has

$$D\Phi(y) = -\frac{1}{n\alpha(n)} \frac{y}{|y|^n} \quad (y \neq 0)$$

and

$$\nu = -\frac{y}{|y|} = -\frac{y}{\varepsilon} \quad (y \in \partial B(0, \varepsilon)).$$

Consequently,

$$\frac{\partial \Phi}{\partial \nu}(y) = \nu \cdot D\Phi(y) = \frac{1}{n\alpha(n)\varepsilon^{n-1}} \quad (y \in \partial B(0, \varepsilon)).$$

But $n\alpha(n)\varepsilon^{n-1}$ is the surface area of the sphere $\partial B(0, \varepsilon)$. Thus, we have

$$\begin{aligned} K_\varepsilon &= -\frac{1}{n\alpha(n)\varepsilon^{n-1}} \int_{\partial B(0, \varepsilon)} f(x-y) dS(y) \\ &= - \oint_{\partial B(x, \varepsilon)} f(y) dS(y) \xrightarrow{\varepsilon \rightarrow 0} -f(x) \end{aligned}$$

- (4) Therefore, combining all of the above computations, one has that in the limit $\varepsilon \rightarrow 0$, $-\Delta u \rightarrow f(x)$.

□

- **Thm. (Mean-Value Formulas for Laplace's Equation)** Fix an open subset $U \subseteq \mathbb{R}^n$. If $u \in C^2(U)$ is harmonic, then

$$u(x) = \oint_{\partial B(x, r)} u dS = \oint_{B(x, r)} u dy,$$

for each ball $B(x, r) \subset U$. I.e., the average of f over the sphere $\partial B(x, r)$ is equal to the average of f over the ball $B(x, r)$ for all such balls completely contained in U , and is equal to $u(x)$.

Proof. Our proof will consist of two steps: (1) in the first step, we will show that the first equality is true; (2) in the second step, we will show that the second equality is true.

- (1) Define

$$\phi(r) := \oint_{\partial B(x, r)} u(y) dS(y) = \oint_{\partial B(0, 1)} u(x + rz) dS(z).$$

Then differentiating with respect to r ,

$$\phi'(r) = \oint_{\partial B(0, 1)} Du(x + rz) \cdot z dS(z).$$

Consequently, by Green's formulas⁴,

$$\begin{aligned}\phi'(r) &= \oint_{\partial B(x,r)} Du(y) \cdot \frac{y-x}{r} \cdot dS(z) \quad ((y-x)/r \text{ is the unit normal}) \\ &= \oint_{\partial B(x,r)} \frac{\partial u}{\partial \nu} dS(y) \\ &= \frac{r}{n} \oint_{B(x,r)} \Delta u(y) dy = 0,\end{aligned}$$

since u is harmonic in U and $B(x,r) \subset U$. Hence, since ϕ is constant,

$$\phi(r) = \lim_{t \rightarrow 0} \phi(t) = \lim_{t \rightarrow 0} \oint_{\partial B(x,t)} u(y) dS(y) = u(x).$$

(2) Using polar coordinates (see, e.g., Theorem 2.49 from Folland), one has

$$\begin{aligned}\int_{B(x,r)} u dy &= \int_0^r \left(\int_{\partial B(x,s)} u dS \right) ds \\ &= u(x) \cdot \int_0^r n \alpha(n) s^{n-1} ds = \alpha(n) r^n u(x) \\ \implies u(x) &= \frac{1}{\alpha(n) r^n} \int_{B(x,r)} u dy = \oint_{B(x,r)} u dy.\end{aligned}$$

□

• **Thm. (Converse to Mean-Value Property)** If $u \in C^2(U)$ satisfies

$$u(x) = \oint_{\partial B(x,r)} u dS$$

for each ball $B(x,r) \subset U$, then u is harmonic.

Proof. Assume to the contrary that u is not harmonic (i.e., assume $\Delta u \neq 0$). Then there exists some ball $B(x,r)$ so that $\Delta u > 0$ within this ball. Define

$$\phi(r) = \oint_{\partial B(x,r)} u(y) dS(y).$$

Then by Green's formulas,

$$0 = \phi'(r) = \frac{r}{n} \oint_{\partial B(x,r)} \Delta u(y) dy > 0,$$

which is a contradiction. □

• **Thm. (Strong Maximum Principle)** Suppose $u \in C^2(U) \cap C(\bar{U})$ is harmonic within U .

- (i) $\max_{\bar{U}} u = \max_{\partial U} u$.
- (ii) If U is connected and there exists $x_0 \in U$ so that

$$u(x_0) = \max_{\bar{U}} u,$$

then u is constant within U .

⁴

By Green's Formulas,

$$\oint_{\partial B(x,r)} \frac{\partial u}{\partial \nu} dS(y) = \frac{1}{|\partial B(x,r)|} \int_{B(x,r)} \Delta u dy = \frac{|B(x,r)|}{|\partial B(x,r)|} \oint_{B(x,r)} \Delta u dy = \frac{r}{n} \oint_{B(x,r)} \Delta u(y) dy.$$

The first result is the *maximum principle* for Laplace's equation, while the second result is the *strong maximum principle*. The first asserts that if u has a maximum in U , it must also be the maximum of u on the closure of U ; the second asserts that if u achieves its maximum in U , then it is constant in U . Replacing u by $-u$ gives us similar assertions with the minimum instead of maximum.

Proof. We will first prove the second statement. The first statement follows from the first.

- (ii) Consider the set $\{x \in U : u(x) = M\}$, where $u(x_0) = M = \max_{\bar{U}} u$. Note that the set is nonempty as it contains (at least) the point x_0 . By the mean-value property,

$$M = u_{x_0} = \int_{B(x_0, r)} u \, dy \leq M,$$

where $0 < r < \text{dist}(x_0, \partial U)$. Equality holds if and only if $u \equiv M$ with $B(x_0, r)$ so that $u(y) = M$ for all $y \in B(x_0, r)$. This proves that the set $\{x \in U : u(x) = M\}$ is open in U . By continuity of M , $\{x \in U : u(x) = M\}$ is closed in U . Thus, since U is connected, $\{x \in U : u(x) = M\} = U$. Therefore, u is constant in U . □

- **Cor. (Positivity Principle)** Let U be a connected open subset of $\mathbb{R}^n$, and assume $u \in C^2(U) \cap C(\bar{U})$ satisfies

$$\begin{cases} \Delta u = 0 & \text{in } U \\ u = g & \text{on } \partial U, \end{cases}$$

where $g \geq 0$. Then u is positive everywhere in U if g is positive somewhere on ∂U .

Proof. Since u satisfies Laplace's equation in U , it follows that u is harmonic in U so that the strong maximum/minimum principles apply to u . By the weak minimum principle, for every $x \in \bar{U}$, $u(x) \geq \min_{\partial U} u = \min_{\partial U} g \geq 0$ so that $g(x)$ is nonnegative everywhere. Define the subset

$$A = \{z \in U : u(z) > 0\}.$$

If A is empty, then $u(x) = 0$ everywhere on U . Thus by continuity of u on $\bar{U}$, $u \equiv 0$ on ∂U so that $g \equiv 0$ on ∂U . But this contradicts our hypothesis that g is positive somewhere on ∂U . Therefore, A is nonempty. By continuity of u , A is open in U . Thus it suffices to show that A is relatively closed in U . Let $\{z_k\} \subset A$ be a sequence that converges to some $z \in U$. We must show that $z \in A$. Since $u(z_k) > 0$ for all k , $u(z) \geq 0$. Assume to the contrary that $u(z) = 0$. Then u achieves its minimum at z so that by the strong minimum principle, u must be zero everywhere in U , which is a contradiction. Thus by contradiction, $u(z) > 0$. I.e., $z \in A$, which completes the proof that A is relatively closed in U . Thus since U is connected, we conclude that $A = U$. □

- **Thm. (Uniqueness of Solutions to Laplace Equation)** Let $g \in C(\partial U)$, $f \in C(U)$. Then there exists at most one solution $u \in C^2(U) \cap C(\bar{U})$ of the boundary-value problem

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U. \end{cases}$$

Proof. Let u and $\tilde{u}$ satisfy the above boundary-value problem. Then we consider the function $w = \pm(u - \tilde{u})$. Since $(u - \tilde{u})|_{\partial U} = 0$, by the weak maximum principle, $\max_{\bar{U}} w = 0$. Likewise, by the weak minimum principle, $\min_{\bar{U}} w = 0$. Thus, $w = 0$ everywhere on U so that $u = \tilde{u}$ everywhere in U . □

- **Thm. (Smoothness of Harmonic Functions)** If $u \in C(U)$ satisfies the mean-value property for each $B(x, r) \subset U$, then $u \in C^\infty(U)$. Note that u may not necessarily be smooth, or even continuous, up to ∂U .

Proof. We will prove this result using mollifiers, and showing that u is equivalent to a smooth function locally. Let η be a standard mollifier (recall that η is a radial function). Set $u^\varepsilon = \eta_\varepsilon * u$ in $U_\varepsilon = \{x \in U : \text{dist}(x, \partial U) > \varepsilon\}$. Note that $u^\varepsilon \in C^\infty(U_\varepsilon)$. Fix some $x \in U^\varepsilon$. Then

$$\begin{aligned} u^\varepsilon(x) &= \int_U \eta_\varepsilon(x-y)u(y) dy \\ &= \frac{1}{\varepsilon^n} \int_{B(x,\varepsilon)} \eta\left(\frac{|x-y|}{\varepsilon}\right)u(y) dy \\ &= \frac{1}{\varepsilon^n} \int_0^\varepsilon \eta\left(\frac{r}{\varepsilon}\right) \left(\int_{\partial B(x,r)} u dS \right) dr \quad (\text{Polar Coordinates}) \\ &= \frac{1}{\varepsilon^n} u(x) \cdot \int_0^\varepsilon \eta\left(\frac{r}{\varepsilon}\right) n\alpha(n)r^{n-1} dr \quad (\text{Mean-Value Property}) \\ &= u(x) \int_{B(0,\varepsilon)} \eta_\varepsilon dy = u(x) \quad (\text{Property of Standard Mollifiers}). \end{aligned}$$

Therefore, $u^\varepsilon \equiv u$ in U_ε so that $u \in C^\infty(U_\varepsilon)$ for every $\varepsilon > 0$. This proves that u is smooth on U . $\square$

- **Thm. (Estimates on Derivatives of Harmonic Functions)** Assume u is harmonic in U . Then

$$|D^\alpha u(x_0)| \leq \frac{C_k}{r^{n+k}} \|u\|_{L^1(B(x_0,r))}$$

for each ball $B(x_0, r) \subset U$ and each multiindex α of order $|\alpha| = k$. Here

$$C_0 = \frac{1}{\alpha(n)}, C_k = \frac{(2^{n+1}nk)^k}{\alpha(n)} \quad (k = 1, \dots).$$

Proof. The proof of these results will follow from induction. We will first focus on the base cases $k = 0, 1$, and then work through the general case $k \geq 2$.

- (1) First assume $k = 0$. Since u is harmonic, it satisfies the mean-value property so that for all $x_0 \in U$ and each ball $B(x_0, r) \subset U$, one has

$$u(x_0) = \int_{B(x_0,r)} u dy \implies |u(x_0)| \leq \frac{1}{\alpha(n)r^n} \int_{B(x_0,r)} |u(y)| dy \leq \frac{1}{\alpha(n)r^n} \|u\|_{L^1(B(x_0,r))}$$

Now suppose $k = 1$. By differentiating Laplace's equation, we observe that u_{x_i} ($i = 1, \dots, n$) is harmonic for each i . This means that each u_{x_i} satisfies the mean-value property. Therefore,

$$\begin{aligned} |u_{x_i}(x_0)| &= \left| \int_{B(x_0,r/2)} u_{x_i} dx \right| \\ &= \left| \frac{2^n}{\alpha(n)r^n} \int_{\partial B(x_0,r/2)} uv_i dS \right| \\ &\leq \frac{2n}{r} \|u\|_{L^\infty(\partial B(x_0,r/2))}. \end{aligned}$$

Now if $x \in \partial B(x_0, r/2)$, then $B(x, r/2) \subset B(x_0, r) \subset U$, and so by the case for $k = 0$,

$$|u(x)| \leq \frac{1}{\alpha(n)} \left(\frac{2}{r}\right)^n \|u\|_{L^1(B(x_0,r))} \implies \|u\|_{L^\infty(\partial B(x_0,r/2))} \leq \frac{1}{\alpha(n)} \frac{2^n}{r^n} \|u\|_{L^1(B(x_0,r))}.$$

Therefore, combining the above inequalities, one has that if $|\alpha| = 1$, then

$$|D^\alpha u(x_0)| \leq \frac{2n}{r} \frac{1}{\alpha(n)} \frac{2^n}{r^n} \|u\|_{L^1(B(x_0,r))} = \frac{2^{n+1}n}{\alpha(n)r^{n+1}} \|u\|_{L^1(B(x_0,r))}.$$

This proves the case for $k = 1$.

- (2) Now we will prove the general case for $k \geq 2$. Assume that the estimates are valid for all balls in U and each multiindex of order less than or equal to $k - 1$. Fix $B(x_0, r) \subset U$ and let α be a multiindex with $|\alpha| = k$. Then $D^\alpha u = (D^\beta u)_{x_i}$ for some $i \in \{1, \dots, n\}$, $|\beta| = k - 1$. By repeatedly differentiating Laplace's equation, we can show that $D^\alpha u$ is harmonic. Thus by the mean-value formula, one has

$$\begin{aligned} |D^\alpha u(x_0)| &= \left| \int_{B(x_0, r/k)} D^\alpha u \, dx \right| \\ &= \left| \frac{k^n}{\alpha(n)r^n} \int_{\partial B(x_0, r/k)} D^\beta u v_i \, dS \right| \\ &\leq \frac{nk}{r} \|D^\beta u\|_{L^\infty(\partial B(x_0, r/k))}. \end{aligned}$$

If $x \in \partial B(x_0, r/k)$, then $B(x, r/k) \subset B(x_0, r) \subset U$. Therefore, by the estimates for $k - 1$, one has

$$|D^\beta u(x)| \leq \frac{(2^{n+1}n(k-1))^{k-1}}{\alpha(n) \left(\frac{k-1}{k}r\right)^{n+k-1}} \|u\|_{L^1(B(x_0, r))}.$$

Thus combining the two estimates gets us,

$$|D^\alpha u(x_0)| \leq \frac{(2^{n+1}nk)^k}{\alpha(n)r^{n+k}} \|u\|_{L^1(B(x_0, r))}.$$

Thus by induction, the estimate holds for all $k \geq 1$.

□

- **Thm. (Liouville's Theorem)** Suppose $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded. Then u is constant.

Proof. Fix $x_0 \in \mathbb{R}^n$, $r > 0$, and let $B(x_0, r)$ be the ball of radius r centered at x_0 . To show that u is constant, we just need to show that $Du \equiv 0$; connectedness of $\mathbb{R}^n$ will then show that u is constant everywhere on $\mathbb{R}^n$. By the above estimates, one has

$$\begin{aligned} |Du(x_0)| &\leq \frac{2^{n+1}n}{\alpha(n)r^{n+1}} \|u\|_{L^1(B(x_0, r))} \\ &= \frac{2^{n+1}n}{\alpha(n)r^{n+1}} \int_{B(x_0, r)} |u(x)| \, dx \\ &\leq \frac{2^{n+1}n}{\alpha(n)r^{n+1}} \int_{B(x_0, r)} \|u\|_{L^\infty(B(x_0, r))} \, dx \\ &= \frac{2^{n+1}n}{r} \|u\|_{L^\infty(B(x_0, r))} \leq \frac{2^{n+1}n}{r} \|u\|_{L^\infty(\mathbb{R}^n)} \xrightarrow{r \rightarrow \infty} 0. \end{aligned}$$

Therefore, $Du \equiv 0$; by connectedness of $\mathbb{R}^n$, we conclude that u is constant. □

- **Thm. (Representation Formula)** Let $f \in C_c^2(\mathbb{R}^n)$, $n \geq 3$. Then any bounded solution of $-\Delta u = f$ in $\mathbb{R}^n$ has the form

$$u(x) = \int_{\mathbb{R}^n} \Phi(x - y) f(y) \, dy + C \quad (x \in \mathbb{R}^n)$$

for some constant C .

Proof. For $n \geq 3$, $\Phi(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Therefore, $\tilde{u}(x) = \int_{\mathbb{R}^n} \Phi(x - y) f(y) \, dy$ is a bounded solution of $-\Delta u = f$ in $\mathbb{R}^n$. If u is another solution, $w = u - \tilde{u}$ is constant according to Liouville's Theorem. As a side remark, for $n = 2$, $\Phi(x)$ is unbounded as $|x| \rightarrow \infty$, so that the above form may also be unbounded. □

- **Thm. (Analyticity of Harmonic Functions)** Assume u is harmonic in U . Then u is analytic in U .

Proof. Fix any point $x_0 \in U$. Our goal is to show that u can be represented by a convergent power series in some neighborhood of x_0 . Our proof will be split into multiple steps.

- (1) Let $r = \frac{1}{4} \text{dist}(x_0, \partial U)$. Then since $u \in C^2(U)$, $u \in L^1(B(x_0, 2r))$ so that the quantity

$$M = \frac{1}{\alpha(n)r^n} \|u\|_{L^1(B(x_0, 2r))} < \infty.$$

- (2) Since $B(x, r) \subset B(x_0, 2r) \subset U$ for each $x \in B(x_0, r)$, one has

$$\begin{aligned} |D^\alpha u(x)| &\leq \left(\frac{2^{n+1}n|\alpha|}{\alpha(n)} \right)^{|\alpha|} \cdot \frac{1}{r^{n+|\alpha|}} \|u\|_{L^1(B(x_0, 2r))} \\ &= \left(\frac{2^{n+1}n}{r} \right)^{|\alpha|} \frac{1}{\alpha(n)^{|\alpha|}r^n} \|u\|_{L^1(B(x_0, 2r))} |\alpha|^{|\alpha|} \\ &= C' M \cdot \left(\frac{2^{n+1}n}{r} \right)^{|\alpha|} |\alpha|^{|\alpha|} \\ \implies \|D^\alpha u\|_{L^\infty(B(x_0, r))} &\leq C' \cdot M \left(\frac{2^{n+1}n}{r} \right)^{|\alpha|} |\alpha|^{|\alpha|}, \end{aligned}$$

where C' is some constant that depends solely on n . Then since $k^k/k! < e^k$ for all positive integers k , one has

$$|\alpha|^{|\alpha|} \leq e^{|\alpha|} |\alpha|!$$

for all multiindices α . By the multinomial theorem, one has

$$n^k = (1 + \dots + 1)^k = \sum_{|\alpha|=k} \frac{|\alpha|!}{\alpha!},$$

so that $|\alpha|! \leq n^{|\alpha|} \alpha!$. Thus,

$$\|D^\alpha u\|_{L^\infty(B(x_0, r))} \leq CM \left(\frac{2^{n+1}n^2 e}{r} \right)^{|\alpha|} \alpha!,$$

where C is some constant that depends on n .

- (3) The Taylor series for u at x_0 is

$$\sum_{\alpha} \frac{D^\alpha u(x_0)}{\alpha!} (x - x_0)^\alpha,$$

where the sum is taken over all multiindices α . We will now show that this power series converges provided $|x - x_0| < \frac{r}{2^{n+2}n^3 e}$. To show this, it suffices to show that the remainder term tends to zero. I.e., we begin by computing,

$$\begin{aligned} R_N(x) &:= u(x) - \sum_{k=0}^{N-1} \sum_{|\alpha|=k} \frac{D^\alpha u(x_0)}{\alpha!} (x - x_0)^\alpha \\ &= \sum_{|\alpha|=N} \frac{D^\alpha u(x_0 + t(x - x_0))}{\alpha!} (x - x_0)^\alpha, \end{aligned}$$

for some $0 \leq t \leq 1$ depending on x , where we obtain the above formula by writing out the first N terms and the error in their Taylor expansion about 0 for the function of one variable

$g(t) = u(x_0 + t(x - x_0))$ at $t = 1$. So using the estimates above and from (2), we have

$$\begin{aligned} |R_N(x)| &\leq CM \sum_{|\alpha|=N} \left(\frac{2^{n+1}n^2e}{r} \right)^N \left(\frac{r}{2^{n+2}n^3e} \right)^N \\ &= CM \sum_{|\alpha|=N} \frac{1}{(2n)^N} \leq CM n^N \frac{1}{(2n)^N} = \frac{CM}{2^N} \xrightarrow{N \rightarrow \infty} 0. \end{aligned}$$

Thus the proof concludes. □

- **Thm. (Harnack's Inequality)** For each connected open set $V \Subset U$ (i.e., $V \subset \bar{V} \subset U$ with $\bar{V}$ compact), there exists a positive constant C , depending only on V such that

$$\sup_V u \leq C \inf_V u,$$

for all nonnegative harmonic functions in U .

Proof. [!! Complete Later !!] □

- **Thm. (Representation Formula for Solution to Poisson's Equation using Green's Function)**
Assume now that $U \subset \mathbb{R}^n$ is open, bounded, and ∂U is C^1 . If $u \in C^2(\bar{U})$ solves the boundary-value problem

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U, \end{cases}$$

then

$$u(x) = - \int_{\partial U} g(y) \frac{\partial G}{\partial \nu}(x, y) dS(y) + \int_U f(y) G(x, y) dy \quad (x \in U),$$

where $G = G(x, y)$ is Green's function for the given domain U .

Proof. Suppose $u \in C^2(\bar{U})$ is an arbitrary function. Fix $x \in U$, and choose $\varepsilon > 0$ so small so that $B(x, \varepsilon) \subset U$, and apply Green's formula on the region $V_\varepsilon := U \setminus B(x, \varepsilon)$ to $u(y)$ and $\Phi(y - x)$. We compute

$$\int_{V_\varepsilon} u(y) \Delta \Phi(y - x) - \Phi(y - x) \Delta u(y) dy = \int_{\partial V_\varepsilon} u(y) \frac{\partial \Phi}{\partial \nu}(y - x) - \Phi(y - x) \frac{\partial u}{\partial \nu}(y) dS(y),$$

where ν denotes the outer unit normal vector on ∂V_ε . For $x \neq y$, $\Delta \Phi(x - y) = 0$. Moreover⁵,

$$\left| \int_{\partial B(x, \varepsilon)} \Phi(y - x) \frac{\partial u}{\partial \nu}(y) dS(y) \right| \leq C \varepsilon^{n-1} \max_{\partial B(0, \varepsilon)} |\Phi| = o(1) \text{ as } \varepsilon \rightarrow 0.$$

Furthermore,

$$\int_{\partial B(x, \varepsilon)} u(y) \frac{\partial \Phi}{\partial \nu}(y - x) dS(y) = \oint_{\partial B(x, \varepsilon)} u(y) dS(y) \rightarrow u(x)$$

as $\varepsilon \rightarrow 0$. Thus in the limit $\varepsilon \rightarrow 0$,

$$u(x) = \int_{\partial U} \Phi(y - x) \frac{\partial u}{\partial \nu}(y) - u(y) \frac{\partial \Phi}{\partial \nu}(y - x) dS(y) - \int_U \Phi(y - x) \Delta u(y) dy.$$

This integral is valid for any $x \in U$ and $u \in C^2(\bar{U})$. If we knew the values of Δu inside U and the values of u , $\partial u / \partial \nu$ along ∂U , then we could use the above formula to solve for $u(x)$. However, we do not know $\partial u / \partial \nu$. Thus we must remove this term from the above result. For a fixed x , we introduce the corrector function $\phi^x(y)$ which solves the boundary value problem

$$\begin{cases} \Delta \phi^x = 0 & \text{in } U \\ \phi^x = \Phi(y - x) & \text{in } \partial U. \end{cases}$$

Applying Green's Formula, one obtains

$$\begin{aligned} - \int_U \phi^x(y) \Delta u(y) dy &= \int_{\partial U} u(y) \frac{\partial \phi^x}{\partial \nu}(y) - \phi^x(y) \frac{\partial u}{\partial \nu}(y) dS(y) \\ &= \int_{\partial U} u(y) \frac{\partial \phi^x}{\partial \nu}(y) - \Phi(y - x) \frac{\partial u}{\partial \nu}(y) dS(y). \end{aligned}$$

5

$$\begin{aligned} \left| \int_{\partial B(x, \varepsilon)} \Phi(y - x) \frac{\partial u}{\partial \nu}(y) dS(y) \right| &\leq \int_{\partial B(x, \varepsilon)} |\Phi(y - x)| \left| \frac{\partial u}{\partial \nu}(y) \right| dS(y) = \int_{\partial B(0, \varepsilon)} |\Phi(t)| \left| \frac{\partial u}{\partial \nu}(t + x) \right| dS(t) \\ &\leq \int_{\partial B(0, \varepsilon)} \|\Phi\|_{L^\infty(\partial B(0, \varepsilon))} \left\| \frac{\partial u}{\partial \nu}(x + \cdot) \right\|_{L^\infty(\partial B(0, \varepsilon))} dS(t) = \int_{\partial B(0, \varepsilon)} \|\Phi\|_{L^\infty(\partial B(0, \varepsilon))} \left\| \frac{\partial u}{\partial \nu} \right\|_{L^\infty(\partial B(x, \varepsilon))} dS(t) \\ &= \|\Phi\|_{L^\infty(\partial B(0, \varepsilon))} \left\| \frac{\partial u}{\partial \nu} \right\|_{L^\infty(\partial B(x, \varepsilon))} n\alpha(n) \varepsilon^{n-1}. \end{aligned}$$

So now suppose that $G(x, y) := \Phi(y - x) - \phi^x(y)$ ($x, y \in U, x \neq y$) (i.e., *Green's function for the region U*). Using this terminology and adding the above result to the original expansion for u , one obtains

$$u(x) = - \int_{\partial U} u(y) \frac{\partial G}{\partial \nu}(x, y) dS(y) - \int_U G(x, y) \Delta u(y) dy \quad (x \in U),$$

where $\partial G / \partial \nu(x, y) = D_y G(x, y) \cdot \nu(y)$ is the outer normal derivative of G with respect to the variable y . Since $\partial u / \partial \nu$ does not appear in this fixed equation, the corrector function ϕ^x was chosen correctly. Suppose now that $u \in C^2(\bar{U})$ satisfies the above boundary-value problem. Then making the necessary substitutions, one obtains the representation formula given above. $\square$

- **Thm. (Symmetry of Green's Function)** For all $x, y \in U, x \neq y$, we have $G(y, x) = G(x, y)$.
- **[!! Fill in Green's Functions for Different Geometries !!]**
- **Thm. (Uniqueness of Solution to Poisson Problem; Energy Method)** Consider the boundary-value problem

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U, \end{cases}$$

where U is open, bounded, and ∂U is C^1 . Then there exists at most one solution $u \in C^2(\bar{U})$ of the above problem.

Proof. Assume $\tilde{u}$ is another solution to the given boundary-value problem. Set $w = u - \tilde{u}$. Then $\Delta w = 0$ in U , which means that by integration by parts,

$$0 = - \int_U w \Delta w dx = \int_U |Dw|^2 dx.$$

Since $|Dw|^2 \geq 0$, it follows that $Dw \equiv 0$ in U . Since $w = 0$ on ∂U , we deduce that $w \equiv 0$ in U so that $u \equiv \tilde{u}$. $\square$

- **Def. (Energy Functional for Laplace's Equation)** Define the *energy functional*,

$$I[w] := \int_U \frac{1}{2} |Dw|^2 - wf dx,$$

where w belongs to the *admissible set*

$$\mathcal{A} := \{w \in C^2(\bar{U}) : w = g \text{ on } \partial U\}.$$

- **Thm. (Dirichlet's Principle)** Assume $u \in C^2(\bar{U})$ solves the Poisson boundary-value problem. Then

$$I[u] = \min_{w \in \mathcal{A}} I[w].$$

Conversely, if $u \in \mathcal{A}$ satisfies the above condition, then u solves the Poisson boundary-value problem. I.e., a solution of the Poisson boundary-value problem can be characterized as the minimizer of the energy functional.

Proof. The proof will proceed in two steps. In the first, we will show that $I[u]$ is indeed the minimum of $I[w]$; in the second, we will show that if u is the minimizer of the energy functional, then it solves Poisson's equation.

(1) **[!! Complete Later !!]**

$\square$

Heat Equation

- **Def. ((Non)homogeneous Heat Equation)** The *(non)homogeneous heat equation* is the following:

$$u_t - \Delta u = 0 \ (f),$$

subject to appropriate initial and boundary conditions, where $t > 0$ and $x \in U^{\text{open}} \subseteq \mathbb{R}^n$. The Laplacian Δ is taken with respect to the spatial variables, and in the second case, f is some given function $f : U \times [0, \infty) \rightarrow \mathbb{R}$.

- **Obs. (Physical Interpretation of the Heat Equation)** The heat equation describes the time evolution of the density u of some quantity. If $V \subset U$ is any smooth subregion, the rate of change of the total quantity within V equals the negative of the net flux through ∂V :

$$\frac{d}{dt} \int_V u \, dx = - \int_{\partial V} \mathbf{F} \cdot \mathbf{v} \, dS,$$

where $\mathbf{F}$ is the flux density. Since V was arbitrary, one has $u_t = -\text{div} \mathbf{F}$. If one assumes that $\mathbf{F} = -a \nabla u$ ($a > 0$), then we obtain the PDE

$$u_t = a \, \text{div}(\nabla u) = a \Delta u,$$

which for $a = 1$ is the heat equation.

- **Prop. (Deriving the Fundamental Solution to the Heat Equation)** We begin our study of the heat equation by trying to find *specific* solutions. We begin by looking for solutions u that are *invariant under dilation scaling*. I.e., we look for solutions u so that for all $\lambda > 0$, $x \in \mathbb{R}^n$, and $t > 0$,

$$u(x, t) = \lambda^\alpha u(\lambda^\beta x, \lambda t),$$

where the constants α and β must be found. This leads to us looking for solutions u having the special structure,

$$u(x, t) = \frac{1}{t^\alpha} v\left(\frac{x}{t^\beta}\right) \quad (x \in \mathbb{R}^n, t > 0).$$

7.2.3 Sobolev Spaces

- **Def. (Lipschitz Continuous Functions)** Assume $U^{\text{open}} \subseteq \mathbb{R}^n$ is open, and let $0 < \gamma \leq 1$. $u : U \rightarrow \mathbb{R}$ is said to be *Lipschitz continuous* if and only if

$$|u(x) - u(y)| \leq C|x - y| \quad (x, y \in U),$$

for some constant C . In particular, it follows that u is uniformly continuous.

- **Def. (Hölder Continuous Functions)** A function $u : U \rightarrow \mathbb{R}$ is said to be *Hölder continuous with exponent γ* if and only if

$$|u(x) - u(y)| \leq C|x - y|^\gamma \quad (x, y \in U),$$

for some constant C and $0 < \gamma \leq 1$.

- **Lem. (Constant Hölder Continuous Functions)** If $u : U \rightarrow \mathbb{R}$ is Hölder continuous with exponent $\gamma > 1$, where U is connected, then u is constant.

Proof. Let U be a connected open subset of $\mathbb{R}^n$ and assume that $u : U \rightarrow \mathbb{R}$ is Hölder continuous with exponent $\gamma > 1$. Fix an arbitrary $x \in U$, and let e_i be a basis vector of $\mathbb{R}^n$. For $h \in \mathbb{R}$ small enough so that $x + he_i \in U$, one has by Hölder continuity

$$|u(x + he_i) - u(x)| \leq C|h|^\gamma \implies 0 \leq \left| \frac{u(x + he_i) - u(x)}{h} \right| \leq C|h|^{\gamma-1}.$$

Since $\gamma - 1 > 0$, the RHS tends to zero. On the other hand, since u is differentiable, the middle term tends to $D_{e_i}u(x)$. Thus since e_i and x were arbitrary, one has that $Du(x) = 0$ for all $x \in U$. Thus since U is connected, we conclude that $Du \equiv 0$, which means that u is constant. $\square$

- **Def. (Hölder Norms)** We define the following:

(i) (UNIFORM NORM) If $u : U \rightarrow \mathbb{R}$ is bounded and continuous, we write

$$\|u\|_{C(\bar{U})} := \sup_{x \in \bar{U}} |u(x)|.$$

(ii) The γ^{th} -Hölder seminorm of $u : U \rightarrow \mathbb{R}$ is

$$[u]_{C^{0,\gamma}(\bar{U})} := \sup_{\substack{x, y \in \bar{U} \\ x \neq y}} \left\{ \frac{|u(x) - u(y)|}{|x - y|^\gamma} \right\},$$

and the γ^{th} -Hölder norm is

$$\|u\|_{C^{0,\gamma}(\bar{U})} := \|u\|_{C(\bar{U})} + [u]_{C^{0,\gamma}(\bar{U})}.$$

- **Def. (Hölder Space, $C^{k,\gamma}(\bar{U})$)** Let $k \geq 1$ be a positive integer. Then we define

$$C^{k,\gamma}(\bar{U}) := \left\{ u \in C^k(\bar{U}) : \|u\|_{C^{k,\gamma}(\bar{U})} := \sum_{|\alpha| \leq k} \|D^\alpha u\|_{C(\bar{U})} + \sum_{|\alpha|=k} [D^\alpha u]_{C^{0,\gamma}(\bar{U})} < \infty \right\}.$$

- **Thm. (Hölder and Banach Spaces)** The space of functions $C^{k,\gamma}(\bar{U})$ is a Banach space under the aforementioned norm.
- **Def. (Test Function)** A function $f \in C_c^\infty(U)$.

- **Def. (Weak Partial Derivatives)** Suppose $u, v \in L^1_{\text{loc}}(U)$ and α is a multiindex. We say that α is the α^{th} -weak partial derivative of u , written

$$D^\alpha u = v,$$

if and only if

$$\int_U u D^\alpha \phi \, dx = (-1)^{|\alpha|} \int_U v \phi \, dx,$$

for all test functions $\phi \in C_c^\infty(U)$.

- **Obs. (Motivation for Weak Derivatives)** Suppose $u \in C^1(U)$. Then if $\phi \in C_c^\infty(U)$, by the integration by parts formula,

$$\int_U u \phi_{x_i} \, dx = - \int_U u_{x_i} \phi \, dx \quad (i = 1, \dots, n).$$

Note that there are no boundary terms since ϕ has compact support in U and vanishes near ∂U . More generally, if k is a positive integer, $u \in C^k(U)$, and $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multiindex of order $|\alpha| = \alpha_1 + \dots + \alpha_n = k$, then

$$\int_U u D^\alpha \phi \, dx = (-1)^{|\alpha|} \int_U D^\alpha u \phi \, dx,$$

obtained by induction. We want to see if this formula is true even if $u \notin C^k(U)$. The left side is defined regardless of the regularity of u so long as it is locally integrable. Thus, we replace the definition of $D^\alpha u$ as suggested above.

- **Lem. (Uniqueness of Weak Derivatives)** A weak α^{th} -partial derivative of u , if it exists, is uniquely defined up to a set of measure zero.

Proof. Assume that $v, \tilde{v} \in L^1_{\text{loc}}(U)$ satisfy

$$\int_U u D^\alpha \phi \, dx = (-1)^{|\alpha|} \int_U v \phi \, dx = (-1)^{|\alpha|} \int_U \tilde{v} \phi \, dx$$

for all $\phi \in C_c^\infty(U)$. Then

$$\int_U (v - \tilde{v}) \phi \, dx = 0$$

for all $\phi \in C_c^\infty(U)$, whence $v - \tilde{v} = 0$ a.e. □

- **Cor. (Weak Derivative in $\mathbb{R}$)** Let $n = 1$, $U = (0, 2)$, and

$$u(x) =$$

- **Def. (Sobolev Space)** Fix $1 \leq p \leq \infty$ and let k be a nonnegative integer. Then we define the Sobolev space

$$W^{k,p}(U) := \left\{ u : U \rightarrow \mathbb{R} \mid u \in L^1_{\text{loc}}(U) \text{ and } \forall |\alpha| \leq k, D^\alpha u \text{ exists in the weak sense and belongs to } L^p(U) \right\}$$

7.3 Semidirect Products

- **Ex. 1 (Classification of Groups of Order pq)** Consider the finite group G of order pq , with p, q distinct primes and without loss of generality, $p < q$. By Sylow's theorems, it follows that G contains a unique (hence normal) Sylow q -subgroup Q . Let P denote a Sylow p -subgroup. Then it follows that $G \cong Q \rtimes_{\varphi} P$, where $\varphi : P \rightarrow \text{Aut}(Q)$ is a group homomorphism. Since P and Q are both of prime order, both subgroups are cyclic. Moreover, $\text{Aut}(Q) \cong (\mathbb{Z}/q\mathbb{Z})^{\times} \cong \mathbb{Z}/(q-1)\mathbb{Z}$. We consider two cases.

(Case I) First consider the case $p \nmid (q-1)$. Then as in Problem 2024-J-I, the only homomorphism from Q to $\text{Aut}(P)$ is the trivial homomorphism. Therefore, it follows that in this case, the only possible semidirect product is the direct product. I.e., $G \cong \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/q\mathbb{Z} \cong \mathbb{Z}/(pq)\mathbb{Z}$. Thus up to isomorphism, there exists exactly one group of order pq , which is cyclic and abelian.

(Case II) Now consider the case $p \mid (q-1)$. Let $P = \langle y \rangle$. Since $\text{Aut}(Q)$ is cyclic of order $q-1$, and $p \mid (q-1)$, by Cauchy's theorem, $\text{Aut}(Q)$ contains a unique subgroup of order p , $\langle \gamma \rangle$. Then any homomorphism $\varphi : P \rightarrow \text{Aut}(Q)$ must map y to a generator of $\langle \gamma \rangle$. Thus in fact, there are p homomorphisms: $\varphi_i : y \mapsto \gamma^i$. Note that when $i = 0$, φ_0 corresponds to the trivial homomorphism so that $G \cong Q \rtimes_{\varphi_0} P \cong Q \times P \cong \mathbb{Z}/(pq)\mathbb{Z}$. On the other hand, all of the other homomorphisms are isomorphic since for every $i > 0$, there exists some generator y_i of P so that $p_i(y_i) = \gamma$. Thus up to isomorphism, there exist exactly 2 groups of order pq with $p < q$; one is cyclic and abelian, and the other is nonabelian.

- **Ex. 2 (Classification of Groups of Order 30)** Let G be a group of order $30 = 2 \cdot 3 \cdot 5$. We begin by showing that at least one of n_3 and n_5 must be one. By Sylow's theorems, one has

$$n_3 \in \{1, 2, 5, 10\} \cap \{1, 4, 7, 10, \dots\} = \{1, 10\}.$$

$$n_5 \in \{1, 2, 3, 6\} \cap \{1, 11, 16, 21, \dots\} = \{1, 6\}.$$

Suppose $n_3 = 10$ and $n_5 = 6$. Since each Sylow 5-subgroup and each Sylow 3-subgroup has prime order, they must all intersect trivially. Moreover, each Sylow 5-subgroup has trivial intersection with each Sylow 3-subgroup by Lagrange's theorem. Thus, there are 30 elements of G with order 5, and 20 elements of G with order 3. This means that $|G| > 30 + 20 = 50 > 30 = |G|$, which is a contradiction. Thus G must have a normal Sylow 3-subgroup or a normal Sylow 5-subgroup. In any case, it then follows that G must have a subgroup H of order 50. Since H has index 2 in G , it must be normal and cyclic in G . Therefore, denoting a Sylow 2-subgroup of G by K , $G \cong H \rtimes_{\varphi} K$ for some homomorphism $\varphi : K \rightarrow \text{Aut}(H)$. Then since $H \cong \mathbb{Z}_{15}$, one has

$$\text{Aut}(\mathbb{Z}_{15}) \cong (\mathbb{Z}/15\mathbb{Z})^{\times} \cong (\mathbb{Z}/5\mathbb{Z})^{\times} \times (\mathbb{Z}/3\mathbb{Z})^{\times} \quad (\text{Chinese Remainder Theorem})$$

Thus, $\text{Aut}(\mathbb{Z}_{15}) \cong \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.