

1 Basic Category Theory

Remark 1.1. The notion of “class” in ZFC requires a definition. This can be done (“parametrically definable classes”), but for now let’s blackbox the concept and just say “classes are a generalization of sets which are allowed to be ‘large’”, e.g. so a “class of all sets” makes sense even though a “set of all sets” does not. This shouldn’t matter ultimately anyways.

Definition 1.1. A **Category**, $\mathcal{C}$, is a triple of classes, $\text{obj}(\mathcal{C})$, $\text{Hom}(\mathcal{C})$, and “ $\circ$ ”, AKA $\text{Comp}(\mathcal{C})$, such that:

- For every $a, b \in \text{obj}(\mathcal{C})$, there is a set $\text{Hom}_{\mathcal{C}}(a, b)$ of elements of $\text{Hom}(\mathcal{C})$.
- For every $a, b, c \in \text{obj}(\mathcal{C})$, composition is a “map” $\circ : \text{Hom}_{\mathcal{C}}(a, b) \times \text{Hom}_{\mathcal{C}}(b, c) \rightarrow \text{Hom}_{\mathcal{C}}(a, c)$
 - Composition is associative, in that $(f \circ g) \circ h = f \circ (g \circ h)$ for any morphisms $f, g, h \in \text{Hom}(\mathcal{C})$ (whenever this composition makes sense).
- For every $a \in \text{obj}(\mathcal{C})$, there is an element $\text{id}_a \in \text{Hom}_{\mathcal{C}}(a, a)$ which is an identity for composition.

In short: we have objects, arrows between objects, a way of composing arrows, and identity arrows on each object. Oftentimes, when working with a category of modules or vector spaces, we will omit the name of the full category and simply denote $\text{Hom}_R(M, N)$ for (say) $\text{Hom}_{R\text{Mod}}(M, N)$.

Definition 1.2. Given a category $\mathcal{C}$, we define its opposite category $\mathcal{C}^{op}$ to be the category with the same objects, but such that for every morphism $f : a \rightarrow b$ in $\text{Hom}(\mathcal{C})$ we have an arrow $f^{op} : b \rightarrow a$ in $\text{Hom}(\mathcal{C}^{op})$. Composition is defined by saying that, for every

$$a \xrightarrow{f} b \xrightarrow{g} c$$

in $\text{Hom}(\mathcal{C})$,

$$c \xrightarrow{(g \circ f)^{op}} a := c \xrightarrow{g^{op}} b \xrightarrow{f^{op}} a$$

in $\text{Hom}(\mathcal{C}^{op})$.

Definition 1.3. Given two categories, $\mathcal{C}$ and $\mathcal{D}$, a **functor** $F : \mathcal{C} \rightarrow \mathcal{D}$ is a map on objects $F_{\text{obj}} : \text{obj}(\mathcal{C}) \rightarrow \text{obj}(\mathcal{D})$ together with a collection of maps on morphisms $F_{\text{Hom}(a,b)} : \text{Hom}_{\mathcal{C}}(a, b) \rightarrow \text{Hom}_{\mathcal{D}}(Fa, Fb)$. By convention we omit any subscripts and, for any $\mathcal{C}$ -object a or any $\mathcal{C}$ -morphism φ , simply denote Fa and $F\varphi$ for the corresponding object/morphism in $\mathcal{D}$.

Now, sometimes we wish to talk about things which don’t *respect* composition, but instead reverse it. These still get to be called functors, but we call those “**contravariant functors**” and (even if we *really*) think of them as “composition reversing functors $\mathcal{C} \rightarrow \mathcal{D}$ ” it’ll end up being nice to think of these quite formally as functors $\mathcal{C}^{op} \rightarrow \mathcal{D}$ —e.g. they “preserve reverse composition of arrows”.

Now we know what a functor is, we should talk about some examples and basic properties thereof.

Definition 1.4. A functor, $F : \mathcal{C} \rightarrow \mathcal{D}$, is said to be:

- **faithful** if the induced map $\text{Hom}_{\mathcal{C}}(a, b) \rightarrow \text{Hom}_{\mathcal{D}}(Fa, Fb)$ is injective for all $a, b \in \text{obj}(\mathcal{C})$
- **full** if the induced map $\text{Hom}_{\mathcal{C}}(a, b) \rightarrow \text{Hom}_{\mathcal{D}}(Fa, Fb)$ is surjective for all $a, b \in \text{obj}(\mathcal{C})$
- **bijective** if the induced map $\text{Hom}_{\mathcal{C}}(a, b) \rightarrow \text{Hom}_{\mathcal{D}}(Fa, Fb)$ is bijective for all $a, b \in \text{obj}(\mathcal{C})$

Now, in many contexts we want to compare two categories, talk about whether or not they’re the “same”. In such circumstances we *could* require that we have a pair of inverse functors—an “isomorphism of categories”—but the problem is that this is way too strong (to the point of coming down to “how strict are we about what sets we use to define our objects”). It is, for instance, perfectly valid linear algebra-wise to imagine all finite $\mathbb{R}$ -vector spaces as just “vector spaces $\mathbb{R}^n$ ”, but the two categories *aren’t* equivalent because one only has a $\mathbb{Z}^+$ worth of sets while the other has way more. The notion of “isomorphism” is thus the “wrong definition”. For a better definition, rather than being “bijective on objects”, we require a sort of weaker notion:

Definition 1.5. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is said to be **essentially surjective** if, for every $d \in \text{obj}(\mathcal{D})$, there exists some $c \in \text{obj}(\mathcal{C})$ such that $Fc \cong d$.

That is to say, F “reaches all of $\mathcal{D}$ (up to abstract isomorphism)” With this in mind, we can formulate the following:

Definition 1.6. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is said to be a **(weak) equivalence of categories** if F is fully faithful and essentially surjective.

This is to say: morphisms in $\mathcal{C}$ and $\mathcal{D}$ can be thought of interchangeably—so long as you’re willing to transport along abstract isomorphisms in $\mathcal{D}$. Subscribing to the philosophy that “objects are but vessels for morphisms” (hey, I don’t wanna bother with the yoneda lemma here), this tells us the two categories are “morally” the same. The reason for the parenthetical “weak” is more or less a “you can choose to be careful about the axiom of choice”—in particular, there’s a natural notion of a “(homotopy) equivalence of categories”—cf. 1.9, which is nice and cool but requires a bit more terminology. With Axiom of choice these are the same, but without axiom of choice this notion is “weaker”. Jason has said he often means “weak equivalence” when he says equivalence though (even though he’s distinguished them in the past?), so mostly it’s a “have the homotopic notion in mind morally, and the weak notion in mind for like 80% of practical examples.”

Definition 1.7. Given two functors, $F, G : \mathcal{C} \rightarrow \mathcal{D}$, a **natural transformation** $\eta_\bullet : F \rightarrow G$ is a collection of maps for each $a \in \text{obj}(\mathcal{C})$, $\eta_a : Fa \rightarrow Ga$ such that for every $a \xrightarrow{\varphi} b$ in $\text{Hom}(\mathcal{C})$ the following square commutes:

$$\begin{array}{ccc} Fa & \xrightarrow{F\varphi} & Fb \\ \eta_a \downarrow & & \downarrow \eta_b \\ Ga & \xrightarrow{G\varphi} & Gb \end{array}$$

Remark 1.2. If the notation was not itself suggestive enough (and perhaps it wasn’t at this high a level of abstraction, alas), a “natural transformation” is a good concept of “map between functors”—this should already be a somewhat familiar notion to those who know some representation theory: A representation of G over k (take $k = \mathbb{R}, \mathbb{C}$ for comfort if you like) is a sort of map (indeed, literally a functor) $G \rightarrow \text{Vect}_k$, and an equivariant map between two representations is analogous to (indeed, literally is) a natural transformation between them. People also sometimes like to illustrate these with a “globular diagram”

$$\begin{array}{ccc} & F & \\ \mathcal{C} & \begin{array}{c} \curvearrowright \\ \Downarrow \eta_\bullet \\ \curvearrowleft \end{array} & \mathcal{D} \\ & G & \end{array}$$

, which in particular should make you imagine a sort of “directed” homotopy from F to G (and indeed, one can kind of formulate homotopies of maps of topological spaces in this “maps between maps” language blabla)—in a bit, we’ll see some concepts which are roughly “homotopy equivalences of categories”, so this intuition is by no means off the mark.

Example 1.1. Denote $(-)^{\times} : \text{CRing} \rightarrow \text{Group}$ the functor that associates any commutative ring R to its group of units. Denote $GL_n(-) : \text{CRing} \rightarrow \text{Group}$ the functor that associates any commutative ring R to the group of invertible $n \times n$ matrices with coefficients in R . Then, the determinant map defines a natural transformation $\det_\bullet : GL_n(-) \rightarrow (-)^{\times}$.

Example 1.2. Denote $(-)^{\vee} : \mathbb{R}\text{FinVect} \rightarrow \mathbb{R}\text{FinVect}$ the functor that takes a vector space, V , to its dual $V^{\vee} := \text{Hom}_{\mathbb{R}}(V, \mathbb{R})$. Now, for each finite dimensional vector space V , one can choose an isomorphism $\eta_V : V \rightarrow V^{\vee}$ by choosing a basis. Nonetheless, this can not be assembled into a natural transformation $\text{id}_{\mathcal{C}} \rightarrow (-)^{\vee}$ —intuitively because there is no way to make the choice of isomorphism “naturally” or “canonically”

“why do we care about this?” well, to curry your question into a different question, we define another thing:

Definition 1.8. A pair of functors $L : \mathcal{C} \rightarrow \mathcal{D}$, $R : \mathcal{D} \rightarrow \mathcal{C}$ are said to be an **adjoint pair** if any of the following equivalent definitions hold:

- (Hom bijection) For every $c \in \text{obj}(\mathcal{C}), d \in \text{obj}(\mathcal{D})$, there is a natural isomorphism $\eta_{c,d} : \text{Hom}_{\mathcal{D}}(Lc, d) \rightarrow \text{Hom}_{\mathcal{C}}(c, Rd)$, where “natural” is taken to mean “for any $c \xrightarrow{\varphi} c' \in \text{Hom}(\mathcal{C}), d \xrightarrow{\psi} d' \in \text{Hom}(\mathcal{D})$, the following commutes:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(Lc, d) & \xrightarrow{\eta_{c,d}} & \text{Hom}_{\mathcal{D}}(c, Rd) \\ \downarrow \text{Hom}_{\mathcal{C}}(F\varphi, \psi) & & \downarrow \text{Hom}_{\mathcal{D}}(\varphi, R\psi) \\ \text{Hom}_{\mathcal{C}}(Lc', d') & \xrightarrow{\eta_{c',d'}} & \text{Hom}_{\mathcal{D}}(c', Rd') \end{array}$$

- (equational) We have natural transformations $\eta : \text{id}_{\mathcal{C}} \rightarrow R \circ L$ (the “unit”) and $\varepsilon : L \circ R \rightarrow \text{id}_{\mathcal{D}}$ (the “counit”) such that the following commutes:

$$\begin{array}{ccccc} & & \text{id}_R & & \\ & \nearrow & & \searrow & \\ R & \xrightarrow{\eta_R(-)} & RLR & \xrightarrow{R\varepsilon(-)} & R \\ & \searrow & & \nearrow & \\ & & \text{id}_L & & \end{array}$$

$$\begin{array}{ccccc} & & L\eta(-) & & \\ & \nearrow & & \searrow & \\ L & \xrightarrow{L\eta(-)} & LRL & \xrightarrow{\varepsilon_L(-)} & L \\ & \searrow & & \nearrow & \\ & & \text{id}_L & & \end{array}$$

(in a quip, “ $\varepsilon \circ \eta = \text{id}$ ”).

- (universal arrow for left adjoint) For every $c \in \mathcal{C}$, there exists $Rc \in \text{obj}(\mathcal{D})$ and a map $\varepsilon_c : L Rc \rightarrow c$ such that, for every $d \in \text{obj}(\mathcal{D})$ and every $\varphi : Ld \rightarrow c$, there exists unique $\tilde{\varphi} : d \rightarrow Rc$ making the following diagram commute

$$\begin{array}{ccc} Ld & & \\ \downarrow \exists! L\tilde{\varphi} & \searrow \varphi & \\ L Rc & \xrightarrow{\varepsilon_c} & c \end{array}$$

- (universal arrow for right adjoint) For every $d \in \mathcal{D}$, there exists $Ld \in \text{obj}(\mathcal{C})$ and a map $\eta_d : d \rightarrow RLd$ such that, for every $c \in \text{obj}(\mathcal{C})$ and every $\psi : d \rightarrow Rc$, there exists unique $\tilde{\psi} : Ld \rightarrow c$ making the following diagram commute

$$\begin{array}{ccc} & Rc & \\ & \uparrow \exists! R\tilde{\psi} & \\ RLd & \xleftarrow{\eta_d} & d \end{array}$$

In any of these cases, L is said to be a **left adjoint** (to R), and R is said to be a **right adjoint** (to L).

Remark 1.3. Each of these definitions gives a slightly different insight into what a “adjunction” really is, namely:

Remark 1.4. Technically, a proof that these definitions are equivalent is required. Practically, it’s pure abstract nonsense and not particularly insightful. To see the proofs, look at the nlab pages on “adjunction” and “adjoint functors”, but you shouldn’t need to remember them nearly so much as (the intuition behind) the definitions.

Now that we have the requisite language, we can finally formulate a “(homotopy) equivalence of categories”:

Definition 1.9. Two categories, $\mathcal{C}, \mathcal{D}$ are said to be **equivalent** if there exists pair of functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ and a pair of natural isomorphisms $\eta : \text{id}_{\mathcal{C}} \rightarrow F \circ G$ and $\varepsilon : G \circ F \rightarrow \text{id}_{\mathcal{D}}$.

Remark 1.5. Any equivalence is an adjunction by the counit-unit definition above, and indeed: an equivalence is precisely an adjunction for which the counit and unit transformations are iso.

Definition 1.10. In a category, $\mathcal{C}$, a “**diagram**” is a functor $F : \mathcal{I} \rightarrow \mathcal{C}$ where $\mathcal{I}$ is small (and thought of as a “index category”).

The term “index category” here is taken in analogy to that of “index set”, e.g. for a product/direct sum.

Example 1.3. Take $\mathcal{I}$ to be the category given by the following digraph (with identity maps implicit):

$$1 \xleftarrow{i} 3 \xrightarrow{j} 2$$

Take $\mathcal{C} = \mathbf{CRing}$. Then, the functor such that $F(1) = \mathbb{Z}[x]$, $F(2) = \mathbb{Z}/\neq\mathbb{Z}$, $F(3) = \mathbb{Z}$ with $F(i), F(j)$ taken to be the natural inclusions gives a diagram, visualized as

$$\mathbb{Z}[x] \longleftarrow \mathbb{Z} \xrightarrow{j} \mathbb{Z}/2\mathbb{Z}$$

Example 1.4. Take $\mathcal{I}$ to be the category given by the following infinite digraph (with identity maps implicit):

$$1 \xrightarrow{i_{12}} 2 \xrightarrow{i_{23}} 3 \xrightarrow{i_{34}} \dots$$

Take $\mathcal{C} = \mathbf{CRing}$. Then, the functor such that, for every positive integer n , we take $F(n) = \mathbb{Z}/2^n\mathbb{Z}$ and $F(i_{n(n+1)})$ is taken to be the natural inclusion $\mathbb{Z}/2^n\mathbb{Z} \rightarrow \mathbb{Z}/2^{n+1}\mathbb{Z}$ is a diagram, visualized as

$$\mathbb{Z}/2\mathbb{Z} \longrightarrow \mathbb{Z}/2^2\mathbb{Z} \longrightarrow \mathbb{Z}/2^3\mathbb{Z} \longrightarrow \dots$$

Example 1.5. Take $\mathcal{I}$ to be the category given by the following digraph (with identity maps implicit):

$$1 \quad 2$$

Take $\mathcal{C} = \mathbf{CRing}$. Then, the functor with $F(1) = \mathbb{R}$ and $F(2) = \mathbb{C}$ gives a diagram, visualized as:

$$\mathbb{R} \quad \mathbb{C}$$

Definition 1.11. Let $F : \mathcal{I} \rightarrow \mathcal{C}$ be a diagram. A “**cone over**” (resp. “**cocone under**”) is a $\mathcal{C}$ -object c , together with arrows $c \xrightarrow{\pi_n} F(n)$ (resp. $c \xleftarrow{\iota_n} F(n)$) for each $n \in \text{obj}(\mathcal{I})$, such that for every $n \xrightarrow{i_{nm}} m \in \text{Hom}(\mathcal{I})$ we have that the following diagram on the left (resp. on the right) commutes:

$$\begin{array}{ccc} c & & F(n) \xrightarrow{F(i_{nm})} F(m) \\ \pi_n \searrow & & \downarrow \iota_n \quad \downarrow \iota_m \\ F(n) & \xrightarrow{F(i_{nm})} & F(m) \end{array}$$

Definition 1.12. Let $F : \mathcal{I} \rightarrow \mathcal{C}$ be a diagram. A **limit** (resp. **colimit**) is a terminal cone over F (resp. initial cocone under F)— e.g. an object c equipped with maps $c \xrightarrow{\pi_n} F(n)$ (resp. $c \xleftarrow{\iota_n} F(n)$) such that, for any other object c' equipped with maps $c' \xrightarrow{\pi'_n} F(n)$ (resp. $c' \xleftarrow{\iota'_n} F(n)$), there exists a unique map $\pi : c' \rightarrow c$ (resp. $\iota : c \rightarrow c'$) such that the following diagram on the left (resp. on the right) commutes:

$$\begin{array}{ccc} c' & & F(n) \xrightarrow{F(i_{nm})} F(m) \\ \downarrow \exists! \pi & & \downarrow \iota_n \quad \downarrow \iota_m \\ c & & c \\ \pi'_n \searrow \quad \pi_n \searrow & & \downarrow \iota'_n \quad \downarrow \iota'_m \\ F(n) & \xrightarrow{F(i_{nm})} & F(m) \end{array}$$

Example 1.6. Let $F : \mathcal{I} \rightarrow \mathcal{C}$ be as in example 1.3. Then, the colimit over the diagram is given by the tensor product as (commutative) $\mathbb{Z}$ -algebras, $\mathbb{Z}[x] \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}[x]$:

$$\begin{array}{ccc} \mathbb{Z}[x] \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} & \longleftarrow & \mathbb{Z}[x] \\ \uparrow & & \uparrow \\ \mathbb{Z}/2\mathbb{Z} & \longleftarrow & \mathbb{Z} \end{array}$$

This is also called a **pushout** (its dual is called a **pullback** or a **fibre product**).

Example 1.7. Let $F : \mathcal{I} \rightarrow \mathcal{C}$ be as in example 1.4. Then, the limit over the diagram is given by the 2-adic integers $\mathbb{Z}_2$ (look them up on wikipedia if you haven't seen them before, they're useful for number theory etc.), together with the natural projections:

$$\begin{array}{ccccc} & & \mathbb{Z}_2 & & \\ & \nearrow & \uparrow & \nwarrow & \\ \mathbb{Z}/2\mathbb{Z} & \longrightarrow & \mathbb{Z}/2^2\mathbb{Z} & \longrightarrow & \mathbb{Z}/2^3\mathbb{Z} \longrightarrow \dots \end{array}$$

This is also called a **projective/inverse limit** (its dual is called a **direct/inductive limit**).

Example 1.8. Let $F : \mathcal{I} \rightarrow \mathcal{C}$ be as in example 1.5. Then, the limit over the diagram is given by the product as rings, and the colimit over the diagram is given by the tensor product as $\mathbb{Z}$ -algebras (note $\mathbb{Z}$ the initial object of $\mathbf{CRing}$, so one has “invisible” maps $\mathbb{Z} \rightarrow \mathbb{R}$ and $\mathbb{Z} \rightarrow \mathbb{C}$):

$$\begin{array}{ccc} & \mathbb{R} \times \mathbb{C} & \\ \pi_{\mathbb{R}} \swarrow & & \searrow \pi_{\mathbb{C}} \\ \mathbb{R} & & \mathbb{C} \end{array} \qquad \begin{array}{ccc} \mathbb{R} & & \mathbb{C} \\ \searrow \iota_1 & & \swarrow \iota_2 \\ & \mathbb{R} \otimes_{\mathbb{Z}} \mathbb{C} & \end{array}$$

The left limit is called a **product**, the right colimit is called a **coproduct** (though indeed, coproducts in $\mathbf{CRing}$ can be thought of as pushouts under $\mathbb{Z}$).

Idea: colimits usually look like disjoint unions modulo identifying some things, limits usually look like subcollections of products. Google examples of (co)products, pushouts, and pullbacks in categories you know about (including $\mathbf{Top}$ by the way) to feel a bit more comfortable with the terminology. Important note: in module categories, finite (but not infinite!) products and coproducts coincide. “coproducts” bear the name of “direct sum” more generally, but finite direct sums *are* products as well (which is part of what makes “abelian categories” so nice).

Theorem 1.1. Suppose $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor. Then:

- If F is a left adjoint (e.g., has a right adjoint), then F preserves colimits.
- If F is a right adjoint (e.g., has a left adjoint), then F preserves limits.

This has a proof by pure formal nonsense given by unraveling the definitions. This is also an incredibly useful fact from which many other properties follow. If it helps to remember, one has the mnemonics “LAPC [left adjoints preserve colimits]” and “RAPL [right adjoints preserve limits].”

Proposition 1.1. Let $F : \mathcal{I} \rightarrow \mathcal{C}$ be a diagram. Then, a limit of F (resp. a colimit of F) is precisely a $\mathcal{C}$ -object $\lim F$ (resp. $\operatorname{colim} F$) such that, for all $c \in \operatorname{obj}(\mathcal{C})$ we have a natural bijection

$$\operatorname{Hom}_{\mathcal{C}}(c, \lim F) \cong \operatorname{Hom}_{\operatorname{Fun}(\mathcal{I}, \mathcal{C})}(\operatorname{const}_c, F)$$

or respectively a natural bijection

$$\operatorname{Hom}_{\mathcal{C}}(\operatorname{colim} F, c) \cong \operatorname{Hom}_{\operatorname{Fun}(\mathcal{I}, \mathcal{C})}(F, \operatorname{const}_c),$$

where by $\mathbf{Fun}(\mathcal{I}, \mathcal{C})$ we mean the category of functors and natural transformations (ignore set-theoretic issues here), and by $\mathbf{const}_c$ we mean the functor that takes every object to c and every morphism to id_c .

Rephrasing: $\lim_{\mathcal{I}}$ and $\mathrm{colim}_{\mathcal{I}}$ are right (resp. left) adjoint to the functor $\mathbf{const} : \mathcal{C} \rightarrow \mathbf{Fun}(\mathcal{I}, \mathcal{C})$.

This above proposition, modulo set-theoretic issues w.r.t. defining a functor category, is a valid alternative definition of “limit/colimit”, and can lead one to various useful ideas (e.g., the 2-category of categories, viewing $\lim$ / colim as functors on their own right, left/right Kan extensions, etc.). It is a fine exercise to try and unravel this definition and see how the universal property (w.r.t. cones/cocones) is equivalent to this statement, but we would suggest not overemphasizing this definition if any of these concepts are new to the reader.

2 Tensor products (even over potentially non-commutative rings) and more miscellaneous module theory

Definition 2.1. *R*-balanced maps Let R be a ring, $M \in \text{Mod}_R, N \in {}_R\text{Mod}$. Then, an R balanced map $\varphi : M \times N \rightarrow A$ an abelian group is a biadditive map (e.g., $\mathbb{Z}$ -bilinear map) such that, for all $r \in R$, $\varphi(mr, n) = \varphi(m, rn)$ for all $m \in M, n \in N$.

Definition 2.2. *tensor product* Let $M \in {}_S\text{Mod}_R, N \in {}_R\text{Mod}_{S'}$ for a ring R . Then, the tensor product $M \otimes_R N$ is an abelian group (in particular, an S - S' -bimodule) together with a map $\otimes_R : M \times N \rightarrow M \otimes_R N$ such that, for any abelian group A together with an R -balanced map $\varphi : M \times N \rightarrow A$, there exists unique abelian group homomorphism $\tilde{\varphi} : M \otimes_R N \rightarrow A$ making the following diagram commute:

$$\begin{array}{ccc} M \times N & \xrightarrow{\otimes} & M \otimes_R N \\ & \searrow \varphi & \downarrow \exists! \tilde{\varphi} \\ & & A \end{array}$$

. If R is commutative, then $M \otimes_R N$ admits the natural structure of an R -module, and we can instead consider the universal R -module such that any R -bilinear map $M \times N \rightarrow A$ factors through an R -module homomorphism $M \otimes_R N \rightarrow A$.

Note: while we take bimodules above, we can simply take S, S' , or R to be $\mathbb{Z}$ and recover “one-sided” modules. That’s a universal property or what have you– really, it’s the same philosophy as viewing the “tensor product of V and W ” as “the multilinear maps from $V \times W$ ”, we’re just being cautious about commutativity and being a smidgeon more careful since not every module is free here. Anyways, “how/why does this universal object actually exist?”, well, we just take a super big free module and the mod out by what it means to be R -balanced, aka by what it means to be (non-commutatively) multilinear. We give the proof of existence as follows:

Lemma 2.1.)(tensor products exist)

The tensor product has, itself, a deep relation to Hom -functors– namely, we have the *tensor-hom adjunction*:

Theorem 2.1.

Definition 2.3. Let R be a ring. An R -**algebra** is a ring, S , together with a ring homomorphism $R \rightarrow S$ (the “structure map”).

Definition 2.4. Let $R \rightarrow S, R \rightarrow S'$ be two R -algebras. A **(homo)morphism of R -algebras** from $S \rightarrow S'$ is a ring homomorphism $\varphi : S \rightarrow S'$ which commutes with the structure maps, e.g. making the following diagram commute:

$$\begin{array}{ccc} & R & \\ & \swarrow \quad \searrow & \\ S & \xrightarrow{\varphi} & S' \end{array}$$

Definition 2.5. Let R be a commutative ring, let $R \rightarrow S$ and $R \rightarrow S'$ be R -algebras. Then, the tensor product of R -algebras is the colimit in CRing over the diagram

$$S' \leftarrow R \rightarrow S,$$

e.g. the pushout in CRing (or, as one can verify, the coproduct in ${}_R\text{CAlg}$).

Remark 2.1. Note that the tensor product of R -algebras can also be taken in steps. Namely, any R -algebra $R \rightarrow S$ admits the natural structure of an R -module by left/right multiplication. Then, for two R -algebras S and S' , one can take their tensor product as R -modules, and then endow $S \otimes_R S'$ with the structure of an ring by defining $(s_1 \otimes s'_1)(s_2 \otimes s'_2) := s_1 s_2 \otimes s'_1 s'_2$, which one can check gives an R -algebra in the natural way.

Definition 2.6. Let R be a commutative ring, and $S \subset R$ a multiplicatively closed subset. The **localization**, $S^{-1}R$ is taken to be a (commutative) R -algebra $R \rightarrow S^{-1}R$ such that, for any ring homomorphism $R \rightarrow T$ taking every element of S to a unit, there exists a unique ring homomorphism $S^{-1}R \rightarrow T$ making the following diagram commute:

$$\begin{array}{ccc} R & \xrightarrow{\quad} & S^{-1}R \\ & \searrow & \downarrow \exists! \\ & & T \end{array}$$

This is to say, $S^{-1}R$ is the “smallest” (e.g., initial) R -algebra for which every element of S is a unit. As a universal property, this requires a construction/proof of existence—this one justifies the common name “ring of fractions” this ring.

Theorem 2.2. (Construction of localizations) *Localizations exist.*

Proof. Let S be a multiplicative subset of R commutative. Define, on $R \times S$, an equivalence relation

$$(a, b) \sim (c, d)$$

if there exists $s \in S$ such that

$$s(ac - bd) = 0.$$

As this forms an ideal, we may take $S^{-1}R$ to be the ring of equivalence classes under this relation; we denote the equivalence class of (r, s) as either $\frac{r}{s}$ or $s^{-1}r$. Then, the map $r \mapsto \frac{r}{1} : R \rightarrow S^{-1}R$ makes $S^{-1}R$ initial among all R -algebras for which S is taken to units by the structure map. $\square$

The relation in question should be taken to be exactly that of fraction addition working “as it should”. The use of an extra $s \in S$ in that definition is mostly just to fudge things a little so that we don’t need to make an extra case for the multiplicative set having zero divisors (but, for S to be a multiplicative set containing a zero divisor, it must also contain 0 and hence the localization must be the zero ring anyways.)

There is some special notation in the following cases:

Definition 2.7. If $f \in R$, then we denote $R_f := \{f, f^2, f^3, \dots\}^{-1}R$ the “**localization away from f** ”. If $\mathfrak{p}$ is a prime ideal of R , then we denote $R_{\mathfrak{p}} := (R - \mathfrak{p})^{-1}R$ the “**localization at $\mathfrak{p}$** .”¹

¹This terminology comes from algebraic geometry, where prime ideals (or, intuitively, maximal ideals) are thought of as “points” of a topological space. Imagining R as the ring of continuous functions on a compact Hausdorff space, $\mathfrak{p}$ as a point of said space, then we have the localization *at* $\mathfrak{p}$ arising as the “ring of germs of functions at $\mathfrak{p}$ ”. The terminology for localization *away from* an element is analogous: the localization at $f \in R$ is precisely the localization (intuitively, restriction to) all prime ideals (“points”) not containing f (that is, where f does not vanish).

3 (co)homological algebra

Definition 3.1. If $F : {}_R\text{Mod} \rightarrow {}_S\text{Mod}$ is a functor of module categories (it can be different sided module categories too, it doesn't matter), F is said to be *additive* if, given any two $\varphi, \psi \in \text{Hom}_R(M, N)$, we have $F(\varphi + \psi) = F(\varphi) + F(\psi)$ —that is, on each Hom it's a homomorphism of additive groups.

I'm really only mentioning the word “additive” here so that I can say “additive” later on without needing to refer you to the stuff on abelian categories (at least, not yet).

Definition 3.2. An R module, P , is said to be **projective** if any of the following equivalent statements are true:

- For any surjective module homomorphism $N \twoheadrightarrow M$, any module homomorphism $P \rightarrow M$ admits a (not-necessarily unique) lift $P \rightarrow N$ such that the following diagram commutes:

$$\begin{array}{ccc} & & N \\ & \nearrow \text{dashed} & \downarrow \\ P & \xrightarrow{\quad} & M \end{array}$$

- Any short exact sequence

$$0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$$

splits.

- The (covariant) functor $\text{Hom}_R(P, -)$ is exact (note left-exactness is always true for Hom functors)
- P is a direct summand of a free module²

Remark 3.1. Any free module is projective. Furthermore, the converse is occasionally (but not always!) true: a projective module is always free over a PID (basic fact), and a projective module over a local ring is always free (straightforward to prove for finitely generated modules via Nakayama's lemma from commutative algebra, but a difficult fact in general—keep the general result in mind only as a restriction for coming up with counterexamples).

Definition 3.3. An R -module, F , is said to be **flat** if any of the following equivalent conditions are true:

- The functor $F \otimes_R - \cong - \otimes_R F$ is exact (note right-exactness is always true for tensoring)
- The functor $F \otimes_R -$ takes injective maps to injective maps.
- For any R -module homomorphism $\varphi : R^n \rightarrow F$, and any K a finitely generated submodule of $\ker \varphi$, φ factors through a map $\psi : R^n \rightarrow F'$ for some free³ R -module F' so that $K \subset \ker \psi$, e.g. such that the following diagram commutes:

$$\begin{array}{ccccc} & & & & F' \\ & & & \nearrow \exists \psi & \searrow \\ K & \xrightarrow{f \cdot g} & \ker \varphi & \hookrightarrow & R^n & \xrightarrow{\varphi} & F \\ & \searrow & \swarrow & & \uparrow & & \\ & & \ker \psi & & & & \end{array}$$

²Note: not every abelian category has a good notion of “free object”, so this particular description is distinct for categories of modules.

³Note: not every abelian category has a good notion of “free object”, so this particular description is distinct for categories of modules.

- (translating the above into a statement about relations– thank you wikipedia) For any R -linear relation

$$\sum_{i=1}^n r_i x_i = 0$$

with $r_i \in R$, $m_i \in F$, there exist some (“change of basis/generator”) elements $y_j \in F$, $a_{ij} \in R$ such that

$$\forall j, \sum_{i=1}^n r_i a_{ij} = 0$$

and

$$\forall i, x_i = \sum_{j=1}^m a_{ij} y_j.$$

Remark 3.2. Any projective module is necessarily flat. The converse is (“clearly”) true for finitely presented flat modules (hence for finitely generated flat modules over Noetherian rings).

Definition 3.4. An R -module, I , is said to be **injective** if any of the following equivalent statements are true:

- For any module homomorphism $M \rightarrow I$, any injective module homomorphism $M \otimes N$ admits an extension $N \rightarrow I$ (“along $M \rightarrow I$ ”) such that the following diagram commutes:

$$\begin{array}{ccc} I & \xleftarrow{\quad} & M \\ & \swarrow \text{dashed} & \downarrow \\ & & N \end{array}$$

- Any short exact sequence

$$0 \rightarrow I \rightarrow M \rightarrow N \rightarrow 0$$

splits.

- The (contravariant) functor $\text{Hom}_R(-, I)$ is exact (note left-exactness [interpreted appropriately for variance] is always true for Hom functors)

Note that projective modules are “dual” to injective modules and vice versa. Here is a useful criterion for checking injectivity:

Theorem 3.1. (Baer’s Criterion) Let I be an R -module. Then, I is injective if and only if any homomorphism $J \rightarrow I$ from J a submodule/ideal of R admits an extension to a homomorphism $R \rightarrow I$.

As an immediate corollary, we have the following:

Corollary 3.1. Any abelian group A is injective (as a $\mathbb{Z}$ -module) if and only if it is divisible (e.g., the map $a \mapsto na$ is a surjection for all non-zero $n \in \mathbb{Z}$).

4 Generalities on “Abelian Categories”

Remark 4.1. *Jason might require us to technically be able to unravel the definition, but, in practice “abelian category” just means “category which we can think of like a category of left/right R -modules”. This is possible to formalize by the 4.1 Freyd-Mitchell Embedding Theorem, which is technically useful. It’s also complete abstract nonsense, and for every almost every single thing we talk about, you can use the words “abelian category” interchangeably with “category of R -modules” (or, if you really want, “category of chain complexes of R -modules”), so just think in terms of whatever’s easiest.*

Theorem 4.1. *(Freyd-Mitchell Embedding) Let $\mathcal{A}$ be a (small) abelian category. Then, there is a ring R (not necessarily commutative) and a full, faithful, exact (hence additive) functor $i : \mathcal{A} \otimes {}_R\text{Mod}$ the category of left R -modules.*

You do *not* need to memorize this (nor will we see a proof), but I feel it necessary to mention it as a name to look up/be referred to.