

Taper and total aerial volumes

Florence Gohon and Amaël Le Squin

29th October 2025

Institut National de l'information géographique et forestière

1. Taper volume

In tree sample...

- trees j , where c, h, h_{dec} are measured

$$V_{\text{bole}}(j) = f_{III}(c, h, h_{\text{dec}})$$

- *simplified or remeasured* trees i , where c is measured

$$\cancel{V_{\text{bole}}(i) = V_{\text{bole}}(j)} \rightarrow \cancel{V_{\text{bole}}(i) = V_{\text{bole}}(j) \frac{c(i)}{c(j)}} \rightarrow V_{\text{bole}}(i) = V_{\text{bole}}(j) \frac{f_I(c(i))}{f_I(c(j))}$$

- Data sets source
- Linear models
- Transformations

$$\begin{cases} V_{\text{bole}} = \frac{c^2 h}{4\pi \left(1 - \frac{1,3}{h}\right)^2} f_{\text{new}} \\ f_{\text{new}} = f_{III}(c, h, h_{\text{dec}}) \end{cases}$$


$$\ln(V_{\text{bole}}) = f_I(\ln(c))$$

- Explanatory variables
 - ▶ h_{dec} turns into h'_{dec}
 - ▶ variables are shared by every species
 - ▶ $f(g)$ is dropped
- Parameters
 - ▶ the three-variable model is no longer cut-specific

$$f_{\text{new}} = \alpha + \beta c + \gamma \frac{\sqrt{c}}{h'_{\text{dec}}} + \delta \frac{\sqrt{h'_{\text{dec}}}}{c^2 h} + \eta \left(1 - \frac{h'_{\text{dec}}}{h} \right)$$


$$V_{\text{bole},I} = e^{\alpha + \beta \ln(c) + \gamma \ln(c)^2 + \frac{\sigma^2}{2}}$$

Performances similar to current models



2. Total aerial volume


A tree in Emerge data




- Bole volume
- Large branches
- Small branches

1. Bole volume (Florence) and independent crown model
2. Multivariate model for both bole and crown
3. Model on the ratio between bole and total volume

Independent crown model – 1




● Small ● Median ● Tall ● Very tall



● Small ● Median ● Tall ● Very tall

Independent crown model – 2



3. Conclusion

Figure that sums up everything!