


Enhancement Due to Photonic Coupling in Nanocavity Structures

Bumbia, Amal;
Abudayyeh, Hamza;
Liu, Zhida;
Li, Xiaoqin


Department of Physics UT
Austin

Background


How do quantum emitters work?



Why can a nanoscale quantum emitter can have emitting behavior?



What is the interplay between quantum emitters and cavities?



Cavities can impact emitters in a few ways:

- decay rate enhancement --- The weak coupling of a 2-level emitter to a cavity leads to the modification of the spontaneous emission rate. The emitter can have multiple radiative modes and some nonradiative processes
- emission redirection --- the intrinsic radiation pattern of a quantum emitter is omnidirectional, making it hard to collect emitted photons. Cavities can redirect the emission to the desired optical mode with an efficiency η_{NA} . Different modes can have different efficiencies.
- absorption enhancement --- cavities alter the absorption cross-section of the emitter which causes absorption enhancement and lowers the saturation power needed from the pump laser.

Coupling parameters:

- emitter's decay in free space Γ_0
- cavity's loss rate κ
- emitter-cavity coupling coefficient g

This is dependent on the overlap of the photonic cavity and the resonances of the 2-level system.

Due to the interplay between the photonic cavity and the system, we are able to enhance some processes at the expense of others.

For instance, if the resonances of the plasmonic cavity are vertical, they selectively enhance dipole emitters with a vertical dipole.

Not only are we increasing the lifetime, but we are also altering our characterization and probability of the emission.

Weak Coupling Regime

We are concerned with the weak coupling regime $g < \kappa$, Γ_0 .

Here, the photon is released by an excited emitter due to spontaneous emission and lost to far-field radiation modes leaving the cavity in a vacuum state.

In this case, the cavity becomes the system's environment and results in a modified decay rate $\Gamma = F \Gamma_0$ where $F = \frac{3}{4\pi^2} Q \frac{\lambda^3}{V}$ (Purcell factor - represents the shortening of lifetimes; losses ensure that it is not directly related to the number of emitted photons), $Q \approx \frac{1}{\kappa}$ (cavity quality factor), and the cavity mode volume $\frac{V}{\lambda^3}$.

The advantage of the weak coupling regime is that for emitted photons, the probability that they will be reabsorbed by the emitter is low compared to the likelihood of the photon escaping the cavity.

What are the effects of weak
coupling between
plasmonic nanocavities and
excitons?

How does plasmon-exciton
coupling effect emission
enhancement?


Background




Sample 1 : Gold Nanocube

Sample 1: Gold Nanocube : Optical Data


The lightning-rod effect causes the corners of the cubes to be the regions of greatest enhancement





Darkfield Spectroscopy


- white light scattering by nanocavities
- measure plasmonic resonances



Where do the plasmonic resonances occur?
 This is a point where the data gets murky...




- 540 nm
- *625 nm !!!*
- 725 nm



Labelled points on
 the darkfield image



Hyperspectral Spatial Image


- well localized points with nanocubes
- intensity coming out of these points is enhanced







Photoluminescence (PL)

- excite sample at different wavelengths
- measure exciton resonances
- longer wavelengths enhanced
- greater, non-uniform intensity enhancement along nanocubes
- find overlap between the wavelengths where plasmonic resonances and enhancement occur








Notice the peaks between 603-609 nm and 612-620 nm and around 625 nm



Slight trough at 608-611 nm and again at 620-625 nm

These were the brightest spots - less uniformity between them but we still have peaks at around 605, 630, 640 and 615 nm

Recall the plasmonic spectrum via darkfield spectroscopy and the peaks around 625 nm — possible overlap between plasmonic and exciton resonances?



Room Temperature Higher Power PL



Peaks are largely concentrated at
610, 625, 640 nm

Background




Sample 1 : Gold Nanocube



Sample 1 : Silver Nanocube


Sample 1: Silver Nanocube : Optical Data


Initially, the silver nanocubes were placed onto the original gold nanocube sample. There was an issue with the trilayer sample so it needs to be remade.





Photoluminescence (PL)


- Focus on each point individually to obtain a spectrum over a range of wavelengths
- This image is focused on a bare portion of the sample for control purposes








Hyperspectral Spatial Image - Cryogenic





Hyperspectral Spatial Image - Cryogenic 20uW







Comparing Results

Do the spectra of the same points peak at the same wavelengths between the samples? How does the presence of silver nanocubes impact the measurements?

Can we identify which points are likely silver nanocubes?

In general, how do the peaks differ between the gold and silver nanocubes?
Does either provide more obvious enhancement?

Sample 1 Before and After Silver Nanocube Placement




Room Temperature PL Gold v. Silver


It appears that the correlations between the new and old points are as follows:

1 - 35
4 - 11
5 - 5
6 - 12
8 - 28
10 - 14
11 - 30
12 - 29
15 - 26


So we will compare the plots. Notice how the spectra will look very different but still preserve the location of certain peaks.




1 RT mAgNC




mAuNC_fine_35





5 RT mAgNC




mAuNC_fine_5







10 RT mAgNC




mAuNC_fine_14



15 RT mAgNC




mAuNC_fine_26






The following comparisons are interesting because the PL spectrum from the points resemble each other, but the correlated points from prior measurements have even more drastically different spectra. They also resemble uncorrelated points...

4 RT mAgNC




mAuNC_fine_11







11 RT mAgNC




mAuNC_fine_30



12 RT mAgNC






mAuNC_fine_29





Point 2 is most likely a silver nanocube – out of all the new point measurements, it was the most isolated from old points.

Notice the peak at around 620 nm corresponding to a very high intensity





Notice how these spectra have a similar structure. Point 16 is notably outside of the general range of intensity for the peak. It is possible that these denote silver nanocubes.



Do the spectra of the same points peak at the same wavelengths between the samples? How does the presence of silver nanocubes impact the measurements?


It seems like some peaks existed in both measurements between the same gold nanocube points, but the spectra of gold cubes from the sample with silver nanocubes had significantly more peaks at higher intensities. There may be some odd interactions between the gold and silver cubes.

Can we identify which points are likely silver nanocubes?

Point 2 is definitely silver, points 7, 13, 14, and 16 are also likely silver. Especially 7.


In general, how do the peaks differ between the gold and silver nanocubes? Does either provide more obvious enhancement?

The gold nanocubes without any influence from the silver caused the highest peaks in the spectrum to be in the 1000-1200 intensity range and the 615-620 nm range. Overall, it appears the silver candidates have cleaner spectra and more obvious peaks than the gold nanocubes did. With the current data, there's no discernable difference in whether or not there's more obvious enhancement between the silver and gold nanocubes.



Gold outlier

Silver outlier



Semester Progress

- Optical data analysis + summary for nanocube project (Sample 1)
- Noise reduction code using DMD algorithm
- Lots of graphene exfoliation + practice transfers with graphene and HBn during slower weeks

Grey points - Cryo Au

Blue points - Cryo Ag

Ag — Au

8 - new silver

10 - 27

17 - 21

21 - 9

22 - 6

23 - new silver

26 - 17

29 - 33

35 - 28

36 - 29

42 - new silver

43 - 3

9 - 40

13 - 25


20 - 11

25 - 12


27 - new silver

31 - new silver


33 - 5




Cryogenic mAgNC cleaned




cryo_mAgNC_8




cryo_mAgNC_23




cryo_mAgNC_27




cryo_mAgNC_31




cryo_mAgNC_31




cryo_mAgNC_13




mAuNC_fine_25




cryo_mAgNC_9




mAuNC_fine_40




cryo_mAgNC_17




mAuNC_fine_21




cryo_mAgNC_43




mAuNC_fine_3





cryo_mAgNC_36




mAuNC_fine_29







cryo_mAgNC_10




mAuNC_fine_27




cryo_mAgNC_21




mAuNC_fine_9




cryo_mAgNC_22




mAuNC_fine_6




cryo_mAgNC_26




mAuNC_fine_17




cryo_mAgNC_29




mAuNC_fine_33




cryo_mAgNC_20




mAuNC_fine_11




cryo_mAgNC_25



mAuNC_fine_12



cryo_mAgNC_33



mAuNC_fine_5

