

# The *Kepler* Smear Campaign I: An Asteroseismic Catalogue of Bright Red Giants

Benjamin J. S. Pope,<sup>1,2,3</sup>★ Guy R. Davies,<sup>4,5</sup> Keith Hawkins,<sup>6,7</sup> Timothy R. White,<sup>5,8</sup> Timothy R. Bedding,<sup>9,5</sup> Allyson Bieryla,<sup>10</sup> David W. Latham,<sup>10</sup>, Conny Aerts,<sup>11,12,13</sup> Suzanne Aigrain,<sup>3</sup> Victoria Antoci,<sup>5</sup> Dominic Bowman,<sup>11,12,13</sup> Ashley Chontos,<sup>9</sup> Gilbert A. Esquerdo,<sup>10</sup> Daniel Huber,<sup>14,15,9</sup> Paula Jofré,<sup>16</sup> Simon Murphy,<sup>5,9</sup> Timothy van Reeth,<sup>5,9</sup> Victor Silva Aguirre,<sup>5</sup> Amalie Stokholm,<sup>5</sup> Jie Yu,<sup>9,5</sup> and friends

<sup>1</sup>Center for Cosmology and Particle Physics, Department of Physics, New York University, 726 Broadway, New York, NY 10003, USA

<sup>2</sup>NASA Sagan Fellow

<sup>3</sup>Oxford Astrophysics, Denys Wilkinson Building, University of Oxford, OX1 3RH, Oxford, UK

<sup>4</sup>School of Physics and Astronomy, University of Birmingham, Birmingham B15 2TT, UK

<sup>5</sup>Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark

<sup>6</sup>Department of Astronomy, The University of Texas at Austin, 2515 Speedway Boulevard, Austin, TX 78712, USA

<sup>7</sup>Department of Astronomy, Columbia University, 550 W 120th St, New York, NY 10027, USA

<sup>8</sup>Research School of Astronomy and Astrophysics, Mount Stromlo Observatory, The Australian National University, Canberra, ACT 2611, Australia

<sup>9</sup>Sydney Institute for Astronomy (SIfA), School of Physics, University of Sydney, NSW 2006, Australia

<sup>10</sup>Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

<sup>11</sup>Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium

<sup>12</sup>Department of Astrophysics, IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands

<sup>13</sup>Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA

<sup>14</sup>Institute for Astronomy, University of Hawai‘i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

<sup>15</sup>SETI Institute, 189 Bernardo Avenue, Mountain View, CA 94043, USA

<sup>16</sup>Núcleo de Astronomía, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Ejército 441, Santiago De, Chile

Accepted XXX. Received YYY; in original form ZZZ

## ABSTRACT

Here we present the first data release of the *Kepler* Smear Campaign, using collateral ‘smear’ data obtained by *Kepler* to reconstruct light curves of 102 stars too bright to have been otherwise targeted. We describe the pipeline developed to extract and calibrate these light curves, and show that we attain photometric precision comparable to stars analyzed by the standard pipeline in the nominal *Kepler* mission. In this Paper, we focus in particular on a subset of these consisting of 64 red giants for which we detect solar-like oscillations. Using high-resolution spectroscopy from the Tillinghast Reflector Échelle Spectrograph (TRES) together with asteroseismic modelling, we obtain the stellar densities and derive masses, radii and ages of 34 of these red giant and red clump stars as benchmarks. All source code, light curves, TRES spectra, and asteroseismic and stellar parameters are publicly available as a *Kepler* legacy sample.

**Key words:** asteroseismology – techniques: photometric – stars: variable: general

## 1 INTRODUCTION

*Kepler* has revolutionized the field of asteroseismology for solar-like oscillations (Gilliland et al. 2010; Chaplin et al. 2010). It has yielded the detection of gravity-mode period spacings in a red giant

(Beck et al. 2011; Mosser et al. 2014), enabling probes of interior rotation of red giants (Beck et al. 2012; Mosser et al. 2012b) and distinguishing between hydrogen- and helium-burning cores (Bedding et al. 2011; Mosser et al. 2012a). It has also permitted the determination of ages and fundamental parameters of main-sequence stars as cool as the Sun and hotter (Silva Aguirre et al. 2013), including planet-hosting stars (Huber et al. 2013; Silva Aguirre et al. 2015;

★ E-mail: benjamin.pope@nyu.edu

Van Eylen et al. 2018), revealing the most ancient known planetary system, dating back to the earliest stages of the galaxy (Campante et al. 2015). By comparing asteroseismic stellar ages to stellar rotation periods, Angus et al. (2015) have shown that gyrochronology models cannot fit the data with a single relation, leading van Saders et al. (2016) to suggest a qualitative change in dynamo mechanism as stars age through the main sequence.

A major outcome of the *Kepler* asteroseismology programme is a legacy sample of extremely well characterized stars that can serve as benchmarks for future work (Lund et al. 2016; Silva Aguirre et al. 2016). Asteroseismological studies with *Kepler* complement other probes of stellar physics well: for example, the APOKASC sample of 1916 spectroscopically- and asteroseismically-characterized red giant stars (Pinsonneault et al. 2014). For this APOKASC sample, Hawkins et al. (2016c) have been able to extract precise elemental abundances by fitting spectroscopic data with  $\log g$  and  $T_{\text{eff}}$  fixed to asteroseismically-determined values. It is necessary to calibrate such a study against benchmark stars with very precisely-determined parameters, which in practice requires nearby bright stars that are amenable to very high signal-to-noise spectroscopy plus asteroseismology (Creevey et al. 2013), parallaxes (Hawkins et al. 2016a), and/or interferometry (Casagrande et al. 2014; Creevey et al. 2015). This is especially important in the context of the *Gaia* mission (Gaia Collaboration et al. 2016), which has recently put out its second data release of 1,692,919,135 sources, including 1,331,909,727 with parallaxes (Gaia Collaboration et al. 2018). These data will form the basis of many large surveys and it is vital that they are calibrated correctly. To this end, 34 FGK stars including both giants and dwarfs have been chosen as *Gaia*-ESO benchmark stars for which metallicities (Jofré et al. 2014), effective temperatures and asteroseismic surface gravities (Heiter et al. 2015), and relative abundances of  $\alpha$  and iron-peak elements (Jofré et al. 2015) have been determined. This includes only four main sequence stars much cooler than the Sun, due to the paucity of such stars with asteroseismology. This has been accompanied by the release of high resolution spectra (Blanco-Cuaresma et al. 2014) and formed the basis of extensions to lower metallicities (Hawkins et al. 2016b), stellar twin studies (Jofré 2016) and comparisons of stellar abundance determination pipelines (Jofré et al. 2017). Furthermore, by combining asteroseismology with optical interferometry, it has been possible to determine fundamental parameters of main-sequence and giant stars with unprecedented precision (Huber et al. 2012; White et al. 2013, 2015).

Brighter *Kepler* stars are therefore ideal benchmark targets, since photometry can be most easily complemented by *Gaia* parallaxes, interferometric diameters, and high resolution spectroscopy. Unfortunately, the *Kepler* field was deliberately placed to minimize overall the number of extremely bright stars on the detectors, so that only a dozen stars brighter than 6th magnitude landed on silicon (Koch et al. 2010). This was because stars brighter than  $Kp \sim 11$  saturate the CCD detectors, with their flux distributed along a bleed column and rendering those pixels otherwise unusable. Furthermore, due to the limited availability of bandwidth to download data from the spacecraft, only  $\sim 5.7\%$  of pixels on the *Kepler* detectors are actually downloaded in any one Quarter (Jenkins et al. 2010). The result of these two target selection constraints is that photometry was obtained for most of the mission for only 35 stars brighter than  $Kp < 7$  in the *Kepler* field, while 17 targets in this range were observed for less than half the mission and 29 targets brighter than this threshold were entirely ignored. The availability of *Kepler* data remains significantly incomplete down to fainter magnitudes, and in this work we consider  $Kp = 9$  to be an arbitrary cutoff for

bright stars of interest. In the *K2* mission (Howell et al. 2014), very saturated stars have been observed with ‘halo photometry’ using unsaturated pixels in a specially-determined region around bright stars, including the Pleiades (White et al. 2017), Aldebaran (Farr et al. 2018), and  $\rho$  Leonis (Aerts et al. 2018). Unfortunately, in the legacy *Kepler* sample, photometry of such saturated stars was rarely attempted.

Kolodziejczak & Caldwell (2011) noted a way to obtain photometry of every target on-silicon in *Kepler* using a data channel normally used for calibration, even if active pixels were not allocated and downloaded. *Kepler* employs an inter-line transfer CCD as its detector, which successively shuffles each row of pixels down to the edges of the chip to be read out. Because the *Kepler* camera lacks a shutter, the detector is exposed to light during the readout process, with the result that fluxes in each pixel are contaminated by light collected from stars in the same column. This is a particularly serious issue for faint stars in the same detector column as brighter stars, and it is important to calibrate this at each readout stage. Six rows of blank ‘masked’ pixels were allocated in each column to measure the smear bias; furthermore, six ‘virtual’ rows were recorded at the end of the readout, with the result that twelve rows of pixels sample the smear bias in each column. Kolodziejczak & Caldwell (2011) realized that these encode the light curves of bright targets in a 1D projection of the star field. The masked and virtual smear registers each receive  $\sim 1/1034$  of the incident flux in each column. If this is dominated by the light from a single star, the flux combining both smear registers is equivalent to that of a star  $\sim 6.8$  times fainter.

In Pope et al. (2016), we demonstrated a method for extracting precise light curves of bright stars in *Kepler* and *K2* from these collateral data, and presented light curves of a small number of variable stars as examples to illustrate this method. In this paper we present smear light curves of all unobserved or significantly under-observed stars brighter than  $Kp = 9$  in the *Kepler* field. This sample mostly consists of red giants and hot stars, containing only one G dwarf. We find no transiting planets, but detect one new eclipsing binary, and measure solar-like oscillations in 34 red giants. We do not model main sequence stars in great detail, but provide some discussion and initial classification of interesting variability. For the oscillating red giants that constitute the bulk of the sample, we determine the asteroseismic parameters  $\nu_{\text{max}}$  and  $\Delta\nu$ , and therefore stellar masses and  $\log g$  measurements. We have also obtained high-resolution optical spectroscopy of 63 stars, predominantly giants, with the Tillinghast Reflector Échelle Spectrograph (TRES). For the 34 stars with both spectroscopy and asteroseismic parameters we derive fundamental stellar parameters and elemental abundances. These asteroseismic constraints can be compared to those from *Gaia*, offering the opportunity both to test asteroseismic scaling relations and combine both datasets to refine the benchmark star properties further.

We have made all new data products and software discussed in this paper publicly available, and encourage interested readers to use these in their own research.

## 2 METHOD

We have obtained smear light curves for our sample of red giant stars with the `keplersmear` pipeline as described in Section 2.2, performed asteroseismology on all of these to extract  $\nu_{\text{max}}$  and therefore  $\log g$  as described in Section 2.3, and combined these with TRES spectra to obtain chemical abundances as described in Section 2.4.

## 2.1 Sample

We selected as our sample all stars on-silicon in *Kepler* with  $Kp < 9$  that were targeted for fewer than an arbitrary 8 quarters, the majority of which were previously entirely missing. Sixteen stars were to some extent observed conventionally: HD 174020 was targeted in LC for Q2, 6, 10, and 14; HD 175841 for Q11-12, 14-16, with SC for Q3; HD 176582 for Q12-13; HD 178090 for Q1, 3, and 10; HD 180682 for Q0, 3, and 7; HD 181069 for Q1, 10, 13, 14, and 17; HD 181878 for Q14-17; HD 182694 for Q2; HD 183124 for even quarters; HD 185351 for Q1-3 and with SC for Q16; HD 186155 for only Q1; HD 187217 for Q14-17; HD 188252 for only Q13; HD 189013 with SC for Q3 as a  $\gamma$  Dor; V380 Cyg for Q11 and with SC for Q7, 9, 10, 12-17; and V819 Cyg for Q14, 16 and 17. A number of these lay at the edge of a detector, with the result that in some cadences the centroid of the star did not lie on the chip; light curves from these targets were found to be of extremely low quality and all of these stars were discarded. After applying these criteria we obtained a list of 102 targets, which are listed in Table 1 with their *Kepler* magnitude  $Kp$  together with their spectral type from SIMBAD, *Gaia* DR2 apparent  $G$  magnitudes and  $Bp - Rp$  colours, *Gaia* DR2 calibrated distances from Bailer-Jones et al. (2018), variability classification and availability of TRES spectroscopy. A Venn diagram of the sample, noting the availability of asteroseismic fits Section 2.3 and spectroscopy (Section 2.4) is displayed in Figure 2. The *Kepler* spacecraft rotates between quarters, so that it cycles through four orientation ‘seasons’ each rotated from the last by  $90^\circ$ . Some stars did not land on silicon for all seasons: we have only one season of HD 179394; two for HD 187277, HD 226754, V554 Lyr, and BD+47 2891; and three for BD+43 3064. The addition of our sample to the conventionally-observed stars makes the *Kepler* survey magnitude-complete down to  $Kp = 9$  for all stars on-silicon.

Figure 1 shows these stars on a colour-magnitude diagram using *Gaia*  $Bp - Rp$  and absolute  $G$  magnitudes and *Gaia* DR2 calibrated distances (Bailer-Jones et al. 2018), overlaid on the *Kepler* sample from the Bedell [gaia-kepler.fun](https://github.com/bedell/gaia-kepler.fun) crossmatch. The smear targets in this diagram appear to have not merely higher apparent brightnesses than the general *Kepler* population, but also higher intrinsic luminosities. While this could simply arise from being selected for their apparent brightness, it is worth considering whether this is because of a bias in their parallax measurements. While *Gaia* parallaxes for very bright stars can be subject to systematic error, we have compared these to those found by *Hipparcos* (van Leeuwen 2007), and found close agreement for the brightest stars, with a scatter that increases with magnitude. We therefore suggest that parallax bias is not the reason for the smear sample sitting above the remainder of the *Kepler* sample.

We identify the evolutionary state of stars in the main sequence versus evolved stars first from the *Gaia* colour-magnitude diagram in Figure 1. Taking a cutoff in *Gaia*  $Bp - Rp > 1$ , we identify 64 of these stars as clearly evolved systems, and the remaining 38 lie apparently on the main sequence.

The coolest main sequence star, BD+43 3068, is a G0 dwarf with a  $G$  magnitude of 8.3 and a distance of  $53.8 \pm 0.1$  pc, and it is therefore surprising that it was not included in the nominal *Kepler* survey as a solar analogue. It is possible that it was previously misidentified as a giant. Regrettably, it is only possible to reconstruct a light curve with the 30 minute long cadence and therefore it is not possible to do asteroseismology on this bright, nearby solar-like star. Its light curve shows neither rotational modulation (as determined by its featureless autocorrelation) nor evidence for transits.

Considering stars lying close to the main sequence, from the *Kepler* power spectrum we identify solar-like oscillations in HD 182354 and HD 176209 at frequencies consistent with them being subgiants or contaminated with flux from red giants. [Get frequencies.](#)


## 2.2 Photometry

In preparing light curves of the *Kepler* smear stars, we have followed the methods described in Pope et al. (2016), with some improvements. We selected our input RA and Dec values from the *Kepler* Input Catalog (KIC) (Brown et al. 2011), and queried MAST to find the corresponding mean pixel position for a given *Kepler* quarter. We then measured the centroid of smear columns in the vicinity, and used these values to do raw aperture photometry. We found that the cosine-bell aperture used for raw photometry in Pope et al. (2016) can in some light curves introduce position-dependent systematics and jumps. We instead in this work have applied a super-Gaussian aperture,  $A \propto \exp \frac{-(x - x_0)^4}{w}$ , where  $x_0$  is the centroid and  $w$  a width in pixels. The very flat top of this function helps avoid significant variation with position, while still smoothly rolling off at the edges to avoid discontinuous artefacts. This is calculated on a grid of  $10 \times$  subsampled points in pixel space so that the sharply varying edge changes column weights smoothly as a function of centroid. We have then extracted photometry using apertures with a range of widths  $w \in \{1.5, 2, 3, 4, 5\}$  pixels.


From this raw photometry a background light curve was subtracted, which corrects for time-varying global systematics. Whereas in Pope et al. (2016) we subtracted a background estimate chosen manually, for this larger set of light curves, we have now chosen the lowest 25% of pixels by median flux as being unlikely to be contaminated by stars, and taken our background level to be the median of this at each time sample. To denoise this, we fit a Gaussian Process with a 30-day timescale squared exponential kernel using GEORGE (Ambikasaran et al. 2015), and our final background light curve is taken to be the posterior mean of this GP.

The dominant source of residual systematic errors in nominal *Kepler* time series is a common-mode variation primarily due to thermal changes on board the spacecraft, an issue which is traditionally dealt with by identifying and fitting a linear combination of systematic modes (Twicken et al. 2010; Stumpe et al. 2012; Smith et al. 2012; Petigura & Marcy 2012). We have adopted the same approach here, using the *Kepler* Pre-search Data Conditioning (PDC) Cotrending Basis Vectors (CBVs) available from MAST, finding least-squares fits of either the first 4 or 8 CBVs to each light curve. This can subtract astrophysical signals on long timescales, such that we use and recommend 4 CBV light curves for stars with variability on timescales longer than  $\sim 5$  days, or indeed raw uncorrected lightcurves for stars variable at high amplitude on  $\sim$  quarter timescales, but otherwise we recommend the 8 CBV light curves. There is some room for improvement here by simultaneously modelling astrophysical and instrumental variations, but this is beyond the scope of this paper. In the following, we will use the light curves with the lowest 6.5 hr Combined Differential Photometric Precision (CDPP) (Christiansen et al. 2012) out of all apertures, as calculated with the  $\kappa 2sc$  implementation (Aigrain et al. 2016). This is not necessarily the optimal choice for all red giants, especially those with oscillations on a 6.5 h timescale, but is a reasonable proxy nevertheless for white noise and leads to satisfactory results upon visual inspection of the present sample.

Because the smear data are collected along an entire CCD col-



**Figure 1.** *Gaia* colour-magnitude diagram of the Smear Campaign stars (orange and teal) overlaid on the sample of *Kepler* stars with *Gaia* parallax SNR > 25 (black), using the Bedell [gaia-kepler.fun](https://github.com/bjaspope/gaia-kepler.fun) crossmatch and *Gaia* DR2 calibrated distances from Bailer-Jones et al. (2018). The smear sample includes giants and hot main-sequence stars. Those giants for which TRES spectroscopy have been obtained are highlighted in teal. An interactive version of this diagram is available as supplementary material from the journal or at [benjaminpope.github.io/data/cmd\\_smear.html](https://benjaminpope.github.io/data/cmd_smear.html).



**Figure 2.** Venn diagram showing the overall structure of the *Kepler* Smear Campaign sample, showing the subsets for which TRES spectroscopy and asteroseismology are available and their overlap.

umn, there is the risk of contamination from other bright stars. This is especially true in doing asteroseismology of red giants, where the low-amplitude stochastically-excited oscillations can be washed

out in a power spectrum by the coherent high amplitude variations of a classical pulsator, even if the background star is much fainter. We can assess the importance of this contamination by considering the differences between odd and even quarters: because the *Kepler* spacecraft rotates  $90^\circ$  between successive quarters, any contaminant will lie in the same column as a smear target only every second quarter, falling in the other quarters in the same row but not necessarily the same column. We have therefore generated Lomb-Scargle periodograms (Lomb 1976; Scargle 1982) of each light curve, clipped for outliers, and considering only odd and even quarters, and visually inspected these for significant differences. In the great majority of cases they closely resemble one another, indicating that contamination is at worst a minor effect. In the case of HD 181878, a red giant, there is clear and significant contamination from an M giant, as is seen in Figure 3. Likewise HD 183383 shows two different stars, depending on the quarter: some parts are likely from an ellipsoidal variable with a period of 6.46 days, other parts are from an RR Lyrae pulsator.

### 2.3 Asteroseismology

Among the 64 red giants identified in this sample, for 30 the timescale of their variability is of the same order as a *Kepler* quarter and they are thus badly affected by systematics and systematics correction. In Table 1 we have noted these as ‘long-period variables’ (LPVs), without specifically meaning these are LPVs by a



**Table 1.** The full set of underobserved and unobserved stars for which new light curves have been produced in this smear catalogue. Calibrated *Gaia* distances are from Bailer-Jones et al. (2018). Some objects, such as HD 185351, were observed in long cadence in some quarters and short cadence in others, and this is noted accordingly. The eclipsing binary V2083 Cyg was detected by *Gaia*, but a parallax could not be obtained in DR2, possibly due to binary motion. Variability classes are determined by inspection, having their usual abbreviations. EV denotes an ellipsoidal variable, and RM rotational modulation, though these two can appear similar.  $\alpha^2$  CVn variables are chemically-peculiar stars with rotational spot modulation, and are noted separately from RM without chemical peculiarity.  $\gamma$  Dor/ $\delta$  Sct denotes a  $\gamma$  Dor/ $\delta$  Sct hybrid, not uncertainty. H+S denotes a ‘hump and spike’ star. Question marks indicate uncertainty, and dashes – that no significant variability is observed.

| Object     | KIC      | Spectral Type<br>(SIMBAD) | <i>Kp</i><br>(mag) | <i>G</i><br>(mag) | <i>Bp</i> – <i>Rp</i><br>(mag) | <i>Gaia</i> Distance<br>(pc)            | TRES | Variability<br>Class       |
|------------|----------|---------------------------|--------------------|-------------------|--------------------------------|---|------|----------------------------|
| 14 Cyg     | 7292420  | G8.5IIbFe-0.5             | 5.490              | 5.370             | -0.055                         | 194.3 <sup>+7.0</sup> <sub>-6.6</sub>   | –    | H+S                        |
| BD+36 3564 | 1575741  | F5II-III                  | 8.128              | 8.041             | 1.544                          | 547.1 <sup>+11.6</sup> <sub>-11.1</sub> | ✓    | RG                         |
| BD+39 3577 | 4989821  | G8III                     | 8.131              | 8.090             | 1.134                          | 311.7 <sup>+2.7</sup> <sub>-2.7</sub>   | ✓    | RG                         |
| BD+39 3882 | 4850372  | A2V                       | 8.259              | 8.159             | 0.616                          | 143.3 <sup>+0.7</sup> <sub>-0.7</sub>   | –    | ?                          |
| BD+42 3150 | 7091342  | B9III                     | 8.350              | 8.315             | 1.206                          | 546.0 <sup>+32.5</sup> <sub>-29.1</sub> | –    | ?                          |
| BD+42 3367 | 7447756  | B5V                       | 7.271              | 6.992             | 2.020                          | 762.0 <sup>+15.8</sup> <sub>-15.2</sub> | ✓    | LPV                        |
| BD+42 3393 | 6870455  | M1III                     | 7.664              | 7.414             | 1.952                          | 929.0 <sup>+25.9</sup> <sub>-24.5</sub> | ✓    | LPV                        |
| BD+43 3064 | 8075287  | G7IIIa                    | 8.284              | 8.203             | 1.599                          | 641.0 <sup>+20.3</sup> <sub>-19.1</sub> | ✓    | RG                         |
| BD+43 3068 | 8006792  | B1.1III+B2.5/3V:          | 8.308              | 8.268             | 0.839                          | 53.8 <sup>+0.1</sup> <sub>-0.1</sub>    | –    | –                          |
| BD+43 3171 | 7810954  | M3III                     | 8.373              | 8.178             | 1.858                          | 751.5 <sup>+17.2</sup> <sub>-16.5</sub> | ✓    | LPV                        |
| BD+43 3213 | 7747499  | A5III                     | 8.311              | 8.139             | 1.876                          | 948.8 <sup>+25.8</sup> <sub>-24.5</sub> | ✓    | LPV                        |
| BD+47 2825 | 10337574 | B2III                     | 8.251              | 8.236             | 1.329                          | 485.8 <sup>+7.3</sup> <sub>-7.1</sub>   | –    | EB                         |
| BD+47 2891 | 10347606 | K1III                     | 8.680              | 8.625             | 1.291                          | 262.8 <sup>+1.1</sup> <sub>-1.6</sub>   | –    | RG                         |
| BD+48 2904 | 11085556 | K5                        | 8.487              | 8.439             | 1.355                          | 400.9 <sup>+5.4</sup> <sub>-5.3</sub>   | –    | RG                         |
| BD+48 2955 | 10988024 | K2                        | 7.961              | 7.899             | 1.549                          | 589.4 <sup>+11.6</sup> <sub>-11.1</sub> | ✓    | RG                         |
| HD 174020  | 7800227  | K2                        | 6.753              | 6.600             | 1.754                          | 433.1 <sup>+4.2</sup> <sub>-4.1</sub>   | ✓    | RG                         |
| HD 174177  | 9630812  | M4-IIIa                   | 6.575              | 6.483             | 0.119                          | 223.9 <sup>+1.7</sup> <sub>-1.6</sub>   | –    | ?                          |
| HD 174676  | 7420037  | K0                        | 7.481              | 7.440             | 2.434                          | 993.3 <sup>+26.7</sup> <sub>-25.4</sub> | ✓    | LPV                        |
| HD 174829  | 7339102  | K1III                     | 6.967              | 6.928             | 1.391                          | 355.0 <sup>+3.5</sup> <sub>-3.4</sub>   | ✓    | RG                         |
| HD 175132  | 6020867  | K0                        | 6.362              | 6.242             | -0.063                         | 333.3 <sup>+5.9</sup> <sub>-5.7</sub>   | –    | $\alpha^2$ CVn             |
| HD 175466  | 7340766  | B3V                       | 6.165              | 5.919             | 1.905                          | 397.8 <sup>+6.8</sup> <sub>-6.6</sub>   | –    | LPV                        |
| HD 175740  | 6265087  | K0                        | 5.212              | 5.152             | 1.171                          | 81.5 <sup>+0.6</sup> <sub>-0.6</sub>    | ✓    | RG                         |
| HD 175841  | 4989900  | B9IIpSi                   | 6.885              | 6.797             | 0.172                          | 241.0 <sup>+2.1</sup> <sub>-2.1</sub>   | –    | $\gamma$ Dor/ $\delta$ Sct |
| HD 175884  | 6584587  | B0.5IIIIn                 | 6.210              | 6.144             | 1.448                          | 238.9 <sup>+1.5</sup> <sub>-1.4</sub>   | ✓    | RG                         |
| HD 176209  | 9327530  | B3Ve                      | 7.437              | 7.365             | 0.091                          | 282.2 <sup>+2.7</sup> <sub>-2.7</sub>   | –    | ?                          |
| HD 176582  | 4136285  | K0                        | 6.510              | 6.383             | -0.232                         | 298.6 <sup>+3.9</sup> <sub>-3.8</sub>   | –    | RM                         |
| HD 176626  | 7943968  | G8II                      | 6.933              | 6.841             | 0.035                          | 224.8 <sup>+1.8</sup> <sub>-1.7</sub>   | –    | RM                         |
| HD 176894  | 6267965  | M0II-III                  | 7.700              | 7.610             | 0.530                          | 82.8 <sup>+0.2</sup> <sub>-0.2</sub>    | –    | $\gamma$ Dor               |
| HD 177697  | 4994443  | K5                        | 7.300              | 6.764             | 2.338                          | 472.0 <sup>+5.4</sup> <sub>-5.3</sub>   | –    | RG                         |
| HD 177781  | 2970780  | B5V                       | 7.744              | 7.701             | 1.024                          | 296.2 <sup>+2.6</sup> <sub>-2.5</sub>   | –    | $\gamma$ Dor/ $\delta$ Sct |
| HD 178090  | 6675338  | A2IV                      | 6.758              | 6.549             | 1.892                          | 583.0 <sup>+8.3</sup> <sub>-8.3</sub>   | –    | LPV                        |
| HD 178797  | 10064283 | K0                        | 7.312              | 7.249             | 1.478                          | 406.1 <sup>+4.8</sup> <sub>-4.7</sub>   | ✓    | RG                         |
| HD 178910  | 11288450 | G5                        | 7.864              | 7.848             | 1.346                          | 291.3 <sup>+2.4</sup> <sub>-2.4</sub>   | ✓    | RG                         |
| HD 179394  | 7105221  | K5                        | 7.575              | 7.475             | -0.100                         | 476.2 <sup>+12.2</sup> <sub>-11.6</sub> | ✓    | –                          |
| HD 179395  | 6593264  | A0V                       | 7.168              | 7.070             | 0.067                          | 233.9 <sup>+1.7</sup> <sub>-1.7</sub>   | –    | RM                         |
| HD 179396  | 3838362  | K5                        | 8.001              | 7.970             | 1.244                          | 321.2 <sup>+2.7</sup> <sub>-2.6</sub>   | ✓    | RG                         |
| HD 179959  | 10265370 | K4III                     | 6.280              | 6.258             | 1.168                          | 499.2 <sup>+7.2</sup> <sub>-7.0</sub>   | ✓    | RG                         |
| HD 180312  | 4551179  | A2                        | 7.970              | 7.834             | 1.162                          | 290.5 <sup>+2.4</sup> <sub>-2.4</sub>   | ✓    | RG                         |
| HD 180475  | 11656042 | A3                        | 7.664              | 7.595             | 1.489                          | 546.1 <sup>+8.0</sup> <sub>-7.8</sub>   | ✓    | RG                         |
| HD 180658  | 6195870  | A2                        | 7.932              | 7.871             | 1.256                          | 282.2 <sup>+2.3</sup> <sub>-2.3</sub>   | ✓    | RG                         |
| HD 180682  | 5177450  | K5                        | 6.617              | 6.532             | 1.486                          | 295.8 <sup>+2.3</sup> <sub>-2.3</sub>   | ✓    | LPV                        |
| HD 181022  | 3946721  | A2V                       | 6.496              | 6.248             | 1.892                          | 317.7 <sup>+2.7</sup> <sub>-2.7</sub>   | ✓    | LPV                        |
| HD 181069  | 4049174  | A0                        | 6.279              | 6.264             | 1.237                          | 144.2 <sup>+0.6</sup> <sub>-0.6</sub>   | ✓    | RG                         |
| HD 181097  | 4149233  | A5                        | 7.920              | 7.848             | 1.434                          | 434.3 <sup>+6.2</sup> <sub>-6.0</sub>   | ✓    | RG                         |
| HD 181328  | 12456737 | A3                        | 7.182              | 6.614             | 2.334                          | 353.9 <sup>+3.3</sup> <sub>-3.3</sub>   | ✓    | LPV                        |
| HD 181521  | 5180075  | K0                        | 6.939              | 6.852             | 0.059                          | 217.8 <sup>+3.4</sup> <sub>-3.3</sub>   | –    | $\gamma$ Dor/ $\delta$ Sct |
| HD 181596  | 11910615 | M3                        | 7.050              | 6.863             | 1.841                          | 591.1 <sup>+8.1</sup> <sub>-7.8</sub>   | ✓    | RG                         |
| HD 181597  | 11555267 | K5III                     | 6.040              | 5.985             | 1.283                          | 135.8 <sup>+0.3</sup> <sub>-0.3</sub>   | ✓    | RG                         |
| HD 181681  | 5092997  | B9                        | 6.864              | 6.696             | 1.798                          | 585.0 <sup>+0.1</sup> <sub>-0.1</sub>   | ✓    | RG                         |
| HD 181778  | 7816792  | B8V                       | 7.545              | 7.514             | 1.315                          | 374.5 <sup>+3.4</sup> <sub>-3.4</sub>   | ✓    | RG                         |
| HD 181878  | 4830109  | M1                        | 6.698              | 6.587             | 1.003                          | 259.5 <sup>+1.8</sup> <sub>-1.8</sub>   | ✓    | RG                         |
| HD 181880  | 3337423  | M5                        | 7.982              | 7.940             | 1.498                          | 541.2 <sup>+10.1</sup> <sub>-9.7</sub>  | ✓    | RG                         |

**Table 1** – *continued* The full set of underobserved and unobserved stars for which new light curves have been produced in this smear catalogue. Calibrated *Gaia* distances are from [Bailer-Jones et al. \(2018\)](#).

| Object     | KIC      | Spectral Type<br>(SIMBAD) | <i>Kp</i><br>(mag) | <i>G</i><br>(mag) | <i>Bp</i> – <i>Rp</i><br>(mag) | <i>Gaia</i> Distance<br>(pc)              | TRES | Variability<br>Class |
|------------|----------|---------------------------|--------------------|-------------------|--------------------------------|---|------|----------------------|
| HD 182354  | 2156801  | B9                        | 6.320              | 6.291             | 1.253                          | 228.9 <sup>+1.7</sup> <sub>-1.7</sub>     | –    | RG                   |
| HD 182531  | 11188366 | B9IV                      | 7.955              | 7.859             | 1.502                          | 599.3 <sup>+9.2</sup> <sub>-8.9</sub>     | ✓    | RG                   |
| HD 182692  | 10728753 | M0                        | 7.310              | 7.247             | 1.227                          | 226.6 <sup>+1.3</sup> <sub>-1.3</sub>     | ✓    | RG                   |
| HD 182694  | 7680115  | K5                        | 5.722              | 5.598             | 1.061                          | 133.1 <sup>+0.7</sup> <sub>-0.7</sub>     | ✓    | RG                   |
| HD 182737  | 1572070  | K0                        | 7.820              | 7.758             | 0.421                          | 460.3 <sup>+6.7</sup> <sub>-6.5</sub>     | –    | RM                   |
| HD 183124  | 8752618  | K0                        | 6.441              | 6.395             | 1.176                          | 160.7 <sup>+0.8</sup> <sub>-0.8</sub>     | ✓    | RG                   |
| HD 183203  | 12208512 | B8                        | 6.928              | 6.530             | 2.116                          | 476.9 <sup>+3.9</sup> <sub>-3.8</sub>     | ✓    | LPV                  |
| HD 183362  | 2715115  | K0                        | 6.394              | 6.208             | -0.041                         | 571.1 <sup>+18.2</sup> <sub>-17.2</sub>   | –    | H+S                  |
| HD 183383  | 6777469  | M3III                     | 7.640              | 7.537             | 0.081                          | 357.1 <sup>+5.5</sup> <sub>-5.3</sub>     | –    | ?                    |
| HD 184147  | 9651435  | A0                        | 7.251              | 7.145             | -0.037                         | 175.5 <sup>+2.6</sup> <sub>-2.5</sub>     | –    | ?                    |
| HD 184215  | 11031549 | –                         | 7.321              | 7.189             | -0.135                         | 361.2 <sup>+6.4</sup> <sub>-6.4</sub>     | –    | SPB                  |
| HD 184483  | 7756961  | M0                        | 7.246              | 6.719             | 2.337                          | 492.9 <sup>+5.4</sup> <sub>-5.4</sub>     | ✓    | LPV                  |
| HD 184565  | 6047321  | K0                        | 7.972              | 7.943             | 1.024                          | 380.9 <sup>+4.3</sup> <sub>-4.2</sub>     | –    | LPV                  |
| HD 184787  | 6528001  | B8                        | 6.757              | 6.658             | -0.003                         | 139.6 <sup>+1.1</sup> <sub>-1.1</sub>     | ✓    | H+S                  |
| HD 184788  | 6129225  | A0                        | 7.249              | 7.143             | -0.055                         | 226.5 <sup>+2.4</sup> <sub>-2.3</sub>     | –    | RM                   |
| HD 184875  | 6954647  | B0III                     | 5.403              | 5.279             | 0.107                          | 172.6 <sup>+3.3</sup> <sub>-3.2</sub>     | –    | γ Dor                |
| HD 185117  | 9094435  | B9                        | 7.696              | 7.472             | 1.921                          | 817.7 <sup>+14.8</sup> <sub>-14.3</sub>   | –    | LPV                  |
| HD 185286  | 7966681  | K2                        | 6.151              | 6.055             | 1.645                          | 263.5 <sup>+3.9</sup> <sub>-3.8</sub>     | ✓    | RG                   |
| HD 185351  | 8566020  | K5                        | 5.034              | 4.882             | 1.091                          | 41.2 <sup>+0.1</sup> <sub>-0.1</sub>      | ✓    | RG                   |
| HD 185397  | 3455268  | K5                        | 6.953              | 6.855             | 0.421                          | 180.0 <sup>+1.0</sup> <sub>-1.0</sub>     | –    | δ Sct                |
| HD 185524  | 8960196  | F0                        | 8.022              | 7.953             | 1.368                          | 753.4 <sup>+15.9</sup> <sub>-15.2</sub>   | ✓    | LPV                  |
| HD 186121  | 7456762  | K5                        | 5.773              | 5.176             | 2.250                          | 475.2 <sup>+35.1</sup> <sub>-30.7</sub>   | ✓    | LPV                  |
| HD 186155  | 9163520  | G5                        | 5.055              | 4.923             | 0.529                          | 50.6 <sup>+0.4</sup> <sub>-0.4</sub>      | –    | H+S                  |
| HD 186255  | 4937492  | A0                        | 6.966              | 6.862             | 0.252                          | 254.5 <sup>+4.1</sup> <sub>-4.0</sub>     | –    | δ Sct                |
| HD 186727  | 12316020 | K2                        | 7.499              | 6.917             | 2.388                          | 581.7 <sup>+9.2</sup> <sub>-8.9</sub>     | ✓    | LPV                  |
| HD 186994  | 8766240  | K2                        | 7.585              | 7.451             | -0.185                         | 1866.1 <sup>+38.1</sup> <sub>-120.6</sub> | –    | EB                   |
| HD 187217  | 11824273 | K0                        | 6.399              | 6.345             | 1.273                          | 243.2 <sup>+1.8</sup> <sub>-1.8</sub>     | ✓    | RG                   |
| HD 187277  | 6967644  | K0                        | 7.579              | 7.464             | 0.282                          | 96.9 <sup>+0.4</sup> <sub>-0.4</sub>      | –    | –                    |
| HD 187372  | 10679281 | K5                        | 5.672              | 5.313             | 2.047                          | 306.4 <sup>+10.3</sup> <sub>-9.6</sub>    | ✓    | LPV                  |
| HD 188252  | 10683303 | K2                        | 6.007              | 5.864             | -0.276                         | 1000.6 <sup>+82.6</sup> <sub>-71.1</sub>  | –    | SPB                  |
| HD 188537  | 9110718  | K0II                      | 7.382              | 7.324             | 1.345                          | 629.9 <sup>+11.4</sup> <sub>-11.0</sub>   | ✓    | RG                   |
| HD 188629  | 8710324  | K0                        | 7.743              | 7.546             | 1.888                          | 651.0 <sup>+12.0</sup> <sub>-11.6</sub>   | ✓    | LPV                  |
| HD 188875  | 5041881  | K                         | 6.164              | 6.091             | 1.584                          | 683.8 <sup>+12.4</sup> <sub>-11.9</sub>   | ✓    | RG                   |
| HD 189013  | 10096499 | K2                        | 6.922              | 6.840             | 0.225                          | 188.8 <sup>+6.4</sup> <sub>-6.0</sub>     | –    | γ Dor                |
| HD 189178  | 5219588  | K2                        | 5.552              | 5.410             | -0.106                         | 347.3 <sup>+13.0</sup> <sub>-12.1</sub>   | –    | H+S                  |
| HD 189636A | 10298067 | –                         | 8.025              | 8.118             | 1.211                          | 384.7 <sup>+6.0</sup> <sub>-5.8</sub>     | –    | ?                    |
| HD 189636B | 10298061 | K0                        | 8.107              | 8.024             | 1.316                          | 376.4 <sup>+3.8</sup> <sub>-4.7</sub>     | –    | ?                    |
| HD 189684  | 9305008  | –                         | 5.982              | 5.881             | 0.246                          | 125.2 <sup>+6.2</sup> <sub>-5.7</sub>     | –    | EV                   |
| HD 189750  | 8521828  | K5                        | 8.052              | 8.061             | 1.207                          | 327.0 <sup>+3.0</sup> <sub>-2.9</sub>     | –    | ?                    |
| HD 190149  | 8262528  | G5                        | 6.488              | 6.171             | 2.031                          | 409.4 <sup>+3.8</sup> <sub>-3.7</sub>     | ✓    | LPV                  |
| HD 226754  | 6234579  | –                         | 7.829              | 7.702             | 1.652                          | 391.8 <sup>+6.1</sup> <sub>-5.9</sub>     | ✓    | RG                   |
| V2079 Cyg  | 8818020  | K0                        | 7.174              | 7.034             | -0.221                         | 321.5 <sup>+3.7</sup> <sub>-3.6</sub>     | –    | α <sup>2</sup> CVn   |
| V2083 Cyg  | 10342012 | F5                        | 6.902              | 6.813             | 0.351                          | –   | –    | EB                   |
| V380 Cyg   | 5385723  | K5                        | 5.771              | 5.632             | -0.062                         | 1044.7 <sup>+116.6</sup> <sub>-95.6</sub> | –    | EB                   |
| V398 Lyr   | 4042516  | G0                        | 7.024              | 5.403             | 3.406                          | 494.7 <sup>+34.9</sup> <sub>-30.6</sub>   | ✓    | RG                   |
| V543 Lyr   | 5429169  | K5                        | 6.299              | 6.160             | -0.217                         | 345.1 <sup>+5.6</sup> <sub>-5.4</sub>     | –    | SPB                  |
| V546 Lyr   | 6267345  | K0                        | 7.385              | 6.784             | 2.443                          | 587.8 <sup>+13.1</sup> <sub>-13.0</sub>   | ✓    | LPV                  |
| V547 Lyr   | 5429948  | M0                        | 6.199              | 5.228             | 2.725                          | 288.9 <sup>+13.0</sup> <sub>-12.0</sub>   | ✓    | LPV                  |
| V554 Lyr   | 5001462  | K0                        | 8.179              | 8.092             | -0.129                         | 335.7 <sup>+4.6</sup> <sub>-4.5</sub>     | –    | α <sup>2</sup> CVn   |
| V819 Cyg   | 10618721 | K0                        | 6.381              | 6.243             | -0.160                         | 1114.0 <sup>+70.9</sup> <sub>-63.0</sub>  | –    | SPB                  |

particular astrophysical definition, and they are discussed further in Section 3.1.3. For the 34 giants for which there is high-SNR shorter-timescale variability, we have attempted to extract the asteroseismic parameters  $\nu_{\max}$  and  $\langle \Delta\nu \rangle$  ([Kjeldsen & Bedding 1995](#); [Chaplin](#)

& [Miglio 2013](#)). These constrain fundamental stellar parameters through the approximate scaling relations:

$$\nu_{\max} \propto \frac{g}{g_{\odot}} \cdot \left( \frac{T_{\text{eff}}}{T_{\text{eff}\odot}} \right)^{\frac{1}{2}} \quad (1)$$



**Figure 3.** Power spectra of odd and even quarters of HD 181778. It is clear from inspection that while odd quarters have the power spectrum expected of a giant star, even quarters have very high amplitude coherent oscillations typical of an M giant.

and

$$\langle \Delta\nu \rangle \propto \sqrt{\langle \rho \rangle} = \sqrt{MR^{-3}} \quad (2)$$

We have followed the method of [Davies & Miglio \(2016\)](#), obtaining a Lomb-Scargle periodogram of the smoothed time series according to the method of [García et al. \(2011\)](#). The posterior distribution of the asteroseismic parameters is obtained with a Markov Chain Monte Carlo fit to the smoothed periodogram, applying the combined granulation and oscillation model of [Kallinger et al. \(2014\)](#). This consists of two Harvey profiles for the granulation ([Harvey 1985](#)), a Gaussian envelope for the stellar oscillations, and a white noise background for instrumental noise. The marginal posterior distribution for the oscillation envelope is well-approximated by a single Gaussian, and we have taken its median and standard deviation to be our estimates for  $\nu_{\max}$  and its uncertainty.

To estimate  $\Delta\nu$ , we have divided the power spectrum through

by the granulation and noise models to obtain a signal-to-noise spectrum, and fit a sum of Lorentzians separated by mean large ( $\Delta\nu$ ) and small ( $\delta\nu$ ) separations to the part of this spectrum in the vicinity of  $\nu_{\max}$ . In practice, for this dataset,  $\delta\nu$  is not constrained, but mean  $\langle \Delta\nu \rangle$  is typically well-constrained and its posterior marginal distribution is well-represented by a single Gaussian as with  $\nu_{\max}$ .

We obtain good estimates of these asteroseismic parameters for 34 targets, presented in Table 2. In the remainder of cases, as noted above, we find that the very-low-frequency ( $\lesssim 2\mu\text{Hz}$ ) oscillations are affected by filter artefacts from detrending, and we are not able to obtain good estimates for these stars.

Once  $\nu_{\max}$  has been estimated, we have used Equation 1, the asteroseismic scaling relation for  $\nu_{\max}$  ([Brown et al. 1991](#); [Kjeldsen & Bedding 1995](#)), to estimate  $\log g$  in order to inform extraction of chemical abundances from spectra. Using the initial spectroscopic estimate of  $T_{\text{eff}}$ , which is not significantly informed by  $\nu_{\max}$ , uncertainties in  $\nu_{\max}$  are propagated with Monte Carlo sampling.

For eight stars, we have found that the asteroseismic fit is unsatisfactory: for BD+39 388 we cannot detect the expected oscillations; for BD+43 3064 there are significant peaks but these are not consistent with the pattern expected from a red giant; for HD 179959 and HD 187217 we suspect contamination with the oscillations of a second giant, which is hard to remove from smear light curves; while for HD 188629, HD 188639 and HD 188875 we can extract a  $\nu_{\max}$  but not a robust  $\Delta\nu$ . One star in our sample, HD 185351, has a mode envelope that is not well fit by our model. The smear light curve for this star has already been published by [Hjørringgaard et al. \(2017\)](#), who showed with detailed asteroseismic modelling that it had a zero-age main sequence mass of  $\sim 1.60M_{\odot}$  (a so-called ‘retired A star’) and used it to calibrate the convective overshoot parameter for low-luminosity red giants. The global asteroseismic modelling presented here should therefore be considered to be superseded by the more detailed model of [Hjørringgaard et al. \(2017\)](#).

## 2.4 Spectroscopy

We have obtained high-resolution spectroscopy with TRES for 63 stars, mainly giants, in order to constrain stellar parameters and elemental abundances. Operating with spectral resolving power  $R = 44000$ , we have obtained spectra with signal-to-noise ratios (SNRs) of tens to hundreds per resolution element. Although this resolution and SNR are sufficient for an exploratory study, for more detailed analysis it will be desirable to use APOGEE or similar instruments to obtain greater spectral coverage. From this observing run we have 34 unique targets with seismic  $\log g$  and spectra, one more star than the *Gaia*-ESO benchmark set and a significant addition to the ensemble of bright red giants with asteroseismic parameter determinations. Due to observing constraints, we were unable to obtain spectra for BD+42 3150, BD+48 2904, HD 176209, HD 182354, HD 189636AB, or HD 189750.

To derive stellar parameters from our TRES spectra, we initially ran the Stellar Parameter Classification (SPC; [Buchhave et al. 2012](#)) code to determine  $T_{\text{eff}}$  and  $\log g$ , using the SPC  $T_{\text{eff}}$  to inform the asteroseismic estimation of  $\log g$  from  $\nu_{\max}$ . For deriving abundances,  $T_{\text{eff}}$  is fixed from the results of an initial SPC fit, while  $\log g$  is fixed to the seismic values. The other stellar atmospheric parameters including the microturbulent velocity ( $v_{\text{mic}}$ ), and broadening (convolution by  $V_{\text{mac}}$ ,  $v_{\text{sin } i}$  and the instrumental line profile) as well as [Fe/H] and chemical abundances for 20 chemical species are derived using the Brussels Automatic Code for Characterizing High accuracy Spectra (BACCHUS; [Masseron et al. 2016](#)), and the results from this calculation are displayed in Table 3. BAC-

**Table 2.** Bulk asteroseismic parameters  $\Delta\nu$ ,  $\nu_{\max}$ , and  $\epsilon$  for the red giant sample as discussed in Section 2.3.

| Object     | $\Delta\nu$<br>( $\mu\text{Hz}$ ) | $\nu_{\max}$<br>( $\mu\text{Hz}$ ) | $\epsilon$      |
|------------|-----------------------------------|------------------------------------|-----------------|
| BD+36 3564 | $0.95 \pm 0.03$                   | $5.08 \pm 0.10$                    | $0.83 \pm 0.20$ |
| BD+39 3577 | $1.68 \pm 0.01$                   | $13.27 \pm 0.32$                   | $0.74 \pm 0.06$ |
| BD+42 3150 | $4.22 \pm 0.03$                   | $38.32 \pm 0.96$                   | $0.70 \pm 0.07$ |
| BD+43 3171 | $0.42 \pm 0.05$                   | $1.98 \pm 0.05$                    | $0.80 \pm 0.17$ |
| BD+43 3213 | $0.49 \pm 0.01$                   | $2.56 \pm 0.06$                    | $1.01 \pm 0.07$ |
| BD+48 2904 | $2.85 \pm 0.01$                   | $23.13 \pm 0.72$                   | $0.86 \pm 0.08$ |
| BD+48 2955 | $0.90 \pm 0.01$                   | $5.44 \pm 0.08$                    | $0.81 \pm 0.05$ |
| HD 174020  | $0.56 \pm 0.02$                   | $2.48 \pm 0.10$                    | $0.89 \pm 0.08$ |
| HD 174829  | $1.28 \pm 0.01$                   | $7.95 \pm 0.16$                    | $0.78 \pm 0.06$ |
| HD 175740  | $5.93 \pm 0.01$                   | $64.33 \pm 0.78$                   | $1.00 \pm 0.02$ |
| HD 175884  | $1.12 \pm 0.01$                   | $7.07 \pm 0.11$                    | $0.96 \pm 0.08$ |
| HD 176209  | $4.22 \pm 0.08$                   | $36.08 \pm 0.77$                   | $0.87 \pm 0.06$ |
| HD 178797  | $1.03 \pm 0.02$                   | $6.34 \pm 0.09$                    | $0.74 \pm 0.29$ |
| HD 178910  | $3.64 \pm 0.02$                   | $32.06 \pm 0.31$                   | $0.83 \pm 0.05$ |
| HD 179396  | $3.76 \pm 0.02$                   | $31.02 \pm 0.44$                   | $0.92 \pm 0.03$ |
| HD 180312  | $4.17 \pm 0.02$                   | $33.84 \pm 0.28$                   | $0.96 \pm 0.04$ |
| HD 180475  | $0.82 \pm 0.00$                   | $4.34 \pm 0.10$                    | $0.68 \pm 0.03$ |
| HD 180658  | $4.00 \pm 0.02$                   | $33.76 \pm 0.50$                   | $0.90 \pm 0.05$ |
| HD 180682  | $0.77 \pm 0.05$                   | $3.68 \pm 0.08$                    | $1.07 \pm 0.15$ |
| HD 181022  | $0.38 \pm 0.01$                   | $1.58 \pm 0.03$                    | $0.70 \pm 0.10$ |
| HD 181069  | $4.43 \pm 0.01$                   | $41.46 \pm 0.32$                   | $0.90 \pm 0.02$ |
| HD 181097  | $1.61 \pm 0.02$                   | $11.16 \pm 0.14$                   | $0.72 \pm 0.36$ |
| HD 181597  | $3.11 \pm 0.01$                   | $25.84 \pm 0.25$                   | $0.97 \pm 0.02$ |
| HD 181778  | $2.56 \pm 0.02$                   | $22.86 \pm 0.29$                   | $0.72 \pm 0.06$ |
| HD 181880  | $1.04 \pm 0.01$                   | $6.54 \pm 0.10$                    | $0.76 \pm 0.05$ |
| HD 182354  | $2.66 \pm 0.01$                   | $24.73 \pm 0.37$                   | $0.74 \pm 0.04$ |
| HD 182531  | $1.03 \pm 0.00$                   | $6.47 \pm 0.09$                    | $0.86 \pm 0.03$ |
| HD 182692  | $4.66 \pm 0.01$                   | $44.38 \pm 0.47$                   | $0.87 \pm 0.02$ |
| HD 182694  | $5.71 \pm 0.01$                   | $69.78 \pm 1.02$                   | $0.94 \pm 0.25$ |
| HD 183124  | $4.39 \pm 0.01$                   | $39.59 \pm 0.29$                   | $0.95 \pm 0.03$ |
| HD 185286  | $0.72 \pm 0.01$                   | $4.23 \pm 0.10$                    | $0.73 \pm 0.08$ |
| HD 188537  | $1.55 \pm 0.01$                   | $13.40 \pm 0.34$                   | $0.72 \pm 0.07$ |
| HD 189636  | $2.91 \pm 0.01$                   | $25.97 \pm 0.74$                   | $0.97 \pm 0.04$ |
| HD 189750  | $4.16 \pm 0.04$                   | $36.14 \pm 0.58$                   | $0.94 \pm 0.08$ |
| HD 226754  | $1.19 \pm 0.01$                   | $7.41 \pm 0.19$                    | $0.74 \pm 0.08$ |

CHUS uses an interpolation scheme through a grid of MARCS model atmospheres (Gustafsson et al. 2008) in combination with TURBOSPECTRUM (Alvarez & Plez 1998; Plez 2012). For the calculation of synthetic spectra, atomic line information has been taken from the fifth version of the Gaia-ESO linelist (Heiter et al., in preparation). Additionally we used the molecular species for CH (Masseron et al. 2014), CN, NH, OH, MgH C<sub>2</sub> (T. Masseron, private communication). The SiH molecular information is adopted from the Kurucz linelists and the information for TiO, ZrO, FeH, CaH from B. Plez (private communication).

Individual elemental abundances are derived by first fixing the stellar atmospheric parameters to those determined above. Spectra are then synthesized in regions centered around an absorption feature of the element we want to derive. The spectra generated will have different  $[X/\text{Fe}]$  values. A  $\chi^2$  minimization procedure is then done to derive the best fitting abundance for each line. The reported abundances are the median  $[X/\text{Fe}]$  value of the various line regions for a given element. **Ben: Maybe remove this if we do not actually implement line-by-line differential; Do we have a TRES Arcturus spectrum.** Abundance uncertainties reported are the standard error in the line-by-line abundance ratios. Where only one line exists for a given element, we conservatively assume

the standard error is 0.10 dex. In principle, these uncertainties are underestimated because they do not include the errors driven by imperfect stellar parameter values and other systematic errors arising, for instance, from incorrect line list data. We do note, however, that use of asteroseismology really reduces the uncertainties caused by the stellar parameters (see Hawkins et al. 2016c, for a longer discussion on this). To achieve the most precise abundances we have derived them using both with and without a line-by-line differential approach with respect to Arcturus ( $\alpha$  Boötis) using the method described by Jofré et al. (2015) and the Arcturus abundances from (Hawkins et al. 2016c). The results of these absolute abundance calculations **without the line-by-line differential analysis implemented?**, are presented in Tables 4, 5 and 6. Because for most elements Arcturus differential abundances are not available, these are provided as supplementary online-only material. No abundances for oxygen could be reliably derived for any of the stars in our spectroscopic sample by either method.

### 3 RESULTS

#### 3.1 Red Giants

##### 3.1.1 Chemical Composition

The chemical composition for each star was measured in the  $\alpha$  (Mg, Ti, Si, Ca), odd-Z (Al, Na, Cu, Sc, V), Fe-peak (Fe, Ni, Zn, Co, Cr, Mn), and neutron capture (Sr, Y, Zr, Ba, La, Nd, Eu) elemental families. This was done to assess the Galactic populations to which these stars belong. The first thing to note is that the metallicities, which are tabulated in Table 3, are too high (with  $-0.51 < [\text{M}/\text{H}] < +0.14$  dex) to be belong to the Galactic halo, whose peak metallicity is around  $\sim -1.50$  (e.g. Chiba & Beers 2000). Furthermore, the distance distribution, noted in Table 1, indicates that all stars are located within a few kpc of the Sun and thus are not apart of the Galactic bulge. Thus, these stars are drawn from only the Galactic thick and thin disks. We provide a detailed chemical abundance analysis below to support this claim.

##### $\alpha$ elements:

The  $\alpha$  elements are largely dispersed into the interstellar medium through Type II supernovae (SNII) (Matteucci & Recchi 2001). It has been shown by many studies (e.g. Edvardsson et al. 1993; Adibekyan et al. 2012; Feltzing & Chiba 2013; Bensby et al. 2014, and references therein), that the Galactic thick disk and thin disk separate in the  $\alpha$  elements, where the thick disk is enhanced in  $[\text{Mg}, \text{Si}, \text{Ca}, \text{Ti}/\text{Fe}]$  compared to the Galactic thin disk at a given metallicity. In Fig. 4, we display the  $[\text{Mg}, \text{Si}, \text{Ca}, \text{Ti}/\text{Fe}]$  abundance ratios as a function of  $[\text{Fe}/\text{H}]$  for our stars (black circles) compared<sup>1</sup> to representative disk stars from (Bensby et al. 2014, open red square) and (Adibekyan et al. 2012, open orange triangles).

For most of the stars in our sample, the  $[\text{Mg}, \text{Ti}, \text{Ca}/\text{Fe}]$  abundance ratios are enhanced. The commonly used  $[\alpha/\text{Fe}]$  abundance ratio is the average of Mg, Ti, Si, Ca (thus it is  $([\text{Mg}/\text{Fe}] + [\text{Ca}/\text{Fe}] + [\text{Si}/\text{Fe}] + [\text{Ti}/\text{Fe}]/4.0)$ ) is also enhanced in most stars. This is consistent with most stars belonging to the Galactic thick disk. Though there are a handful of stars where the typical  $[\alpha/\text{Fe}]$  ratio is solar.

##### Odd-Z elements:

The odd-Z elements Na and Al are also shown in Fig. 4. These elements are significantly enhanced

<sup>1</sup> There are likely to be systematics between our  $[X/\text{Fe}]$  abundance scale and those of our comparison samples.



**Table 3.** Fundamental stellar parameters for the red giant sample as determined jointly by asteroseismology (asteroseismic  $\log g$ ; Section 2.3) and spectroscopy (RV,  $T_{\text{eff}}$ ,  $\log g$ , [M/H],  $V \sin i$ , and SNR; Section 2.4.)

| Object     | RV<br>(km/s)      | $T_{\text{eff}}$<br>(K) | $\log g$        | [M/H]            | $V \sin i$<br>(km/s) | SNR   |
|------------|-------------------|-------------------------|-----------------|------------------|----------------------|-------|
| BD+36 3564 | $-77.84 \pm 0.05$ | $4301 \pm 50$           | $1.58 \pm 0.01$ | $-0.34 \pm 0.08$ | $5.14 \pm 0.50$      | 71.8  |
| BD+39 3577 | $-14.81 \pm 0.07$ | $5079 \pm 50$           | $2.03 \pm 0.01$ | $-0.11 \pm 0.08$ | $3.98 \pm 0.50$      | 92.8  |
| BD+43 3171 | $-16.32 \pm 0.11$ | $4072 \pm 50$           | $1.16 \pm 0.01$ | $-0.17 \pm 0.08$ | $5.68 \pm 0.50$      | 68.6  |
| BD+43 3213 | $-14.16 \pm 0.16$ | $4131 \pm 50$           | $1.27 \pm 0.01$ | $0.07 \pm 0.08$  | $6.24 \pm 0.50$      | 57.3  |
| BD+48 2955 | $1.66 \pm 0.04$   | $4344 \pm 50$           | $1.61 \pm 0.01$ | $-0.32 \pm 0.08$ | $4.78 \pm 0.50$      | 31.7  |
| HD 174020  | $-14.84 \pm 0.08$ | $4162 \pm 50$           | $1.26 \pm 0.02$ | $-0.10 \pm 0.08$ | $5.81 \pm 0.50$      | 120.1 |
| HD 174829  | $10.15 \pm 0.03$  | $4482 \pm 50$           | $1.78 \pm 0.01$ | $-0.40 \pm 0.08$ | $4.41 \pm 0.50$      | 112.2 |
| HD 175740  | $-8.82 \pm 0.05$  | $4973 \pm 50$           | $2.71 \pm 0.01$ | $-0.05 \pm 0.08$ | $3.66 \pm 0.50$      | 264.0 |
| HD 175740  | $-8.82 \pm 0.05$  | $4973 \pm 50$           | $2.71 \pm 0.01$ | $-0.05 \pm 0.08$ | $3.66 \pm 0.50$      | 264.0 |
| HD 175740  | $-8.82 \pm 0.05$  | $4973 \pm 50$           | $2.71 \pm 0.01$ | $-0.05 \pm 0.08$ | $3.66 \pm 0.50$      | 264.0 |
| HD 175740  | $-8.82 \pm 0.05$  | $4973 \pm 50$           | $2.71 \pm 0.01$ | $-0.05 \pm 0.08$ | $3.66 \pm 0.50$      | 264.0 |
| HD 175884  | $-34.39 \pm 0.07$ | $4466 \pm 50$           | $1.73 \pm 0.01$ | $-0.27 \pm 0.08$ | $4.46 \pm 0.50$      | 144.4 |
| HD 178797  | $6.35 \pm 0.05$   | $4406 \pm 50$           | $1.68 \pm 0.01$ | $-0.37 \pm 0.08$ | $4.18 \pm 0.50$      | 77.1  |
| HD 178910  | $-14.28 \pm 0.05$ | $4589 \pm 50$           | $2.39 \pm 0.00$ | $0.14 \pm 0.08$  | $4.26 \pm 0.50$      | 76.9  |
| HD 179396  | $24.80 \pm 0.04$  | $4781 \pm 50$           | $2.39 \pm 0.01$ | $-0.21 \pm 0.08$ | $3.99 \pm 0.50$      | 82.7  |
| HD 180312  | $-21.94 \pm 0.05$ | $4916 \pm 50$           | $2.43 \pm 0.00$ | $-0.44 \pm 0.08$ | $4.05 \pm 0.50$      | 73.5  |
| HD 180312  | $-21.94 \pm 0.05$ | $4916 \pm 50$           | $2.43 \pm 0.00$ | $-0.44 \pm 0.08$ | $4.05 \pm 0.50$      | 73.5  |
| HD 180475  | $-45.90 \pm 0.08$ | $4398 \pm 50$           | $1.52 \pm 0.01$ | $-0.44 \pm 0.08$ | $4.39 \pm 0.50$      | 58.4  |
| HD 180658  | $2.97 \pm 0.06$   | $4802 \pm 50$           | $2.43 \pm 0.01$ | $-0.12 \pm 0.08$ | $3.81 \pm 0.50$      | 72.3  |
| HD 180682  | $30.99 \pm 0.07$  | $4410 \pm 50$           | $1.45 \pm 0.01$ | $-0.51 \pm 0.08$ | $4.88 \pm 0.50$      | 80.1  |
| HD 181022  | $-80.39 \pm 0.16$ | $4045 \pm 50$           | $1.06 \pm 0.01$ | $-0.28 \pm 0.08$ | $5.75 \pm 0.50$      | 108.8 |
| HD 181069  | $9.99 \pm 0.05$   | $4842 \pm 50$           | $2.52 \pm 0.00$ | $-0.05 \pm 0.08$ | $3.53 \pm 0.50$      | 90.0  |
| HD 181097  | $-5.60 \pm 0.08$  | $4520 \pm 50$           | $1.93 \pm 0.01$ | $-0.28 \pm 0.08$ | $4.08 \pm 0.50$      | 69.7  |
| HD 181597  | $-13.06 \pm 0.04$ | $4751 \pm 50$           | $2.31 \pm 0.00$ | $-0.23 \pm 0.08$ | $2.23 \pm 0.50$      | 161.8 |
| HD 181778  | $-22.04 \pm 0.06$ | $4664 \pm 50$           | $2.25 \pm 0.01$ | $-0.19 \pm 0.08$ | $4.23 \pm 0.50$      | 87.6  |
| HD 181880  | $0.56 \pm 0.08$   | $4405 \pm 50$           | $1.70 \pm 0.01$ | $-0.30 \pm 0.08$ | $4.44 \pm 0.50$      | 71.2  |
| HD 182531  | $-7.34 \pm 0.05$  | $4413 \pm 50$           | $1.69 \pm 0.01$ | $-0.24 \pm 0.08$ | $4.39 \pm 0.50$      | 71.4  |
| HD 182692  | $-8.01 \pm 0.05$  | $4965 \pm 50$           | $2.55 \pm 0.00$ | $0.09 \pm 0.08$  | $3.40 \pm 0.50$      | 72.8  |
| HD 182694  | $-0.87 \pm 0.06$  | $5178 \pm 50$           | $2.76 \pm 0.01$ | $-0.12 \pm 0.08$ | $5.12 \pm 0.50$      | 187.2 |
| HD 183124  | $14.96 \pm 0.01$  | $4911 \pm 50$           | $2.50 \pm 0.00$ | $-0.15 \pm 0.08$ | $5.19 \pm 0.50$      | 114.3 |
| HD 185286  | $-13.70 \pm 0.08$ | $4301 \pm 50$           | $1.50 \pm 0.01$ | $-0.14 \pm 0.08$ | $5.16 \pm 0.50$      | 135.6 |
| HD 188537  | $-18.03 \pm 0.15$ | $4961 \pm 50$           | $2.03 \pm 0.01$ | $-0.08 \pm 0.08$ | $10.68 \pm 0.50$     | 67.0  |
| HD 226754  | $18.66 \pm 0.10$  | $4370 \pm 50$           | $1.75 \pm 0.01$ | $0.08 \pm 0.08$  | $4.78 \pm 0.50$      | 62.5  |

**Fe-peak elements:**

XX

**Neutron capture elements:**

XX

Two of the stars in our sample also appear in the Hypatia catalogue of stellar abundances (Hinkel et al. 2014): HD 185351 and HD 175740. The abundances reported here for HD 185351 are consistent within the large errorbars of both surveys with those reported in Hypatia, while for HD 175740 they are not. **Keith - what's going on here? Check this?**

**3.1.2 Red Clump Stars**

Red clump stars, which burn helium in their cores, can be distinguished from hydrogen-shell burning giants asteroseismologically, via their much higher  $g$ -mode period spacings (Bedding et al. 2011). The term ‘red clump’ arises from the fact that such stars can have a very narrow range of luminosities, so that they appear as a clump in the HR diagram (Girardi 2016). This property makes them useful standard candles to which distances can be accurately computed from photometry. Red clump stars have been used to calibrate the *Gaia* survey’s parallaxes at long distances (Davies et al. 2017; Hawkins et al. 2017; Ruiz-Dern et al. 2018). *Gaia* DR2 parallaxes

have a zero-point offset of  $\sim 0.03$  mas (Lindegren et al. 2018), and in particular hierarchical models of the ensemble of *Gaia* clump stars can be used to accurately estimate this and thereby improve the accuracy of *Gaia* distances greater than a few kpc (Hawkins et al., in prep.).

From inspection of the power spectra, HD 181069, HD 183124, HD 182354, HD 182692, and HD 180658 are seen to be red clump stars. A power spectrum of the best example of these, HD 183124, together with an échelle diagram used to estimate its  $g$ -mode period spacing, is shown in Figure 5. While precise characterization of these stars is beyond the scope of this paper, they are ideal candidates for anchoring models of the mass and metallicity dependence of red clump properties for calibrating *Gaia* and other distance measures.

**3.1.3 Long Period Variables****3.2 Main Sequence Stars**

For all the main sequence stars in our sample, we inspected light curves and power spectra to determine their variability class. In the following subsections, we will briefly comment on some of the findings. Since main sequence variables are so diverse, and the relevant scientific questions so varied, we have attempted only a

**Table 4.** Chemical abundances relative to iron for stars in the red giant sample as determined by BACCHUS, without differential line-by-line comparison to Arcturus, as described in Section 2.4, for the elements Ca, Mg, Si, Ti, Al, Ba, and Na. Dashes indicate elements for which abundances could not be reliably computed. The catalogue of abundances for more elements continues in Tables 5 and 6.

| Object     | [Ca/Fe]     | [Mg/Fe]     | [Si/Fe]      | [Ti/Fe]     | [Al/Fe]     | [Ba/Fe]     | [Na/Fe]     |
|------------|-------------|-------------|--------------|-------------|-------------|-------------|-------------|
| BD+36 3564 | 0.21 ± 0.02 | 0.33 ± 0.03 | 0.10 ± 0.03  | 0.34 ± 0.04 | 0.40 ± 0.01 | –           | 0.26 ± 0.08 |
| BD+39 3577 | 0.13 ± 0.02 | 0.22 ± 0.04 | −0.11 ± 0.02 | 0.08 ± 0.04 | 0.21 ± 0.01 | 0.35 ± 0.10 | 0.42 ± 0.00 |
| BD+43 3064 | 0.19 ± 0.04 | 0.21 ± 0.03 | −0.01 ± 0.03 | 0.28 ± 0.04 | 0.36 ± 0.01 | –           | 0.48 ± 0.06 |
| BD+43 3171 | 0.29 ± 0.03 | 0.26 ± 0.06 | −0.00 ± 0.07 | 0.21 ± 0.06 | 0.42 ± 0.01 | 0.33 ± 0.18 | 0.18 ± 0.25 |
| BD+43 3213 | 0.19 ± 0.03 | 0.23 ± 0.07 | −0.18 ± 0.11 | 0.27 ± 0.07 | 0.37 ± 0.04 | –           | 0.62 ± 0.37 |
| BD+48 2955 | 0.22 ± 0.05 | 0.20 ± 0.03 | 0.08 ± 0.04  | 0.30 ± 0.04 | 0.30 ± 0.07 | –           | 0.23 ± 0.14 |
| HD 174020  | 0.33 ± 0.03 | 0.23 ± 0.04 | −0.07 ± 0.06 | 0.29 ± 0.07 | 0.39 ± 0.03 | –           | 0.26 ± 0.33 |
| HD 174829  | 0.16 ± 0.04 | 0.20 ± 0.06 | 0.05 ± 0.05  | 0.19 ± 0.03 | 0.29 ± 0.01 | –           | 0.31 ± 0.04 |
| HD 175740  | 0.12 ± 0.02 | 0.07 ± 0.05 | −0.05 ± 0.02 | 0.14 ± 0.03 | 0.21 ± 0.01 | 0.30 ± 0.07 | 0.34 ± 0.03 |
| HD 175884  | 0.23 ± 0.02 | 0.20 ± 0.03 | −0.01 ± 0.03 | 0.32 ± 0.03 | 0.34 ± 0.01 | –           | 0.46 ± 0.06 |
| HD 178797  | 0.22 ± 0.02 | 0.32 ± 0.03 | 0.06 ± 0.03  | 0.40 ± 0.04 | 0.42 ± 0.01 | 0.39 ± 0.22 | 0.45 ± 0.03 |
| HD 178910  | 0.20 ± 0.03 | 0.20 ± 0.03 | 0.15 ± 0.05  | 0.20 ± 0.03 | 0.39 ± 0.04 | 0.25 ± 0.08 | 0.36 ± 0.98 |
| HD 179396  | 0.09 ± 0.02 | 0.19 ± 0.03 | 0.04 ± 0.05  | 0.13 ± 0.02 | 0.27 ± 0.02 | 0.31 ± 0.03 | 0.28 ± 0.04 |
| HD 179959  | 0.04 ± 0.04 | 0.06 ± 0.04 | 0.01 ± 0.03  | 0.03 ± 0.03 | 0.15 ± 0.02 | –           | 0.38 ± 0.02 |
| HD 180312  | 0.09 ± 0.02 | 0.21 ± 0.03 | 0.06 ± 0.03  | 0.09 ± 0.03 | 0.31 ± 0.01 | 0.37 ± 0.08 | 0.19 ± 0.01 |
| HD 180475  | 0.23 ± 0.03 | 0.33 ± 0.03 | 0.03 ± 0.01  | 0.36 ± 0.04 | 0.41 ± 0.02 | 0.30 ± 0.20 | 0.40 ± 0.03 |
| HD 180658  | 0.15 ± 0.03 | 0.19 ± 0.04 | −0.01 ± 0.03 | 0.21 ± 0.03 | 0.35 ± 0.01 | 0.21 ± 0.09 | 0.39 ± 0.04 |
| HD 180682  | 0.25 ± 0.02 | 0.45 ± 0.03 | 0.13 ± 0.02  | 0.47 ± 0.04 | 0.51 ± 0.05 | 0.19 ± 0.05 | 0.32 ± 0.01 |
| HD 181022  | 0.34 ± 0.02 | 0.34 ± 0.06 | 0.01 ± 0.08  | 0.49 ± 0.06 | –           | 0.31 ± 0.23 | 0.09 ± 0.48 |
| HD 181069  | 0.13 ± 0.02 | 0.17 ± 0.04 | −0.03 ± 0.05 | 0.19 ± 0.03 | 0.28 ± 0.02 | 0.26 ± 0.09 | 0.45 ± 0.06 |
| HD 181097  | 0.25 ± 0.02 | 0.27 ± 0.03 | −0.02 ± 0.03 | 0.35 ± 0.03 | 0.34 ± 0.02 | –           | 0.46 ± 0.06 |
| HD 181597  | 0.19 ± 0.02 | 0.20 ± 0.05 | −0.03 ± 0.02 | 0.27 ± 0.04 | 0.28 ± 0.00 | 0.28 ± 0.05 | 0.42 ± 0.04 |
| HD 181778  | 0.06 ± 0.03 | 0.12 ± 0.03 | 0.00 ± 0.03  | 0.09 ± 0.03 | 0.28 ± 0.02 | 0.47 ± 0.05 | 0.42 ± 0.12 |
| HD 181880  | 0.26 ± 0.02 | 0.30 ± 0.03 | 0.06 ± 0.04  | 0.35 ± 0.03 | 0.42 ± 0.01 | –           | 0.40 ± 0.05 |
| HD 182531  | 0.22 ± 0.02 | 0.21 ± 0.05 | −0.07 ± 0.03 | 0.37 ± 0.04 | 0.39 ± 0.01 | –           | 0.48 ± 0.06 |
| HD 182692  | 0.19 ± 0.03 | 0.18 ± 0.04 | −0.12 ± 0.03 | 0.22 ± 0.04 | 0.35 ± 0.03 | 0.13 ± 0.05 | 0.38 ± 0.12 |
| HD 182694  | 0.10 ± 0.02 | 0.11 ± 0.04 | −0.04 ± 0.02 | 0.05 ± 0.02 | 0.14 ± 0.01 | –           | 0.32 ± 0.01 |
| HD 183124  | 0.17 ± 0.02 | 0.21 ± 0.04 | −0.02 ± 0.04 | 0.19 ± 0.03 | 0.29 ± 0.00 | 0.25 ± 0.05 | 0.35 ± 0.02 |
| HD 185286  | 0.34 ± 0.02 | 0.22 ± 0.04 | −0.04 ± 0.04 | 0.40 ± 0.06 | 0.42 ± 0.02 | –           | 0.55 ± 0.53 |
| HD 185351  | 0.13 ± 0.03 | 0.08 ± 0.05 | −0.08 ± 0.02 | 0.20 ± 0.03 | 0.22 ± 0.00 | 0.21 ± 0.09 | 0.38 ± 0.01 |
| HD 187217  | 0.16 ± 0.04 | 0.28 ± 0.02 | −0.09 ± 0.03 | 0.14 ± 0.04 | 0.32 ± 0.03 | 0.21 ± 0.14 | –           |
| HD 188537  | 0.11 ± 0.04 | 0.27 ± 0.04 | 0.02 ± 0.03  | 0.11 ± 0.04 | 0.25 ± 0.05 | 0.24 ± 0.07 | –           |
| HD 188629  | 0.30 ± 0.03 | 0.21 ± 0.03 | −0.04 ± 0.07 | 0.37 ± 0.07 | 0.41 ± 0.04 | –           | 0.46 ± 0.32 |
| HD 188875  | 0.18 ± 0.04 | 0.22 ± 0.03 | −0.07 ± 0.03 | 0.29 ± 0.04 | 0.33 ± 0.02 | –           | 0.61 ± 1.09 |
| HD 226754  | 0.30 ± 0.02 | 0.31 ± 0.04 | 0.03 ± 0.04  | 0.40 ± 0.06 | 0.48 ± 0.07 | 0.43 ± 0.00 | 0.47 ± 0.18 |

very preliminary study of these stars in this paper, leaving detailed analysis to future work.

### 3.2.1 Classical Pulsators

Our sample includes pulsating stars of spectral type B, A, and F, with their names, properties and variability class listed in Table 1.

Among the hot-star sample are 5  $\delta$  Sct stars, which show p-mode pulsation. These oscillation modes have particularly long lifetimes and stable frequencies, making them precise stellar clocks with periods of  $\sim 2$  hr. These can be used to search for binarity and to obtain orbital parameters from photometry alone (Shibahashi & Kurtz 2012). We used the phase-modulation method of Murphy et al. (2014) to investigate whether any of these  $\delta$  Sct stars were binaries. Any phase modulation is converted into a light arrival-time delay, and if a star is a binary, the time delays of each mode should vary in unison. Nearly 350 PM binaries are known in the full *Kepler* dataset (Murphy et al. 2018).

In four of the five targets we found evidence for binarity, while in the fifth (HD 185397) there was some time-delay variation but there was no agreement between different modes so it is not of binary origin. Of the others, HD 175841 and HD 177781 are likely

very long-period binaries, with periods far exceeding the Kepler datasets of  $\sim 1470$  d. HD 181521 appears to be an eccentric binary with a period of at least 1000 d, but there is only 1 maximum and 1 minimum in the time-delay curve (cf. (Murphy & Shibahashi 2015)), so a unique orbital solution was unobtainable. Finally, HD 186255 is likely a binary with a period of  $\sim 415$  d, but there is a slight aperiodicity in the time delays, likely caused by beating between pulsation modes that are not well-separated in the frequency. That, coupled with the fact that this star is on the failed Module 3 and is therefore missing data every 4th quarter (i.e.  $\sim 93$  of every 372.5 d), makes the binary classification uncertain. If indeed this is a 415-d binary, the time delays are consistent with a companion of minimum mass  $\sim 0.45 M_{\odot}$  in an orbit of moderate eccentricity ( $\sim 0.15$ ).

Several stars have a more complex classification than can be adequately noted in Table 1: HD 189684 is listed as an ellipsoidal variable, but also shows evidence for  $\gamma$  Dor variability. HD 185397 and HD 186255 are listed as  $\gamma$  Dor/ $\delta$  Sct hybrids, but may in fact simply be  $\delta$  Sct variables with nonlinear combination frequencies, and a detailed frequency analysis will be required to distinguish between these possibilities. HD 184788 shows a combination of two rotational modulation signals with base frequencies: 0.0885

**Table 5.** Chemical abundances relative to iron for stars in the red giant sample as determined by BACCHUS, without differential line-by-line comparison to Arcturus, as described in Section 2.4, for the elements Ni, Mn, Co, Eu, La, Zr, and Sr. Dashes indicate elements for which abundances could not be reliably computed. The catalogue of abundances for more elements continues in Table 6.

| Object     | [Ni/Fe]      | [Mn/Fe]      | [Co/Fe]      | [Eu/Fe]      | [La/Fe]      | [Zr/Fe]     | [Sr/Fe]     |
|------------|--------------|--------------|--------------|--------------|--------------|-------------|-------------|
| BD+36 3564 | 0.01 ± 0.04  | 0.08 ± 0.00  | 0.13 ± 0.02  | 0.25 ± 0.03  | −0.02 ± 0.07 | 0.10 ± 0.02 | 0.34 ± 0.12 |
| BD+39 3577 | −0.05 ± 0.03 | −0.03 ± 0.06 | −0.02 ± 0.02 | −0.22 ± 0.04 | −0.25 ± 0.02 | 0.13 ± 0.08 | –           |
| BD+43 3064 | 0.05 ± 0.04  | 0.21 ± 0.02  | 0.13 ± 0.02  | 0.28 ± 0.06  | 0.15 ± 0.02  | 0.32 ± 0.04 | 0.25 ± 0.12 |
| BD+43 3171 | 0.04 ± 0.05  | 0.11 ± 0.09  | 0.14 ± 0.05  | 0.21 ± 0.05  | −0.06 ± 0.11 | 0.36 ± 0.07 | –           |
| BD+43 3213 | 0.06 ± 0.10  | 0.33 ± 0.07  | 0.03 ± 0.05  | 0.06 ± 0.04  | −0.11 ± 0.05 | 0.49 ± 0.11 | 0.64 ± 0.47 |
| BD+48 2955 | 0.05 ± 0.04  | 0.10 ± 0.02  | 0.12 ± 0.04  | 0.28 ± 0.04  | 0.24 ± 0.05  | 0.34 ± 0.05 | –           |
| HD 174020  | 0.05 ± 0.05  | 0.23 ± 0.02  | 0.10 ± 0.04  | 0.11 ± 0.04  | 0.02 ± 0.07  | –           | 0.37 ± 0.89 |
| HD 174829  | −0.06 ± 0.04 | −0.02 ± 0.07 | 0.05 ± 0.02  | 0.15 ± 0.01  | 0.12 ± 0.05  | 0.08 ± 0.03 | –           |
| HD 175740  | 0.03 ± 0.04  | 0.06 ± 0.01  | 0.08 ± 0.02  | 0.09 ± 0.07  | 0.12 ± 0.01  | 0.18 ± 0.02 | –           |
| HD 175884  | 0.04 ± 0.05  | 0.14 ± 0.02  | 0.10 ± 0.02  | 0.19 ± 0.02  | 0.14 ± 0.03  | 0.26 ± 0.02 | –           |
| HD 178797  | 0.05 ± 0.04  | 0.13 ± 0.11  | 0.18 ± 0.03  | 0.26 ± 0.02  | 0.14 ± 0.02  | 0.23 ± 0.03 | –           |
| HD 178910  | 0.28 ± 0.07  | 0.21 ± 0.05  | 0.17 ± 0.03  | −0.02 ± 0.06 | −0.13 ± 0.06 | 0.00 ± 0.03 | –           |
| HD 179396  | −0.02 ± 0.04 | 0.09 ± 0.02  | 0.08 ± 0.03  | −0.05 ± 0.03 | 0.05 ± 0.03  | 0.04 ± 0.02 | –           |
| HD 179959  | −0.08 ± 0.04 | −0.15 ± 0.04 | −0.05 ± 0.02 | 0.16 ± 0.06  | 0.18 ± 0.01  | 0.14 ± 0.07 | –           |
| HD 180312  | 0.02 ± 0.03  | −0.09 ± 0.03 | 0.07 ± 0.01  | 0.34 ± 0.05  | 0.04 ± 0.07  | 0.08 ± 0.02 | –           |
| HD 180475  | 0.03 ± 0.05  | 0.16 ± 0.04  | 0.19 ± 0.02  | 0.19 ± 0.07  | 0.18 ± 0.03  | 0.25 ± 0.03 | –           |
| HD 180658  | 0.03 ± 0.06  | 0.13 ± 0.03  | 0.11 ± 0.02  | –            | 0.04 ± 0.04  | 0.16 ± 0.07 | –           |
| HD 180682  | 0.06 ± 0.04  | −0.03 ± 0.08 | 0.20 ± 0.02  | 0.26 ± 0.03  | −0.03 ± 0.02 | 0.22 ± 0.03 | –           |
| HD 181022  | 0.02 ± 0.07  | 0.05 ± 0.11  | 0.14 ± 0.05  | 0.26 ± 0.03  | −0.03 ± 0.21 | 0.36 ± 0.14 | –           |
| HD 181069  | 0.08 ± 0.05  | 0.16 ± 0.03  | 0.12 ± 0.02  | 0.09 ± 0.03  | 0.02 ± 0.04  | 0.10 ± 0.03 | –           |
| HD 181097  | 0.01 ± 0.04  | 0.02 ± 0.11  | 0.14 ± 0.03  | 0.28 ± 0.04  | 0.17 ± 0.02  | 0.23 ± 0.03 | –           |
| HD 181597  | 0.03 ± 0.04  | 0.14 ± 0.01  | 0.13 ± 0.02  | 0.18 ± 0.03  | 0.13 ± 0.01  | 0.26 ± 0.03 | –           |
| HD 181778  | −0.00 ± 0.05 | 0.13 ± 0.02  | 0.04 ± 0.02  | 0.16 ± 0.01  | 0.08 ± 0.03  | 0.11 ± 0.03 | –           |
| HD 181880  | 0.04 ± 0.04  | 0.10 ± 0.01  | 0.18 ± 0.03  | 0.32 ± 0.04  | 0.17 ± 0.02  | 0.33 ± 0.04 | –           |
| HD 182531  | 0.06 ± 0.04  | 0.17 ± 0.06  | 0.11 ± 0.02  | 0.16 ± 0.05  | 0.15 ± 0.03  | 0.36 ± 0.03 | 0.35 ± 0.14 |
| HD 182692  | 0.03 ± 0.05  | 0.22 ± 0.02  | 0.15 ± 0.02  | 0.01 ± 0.05  | 0.06 ± 0.04  | 0.21 ± 0.03 | –           |
| HD 182694  | −0.07 ± 0.04 | −0.08 ± 0.02 | 0.03 ± 0.03  | 0.16 ± 0.02  | 0.16 ± 0.02  | 0.16 ± 0.04 | –           |
| HD 183124  | −0.00 ± 0.05 | 0.01 ± 0.04  | 0.11 ± 0.02  | 0.17 ± 0.05  | 0.04 ± 0.06  | 0.14 ± 0.04 | –           |
| HD 185286  | 0.12 ± 0.04  | 0.25 ± 0.01  | 0.13 ± 0.03  | 0.18 ± 0.03  | 0.12 ± 0.05  | 0.52 ± 0.05 | 0.30 ± 0.05 |
| HD 185351  | 0.01 ± 0.04  | 0.11 ± 0.02  | 0.15 ± 0.03  | −0.06 ± 0.06 | 0.13 ± 0.03  | 0.29 ± 0.04 | –           |
| HD 187217  | −0.03 ± 0.06 | −0.10 ± 0.10 | −0.03 ± 0.02 | –            | −0.07 ± 0.03 | 0.22 ± 0.04 | –           |
| HD 188537  | 0.05 ± 0.07  | 0.10 ± 0.03  | 0.12 ± 0.04  | 0.20 ± 0.04  | 0.15 ± 0.10  | 0.30 ± 0.04 | –           |
| HD 188629  | 0.10 ± 0.06  | 0.22 ± 0.01  | 0.10 ± 0.02  | 0.15 ± 0.03  | 0.06 ± 0.07  | 0.43 ± 0.01 | 0.34 ± 0.22 |
| HD 188875  | −0.02 ± 0.05 | 0.23 ± 0.02  | 0.09 ± 0.03  | 0.19 ± 0.07  | 0.20 ± 0.05  | 0.30 ± 0.03 | –           |
| HD 226754  | 0.19 ± 0.05  | 0.33 ± 0.03  | 0.23 ± 0.03  | 0.28 ± 0.07  | −0.05 ± 0.07 | 0.34 ± 0.04 | 0.26 ± 0.13 |

and 0.1966c/d. HD 184875 is a  $\gamma$  Dor but also shows evidence for an unknown contaminant. V554 Lyr and V2079 Cyg are both known  $\alpha^2$  CVn variables, which are chemically peculiar stars with strong magnetic fields that show rotational modulation. V2079 Cyg also shows a weak  $\delta$  Sct signal. The detection of rotational modulation in the chemically-peculiar HD 175132 suggests its reclassification as an  $\alpha^2$  CVn variable.

The coherent g-mode pulsations in samples of B, A, and F stars observed by *Kepler* previously showed these stars to be near-rigid rotators (Kurtz et al. 2014; Saio et al. 2015; Triana et al. 2015; Van Reeth et al. 2015, 2016, 2018; Murphy et al. 2016; Schmid & Aerts 2016; Moravveji et al. 2016; Ouazzani et al. 2017; Pápics et al. 2017; Aerts et al. 2017; Szewczuk & Daszyńska-Daszkiewicz 2018). These studies cover about 70 stars so far. However, the vast majority of intermediate-mass stars observed by *Kepler* have yet to be subjected to in-depth asteroseismic analyses and modelling of their interior properties. One of the valuable outputs of our current work includes the reduced light curves of several early-B stars, which were only scarcely targeted in the nominal *Kepler* mission. The few that were monitored did not reveal suitable oscillation frequency patterns to achieve a unique mode identification, which is a requirement to perform asteroseismic modelling. The investigation

of pulsation modes in high-mass stars using high-quality *Kepler* smear data combined with high-precision spectroscopy to identify the modes (Aerts et al. 2010, Chapter 6) is an exciting prospect for asteroseismology, as the interior physics of these stars are largely unknown, yet they play a pivotal role in stellar and galactic evolution. The in-depth asteroseismic analysis of the smear data for the B stars in this work is beyond the scope of the current paper, as it requires additional ground-based follow-up spectroscopy. Such studies will be the subject of future work.

### 3.2.2 Hump and Spike Stars

Several stars in the sample show the ‘hump-and-spike’ morphology in their power spectra (a broad ‘hump’ of low-amplitude oscillations dominated by one high amplitude coherent oscillation toward the high frequency end of this band): HD 186155 (HR 7495), 14 Cyg (HD 185872, HR 7483), HD 189178 (HR 7628), HD 183362 (HR 7403), and HD 184787. Of these, HD 186155 and 14 Cyg are the third- and sixth-brightest stars on silicon, making these the brightest stars that show this effect. The identification for HD 189178 is tentative, as the spectrum also resembles that of an SPB star, while for HD 184787 there is long term variability

**Table 6.** Chemical abundances relative to iron for stars in the red giant sample as determined by BACCHUS, without differential line-by-line comparison to Arcturus, as described in Section 2.4, for the elements Zn, Y, Cr, V, Cu, and Sc. Dashes indicate elements for which abundances could not be reliably computed.

| Object     | [Zn/Fe]          | [Y/Fe]           | [Cr/Fe]          | [V/Fe]           | [Cu/Fe]          | [Sc/Fe]          |
|------------|------------------|------------------|------------------|------------------|------------------|------------------|
| BD+36 3564 | $-0.29 \pm 0.20$ | $-0.27 \pm 0.02$ | $0.23 \pm 0.00$  | $0.15 \pm 0.03$  | $-0.04 \pm 0.06$ | $0.17 \pm 0.02$  |
| BD+39 3577 | $-0.24 \pm 0.71$ | $-0.40 \pm 0.04$ | $0.16 \pm 0.10$  | $0.01 \pm 0.02$  | $-0.21 \pm 0.01$ | $-0.12 \pm 0.05$ |
| BD+43 3064 | –                | $-0.14 \pm 0.05$ | $0.32 \pm 0.01$  | $0.24 \pm 0.03$  | $-0.16 \pm 0.10$ | $0.14 \pm 0.02$  |
| BD+43 3171 | $-0.40 \pm 0.05$ | $-0.31 \pm 0.03$ | $0.29 \pm 0.04$  | $0.12 \pm 0.06$  | $0.02 \pm 0.11$  | $0.14 \pm 0.03$  |
| BD+43 3213 | –                | $-0.06 \pm 0.09$ | $0.39 \pm 0.01$  | $0.08 \pm 0.09$  | $-0.28 \pm 0.11$ | $0.18 \pm 0.04$  |
| BD+48 2955 | –                | $-0.15 \pm 0.05$ | $0.23 \pm 0.04$  | $0.20 \pm 0.03$  | $-0.05 \pm 0.04$ | $0.15 \pm 0.03$  |
| HD 174020  | $-0.48 \pm 1.11$ | $-0.19 \pm 0.06$ | $0.41 \pm 0.06$  | $0.26 \pm 0.03$  | $-0.20 \pm 0.11$ | $0.18 \pm 0.03$  |
| HD 174829  | $-0.12 \pm 0.13$ | $-0.25 \pm 0.06$ | $0.16 \pm 0.02$  | $0.01 \pm 0.02$  | $-0.23 \pm 0.03$ | $0.12 \pm 0.03$  |
| HD 175740  | $-0.16 \pm 0.16$ | $-0.09 \pm 0.07$ | $0.13 \pm 0.04$  | $0.09 \pm 0.02$  | $-0.16 \pm 0.04$ | $0.08 \pm 0.03$  |
| HD 175884  | $-0.15 \pm 0.17$ | $-0.21 \pm 0.07$ | $0.26 \pm 0.04$  | $0.21 \pm 0.02$  | $-0.10 \pm 0.05$ | $0.13 \pm 0.02$  |
| HD 178797  | –                | $-0.08 \pm 0.05$ | $0.26 \pm 0.04$  | $0.19 \pm 0.02$  | $-0.11 \pm 0.04$ | $0.23 \pm 0.03$  |
| HD 178910  | $-0.29 \pm 0.74$ | $-0.18 \pm 0.05$ | $0.29 \pm 0.01$  | $0.17 \pm 0.02$  | $0.21 \pm 0.14$  | $0.14 \pm 0.02$  |
| HD 179396  | $-0.07 \pm 0.15$ | $-0.27 \pm 0.07$ | $0.12 \pm 0.03$  | $0.03 \pm 0.02$  | $-0.16 \pm 0.06$ | $0.10 \pm 0.03$  |
| HD 179959  | $0.05 \pm 1.84$  | $-0.08 \pm 0.06$ | $-0.00 \pm 0.03$ | $-0.11 \pm 0.02$ | $-0.29 \pm 0.05$ | $0.10 \pm 0.05$  |
| HD 180312  | $-0.18 \pm 0.01$ | $-0.23 \pm 0.05$ | $-0.06 \pm 0.06$ | $-0.05 \pm 0.02$ | $-0.15 \pm 0.04$ | $0.15 \pm 0.05$  |
| HD 180475  | $-0.09 \pm 0.11$ | $-0.25 \pm 0.08$ | $0.24 \pm 0.04$  | $0.20 \pm 0.02$  | $-0.00 \pm 0.04$ | $0.21 \pm 0.03$  |
| HD 180658  | $0.16 \pm 1.25$  | $-0.20 \pm 0.01$ | $0.19 \pm 0.04$  | $0.15 \pm 0.02$  | $-0.05 \pm 0.06$ | $0.12 \pm 0.03$  |
| HD 180682  | $-0.23 \pm 0.14$ | $-0.29 \pm 0.04$ | $0.23 \pm 0.03$  | $0.26 \pm 0.02$  | $-0.06 \pm 0.04$ | $0.27 \pm 0.02$  |
| HD 181022  | $-0.27 \pm 0.03$ | $-0.23 \pm 0.02$ | $0.19 \pm 0.08$  | $0.10 \pm 0.08$  | $-0.01 \pm 0.12$ | $0.25 \pm 0.04$  |
| HD 181069  | $-0.02 \pm 0.19$ | $-0.11 \pm 0.08$ | $0.22 \pm 0.03$  | $0.15 \pm 0.02$  | $-0.10 \pm 0.05$ | $0.13 \pm 0.03$  |
| HD 181097  | $-0.08 \pm 0.41$ | $-0.21 \pm 0.03$ | $0.25 \pm 0.02$  | $0.19 \pm 0.03$  | $-0.12 \pm 0.03$ | $0.22 \pm 0.03$  |
| HD 181597  | $-0.14 \pm 0.15$ | $-0.19 \pm 0.08$ | $0.19 \pm 0.05$  | $0.21 \pm 0.02$  | $-0.18 \pm 0.04$ | $0.16 \pm 0.02$  |
| HD 181778  | $-0.03 \pm 0.18$ | $-0.13 \pm 0.04$ | $0.18 \pm 0.02$  | $-0.02 \pm 0.02$ | $-0.25 \pm 0.07$ | $0.05 \pm 0.02$  |
| HD 181880  | $-0.04 \pm 0.22$ | $-0.20 \pm 0.07$ | $0.27 \pm 0.03$  | $0.22 \pm 0.02$  | $-0.07 \pm 0.03$ | $0.23 \pm 0.03$  |
| HD 182531  | $0.03 \pm 0.78$  | $-0.19 \pm 0.07$ | $0.29 \pm 0.05$  | $0.24 \pm 0.03$  | $-0.08 \pm 0.05$ | $0.18 \pm 0.02$  |
| HD 182692  | $-0.24 \pm 1.34$ | $-0.21 \pm 0.10$ | $0.15 \pm 0.07$  | $0.24 \pm 0.02$  | $-0.11 \pm 0.06$ | $0.18 \pm 0.03$  |
| HD 182694  | $-0.24 \pm 0.07$ | $-0.12 \pm 0.05$ | $0.04 \pm 0.03$  | $-0.05 \pm 0.02$ | $-0.26 \pm 0.04$ | $0.09 \pm 0.05$  |
| HD 183124  | $-0.18 \pm 0.17$ | $-0.24 \pm 0.03$ | $0.12 \pm 0.04$  | $0.10 \pm 0.02$  | $-0.22 \pm 0.02$ | $0.10 \pm 0.03$  |
| HD 185286  | –                | $-0.19 \pm 0.08$ | $0.46 \pm 0.01$  | $0.34 \pm 0.02$  | $-0.11 \pm 0.10$ | $0.27 \pm 0.03$  |
| HD 185351  | $-0.31 \pm 0.10$ | $-0.16 \pm 0.05$ | $0.16 \pm 0.04$  | $0.18 \pm 0.02$  | $-0.17 \pm 0.03$ | $0.12 \pm 0.04$  |
| HD 187217  | –                | $-0.37 \pm 0.05$ | $0.28 \pm 0.03$  | $0.11 \pm 0.03$  | $-0.23 \pm 0.02$ | $0.04 \pm 0.05$  |
| HD 188537  | $0.32 \pm 0.78$  | $-0.27 \pm 0.09$ | $0.17 \pm 0.01$  | $0.11 \pm 0.02$  | $-0.17 \pm 0.04$ | $0.06 \pm 0.05$  |
| HD 188629  | –                | $-0.04 \pm 0.10$ | $0.30 \pm 0.06$  | $0.31 \pm 0.04$  | $-0.15 \pm 0.09$ | $0.22 \pm 0.04$  |
| HD 188875  | $0.31 \pm 1.71$  | $-0.04 \pm 0.07$ | $0.33 \pm 0.07$  | $0.18 \pm 0.02$  | $-0.25 \pm 0.07$ | $0.13 \pm 0.03$  |
| HD 226754  | $-0.22 \pm 1.07$ | $-0.33 \pm 0.04$ | $0.38 \pm 0.07$  | $0.45 \pm 0.04$  | $-0.02 \pm 0.07$ | $0.30 \pm 0.04$  |

consistent with contamination. The other hump-and-spike identifications seem secure. [Saio et al. \(2018\)](#) have recently interpreted the hump-and-spike power spectra as evidence for Rossby modes. The F5 star HD 186155, identified by SIMBAD as having a giant spectral type of F5II-III, is shown by its *Gaia* distance to in fact lie on the main sequence. A detailed study of these stars will be presented by Antoci et al., in prep.

Another star with a hump-and-spike spectrum is Boyajian’s Star (KIC 8462852), which shows deep enigmatic dips in brightness ([Boyajian et al. 2016](#)), and has faded both throughout the *Kepler* mission ([Montet & Simon 2016](#)) and in relation to Harvard photographic plates from 1890 onwards ([Schaefer 2016](#)). The dimming, which is chromatic in the manner expected of heterogeneous clouds of circumstellar dust in the line of sight ([Davenport et al. 2018; Bodman et al. 2018](#)), has been ascribed to various causes (reviewed in [Wright 2018](#)), most notably a cloud of exocomets surrounding the star (e.g. [Wyatt et al. 2018](#)). It is unclear whether the explanation of the hump-and-spike phenomenon will shed light on the strange behaviour of Boyajian’s Star, but it may be relevant.

Ashley/Dan/Vichi?


### 3.2.3 Binaries

We detect BD+47 2825 as a new eclipsing binary system, and recover light curves for the previously-known eclipsing binaries HD 186994, V2083 Cyg, and V380 Cyg. The known binary system HD 189684 is newly identified as showing ellipsoidal variability, but does not show evidence of eclipses. We do not attempt detailed analysis of their variability in this paper.

## 4 OPEN SCIENCE

We believe in open science, and have therefore made all substantive products of this research available to the interested reader. All code used to produce smear light curves is available under a GPL v3 license at [github.com/benjaminpope/keplersmear](https://github.com/benjaminpope/keplersmear). All smear light curves, both including the red giant sample studied in detail in Section 3.1, and main sequence stars as discussed in Sections 3.2.1 and 3.2.3, can be downloaded from the Mikulski Archive for Space Telescopes (MAST) as a High-Level Science Product. TRES spectra will be made available from the ExoFOP-TESS website, and all asteroseismic parameters and derived stellar parameters for the red giants in Section 3.1 are provided in an online-only table as Supplementary Material to this paper.





**Figure 4.** The  $[\text{Mg}/\text{Fe}]$ ,  $[\text{Si}/\text{Fe}]$ ,  $[\text{Ca}/\text{Fe}]$ , and  $[\text{Ti}/\text{Fe}]$ ,  $[\text{Al}/\text{Fe}]$ ,  $[\text{Na}/\text{Fe}]$  abundance ratios as a function of iron on the top and bottom rows, respectively for our stars (black circles). We also illustrate a representative sample of Galactic disk stars from Bensby et al. (open red square, 2014) and Adibekyan et al. (open orange triangles, 2012).



**Figure 5.** Power spectrum (left) and échelle diagram (right) of the solar-like oscillations of the red clump star HD 183124. The modes in the power spectrum used for the échelle diagram are highlighted with blue dots. In the échelle diagram we see the characteristic pattern of ‘bumped’ modes from avoided crossings between the comb of  $p$ -modes and  $g$ -mode oscillations with a period spacing of  $\Delta\Pi = 300.1$  s.

All smear light curves in this paper, as well as the  $\text{\LaTeX}$  source code used to produce this document, can be found at [github.com/benjaminpope/smearcampaign](https://github.com/benjaminpope/smearcampaign).

## 5 CONCLUSIONS

The *Kepler* Smear Campaign establishes a legacy sample of 102 very bright stars, with *Kepler* light curves that in almost all cases reveal astrophysically interesting variability. The virtue of these bright

stars is that they can be studied with interferometry, and more easily with spectroscopy than fainter targets, permitting especially detailed characterization. We have therefore obtained detailed abundances of a subset of the red giants in this sample, particularly with a view to determining their provenance in the Galactic thick and thin disk populations. The science that can be done both with this sample and with this method are, however, considerably broader: while we have not attempted it in this paper, a compelling next step is to use interferometric diameter measurements and to further constrain the red

giant parameters, and compare these to the constraints from *Gaia*. Any tension between these measurements will help test and refine the asteroseismic scaling relations, and better models will propagate through to smaller systematic uncertainties in large samples of stars too faint for interferometry. Further improvements will be revealed by the detailed modelling of individual oscillation frequencies in these giants to infer interior structure such as convective overshoot, which is at the time of writing an active topic of research. For the lower-frequency M giants classed as LPVs in this paper, extending the systematics correction and quarter-stitching algorithms to more robustly correct their light curves without removing real signal will allow similar asteroseismic analysis, for a sample of stars that are much less well understood than their higher-frequency counterparts.

The *Kepler* Smear Campaign has another natural extension: while many saturated stars in *K2* have now been observed with ‘halo’ apertures including their unsaturated pixels, many were not, either because they were fainter than the typical  $Kp \lesssim 6.5$  limit, or because in Campaigns 0-3 and 5 no such apertures were selected. There is therefore the potential for a *K2* Smear Campaign to complete the *K2* sample down to fainter magnitudes, complementing the very brightest stars studied with halo photometry.

## ACKNOWLEDGEMENTS

This work was performed in part under contract with the Jet Propulsion Laboratory (JPL) funded by NASA through the Sagan Fellowship Program executed by the NASA Exoplanet Science Institute. B.P. also acknowledges support from Balliol College and the Clarendon Fund. D.H. acknowledges support by the Australian Research Council’s Discovery Projects funding scheme (project number DE140101364) and support by the NASA Grant NNX14AB92G issued through the *Kepler* Participating Scientist Program. DWL acknowledges partial support from the Kepler Extended Mission under NASA Cooperative Agreement NNX13AB58A with the Smithsonian Astrophysical Observatory.

BP acknowledges being on the traditional territory of the Lenape Nations and, today, we recognize that Manhattan continues to be the home to many Algonkian peoples. We thank the Lenape peoples for allowing us to carry out this work on the Lenape original homelands at New York University. BP and TW would like to acknowledge the Gadigal people of the Eora Nation and the Norongerragal and Gweagal peoples of the Tharawal Nation as the traditional owners of the land at the University of Sydney and the Sutherland Shire on which some of this work was carried out, and pay their respects to their knowledge, and their elders past, present and future.

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. This work has in particular made use of the [gaia-kepler.fun](https://github.com/gaia-kepler-fun) crossmatch database created by Megan Bedell.

This research made use of NASA’s Astrophysics Data System; the SIMBAD database, operated at CDS, Strasbourg, France; the IPython package (Pérez & Granger 2007); SciPy (Jones et al. 2001); and Astropy, a community-developed core Python package for Astronomy (Collaboration et al. 2018). Some of the data presented in this paper were obtained from the Mikulski Archive for

Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NNX13AC07G and by other grants and contracts. We acknowledge the support of the Group of Eight universities and the German Academic Exchange Service through the Go8 Australia-Germany Joint Research Co-operation Scheme.

## REFERENCES

- Adibekyan V. Z., Sousa S. G., Santos N. C., Delgado Mena E., González Hernández J. I., Israelian G., Mayor M., Khachatryan G., 2012, *A&A*, **545**, A32
- Aerts C., Christensen-Dalsgaard J., Kurtz D. W., 2010, *Asteroseismology*
- Aerts C., Van Reeth T., Tkachenko A., 2017, *ApJ*, **847**, L7
- Aerts C., et al., 2018, *MNRAS*, **476**, 1234
- Aigrain S., Parviainen H., Pope B. J. S., 2016, *MNRAS*, **459**, 2408
- Alvarez R., Plez B., 1998, *A&A*, **330**, 1109
- Ambikasaran S., Foreman-Mackey D., Greengard L., Hogg D. W., O’Neil M., 2015, *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **38**
- Angus R., Aigrain S., Foreman-Mackey D., McQuillan A., 2015, *MNRAS*, **450**, 1787
- Bailer-Jones C. A. L., Rybizki J., Fousneau M., Mantelet G., Andrae R., 2018, preprint, ([arXiv:1804.10121](https://arxiv.org/abs/1804.10121))
- Beck P. G., et al., 2011, *Science*, **332**, 205
- Beck P. G., et al., 2012, *Nature*, **481**, 55
- Bedding T. R., et al., 2011, *Nature*, **471**, 608
- Bensby T., Feltzing S., Oey M. S., 2014, *A&A*, **562**, A71
- Blanco-Cuaresma S., Soubiran C., Jofré P., Heiter U., 2014, *A&A*, **566**, A98
- Bodman E., Wright J., Boyajian T., Ellis T., 2018, preprint, ([arXiv:1806.08842](https://arxiv.org/abs/1806.08842))
- Boyajian T. S., et al., 2016, *MNRAS*, **457**, 3988
- Brown T. M., Gilliland R. L., Noyes R. W., Ramsey L. W., 1991, *ApJ*, **368**, 599
- Brown T. M., Latham D. W., Everett M. E., Esquerdo G. A., 2011, *AJ*, **142**, 112
- Buchhave L. A., et al., 2012, *Nature*, **486**, 375
- Campante T. L., et al., 2015, *ApJ*, **799**, 170
- Casagrande L., et al., 2014, *MNRAS*, **439**, 2060
- Chaplin W. J., Miglio A., 2013, *ARA&A*, **51**, 353
- Chaplin W. J., et al., 2010, *ApJ*, **713**, L169
- Chiba M., Beers T. C., 2000, *AJ*, **119**, 2843
- Christiansen J. L., et al., 2012, *PASP*, **124**, 1279
- Collaboration T. A., et al., 2018, *The Astronomical Journal*, **156**, 123
- Creevey O. L., et al., 2013, *MNRAS*, **431**, 2419
- Creevey O. L., et al., 2015, *A&A*, **575**, A26
- Davenport J. R. A., et al., 2018, *ApJ*, **853**, 130
- Davies G. R., Miglio A., 2016, *Astronomische Nachrichten*, **337**, 774
- Davies G. R., et al., 2017, *A&A*, **598**, L4
- Edvardsson B., Andersen J., Gustafsson B., Lambert D. L., Nissen P. E., Tomkin J., 1993, *A&A*, **275**, 101
- Farr W. M., et al., 2018, preprint, ([arXiv:1802.09812](https://arxiv.org/abs/1802.09812))
- Feltzing S., Chiba M., 2013, *New Astron. Rev.*, **57**, 80
- Gaia Collaboration et al., 2016, *A&A*, **595**, A1
- Gaia Collaboration Brown A. G. A., Vallenari A., Prusti T., de Bruijne J. H. J., Babusiaux C., Bailer-Jones C. A. L., 2018, preprint, ([arXiv:1804.09365](https://arxiv.org/abs/1804.09365))
- García R. A., et al., 2011, *MNRAS*, **414**, L6
- Gilliland R. L., et al., 2010, *PASP*, **122**, 131
- Girardi L., 2016, *ARA&A*, **54**, 95
- Gustafsson B., Edvardsson B., Eriksson K., Jørgensen U. G., Nordlund Å., Plez B., 2008, *A&A*, **486**, 951
- Harvey J., 1985, in Rolfe E., Battick B., eds, *ESA Special Publication Vol. 235, Future Missions in Solar, Heliospheric & Space Plasma Physics*.

- Hawkins K., et al., 2016a, *A&A*, **592**, A70
- Hawkins K., et al., 2016b, *A&A*, **592**, A70
- Hawkins K., Masseron T., Jofré P., Gilmore G., Elsworth Y., Hekker S., 2016c, *A&A*, **594**, A43
- Hawkins K., Leistedt B., Bovy J., Hogg D. W., 2017, *MNRAS*, **471**, 722
- Heiter U., Jofré P., Gustafsson B., Korn A. J., Soubiran C., Thévenin F., 2015, *A&A*, **582**, A49
- Hinkel N. R., Timmes F. X., Young P. A., Pagano M. D., Turnbull M. C., 2014, *AJ*, **148**, 54
- Hjorringgaard J. G., Silva Aguirre V., White T. R., Huber D., Pope B. J. S., Casagrande L., Justesen A. B., Christensen-Dalsgaard J., 2017, *MNRAS*, **464**, 3713
- Howell S. B., et al., 2014, *PASP*, **126**, 398
- Huber D., et al., 2012, *ApJ*, **760**, 32
- Huber D., et al., 2013, *ApJ*, **767**, 127
- Jenkins J. M., et al., 2010, *ApJ*, **713**, L87
- Jofré P., 2016, *Astronomische Nachrichten*, **337**, 859
- Jofré P., et al., 2014, *A&A*, **564**, A133
- Jofré P., et al., 2015, *A&A*, **582**, A81
- Jofré P., et al., 2017, *A&A*, **601**, A38
- Jones E., Oliphant T., Peterson P., Others 2001, SciPy: Open source scientific tools for Python, <http://www.scipy.org/>
- Kallinger T., et al., 2014, *A&A*, **570**, A41
- Kjeldsen H., Bedding T. R., 1995, *A&A*, **293**, 87
- Koch D. G., et al., 2010, *ApJ*, **713**, L79
- Kolodziejczak J., Caldwell D., 2011, Technical Report 20120003045, Science from Kepler Collateral Data: 150 ksec/year from 13 Million Stars?, <http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20120003045.pdf>. NASA Marshall Space Flight Centre, <http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20120003045.pdf>
- Kurtz D. W., Saio H., Takata M., Shibahashi H., Murphy S. J., Sekii T., 2014, *MNRAS*, **444**, 102
- Lindgren L., et al., 2018, preprint, ([arXiv:1804.09366](https://arxiv.org/abs/1804.09366))
- Lomb N. R., 1976, *Ap&SS*, **39**, 447
- Lund M. N., et al., 2016, preprint, ([arXiv:1612.00436](https://arxiv.org/abs/1612.00436))
- Masseron T., et al., 2014, *A&A*, **571**, A47
- Masseron T., Merle T., Hawkins K., 2016, BACCHUS: Brussels Automatic Code for Characterizing High accuracy Spectra, Astrophysics Source Code Library (ascl:1605.004), [doi:10.20356/C4TG6R](https://doi.org/10.20356/C4TG6R)
- Matteucci F., Recchi S., 2001, *ApJ*, **558**, 351
- Montet B. T., Simon J. D., 2016, *ApJ*, **830**, L39
- Moravveji E., Townsend R. H. D., Aerts C., Mathis S., 2016, *ApJ*, **823**, 130
- Mosser B., et al., 2012a, *A&A*, **540**, A143
- Mosser B., et al., 2012b, *A&A*, **548**, A10
- Mosser B., et al., 2014, *A&A*, **572**, L5
- Murphy S. J., Shibahashi H., 2015, *MNRAS*, **450**, 4475
- Murphy S. J., Bedding T. R., Shibahashi H., Kurtz D. W., Kjeldsen H., 2014, *MNRAS*, **441**, 2515
- Murphy S. J., Fossati L., Bedding T. R., Saio H., Kurtz D. W., Grassitelli L., Wang E. S., 2016, *MNRAS*, **459**, 1201
- Murphy S. J., Moe M., Kurtz D. W., Bedding T. R., Shibahashi H., Boffin H. M. J., 2018, *MNRAS*, **474**, 4322
- Ouazzani R.-M., Salmon S. J. A. J., Antoci V., Bedding T. R., Murphy S. J., Roxburgh I. W., 2017, *MNRAS*, **465**, 2294
- Pápics P. I., et al., 2017, *A&A*, **598**, A74
- Pérez F., Granger B. E., 2007, *Computing in Science and Engineering*, **9**, 21
- Petigura E. A., Marcy G. W., 2012, *PASP*, **124**, 1073
- Pinsonneault M. H., et al., 2014, *ApJS*, **215**, 19
- Plez B., 2012, Turbospectrum: Code for spectral synthesis, Astrophysics Source Code Library (ascl:1205.004)
- Pope B. J. S., et al., 2016, *MNRAS*, **455**, L36
- Ruiz-Dern L., Babusiaux C., Arenou F., Turon C., Lallement R., 2018, *A&A*, **609**, A116
- Saio H., Kurtz D. W., Takata M., Shibahashi H., Murphy S. J., Sekii T., Bedding T. R., 2015, *MNRAS*, **447**, 3264
- Saio H., Kurtz D. W., Murphy S. J., Antoci V. L., Lee U., 2018, *MNRAS*, **474**, 2774
- Scargle J. D., 1982, *ApJ*, **263**, 835
- Schaefer B. E., 2016, *ApJ*, **822**, L34
- Schmid V. S., Aerts C., 2016, *A&A*, **592**, A116
- Shibahashi H., Kurtz D. W., 2012, *MNRAS*, **422**, 738
- Silva Aguirre V., et al., 2013, *ApJ*, **769**, 141
- Silva Aguirre V., et al., 2015, *MNRAS*, **452**, 2127
- Silva Aguirre V., et al., 2016, preprint, ([arXiv:1611.08776](https://arxiv.org/abs/1611.08776))
- Smith J. C., et al., 2012, *PASP*, **124**, 1000
- Stumpe M. C., et al., 2012, *PASP*, **124**, 985
- Szewczuk W., Daszyńska-Daszkiewicz J., 2018, *MNRAS*, **478**, 2243
- Triana S. A., Moravveji E., Pápics P. I., Aerts C., Kawaler S. D., Christensen-Dalsgaard J., 2015, *ApJ*, **810**, 16
- Twicken J. D., Chandrasekaran H., Jenkins J. M., Gunter J. P., Girouard F., Klaus T. C., 2010, in Software and Cyberinfrastructure for Astronomy. p. 77401U, [doi:10.1117/12.856798](https://doi.org/10.1117/12.856798)
- Van Eylen V., Agentoft C., Lundkvist M. S., Kjeldsen H., Owen J. E., Fulton B. J., Petigura E., Snellen I., 2018, *MNRAS*, **479**, 4786
- Van Reeth T., et al., 2015, *ApJS*, **218**, 27
- Van Reeth T., Tkachenko A., Aerts C., 2016, *A&A*, **593**, A120
- Van Reeth T., et al., 2018, preprint, ([arXiv:1806.03586](https://arxiv.org/abs/1806.03586))
- White T. R., et al., 2013, *MNRAS*, **433**, 1262
- White T. R., et al., 2015, in European Physical Journal Web of Conferences. p. 06068, [doi:10.1051/epjconf/201510106068](https://doi.org/10.1051/epjconf/201510106068)
- White T. R., et al., 2017, *MNRAS*, **471**, 2882
- Wright J. T., 2018, *Research Notes of the American Astronomical Society*, **2**, 16
- Wyatt M. C., van Lieshout R., Kennedy G. M., Boyajian T. S., 2018, *MNRAS*, **473**, 5286
- van Leeuwen F., 2007, *A&A*, **474**, 653
- van Saders J. L., Ceillier T., Metcalfe T. S., Silva Aguirre V., Pinsonneault M. H., García R. A., Mathur S., Davies G. R., 2016, *Nature*, **529**, 181

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.