

Report of Sri Lanka Dengue Dashboard

Y.M. Amali P. Rajapaksha

2026-07-08

This dashboard provides an interactive overview of the dengue situation in Sri Lanka using the latest available surveillance data. Built with R using the Quarto Dashboard framework, it presents recent dengue trends, district-level analyses, historical patterns, comparisons with the global dengue situation, and explores the relationship between dengue transmission and key climate variables. The dashboard offers valuable insights for monitoring, understanding, and supporting evidence-based decision-making on dengue dynamics in Sri Lanka.

Dashboard Link

https://amalirajapaksha.github.io/denguedashboard_SriLanka/

Created By

Name : Y.M.A.P. Rajapaksha

Index No : AS2022642

Data Sources

- Local Data up to December 2025 :

Talagala, T. (2026). *denguedatahub: A Tidy Format Datasets of Dengue by Country*. R package version 4.1.1. <https://thiyanagt.github.io/denguedatahubweb/>

- Local Data for 2026 :

Epidemiology Unit. (2026). *Weekly Epidemiological Reports on Dengue Surveillance*. Ministry of Health, Sri Lanka. Available at: <https://www.epid.gov.lk/>

National Dengue Control Unit. (2026). *Weekly Dengue Updates*. Ministry of Health, Sri Lanka. Available at: <https://www.dengue.health.gov.lk/updates/>

- Global Data :

World Health Organization. (2026). *Global Dengue Surveillance Dashboard*. Available at: https://worldhealthorg.shinyapps.io/dengue_global/

- Climate Data :

NASA Langley Research Center. (2026). *NASA POWER Data Access Viewer*. National Aeronautics and Space Administration. Available at: <https://power.larc.nasa.gov/data-access-viewer/>

Data Processing and Availability

- All local data up to December 2025 used in this dashboard are obtained from the weekly epidemiological reports published by the Epidemiology Unit, Ministry of Health, Sri Lanka, through the `denguedatahub` R package, available at <https://github.com/thiyangt/denguedatahub>.
- Data for 2026 are obtained from the weekly epidemiological reports published by the Epidemiology Unit, Ministry of Health, Sri Lanka, and the Weekly Dengue Updates published by the National Dengue Control Unit, Ministry of Health, Sri Lanka. These data are extracted through web scraping using the `convert_slwer_to_tidy()` function available in the `denguedatahub` package. The processed datasets are available in `data/sri_lanka_weekly_dengue_cases_2026.csv` and `data/last_week_comparisson.xlsx`.
- The global data used in this dashboard are obtained from the World Health Organization's [Global Dengue Surveillance Dashboard](#). The corresponding dataset is available in `data/dengue-global-data.xlsx`.
- The climate data used in this study were downloaded from the [NASA POWER Data Access Viewer](#), provided by the National Aeronautics and Space Administration (NASA). The processed weekly climate dataset is available in `data/weekly_climate.csv`.
- All raw data files are available in the `raw-data` directory.
- All R scripts used for data preprocessing are available in the `pre_processing` directory.

Methodology

Here, several visualization techniques have been used for different purposes. To explore new visualization techniques that can be applied for disease surveillance and epidemiological analysis, the [Sri Lanka COVID-19 Dashboard](#) (Talagala & De Alwis, 2021) was referred to. To get an overall idea of the methodologies, see Table 1.

Table 1: Summary of methodologies used in the dashboard

Information Presented	Visualization Technique(s)
Key Information about the Present Dengue Situation in Sri Lanka	KPIs
Distribution of Total Dengue Cases in Sri Lanka in 2026	Choropleth Map
Cumulative Dengue Deaths in Sri Lanka in 2026	Choropleth Map
Anomalous Dengue Behaviour in the Current Period Compared with the Previous Two Years in Sri Lanka	Time Series Plot
Last Week’s Dengue Situation Analysis	Choropleth Maps
District-wise Weekly Dengue Case Analysis Over Time until 2025	Faceted and Individual Time Series Plots, Faceted and Individual Seasonal Plots
Distribution of Dengue Patients by District and Year until 2025	Tree map, Faceted Choropleth Map, Heatmap
Relationship Between Climatic Variables and Dengue Cases	Multi-panel Time Series Plot, Scatterplots
Global Distribution of Dengue Patients/Cases in 2025	Choropleth Map
Sri Lanka Compared with the Top 10 Dengue-Affected Countries Worldwide and Other South-East Asian countries in 2025	Stacked Bar Chart

One of the additional analyses included in this dashboard is the climate-dengue relationship analysis. In addition to time series plots (Figure 6.a) and scatter plots (Figure 6.b) of climatic variables vs. dengue cases, I explored the potential lag effect using two-week and four-week lag scatter plots (Figure 6.c , Figure 6.d). Dengue transmission does not respond immediately to climatic changes such as rainfall and humidity. Instead, favourable climatic conditions may enhance mosquito breeding and contribute to increased dengue cases after a certain time lag. Based on this idea, I explored the lag relationship between climatic factors and dengue cases. Furthermore, to obtain a comprehensive understanding of the relationship between climatic variables and dengue cases, the Detrended Cross-Correlation Coefficient (ρ_{DCCA}) was calculated across different time scales (Figueredo et al., 2023). This method is used to study the relationship between multiple time series that change over time, even when the data are not stationary (Zebende et al., 2018). The calculated ρ_{DCCA} values were plotted over different time scales to investigate how climatic variables influence dengue occurrence across short, medium and long-term periods.

In this dashboard, several colour palettes were used ensuring that the visualizations are colour-blind friendly. These include the Viridis palette from the viridis package (Garnier et al., 2024), the Paired palette from the ColorBrewer collection (Brewer, 2003), and the Magma palette from the Viridis colour map family (van der Walt & Smith, 2015). In addition, some colour

themes were manually defined according to the context of the visualizations.

In this dashboard, multiple visualization techniques have sometimes been used for the same data. Although the data are the same, each visualization provides different insights that can be directly obtained from the data. Therefore, using multiple plots for the same data helps explore different aspects of the actual situation more comprehensively.

Results

Table 2: Description of Dashboard Tabs

Tab No.	Description
Tab 1	This provides the latest information about the dengue epidemic in Sri Lanka. It gives a brief overview of the current dengue situation in Sri Lanka.
Tab 2	This compares the dengue situation in the 26th week of this year with the same week of the previous year. This is the latest data we have.
Tab 3	There are two sub-tabs. The first contains faceted time series plots of weekly dengue cases from 2006 to 2025 for each district separately. The second tab allows users to explore small changes in dengue cases for each district individually using dygraphs.
Tab 4	There are two sub-tabs. The first contains faceted seasonal plots of weekly dengue cases from 2006 to 2025 for each district separately. The second tab allows users to explore small changes in seasonal patterns for each district individually using Plotly graphs.
Tab 5	There are three sub-tabs. The first contains a treemap showing the distribution of total dengue patients in 2025. The second and third sub-tabs contain the distribution of dengue patients over the last 20 years by district and year.
Tab 6	This section contains five sub-tabs and provides a comprehensive analysis of the relationship between climatic variables and dengue cases. The first four sub-tabs present time series plots and scatter plots to explore temporal patterns and associations between climatic variables and dengue incidence. The fifth sub-tab presents a map displaying the calculated raw DCCA values across different time scales.
Tab 7	This section has three sub-tabs. The first shows the distribution of dengue patients worldwide in 2025. The second and third sub-tabs compare the dengue situation in Sri Lanka with the global and South-East Asia region levels, respectively.
Tab 8	This contains detailed information about the dashboard, data sources, references, and other related information.

The dashboard is organized into 8 navigation tabs, including Overview, Last Week Dengue Situation Analysis, District-wise Time Series Trends, District-wise Seasonal Plots, Dengue Patients

Distribution, Climate-Dengue Relationship Analysis, Global Comparison, and About. Each tab contains interactive visualizations and analytical outputs. Table 2 provides a description of each tab and summarizes its key content.




Figure 1: Screenshot of Tab 1 (Overview)

According to Figure 1, a rapid increase in dengue cases can be observed in 2026 from around the 21st week compared with 2024 and 2025. Public sources suggest that this increase may be associated with enhanced *Aedes* mosquito breeding due to favourable climatic conditions, post-flood environmental changes following Cyclone Ditwah, urbanization-related breeding habitats, and ongoing dengue virus transmission dynamics.




Figure 2: Screenshot of Tab 2 (Last Week Dengue Situation Analysis)

Figure 2 also proves the severity of the situation Sri Lanka faces today. Approximately, the

number of cases reported in the 26th week this year is nearly twice that of the same week last year.




Figure 3: Screenshot of Tab 3 (District-wise Time Series Trends)

Figure 3.a shows that many districts experienced notable peaks in dengue cases during 2017, 2019, and 2023. Among these, the 2017 outbreak was the most severe, with consistently high case counts across nearly all districts. According to published reports, Sri Lanka recorded its largest dengue outbreak in 2017, with 186,101 suspected cases and 440 deaths nationwide. Preliminary laboratory investigations identified Dengue virus serotype 2 (DENV-2) as the predominant circulating strain associated with this outbreak. Heavy monsoon rains and flooding also contributed to the outbreak by creating favourable breeding conditions for *Aedes* mosquitoes through standing water and inadequate waste management, leading to increased transmission, particularly in urban and suburban areas.




Figure 4: Screenshot of Tab 4 (District-wise Seasonal Plots)

Dengue transmission in Sri Lanka shows a clear seasonal pattern, with two annual peaks corresponding to the Southwest (May to September) and Northeast (October to January) monsoon periods. However, this pattern is not observed uniformly across all districts.




Figure 5: Screenshot of Tab 5 (Dengue Patients Distribution)

The Southwest monsoon mainly affects districts such as Colombo, Gampaha, Kalutara, Galle, Matara, Hambantota, Kandy, Matale, Nuwara Eliya, Ratnapura, Kegalle, Puttalam, and Kurunegala. In contrast, districts such as Trincomalee, Batticaloa, Ampara, Jaffna, Kilinochchi, Mannar, Mullaitivu, Vavuniya, Badulla, and Monaragala are not significantly affected by the Southwest monsoon. This difference is clearly visible in Figure 4.a. While the Southwest monsoon affected districts show annual peaks in dengue cases during the period from May to September, districts such as Trincomalee, Batticaloa, Ampara, Jaffna, Kilinochchi, Mannar, Vavuniya, Badulla, and Monaragala do not exhibit a similar seasonal peak during this period.

Figure 5.c shows that the Western Province, particularly the districts of Colombo, Gampaha, and Kalutara, has consistently reported the highest dengue burden in the country. As shown in Figure 5.b, these districts are geographically very small. However, the Western Province serves as Sri Lanka's administrative, commercial, and economic hub, attracting a large number of people to live and work in these areas.

Consequently, the population density in these districts is exceptionally high. In addition, rapid urbanization, extensive construction activities, environmental conditions favourable for mosquito breeding, and increased human mobility have further contributed to the high transmission of dengue in the region. Figure 5.a clearly illustrates the hierarchical distribution of dengue cases across all districts in 2025 using a treemap.



Figure 6: Screenshot of Tab 6 (Climate-Dengue Relationship Analysis)

Figure 6.a shows a clearly visible relationship between dengue cases and climatic variables, particularly relative humidity, temperature, and rainfall. According to the figure, dengue cases increase when relative humidity is high and temperature is low. During periods with high dengue cases, rainfall is also higher than during other periods.

When Figures 6.b, 6.c, and 6.d are examined, the above-mentioned lag effect can be clearly seen. Among them, the scatter plots in Figure 6.b at a lag of 4 weeks show that the points are more closely clustered and move in the same direction. This indicates a positive relationship between relative humidity and dengue cases and a negative relationship between temperature and dengue cases. However, the relationship between rainfall and dengue cases is not as strong.

However, scatter plots and Pearson correlation coefficients are not very suitable methods for quantifying the strength and direction of relationships in time series data. Therefore, DCCA

was used for this purpose. According to the DCCA results shown in Figure 6.e, there is no strong relationship in the short run. However, in the long run, there is a strong positive relationship between relative humidity and dengue cases, while temperature has a strong negative relationship with dengue cases. The relationship between rainfall and dengue cases is moderate in the long run.




Figure 7: Screenshot of Tab 7 (Global Comparison)

These findings may be highly useful for policy-making purposes and for promoting personal precautionary activities. For example, when a period of high relative humidity is observed for a certain duration, people can be made aware that there may be an increased risk of dengue transmission. Based on such early indications, health authorities can organize public awareness programs, strengthen healthcare facilities, and ensure the availability of necessary medical resources. At the individual level, people can take preventive measures such as cleaning their surroundings to eliminate mosquito breeding sites, using mosquito repellents, and adopting other protective practices to reduce the risk of dengue infection.

Figure 7.a highlights the geographical variation of dengue cases worldwide in 2025, while Figure 7.b compares the global situation with that of Sri Lanka. According to the figure, although Sri Lanka has a lower number of dengue cases, its mortality is comparatively higher, even exceeding that of some of the top 10 countries. However, when considering the South-East Asian countries, Sri Lanka performs better than several regional peers, suggesting comparatively lower mortality. This is shown in Figure 7.c.




Figure 8: Screenshot of Tab 8 (About)

References

1. Talagala, T. S; De Alwis R. S (2021, December 6). Covid-19 Dashboard Visualisation: Sri Lanka <https://thiyangt.github.io/coviddashboard/>
2. Figueredo, M. B., Monteiro, R. L. S., do Nascimento Silva, A., de Araújo Fontoura, J. R., da Silva, A. R., & Alves, C. A. P. (2023). Analysis of the correlation between climatic variables and Dengue cases in the city of Alagoinhas/BA. *Scientific Reports*, 13(1), 7512.
3. Duque-Lee, C. D., Yu, A. K. D., Ytienza, S. I. E., Yu, A. M. D., Yu, V. C. S., Wangkay, K. A. K., ... & Ysip, G. B. (2020). Correlation between incidence of dengue and climatic factors in the Philippines: An ecological study. *Health Sciences Journal*, 9(2), 1-1.
4. Zebende, G. F., Brito, A. A., Silva Filho, A. M., & Castro, A. P. (2018). DCCA applied between air temperature and relative humidity: An hour/hour view. *Physica A: Statistical Mechanics and its Applications*, 494, 17-26.
5. Brewer, C. A. (2003). *ColorBrewer.org: An online tool for selecting colour schemes for maps*.
6. Garnier, S., Ross, N., Rudis, B., Camargo, A. P., Sciaini, M., & Scherer, C. (2024). *viridis: Colorblind-Friendly Color Maps for R*. R package version 0.6.5.
7. van der Walt, S., & Smith, N. (2015). *mpl-colormaps: Viridis and other perceptually uniform colour maps*.