

Intermediate Anabolic Cycle Blueprint

Multi-Compound Protocol — Advanced Management & Periodisation

14–16 Weeks	3–4 Compounds	Advanced Management	68+ pg Pages
-----------------------	-------------------------	-------------------------------	------------------------

■ **MEDICAL DISCLAIMER:** This guide is for educational and harm-reduction purposes only. The compounds discussed are prescription medications in many countries. Always consult a qualified physician before using any pharmacological substance. Tiger Fitness Pro does not encourage illegal activity.

Intermediate Anabolic Cycle Blueprint

After completing at least 1–2 beginner cycles, intermediate users can introduce a second compound. The most effective and safest intermediate stack is Testosterone + Nandrolone (Deca) with optional oral kickstart. This guide covers the full protocol, estrogen management, and prolactin control.

Intermediate Cycle Options

Stack	Weeks	Compounds	Best For	Complexity
Test + Deca	14–16	Test-E 500 + Deca 400	Mass & strength	Moderate
Test + EQ	16	Test-E 500 + EQ 400	Lean mass, vascularity	Moderate
Test + NPP	10–12	Test-P 400 + NPP 300	Mass, faster recovery	Moderate-High
Test + Deca + Dbol	16	Test + Deca + Dbol 4wks	Maximum mass	High
Test + Primo	14–16	Test-E 400 + Primo 400	Lean mass, no water	Low (mild compounds)

Recommended Protocol — Test + Deca

Compound	Dose	Frequency	Weeks	Notes
Testosterone Enanthate	500 mg	2x/week	1–16	Backbone of cycle
Nandrolone Decanoate (Deca)	400 mg	1x/week	1–14	Add weeks 1–14 only
Dianabol (kickstart)	30–40 mg/day	Daily split	1–4	Oral kickstart
Arimidex (AI)	0.5 mg	EOD	1–16	Estrogen control
Cabergoline	0.25 mg	2x/week	1–14	Prolactin control (Deca)
HCG	500 IU	2x/week	1–14	Testicular maintenance
TUDCA	500 mg	Daily	1–4 (Dbol weeks)	Liver support

Prolactin Management — Essential with Deca

Nandrolone (Deca) is a progestin and elevates prolactin in some users, causing reduced libido and erectile dysfunction ('Deca dick'). Cabergoline is a dopamine agonist that suppresses prolactin.

Compound	Dose	Timing	Purpose
Cabergoline	0.25 mg	2x/week (e.g. Mon/Thu)	Primary prolactin control
Pramipexole (alt)	0.25 mg	Daily	Alternative if Caber unavailable
Vitamin B6 (P5P)	200 mg	Daily	Mild prolactin support

Bloodwork Schedule

Timepoint	Tests	Target
Pre-cycle	Full panel + prolactin baseline	All normal
Week 4	E2, prolactin, liver (Dbol), CBC	E2 20–40, prolactin <15 ng/mL
Week 8	Total T, E2, prolactin, hematocrit	Hct <52%
Week 14 (drop Deca)	Full panel	Prepare for PCT
4 weeks post-PCT	T, LH, FSH, E2	Recovery assessment

Advanced Training Periodisation

Phase	Weeks	Volume	Intensity	Recovery
Strength phase	1–6	12–16 sets/muscle	85–90% 1RM	48 hrs between sessions
Hypertrophy phase	7–12	18–22 sets/muscle	70–80% 1RM	24–36 hrs
Peak phase	13–16	Reduce 20%	90%+ 1RM	72 hrs — quality over quantity