

Advanced Cutting Cycle

12-Week Protocol for Elite Fat Loss

12 Weeks	3 Compounds	500 kcal Deficit	48+ pg Pages
--------------------	-----------------------	----------------------------	------------------------

TIGER FITNESS PRO

■ **MEDICAL DISCLAIMER:** This guide is for educational and harm-reduction purposes only. The compounds discussed are prescription medications in many countries. Always consult a qualified physician before using any pharmacological substance. Tiger Fitness Pro does not encourage illegal activity.

Advanced Cutting Cycle — 12-Week Protocol

This advanced protocol combines testosterone, anavar, and clenbuterol in a calculated 12-week cutting cycle designed for experienced athletes with 3+ cycles of prior experience. Always obtain bloodwork before, during (week 6), and after the cycle.

Pre-Cycle Requirements

Before starting, confirm: baseline testosterone levels, LH/FSH, estradiol, hematocrit, lipid panel, liver enzymes (AST/ALT), and PSA (males 35+). Body fat should be ≤18% for optimal results. Minimum 16 weeks since last cycle.

Compound Stack & Dosing

Compound	Weeks	Dose	Frequency	Notes
Testosterone Propionate	1–12	100 mg	EOD	Maintains muscle, libido
Anavar (Oxandrolone)	1–8	50 mg/day	Daily (split AM/PM)	Dry, lean gains
Clenbuterol	1–12	40→120 mcg	2-week on/off	Beta-2 agonist fat burner
Aromasin (AI)	1–12	12.5 mg	E3D or as needed	Controls estrogen
HCG	1–10	500 IU	2x/week	Testicular function

Week-by-Week Breakdown

Phase	Weeks	Key Focus	Clen Dose	Expected Result
Ramp-Up	1–2	Adjust to compounds	40→60 mcg	Body adapts, slight fat loss
Active Cut	3–6	Max deficit, high intensity	80 mcg	1–1.5 kg fat/week
Peak Cutting	7–10	Carb cycling, posing	100 mcg	Vascularity emerging
Finishing	11–12	Taper, water management	120→80 mcg	Final conditioning
PCT Prep	13	Last HCG shot, stop all	Off	Prepare for SERM PCT

Nutrition Protocol — Cutting Phase

Macro	Intake	Timing	Food Sources
Protein	2.5 g/kg BW	Every 3 hrs	Chicken, eggs, whey, cottage cheese
Carbohydrates	1.5–2 g/kg (training)	Pre/post workout only	Rice, oats, sweet potato
Fats	0.8 g/kg	With non-workout meals	Eggs, nuts, olive oil, fish
Calories	BMR × 1.4 – 500	Deficit maintained	Track with MyFitnessPal
Water	4–5 L/day	Throughout	Electrolytes essential on Clen

Training Split

Day	Focus	Volume	Intensity	Notes
Mon	Chest + Triceps	16 sets	75–80% 1RM	Add supersets
Tue	Back + Biceps	18 sets	70–80% 1RM	Pull-ups essential
Wed	Legs	20 sets	60–75% 1RM	Squats, RDL, leg press
Thu	Shoulders + Core	14 sets	70% 1RM	Lateral raises focus
Fri	Arms + Weak points	12 sets	65–75% 1RM	High reps
Sat	LISS Cardio	45–60 min	60% max HR	Fasted preferred
Sun	Rest/Stretching	—	—	Recovery priority

Side Effect Management

Side Effect	Cause	Prevention	Treatment
Elevated estrogen	Test aromatization	Aromasin E3D	Increase AI dose
Insomnia	Clenbuterol stimulation	Take Clen before noon	Melatonin 3–5 mg
Muscle cramps	Clen depletes taurine	Taurine 3–5 g/day	Potassium + magnesium
Hair thinning	DHT elevation	Nizoral shampoo	Finasteride (optional)
Liver stress	Oral Anavar	TUDCA 500 mg/day	Milk thistle, NAC

Post-Cycle Therapy (PCT)

Begin PCT 3 days after last Testosterone Propionate injection. The goal is HPTA restoration within 4–6 weeks.

SERM	Weeks	Dose	Notes
Nolvadex (Tamoxifen)	1–2	40 mg/day	Kickstart LH/FSH
Nolvadex	3–4	20 mg/day	Taper down
Clomid (optional)	1–2	50 mg/day	Stack with Nolvadex if needed
Zinc + D3	PCT + 4 weeks	30 mg / 5000 IU	Natural testosterone support

Bloodwork Checkpoints

Timepoint	Tests Required	Target Values
Pre-cycle	Total T, E2, LH, FSH, CBC, lipids, liver	All within normal range
Week 6	Total T, E2, RBC, hematocrit, liver	Hct <52%, E2 20–40 pg/mL
4 wks post-PCT	Total T, LH, FSH, E2	T recovering toward baseline
8 wks post-PCT	Full panel	All normalized before next cycle