

COMPLETE PEPTIDE PROTOCOL BIBLE

Ch 1: Peptide Fundamentals

Peptides are short chains of amino acids (2–50 AAs) that act as signalling molecules. Unlike anabolic steroids, most peptides work by stimulating the body's own hormone production rather than replacing it. This makes them a unique category of performance compounds.

Peptide Category	Examples	Primary Action	Main Application
GHRPs (GH-releasing peptides)	GHRP-2, GHRP-6, Ipamorelin	Stimulate GH pulse from pituitary	Mass gain, fat loss, recovery
GHRHs (GH-releasing hormones)	CJC-1295, Sermorelin, Tesamorelin	Extend and amplify GH pulse	Synergy with GHRPs
Repair peptides	BPC-157, TB-500	Tissue healing, angiogenesis	Injury recovery, gut repair
Metabolic peptides	AOD-9604, Fragment 176-191	Fat cell lipolysis stimulation	Fat loss without muscle loss
Melanocortin peptides	PT-141, Melanotan II	CNS and melanocortin activation	Libido enhancement, tanning
Cognitive peptides	Semax, Selank	Neuroprotection, BDNF increase	Focus, anxiety, recovery

Ch 2: GH-Releasing Peptide Profiles

Peptide	Effective Dose	Frequency	Half-life	Key Side Effects	Best Use Case
GHRP-6	100–200 mcg	3x/day	< 30 min	Hunger spike, cortisol rise	Mass gain; appetite stimulation
GHRP-2	100–200 mcg	3x/day	< 30 min	Cortisol, prolactin elevation	GH pulse without hunger
Ipamorelin	200–300 mcg	2–3x/day	2 hours	Minimal	Cleanest GHRP; best for beginners
Hexarelin	200 mcg	2x/day	< 30 min	Desensitisation, cardiac effects	Short cycles only; powerful
CJC-1295 (DAC)	2 mg/week	1–2x/week	8 days	Blunted natural GH pulses	Steady GH elevation; anti-aging
CJC-1295 (no DAC)	100 mcg	3x/day with GHRP	30 min	Minimal	Best synergy with GHRPs
Sermorelin	200–500 mcg	Before sleep	12 min	Injection site reaction	Anti-aging; sleep GH optimisation

Ch 3: BPC-157 Complete Guide

Body Protection Compound 157 (BPC-157) is a 15-amino acid peptide derived from a protein in gastric juice. It shows extraordinary tissue-healing properties and is the most researched peptide for injury recovery.

Condition	Protocol	Route	Duration	Expected Outcome
Tendon/ligament injury	250 mcg 2x/day near injury site	Subcutaneous or oral	4–8 weeks	Accelerated healing; reduced pain
Muscle tear	250–500 mcg/day near tear	Subcutaneous	2–4 weeks	Faster fibre repair; less scar tissue
GI issues (IBS, leaky gut)	250 mcg 2x/day	Oral (capsule in water)	8–12 weeks	Gut lining repair; reduced inflammation
General recovery (on cycle)	200 mcg/day	Subcutaneous any site	Duration of cycle	Prevents joint damage; recovery speed

Ch 4: Peptide Stacking Protocols

Goal	Peptide Stack	Daily Dose	Timing	Duration
Maximum GH pulse	Ipamorelin + CJC-1295 no DAC	300 + 100 mcg each	Fasted AM + pre-sleep	12 weeks on, 4 off
Injury recovery	BPC-157 + TB-500	250 mcg + 2mg/wk	BPC 2x/day; TB weekly	Until healed + 2 wks
Body recomposition	AOD-9604 + Ipamorelin	300 + 300 mcg	AM fasted + pre-sleep	16 weeks
Anti-aging protocol	Sermorelin + BPC-157	500 + 250 mcg	Pre-sleep + AM	Ongoing (12 on/4 off)
Bulking assist	GHRP-6 + CJC no DAC + BPC-157	200+100+250 mcg	3x/day (GHRP/CJC); BPC 2x	Duration of cycle

Ch 5: Reconstitution, Storage & Safety

Peptide Handling Protocol

- Reconstitute with bacteriostatic water (BacWater) — NOT regular sterile water
- Standard dilution: 1 mL BacWater per 2 mg peptide = 2000 mcg/mL concentration
- Inject BacWater slowly down the SIDE of the vial — never directly on the powder
- Gently swirl to mix — never shake (shaking denatures the peptide)
- Unreconstituted: store in freezer up to 2 years; Reconstituted: refrigerator, use within 4 weeks
- Syringes: 29–31G insulin syringes; 0.5–1 mL volume
- Always allow peptide solution to reach room temperature before injection

Red Flag When Buying Peptides	Why It Matters
No HPLC purity certificate from third party	Cannot verify actual peptide content or purity
Price far below market average	Likely diluted, counterfeit, or improperly stored
No bacteriostatic water provided or sold	Vendor does not understand sterile reconstitution requirements
No labelling of amino acid sequence and percentage purity	Cannot confirm correct peptide was synthesised
Vendor cannot answer basic reconstitution questions	Indicates lack of genuine product knowledge

Ch 6: Peptide Dosing Reference Card

Peptide	Storage	Reconstitution	Injection Site	Injection Time
BPC-157	Freezer (unrecon); Fridge (recon)	250 mcg/0.125 mL if 2000 mcg/mL	Near injury or any subcut site	AM fasted + evening
TB-500	Freezer (unrecon); Fridge (recon)	2 mg/1 mL	Any subcutaneous site (abdomen easy)	Once weekly (or split 2x)
Ipamorelin	Freezer (unrecon); Fridge (recon)	300 mcg/0.15 mL if 2000 mcg/mL	Abdomen subcutaneous	AM fasted; pre-sleep
CJC-1295 no DAC	Freezer (unrecon); Fridge (recon)	100 mcg/0.05 mL if 2000 mcg/mL	Abdomen subcutaneous	Same time as Ipamorelin
AOD-9604	Freezer (unrecon); Fridge (recon)	300 mcg/0.15 mL if 2000 mcg/mL	Abdomen subcutaneous	AM fasted only