

ULTIMATE FAT LOSS MASTERCLASS

Ch 1: The Fat Loss Equation

Fat loss occurs when energy expenditure exceeds energy intake — a caloric deficit. All diets that produce fat loss do so by creating this deficit. The challenge is maintaining the deficit while preserving muscle, managing hunger, and sustaining consistency over weeks and months.

Deficit Size	Weekly Fat Loss	Risk Level	Best For
250 kcal/day	~250g/week	Minimal; highly sustainable	Slow recomp; advanced trainees; long-term dieting
500 kcal/day	~500g/week	Low; optimal for most people	Standard cut; best balance of speed and muscle preservation
750 kcal/day	~750g/week	Moderate; some muscle loss risk	Aggressive cut with very high protein (2.5g/kg+)
1000+ kcal/day	1 kg+/week	High; muscle loss and hormonal disruption likely	Short-term only; obese individuals with medical supervision

Ch 2: Macros & Calorie Calculation

Calculate your TDEE (Total Daily Energy Expenditure) using the Mifflin-St Jeor equation multiplied by your activity factor. Subtract 500 kcal for standard fat loss. Adjust every 3–4 weeks as body weight decreases.

Activity Level	Multiplier	Example (70 kg person, BMR ~1650 kcal)
Sedentary (desk job, no exercise)	x 1.2	$1650 \times 1.2 = 1,980$ kcal TDEE
Lightly active (1–3 workouts/week)	x 1.375	$1650 \times 1.375 = 2,269$ kcal TDEE
Moderately active (4–5 sessions/week)	x 1.55	$1650 \times 1.55 = 2,558$ kcal TDEE
Very active (hard training + physical job)	x 1.725	$1650 \times 1.725 = 2,846$ kcal TDEE

Macro	Cutting Target	Why It Matters	Indian Sources
Protein	2.2–2.5 g/kg BW	Muscle preservation; most satiating macro	Paneer, dal, eggs, chicken, soya chunks, curd
Fat	0.7–1.0 g/kg BW	Hormones; satiety; fat-soluble vitamins	Ghee, nuts, olive oil, egg yolks, fish oil
Carbs	Fill remaining calories	Training performance; glycogen; mood	Rice, oats, roti, sweet potato, banana, fruit

Ch 3: Evidence-Based Strategies

Strategy	Mechanism	How to Implement	Evidence Rating
High protein diet	Thermic effect 25–30%; strongest satiety; muscle preservation	2g+/kg every single day; distribute across 4–5 meals	5/5
Resistance training	Preserves muscle during deficit; raises RMR long-term	3–4x/week compound movements; progressive overload	5/5
Cardio (LISS + HIIT combo)	Burns additional calories; improves cardiovascular health	LISS 4x45 min + HIIT 2x20 min per week	4/5
Sleep 7–9 hours	Controls ghrelin and leptin; reduces cravings by 30%+	Fixed sleep/wake schedule; dark, cool room	4/5
Calorie tracking (6 weeks min)	Eliminates unconscious caloric overconsumption	HealthifyMe or MyFitnessPal; weigh food for 1 week	4/5
High-volume low-calorie foods	Low calorie density allows larger meal volume	Start meals with salad/soup; eat cucumber and watermelon	3/5
Intermittent fasting (16:8)	Reduces eating window; helps some with caloric adherence	Not magic; works only if it reduces total intake	3/5

Ch 4: Breaking Plateaus

Plateau-Breaking Protocol (Apply in Order)

- 1. Recalculate TDEE — it drops as you lose weight; recalculate every 5 kg of weight loss
- 2. Refeed day — 1 day at maintenance calories (high carb) weekly to restore leptin
- 3. Diet break — 1–2 weeks at maintenance; alleviates hormonal adaptation; resume refreshed
- 4. Increase NEAT — 2000 extra steps per day; stand more; take stairs; park far from entrance
- 5. Change cardio stimulus — if doing LISS only, add HIIT; if doing HIIT only, add fasted LISS
- 6. Re-audit food logging — weigh all food for 7 days; eyeballing creates 20–40% underestimation
- 7. Check hormones — if plateau persists over 3 weeks: thyroid, cortisol, insulin sensitivity

Ch 5: Indian Cutting Meal Plans

Indian Vegetarian Cutting Plan (1800 kcal, 130g protein)

Meal	Food	kcal	Protein
Breakfast 7 AM	Moong dal chilla x3 + green chutney + chai (no sugar)	340	20g
Mid-morning 10 AM	100g soya chunks (boiled, spiced) + cucumber + lemon	200	26g
Lunch 1 PM	150g paneer bhurji + 1 roti + mixed sabzi + salad	420	28g
Snack 4 PM	1 cup low-fat curd + 10 almonds	200	12g
Post-workout 6 PM	Whey shake 30g + 1 banana	280	26g
Dinner 8 PM	100g tofu stir-fry (Indian spices) + 1 roti + dal soup	360	24g
TOTAL		1,800 kcal	136g

Indian Non-Vegetarian Cutting Plan (1800 kcal, 160g protein)

Meal	Food	kcal	Protein
Breakfast 7 AM	4 egg whites + 1 whole egg bhurji + 1 roti	310	28g
Mid-morning 10 AM	30g whey + 100g curd	210	32g
Lunch 1 PM	200g grilled chicken + 3/4 cup brown rice + sabzi	460	45g
Snack 4 PM	Boiled eggs x2 + cucumber + lemon	160	14g
Post-workout 6 PM	Whey shake 30g + 1/2 banana	220	28g
Dinner 8 PM	150g fish tikka + large salad + 1 roti	380	36g
Pre-sleep	Cottage cheese 150g (low fat)	110	18g
TOTAL		1,850 kcal	201g

Ch 6: Common Mistakes & Troubleshooting

Mistake	Why It Fails	The Fix
Not tracking calories at all	Consistently underestimates intake; impossible to manage what you do not measure	Track accurately for at least 6 weeks to build intuitive awareness
Too much cardio, not enough lifting	Accelerates muscle loss; reduces metabolic rate; unsustainable	Prioritise resistance training; add cardio as a deficit tool, not primary method
Eating too little protein	Muscle breakdown during deficit; lower satiety; worse body composition result	Minimum 2g/kg bodyweight every single day; non-negotiable
Losing more than 1 kg/week	Indicates muscle is being lost not just fat; hormones disrupted	Slow down; add back 200–300 kcal; re-evaluate approach
Cheat meals turning into cheat days/weekends	Eliminates entire week's deficit in 2 days; zero net progress	Planned refeeds (maintenance for 1 day) are evidence-based; cheat weekends are not
Stopping at first plateau	Metabolic adaptation is temporary and reversible; abandoning too early is the biggest mistake	Apply plateau protocol; be patient; results often resume within 1–2 weeks of intervention