

ROYAL FITNESS CLUB

RFC PREMIUM GUIDE #20

ANABOLIC CYCLE NUTRITION & SUPPLEMENTATION BIBLE

**Complete Nutritional Strategy for Before, During & After Every
Cycle**

6 Chapters · Essential Reference · Bulking & Cutting · Organ Protection

© Royal Fitness Club — For Members Only

TABLE OF CONTENTS

Chapter 1 — Nutrition Principles for Enhanced Athletes	3
Chapter 2 — Bulking Nutrition: Maximising Anabolic Environment	4
Chapter 3 — Cutting Nutrition on Cycle	5
Chapter 4 — Organ Protection & Support Supplements	6
Chapter 5 — Intra-Cycle Supplementation Strategy	7
Chapter 6 — Post-Cycle Nutrition for Recovery	8

DISCLAIMER

For educational and harm-reduction purposes only. Not medical advice. Consult a qualified healthcare professional before starting any protocol. Individual results vary.

CHAPTER 1 — NUTRITION FOR ENHANCED ATHLETES

"Anabolic compounds are nutrient partitioning agents — they do not create muscle from nothing. They direct protein and calories toward muscle growth at a dramatically higher efficiency. Supply the raw materials."

Enhanced athletes have fundamentally different nutritional requirements from natural athletes. The increased protein synthesis rate, nitrogen retention, and anabolic signalling during a cycle creates a dramatically elevated demand for dietary substrate. Insufficient protein or calories while "on cycle" is the most common reason for below-average results — the compounds cannot build what the nutrition does not provide.

Nutritional Variable	Natural Athlete Target	Enhanced Athlete Target	Reason for Difference
Protein (g/kg BW/day)	1.6–2.0g	2.2–3.0g	Enhanced protein synthesis requires more substrate
Total calories (surplus)	+200–300 kcal	+300–600 kcal	Greater anabolic capacity utilises larger surplus cleanly
Carbohydrates (g/kg)	3–5g	4–7g (bulking)	Increased glucose uptake and glycogen synthesis rate
Meal frequency	3–4 meals	5–6 meals	Sustained amino acid availability for enhanced MPS
Protein per meal	30–40g	40–60g	Higher leucine threshold with elevated MPS

CHAPTER 2 — BULKING NUTRITION ON CYCLE

Bodyweight	TDEE	Cycle Surplus	Target Kcal	Protein	Carbs	Fat
70kg	2800	+400	3200	175g	450g	85g
80kg	3100	+450	3550	200g	510g	95g
90kg	3400	+500	3900	225g	570g	105g
100kg	3700	+550	4250	250g	630g	115g

6-Meal Bulking Day Template

Meal #	Timing	Foods	Protein	Kcal
1	7:00 AM	6 eggs + 3 roti + 1 cup full-fat milk + banana	50g	730
2	10:30 AM	200g chicken + 1.5 cups rice + sabzi	45g	600
3 (pre-workout)	1:30 PM	50g oats + 1.5 scoops whey + 1 tbsp PB	45g	490
4 (post-workout)	4:30 PM	2 scoops whey + 60g dextrose + creatine	55g	470
5	7:30 PM	250g paneer or fish + 2 roti + 1 cup daal + veg	60g	740
6	10:30 PM	300ml full-fat milk + 30g casein + 30g almonds	45g	520

CHAPTER 3 — CUTTING NUTRITION ON CYCLE

"The advantage of cutting on cycle is that muscle preservation is dramatically enhanced — you can run a larger deficit than a natural athlete without the same muscle loss penalty."

Bodyweight	TDEE	On-Cycle Deficit	Target Kcal	Protein	Carbs	Fat
80kg	3100	−600	2500	220g	240g	75g
90kg	3400	−650	2750	248g	270g	82g
100kg	3700	−700	3000	275g	300g	90g

■ ***On-cycle cutting deficit can be 500–700 kcal vs the 300–400 kcal recommended for naturals. The anabolic environment prevents muscle catabolism at this deficit. Do not exceed 700 kcal — cortisol elevation is still a risk.***

CHAPTER 4 — ORGAN PROTECTION & SUPPORT SUPPLEMENTS

Non-Negotiable On-Cycle Supplements

Supplement	Dose	Primary Protection	Take With
TUDCA (Tauroursodeoxycholic Acid)	500mg/day	Bile acid transport, hepatocyte protection	Any meal
UDCA or Liv52 DS (alternative)	2 tabs twice daily	Liver enzyme normalisation	Meals
Omega-3 (EPA/DHA)	4–6g/day combined	LDL reduction, anti-inflammatory, cardiovascular	Meals
CoQ10	200–400mg/day	Mitochondrial function, cardiac protection	Morning, with fat
NAC (N-Acetyl Cysteine)	600mg twice daily	Glutathione precursor, liver + kidney	Between meals
Hawthorn Berry	500mg twice daily	Blood pressure management, cardiac	Morning + evening
Vitamin K2 (MK-7)	200mcg/day	Arterial calcification prevention (with D3)	With fat

CHAPTER 5 — INTRA-CYCLE SUPPLEMENTATION

Supplement	On-Cycle Role	Dose	Timing
Creatine Monohydrate	Cell volume, strength, recovery	5g/day	Any time consistently
Whey Isolate	Rapid protein synthesis substrate	1–2 scoops	Post-workout + between meals
Casein Protein	Sustained overnight MPS	30–40g	Pre-sleep
BCAA (2:1:1)	Anti-catabolic, intra-workout	10g	During training
Vitamin C	Cortisol reduction, collagen	1000mg twice daily	Morning + post-workout
Magnesium Glycinate	Sleep quality, CNS, insulin sensitivity	400mg	Before bed
Zinc	Testosterone co-factor, immune	25–30mg	Before bed (away from calcium)
Probiotics	Gut health, nutrient absorption	10B CFU daily	With breakfast

CHAPTER 6 — POST-CYCLE NUTRITION

"Post-cycle nutrition is the final lever between keeping your gains and losing them. PCT drugs restart hormones; nutrition provides the substrate that holds muscle while testosterone is suboptimal."

The post-cycle period (including PCT) is a catabolic risk window. Testosterone is low, cortisol may be elevated, and anabolic signalling is suppressed. Aggressive nutrition counteracts the catabolic environment.

PCT Phase	Caloric Strategy	Protein Target	Carb Approach	Fat Strategy
Weeks 1–2 (SERM initiation)	Maintenance + 100	2.5g/kg BW	Moderate — 3g/kg	0.8g/kg
Weeks 3–4 (Mid PCT)	Maintenance	2.5g/kg BW	Timed peri-workout	0.8g/kg
Post-PCT consolidation	Small surplus (+150)	2.2g/kg BW	Normal cycling	0.8–1.0g/kg

- Maintain training intensity and volume through PCT — do not reduce training out of fear of loss
- Prioritise sleep during PCT — it is the single most important recovery intervention available
- No alcohol during PCT — the HPTA is fragile; do not add pharmacological stress
- Continue all on-cycle liver/organ support through PCT and 4 weeks beyond
- Avoid large caloric deficits for at least 12 weeks after PCT completion